



OPEN ACCESS

EDITED AND REVIEWED BY
Candan Tamerler,
University of Kansas, United States

*CORRESPONDENCE

Yumin Yang,
✉ yangym@ntu.edu.cn
Xiaoming Yang,
✉ sammy@ntu.edu.cn
Yahong Zhao,
✉ Zhaoyh108@ntu.edu.cn

RECEIVED 24 April 2025

ACCEPTED 26 May 2025

PUBLISHED 12 June 2025

CITATION

Wang Y, Liang Y, Huang J, Gao Y, Xu Z, Ni X, Yang Y, Yang X and Zhao Y (2025) Correction: Proteomic analysis of silk fibroin reveals diverse biological function of different degumming processing from different origin. *Front. Bioeng. Biotechnol.* 13:1612862. doi: 10.3389/fbioe.2025.1612862

COPYRIGHT

© 2025 Wang, Liang, Huang, Gao, Xu, Ni, Yang, Yang and Zhao. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Correction: Proteomic analysis of silk fibroin reveals diverse biological function of different degumming processing from different origin

Yaling Wang^{1,2}, Yunyun Liang¹, Jiacen Huang³, Yisheng Gao¹, Zhixin Xu¹, Xuejun Ni⁴, Yumin Yang^{1*}, Xiaoming Yang^{1,3*} and Yahong Zhao^{1,3*}

¹Key Laboratory of Neuroregeneration of Jiangsu and Ministry of Education, Co-Innovation Center of Neuroregeneration, NMPA Key Laboratory for Research and Evaluation of Tissue Engineering Technology Products, Nantong University, Nantong, China, ²School of Pharmacy, Nantong University, Nantong, China, ³School of Public Health, Nantong University, Nantong, China, ⁴Affiliated Hospital of Nantong University, Nantong University, Nantong, China

KEYWORDS

silk, silk fibroin, silk sericin, degum, proteomic

A Correction on

Proteomic analysis of silk fibroin reveals diverse biological function of different degumming processing from different origin

by Wang Y, Liang Y, Huang J, Gao Y, Xu Z, Ni X, Yang Y, Yang X and Zhao Y (2022). *Front. Bioeng. Biotechnol.* 9:777320. doi: 10.3389/fbioe.2021.777320

In the published article, there was an error in [Figure 4](#) as published. The images were mistakenly selected from a different experimental condition than the one described in the corresponding figure legend and main text. The error occurred during the figure assembly process, as the original image files for multiple experimental conditions were stored in the same folder without distinct subcategorization. The corrected [Figure 4](#) and its caption appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.




FIGURE 4

Viability of Schwann cells on different samples. Images of calcein-AM/propidium iodide (PI) double-staining: (A) control, (B) JS-0.05, (C) JS-0.5, (D) ZJ-0.05, and (E) ZJ-0.5. Scale bar, 50 μ m. Green fluorescence indicates live cells stained with calcein-AM and red fluorescence indicates dead cells stained with PI. Scale bar, 200 μ m. (F) CCK-8 test of Schwann cells.