



Corrigendum: Novel Porcine Model of Coronary Dissection Reveals the Impact of Impella on Dissected Coronary Arterial Hemodynamics

OPEN ACCESS

Edited and reviewed by:

Hendrik Tevæearai Stahel,
Bern University Hospital, Switzerland

*Correspondence:

Kiyotake Ishikawa
kiyotake.ishikawa@mssm.edu

Specialty section:

This article was submitted to
Heart Failure and Transplantation,
a section of the journal
Frontiers in Cardiovascular Medicine

Received: 27 December 2020

Accepted: 17 March 2021

Published: 08 April 2021

Citation:

Kariya T, Yamada KP, Bikou O,
Tharakan S, Miyashita S and
Ishikawa K (2021) Corrigendum: Novel
Porcine Model of Coronary Dissection
Reveals the Impact of Impella on
Dissected Coronary Arterial
Hemodynamics.
Front. Cardiovasc. Med. 8:646675.
doi: 10.3389/fcvm.2021.646675

Taro Kariya, Kelly P. Yamada, Olympia Bikou, Serena Tharakan, Satoshi Miyashita and Kiyotake Ishikawa*

Cardiovascular Research Center, Icahn School of Medicine at Mount Sinai, New York, NY, United States

Keywords: coronary arterial dissection, spontaneous coronary artery dissection, impella, large animal, LV unloading, coronary angiography, intimal flap

A Corrigendum on

Novel Porcine Model of Coronary Dissection Reveals the Impact of Impella on Dissected Coronary Arterial Hemodynamics

by Kariya, T., Yamada, K. P., Bikou, O., Tharakan, S., Miyashita, S., and Ishikawa, K. (2020). *Front. Cardiovasc. Med.* 7:162. doi: 10.3389/fcvm.2020.00162

In the original article, there was a mistake in **Table 2**, as published. The correct heart rate values appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2021 Kariya, Yamada, Bikou, Tharakan, Miyashita and Ishikawa. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

TABLE 2 | Effect of Impella on LV parameters after coronary dissection in animals with large coronary flap.

	Impella off (N = 6)	Impella max flow (N = 6)	P-value
LV end-diastolic pressure, mmHg	20.6 ± 6.6	12.0 ± 3.4	0.032
LV end-systolic pressure, mmHg	112.2 ± 28.6	126.9 ± 22.9	0.066
LV end-diastolic volume, ml	127 ± 32	97 ± 26	0.015
LV end-systolic volume, ml	71 ± 32	63 ± 27	0.14
Stroke volume, ml	68 ± 16	48 ± 14	0.003
Stroke work, mmHg·ml	5,744 ± 1,866	4,424 ± 1,650	0.003
Heart rate, bpm	71.4 ± 6.6	64.9 ± 9.3	0.014
Cardiac output, L/min	4.78 ± 0.83	5.70 ± 0.46	0.03

*Cardiac output is a summation of one from left ventricular contraction and one from Impella.
LV, left ventricle.*