



Genetic Imaging of Neuroinflammation in Parkinson's Disease: Recent Advancements

Longping Yao^{1*}, Jiayu Wu¹, Sumeyye Koc² and Guohui Lu^{1*}

¹ Department of Neurosurgery, The First Affiliated Hospital of Nanchang University, Nanchang, China, ² Department of Neuroscience, Institute of Health Sciences, Ondokuz Mayıs University, Samsun, Turkey

OPEN ACCESS

Edited by:

Chencheng Zhang,
Shanghai Jiao Tong University, China

Reviewed by:

Sachchida Nand Rai,
University of Allahabad, India
Huajian Chen,
Southern Medical University, China

*Correspondence:

Longping Yao
loupe_yao@163.com
Guohui Lu
guohui-lu@163.com

Specialty section:

This article was submitted to
Molecular Medicine,
a section of the journal
*Frontiers in Cell and Developmental
Biology*

Received: 19 January 2021

Accepted: 14 June 2021

Published: 15 July 2021

Citation:

Yao L, Wu J, Koc S and Lu G
(2021) Genetic Imaging
of Neuroinflammation in Parkinson's
Disease: Recent Advancements.
Front. Cell Dev. Biol. 9:655819.
doi: 10.3389/fcell.2021.655819

Parkinson's disease (PD) is one of the most prevalent neurodegenerative aging disorders characterized by motor and non-motor symptoms due to the selective loss of midbrain dopaminergic (DA) neurons. The decreased viability of DA neurons slowly results in the appearance of motor symptoms such as rigidity, bradykinesia, resting tremor, and postural instability. These symptoms largely depend on DA nigrostriatal denervation. Pharmacological and surgical interventions are the main treatment for improving clinical symptoms, but it has not been possible to cure PD. Furthermore, the cause of neurodegeneration remains unclear. One of the possible neurodegeneration mechanisms is a chronic inflammation of the central nervous system, which is mediated by microglial cells. Impaired or dead DA neurons can directly lead to microglia activation, producing a large number of reactive oxygen species and pro-inflammatory cytokines. These cytotoxic factors contribute to the apoptosis and death of DA neurons, and the pathological process of neuroinflammation aggravates the primary morbid process and exacerbates ongoing neurodegeneration. Therefore, anti-inflammatory treatment exerts a robust neuroprotective effect in a mouse model of PD. Since discovering the first mutation in the α -synuclein gene (SNCA), which can cause disease-causing, PD has involved many genes and loci such as LRRK2, Parkin, SNCA, and PINK1. In this article, we summarize the critical descriptions of the genetic factors involved in PD's occurrence and development (such as LRRK2, SNCA, Parkin, PINK1, and inflammasome), and these factors play a crucial role in neuroinflammation. Regulation of these signaling pathways and molecular factors related to these genetic factors can vastly improve the neuroinflammation of PD.

Keywords: Parkinson's disease, microglia, genetics, neuroinflammation, dopaminergic neurons, neurotoxins

INTRODUCTION

Parkinson's disease (PD) is an age-related neurodegenerative disease characterized by motor and non-motor symptoms (Brundin et al., 2018). It is the second most common neurodegenerative disorder after Alzheimer's disease, and now, it has become a significant public health problem worldwide (Fan, 2020). One characteristic pathological change is the loss of midbrain dopaminergic

(DA) neurons (Charvin et al., 2018). Parkinson's disease is usually characterized by the deposition of protein aggregates containing α -synuclein (Lewy bodies) in multiple brain regions (Rey et al., 2018). The decreased viability of DA neurons slowly results in the appearance of motor symptoms, which largely depend on dopaminergic nigrostriatal denervation. With the progression of neurodegeneration and advancing disease, people with PD also experience sleep disturbances, fatigue, altered mood, cognitive changes, autonomic dysfunction, and pain (Schapira et al., 2017). These "non-motor" symptoms dominate the clinical picture and are the main determinants of quality of life. Besides, some endophenotypes dominated by non-motor symptoms also have been reported in recent studies (Ehgoetz Martens and Shine, 2018). Nowadays, pharmacological and surgical interventions are the main treatments for improving clinical symptoms. However, preventing or curing the disease is not possible at present (Sarrafchi et al., 2016).

Although various possible pathogenetic mechanisms have been proposed in recent years, the cause of neurodegeneration remains unknown. One possible mechanism is inflammation mediated by microglial cells (Yao et al., 2018). Emerging evidence indicates that sustained inflammatory stimulation plays a vital role in the degeneration of DA neurons and is a common feature in both human PD patients and animal models of PD (Ikeda-Matsuo et al., 2019). The neuroinflammatory response may also lead to a cascade of events leading to neuronal degeneration. Anti-inflammatory treatment showed beneficial effects in preventing neurodegeneration mediated by inflammatory damage (Singh et al., 2020). For example, ursolic acid acted as a therapeutic drug targeted for protecting from neuroinflammation-induced neurodegeneration (Rai et al., 2019; Singh et al., 2020; Zahra et al., 2020). Besides, the supplement of tyrosine hydroxylase in MPTP-intoxicated mice could protect dopaminergic neurons by suppressing neuroinflammation (Birla et al., 2019). Thus, regulating neuroinflammation is a therapeutic strategy by reducing oxidative stress (Rai et al., 2017; Yadav et al., 2017; Singh et al., 2018). In recent years, research has shown that many genetic factors are associated with neurodegeneration

Abbreviations: 6-mer, Six monosaccharides; 6-OHDA, 6-hydroxydopamine; AIM2, Absent in melanoma 2; ATP, Adenosine triphosphate; BHB, β -hydroxy butyrate; c-Abl, c-Abelson murine leukemia viral oncogene homolog; CNS, Central nervous system; COX2, Cyclooxygenase 2; DA, Dopaminergic; DAMPs, damage associated proteins; EAE, Experimental autoimmune encephalomyelitis; FAK, Focal adhesion kinase; GFAP, Glial fibrillary acidic protein; IL-1 β , Interleukin-1 β ; IL-6, Interleukin-6; iNOS, Inducible nitric oxide synthase; IRFs, Interferon regulatory factors; JAK/STAT, Janus kinase/signal transducers and activators of transcription; LPC, Lysophosphatidylcholine; LPS, Lipopolysaccharide; LRRK2, Leucine-rich repeat kinase 2; MAPK-NF- κ B, p38 mitogen-activated protein kinase-NF- κ B; MPTP, 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine; MSU, Monosodium urate; NADPH, Nicotinamide adenine dinucleotide phosphate; NFA, Nuclear factor of activated T cell; NF- κ B, Nuclear factor-kappa B; NLRP4, Nod-like receptor family CARD domain containing protein 4; NO, Nitric oxide; Nrf2/HO-1, Nuclear factor erythroid 2-related factor 2/heme oxygenase-1; PAMPs, Pathogen specific proteins; PD, Parkinson's disease; p-ERK, Phosphorylated extracellular signal-regulated kinase; PINK1, PTEN-induced putative kinase 1; PKA, Protein kinase A; PPAR- γ , Peroxisome proliferator-activated receptor γ ; PRGF, Plasma rich in growth factors; PRRs, Pattern recognition receptors; ROS, Reactive oxygen species; ROT, LPS-primed rotenone; SNCA, α -synuclein gene; SNpc, Substantia nigra pars compacta; TNF- α , Tumor necrosis factor- α ; TrxR, Thioredoxin reductase; UBL, Ubiquitin-like; α -syn, Alpha-synuclein.

of the nigrostriatal DA, which includes apoptosis or death caused by neuroinflammation (Zhang et al., 2018). These factors are potential targets to interfere with the disease process in PD. This review discusses the details of genetic imaging of neuroinflammation in PD.

INFLAMMATION AND PARKINSON'S DISEASE

The presence of activated microglial cells in the substantia nigra has been shown in postmortem studies and in the 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine (MPTP)-animal models (both mice and non-human primates) of PD (Zella et al., 2019). Microglia are the macrophages that reside in the central nervous system (CNS) and are the brain's primary innate immune effector cells (Yin et al., 2017). Besides, they are the main generator of inflammatory cytokines and reactive oxygen species (ROS), acting as the central active immune defense under normal conditions (Takeda et al., 2018). In pathological conditions, microglia are activated and releases anti-inflammatory cytokines and neurotrophic factors, contributing to tissue repair and the protection of neurons against apoptosis or death (Hilla et al., 2017). However, when pathological factors are continuously present, the number of noxious phenotypes of microglia increase and release a large amount of ROS, proteinases, and inflammatory cytokines contributing to neuronal damage. In PD, the inflammatory mediators such as tumor- α (TNF- α), interleukin-1 β (IL-1 β), inducible nitric oxide synthase (iNOS), and interleukin-6 (IL-6) have been found to regulate the progression of PD (Yao et al., 2019b). These pro-inflammatory mediators have been found to increase significantly in the midbrain of PD patients and animal models. Furthermore, numerous studies have shown that impaired or dead DA neurons can directly induce the activation of microglia, increasing the production of ROS and pro-inflammatory cytokines. Therefore, as mentioned above, the activation of microglia and DA neuronal damage form a self-propelled degeneration cycle in PD; thus, microglia are more likely to play critical roles in establishing and maintaining inflammatory responses in PD (Figure 1).

The anti-inflammatory treatment has been found to exert a strong neuroprotective effect in a mouse model of PD. For example, specially designed liposomes targeted for the CD163 receptor were loaded with glucocorticoids involved in PD-like neurodegeneration. The results showed that glucocorticoids protect DA neurons in the 6-hydroxydopamine (6-OHDA) PD model via anti-inflammatory modulation (Tentillier et al., 2016). APSE-Aq (fraction isolated from *A. pyrifolium* seeds) may have antioxidant and anti-inflammatory properties, which may offer neuroprotection in a model of PD (de Araujo et al., 2018). Tetramethylpyrazine exhibits its anti-apoptotic, anti-inflammatory, and antioxidant actions via attenuating rotenone-induced upregulation of the transcription factor nuclear factor erythroid 2-related factor 2/heme oxygenase-1 (Nrf2/HO-1) pathway and inflammation markers: nuclear factor-kappa B (NF- κ B), iNOS, cyclooxygenase 2 (COX2), and glial fibrillary acidic protein (GFAP) expression (Lu et al., 2014; Michel et al., 2017).




FIGURE 1 | A self-propelled degeneration cycle in PD. In PD's pathological conditions, microglia are activated and release anti-inflammatory cytokines to repair the tissues, protecting neurons against apoptosis or death. However, the continuous stimulation of pathological factors increases the number of toxic phenotypes of microglia, releasing a large number of inflammatory cytokines, such as TNF- α , IL-1 β , iNOS, IL-6, and ROS, which contribute to neuronal damage. Besides, the impaired or dead DA neurons can directly induce microglial activation, increasing ROS and pro-inflammatory cytokines. Thus, the activation of microglia and DA neuronal damage form a self-propelled degeneration cycle in PD. PD, Parkinson's disease; DA, dopaminergic; ROS, reactive oxygen species; iNOS, inducible nitric oxide synthase; IL-6, interleukin-6; TNF- α , tumor necrosis factor- α ; IL-1 β , interleukin-1 β .

The isothiocyanate is mainly found in Brassica vegetables (Brassicaceae) and Moringaceae plants. It shows potent anti-inflammatory activity in the treatment of murine subacute PD and promises compounds against oxidative stress and neuroinflammation (Sita et al., 2016; Giacoppo et al., 2017). The neuroprotective properties of securinine may be due

to the inhibition of glial activation and the subsequent generation of pro-inflammatory factors via inhibiting the p38 mitogen-activated protein kinase-NF- κ B (MAPK-NF- κ B) pathway (Leonoudakis et al., 2017). A novel compound (VSC2) has anti-inflammatory and antioxidant properties in microglia and in an animal model of PD by preventing NF- κ B activation

and the production of TNF- α , iNOS, IL-1 β , NO, and COX-2 (Lee et al., 2015). Morin effectively prevents MPTP-induced PD-like pathologies in mice and protects primary neurons against MPP + -induced toxicity by ameliorating oxidative stress and inflammation (Zhang et al., 2010; Lee et al., 2016). The anti-inflammatory effect of β -Hydroxybutyric acid on microglia is mediated by G-protein-coupled receptor 109A. This process involves the NF- κ B signaling pathway in both *in vivo* and *in vitro* PD models, causing the inhibition of the production of pro-inflammatory enzymes (iNOS and COX-2) and pro-inflammatory cytokines (TNF- α , IL-1 β , and IL-6) (Fu et al., 2015). The synthesis and biological evaluation of clovamide analogs show that they have potent anti-neuroinflammatory effects by reducing the expression of GFAP in a model of PD (Hu et al., 2018). Compound 21 is obtained from 2,4,6-trimethoxybenzaldehyde by adding amines or alkyl (methyl or ethyl) ester of the amino acid hydrochloride salts. Compound 21 may be a potential candidate for PD treatment because of its potent anti-neuroinflammatory activity, novel mechanism, impressive penetration of the blood-brain barrier, and low toxicity *in vitro* and *in vivo* (Wang et al., 2016). The behavioral and neurochemical alterations in a rat model of PD are partially reversed by *Spirulina platensis*, which is primarily related to its anti-inflammatory effects (Lima et al., 2017). Phytic acid shows a solid neuroprotective effect in an MPTP-induced PD model correlated with its anti-inflammatory effect by suppressing the NF- κ B and phosphorylated extracellular signal-regulated kinase (p-ERK) pathways (Lv et al., 2015).

Based on the hypothesis that neuroinflammation is involved in the pathophysiology of PD, scientists have evaluated the feasibility of using non-steroidal anti-inflammatory drugs (NSAIDs) to cure PD patients. Interestingly, the results of a prospective cohort study showed that NSAIDs might delay or prevent the onset of PD (Chen et al., 2003). Ibuprofen users had a lower risk of PD than non-users, suggesting that ibuprofen use may delay or prevent PD onset (Poly et al., 2019). However, the same anti-inflammatory effect was not observed for aspirin, other NSAIDs, or acetaminophen in PD patients. The inconsistent results between ibuprofen and other NSAIDs indicate that ibuprofen has specific protective properties of the anti-inflammatory response in PD.

THE ROLE OF NF- κ B IN NEUROINFLAMMATION IN PD

Transcription factors such as NF- κ B, STAT1, STAT3, and SMAD7 are the proteins that can bind to a specific sequence of DNA to regulate the transcription of genes. Recent literature shows that the upregulation of transcription factors can cause microglial activation, leading to a self-sustaining neuroinflammation environment via various mechanisms in PD. In this article, we will describe how NF- κ B is associated with different pathways and molecular regulation. For now, we will give a clear description of the role of NF- κ B in neuroinflammation in PD.

NF- κ B is crucial for neuroinflammation responses; NF- κ B is a group of transcription factors including RelA, RelB,

c-Rel, NF- κ B1/p50, and NF- κ B2/p52 (Sokolova and Naumann, 2017). The transcription factors can form homodimers and heterodimers to regulate the expression of genes. The abundant basal expression of NF- κ B in the brain is much higher than in peripheral tissues (Shih et al., 2015). NF- κ B is largely involved in the progression of PD. For example, it reveals that NF- κ B is activated in an MPTP model of PD (Dehmer et al., 2004), following the microglia activation (Aoki et al., 2009). One research study showed that NF- κ B increased more than 70-fold in the brain tissue of PD patients and exhibited strong nuclear p65 immunoreactivity of DA neurons in the substantia nigra (Mattson and Camandola, 2001). The activated NF- κ B leading to DA neuron degeneration has been demonstrated in PD (Phani et al., 2012). Selective inhibition NF- κ B has been found to protect against DA neurons' death from MPTP toxicity in a PD model (Bassani et al., 2015).

In neuroinflammation, NF- κ B can regulate the production of pro-inflammatory cytokines, such as IL-6, TNF- α , G-CSF, iNOS, and IL-1 β . The sustained inflammatory stimulus is related to the NF- κ B pathway in the activation of uncontrolled microglia, resulting in ROS production, neurotoxic factors, interferon- γ (INF- γ), and glutamate, whose excessive formation induces neuronal damage (Spencer et al., 2012). Toll-like receptors are a vital class of membrane proteins that can activate microglial cells—the predominant pathways that TLRs trigger are associated with the NF- κ B pathway (Kawai and Akira, 2007). Inhibition of the NF- κ B pathway can sufficiently suppress the activation of microglia and neuroinflammation.

Recent studies show that NF- κ B is involved in the inflammatory response of microglia in the progression of PD. Therefore, the regulation of the abnormal expression of NF- κ B exerts fortissimo neuroprotection and inhibition of inflammation in PD. For example, our previous study found that miR-124 can prevent DA neuronal death and suppress microglia activation via suppressing the MEKK3/NF- κ B pathway in a mouse model of PD (Yao et al., 2018). Intranasal plasma rich in growth factors (PRGF)-Endoret provides a novel neuroprotective strategy for DA and attenuates NF- κ B-dependent inflammation processes in a PD model (Anitua et al., 2015). Inhibition of NF- κ B activation leads to the suppression of pro-inflammatory molecules, improvement in locomotor activity, and DA neurons' protection in the substantia nigra pars compacta (SNpc) (Mondal et al., 2012). In DA neuron-glia co-cultures, pioglitazone prevents DA cells' death from lipopolysaccharide (LPS)-induced exacerbation of microglia activation by interfering with the NF- κ B pathway (Esposito et al., 2007). Besides, schisandrol A could enhance the PI3K/AKT pathway and inhibit the IKK/I κ B α /NF- κ B pathway to reduce neuronal inflammation, oxidative stress and enhance the survival of DA neurons in the brains of PD mice (Yan et al., 2019). Rosmarinic acid could attenuate inflammatory responses by suppressing the HMGB1/TLR4/NF- κ B signaling pathways, which may contribute to its anti-PD activity (Lv et al., 2019). Cordycepin mitigates MPTP-induced inflammatory response in PD by inhibiting the TLR/NF- κ B signaling pathway (Cheng and Zhu, 2019). The knockdown of cathepsin D can protect dopaminergic neurons from neuroinflammation-mediated neurotoxicity via inhibition of the NF- κ B signaling pathway in

a PD model (Gan et al., 2018). Polydatin treatment protects DA neurons and ameliorates motor dysfunction by inhibiting microglial activation and the release of pro-inflammatory mediators via regulation of the AKT/GSK3 β -Nrf2/NF- κ B signaling axis (Huang et al., 2018).

Meanwhile, several NSAIDs, such as sodium salicylate, celecoxib, aspirin, and diclofenac, have been found to exert a neuroprotective role by decreasing NF- κ B expression in an MPTP-induced model of PD (Bassani et al., 2015). The peroxisome proliferator-activated receptor γ (PPAR- γ) agonist pioglitazone mediates microglial activation and NF- κ B expression in the 6-hydroxydopamine model of PD (Goes et al., 2018). A20 enzyme, which inhibits NF- κ B by restricting the duration and intensity of its action, has been found to significantly decrease in a blood sample of patients with PD (Mazo et al., 2017). The topics which are discussed next are widely associated with NF- κ B.

THE ROLE OF GENETICS IN THE NEUROINFLAMMATION OF PD

Since discovering the first mutation in the α -synuclein gene (SNCA) can cause disease-causing, PD has involved many genes and loci. For example, the deficiencies of genes such as LRRK2, Parkin, SNCA, and PINK1 are risk factors for PD (including family and sporadic PD). The genetic discoveries clearly illustrate the cellular pathways and functions that are involved in the development of PD. To date, at least 23 loci and 19 genes (Table 1) have been identified and designated as both 10 autosomal dominant and 9 autosomal recessive PD.

Recent epidemiological and genetic studies have indicated that some PD-associated genes are involved in regulating neuroinflammation in the CNS. The discoveries of genetic factors highlight the biological mechanism of PD. According to the literature, we summarized whether the 19 genes associated with PD are also associated with neuroinflammation (Table 1). Understanding how genetic factors influence the inflammatory pathogenesis of PD can help decipher the disease's etiology.

Leucine-Rich Repeat Kinase 2 (LRRK2)

Missense mutations in the LRRK2 gene are the most common cause of autosomal-dominant inherited PD (Chan and Tan, 2017); the standard variants of the LRRK2 gene have also been associated with sporadic PD (Kluss et al., 2019). LRRK2 has been a therapeutic target for family and sporadic PD (Tufekci et al., 2012). The penetrance of LRRK2 mutations is incomplete in PD because the lifetime risk is estimated to be 22–32% in clinical populations, suggesting strong modifiers of LRRK2 disease (Goldwurm et al., 2007). Recently, genome-wide association studies show that LRRK2 is also involved in modifying immunogenic responses in PD (Moehle et al., 2012). Injecting LPS can strongly induce LRRK2 expression in SNpc in mice. Idiopathic PD patients and those with an LRRK2 mutation have increased levels of pro-inflammatory serum markers (Brockmann et al., 2016). However, no activated inflammatory profiles are observed in PD patients with a

TABLE 1 | The genes associated with the pathogenesis of PD.

Gene	Full name	Locus	Location
SNCA	Synuclein alpha	PARK1	4q22.1 Azizi and Azizi (2018)
Parkin	Parkin RBR E3 ubiquitin protein ligase	PARK2	6q26 Morato Torres et al. (2020)
PARK3	Parkinson disease 3	PARK3	2p13 Kachidian and Gubellini (2021)
SNCA	Synuclein alpha	PARK4	4q22 Forero et al. (2020)
UCHL1	Ubiquitin C-terminal hydrolase L1	PARK5	4p13 Kachidian and Gubellini (2021)
PINK1	PTEN induced putative kinase 1	PARK6	1p36 Li D. et al. (2020)
PARK7	Parkinsonism associated deglycase	PARK7	1p36.23 Li Y. et al. (2020)
LRRK2	Leucine rich repeat kinase 2	PARK8	12q12 Li D. et al. (2020)
ATPase 13A2	ATPase 13A2	PARK9	1p36.13 Gago et al. (2020)
PARK10	Parkinson disease 10	PARK10	1p32 Kachidian and Gubellini (2021)
GIGYF2	GRB10 interacting GYF protein	PARK11	2q37.1 Calatayud et al. (2017)
PARK12	Parkinson disease 12	PARK12	Xq21-q25 Zanon (2020)
HTRA2	HtrA serine peptidase 2	PARK13	2p13.1 Zanon (2020)
PLA2G6	Phospholipase A2 group VI	PARK14	22q13.1 Sudira et al. (2018)
FBXO7	F-box protein 7	PARK15	22q12.3 Yuan et al. (2017)
PARK16	Parkinson disease 16	PARK16	1q32 Han et al. (2019)
VPS35	VPS35, retromer complex component	PARK17	16q11.2 Cutillo et al. (2020)
EIF4G1	Eukaryotic translation initiation factor 4 gamma 1	PARK18	3q27.1 Li D. et al. (2020)
DNAJC6	DnaJ heat shock protein family (Hsp40) member C6	PARK19	1p31.3 Velez-Pardo and Jimenez-Del-Rio (2020)
SYNJ1	Synaptojanin 1	PARK20	21q22.1 Zanon (2020)
TMEM230	Transmembrane protein 230	PARK21	20p13 Deng et al. (2018)
CHCHD2	Coiled-coil-helix-coiled-coil-helix domain containing 2	PARK22	7p11.2 Velez-Pardo and Jimenez-Del-Rio (2020)
VPS13C	Vacuolar protein sorting 13	PARK23	15q22.2
RIC3	homolog C acetylcholine receptor chaperone RIC3		11p15.4 Zanon (2020)

non-manifesting LRRK2 mutation (Brockmann et al., 2016). A recent study showed that the kinase activity of LRRK2 increased in microglia cells in sporadic PD postmortem tissue (Di Maio et al., 2018). Besides, the accumulation of α -syn results in increased ROS expression via inducing mitophagy in neurons, a process linked to LRRK2 activity (Choubey et al., 2011; Saez-Atienzar et al., 2014). LRRK2 can modulatesmokine (C–X3–C) receptor 1-mediated signaling pathways to modulate microglial activity (Ma et al., 2016). LRRK2 kinase activity contributes to neuroinflammation via phosphorylating p53 in PD, and the phosphorylation of p53 induces the expression of TNF- α (Muda et al., 2014).

LRRK2 is upstream of protein kinase A (PKA) and can negatively control PKA activity, thus modulating neuronal functions (Greggio et al., 2017). Meanwhile, it has been found

that LRRK2 can control microglial inflammation by regulating PKA-mediated NF- κ B p50 phosphorylation in microglia cells (Russo et al., 2015). The mutant variants of LRRK2 can vastly enhance the transcriptional activity of NF- κ B in microglia (Kim et al., 2012). LRRK2 knockdown increases the levels of phosphorylated NF- κ B p50 in primary microglial cells (Russo et al., 2015). Further, phosphorylated NF- κ B translocates into the nucleus, competes with, and displaces DNA-bound p50:p50 to initiate mRNAs transcription (Zhong et al., 2002). Thus, LRRK2 may control its activation to affect the consequent transcription of pro-inflammatory mediators via NF- κ B in microglia. A recent study demonstrated that LRRK2 acts as a negative regulator of the nuclear factor of activated T cell (NFAT) transcription factors which are associated with the inflammatory response in a large set of immune cells (Liu et al., 2011). Moreover, the abnormal activity of LRRK2 modulates the activation and phagocytosis of microglia cells by the hyperpolymerization of cytoskeleton components such as actin and β -tubulin (Russo et al., 2014). Moreover, LRRK2 suppresses focal adhesion kinase (FAK) Y397 phosphorylation through the phosphorylation of Thr-X-Arg/Lys (TXR) motif(s) in FAK in microglial cells (Choi et al., 2015). LRRK2 has a negative regulatory role in α SYN clearance through down-regulation of the endocytosis pathway in microglia (Maekawa et al., 2016). Besides, LRRK2 promotes mitochondrial alteration in microglia via Drp1 in a kinase-dependent manner, contributing to pro-inflammatory responses, which is regarded as a potential therapeutic target in PD (Ho et al., 2018). LRRK2 has been found to modulate neuroinflammation and neurotoxicity in models of human immunodeficiency virus 1-associated neurocognitive disorders (Puccini et al., 2015).

The inhibition of LRRK2 kinase activity attenuates the expression of pro-inflammatory microglial signaling to modulate neuroinflammation. In this context, several studies have identified that LRRK2 inhibitors show good physicochemical and pharmacokinetic properties and good selectivity and blood-brain barrier permeability against both kinases (Koshibu et al., 2015). Disruption of LRRK2 activity prevents a complete inflammatory response and microglial morphological remodeling (Moehle et al., 2012). As of now, some inhibitors of the LRRK2 gene have been found, showing a potential new neuroprotective role in PD (Lee et al., 2010). Manganese could induce neuroinflammation and the up-regulation of LRRK2 *in vitro* and *in vivo*, and the knock down of LRRK2 can attenuate manganese-induced autophagy dysfunction and inflammation in microglia (Chen J. et al., 2018). Wave2, an actin-cytoskeletal regulator which can directly couple to LRRK2, mediates Lrrk2-G2019S-induced DA neuronal death in both macrophage-midbrain cocultures and *in vivo* in PD (Kim K. S. et al., 2018).


Meanwhile, LRRK2 can phosphorylate Wave2 at the spot of Thr470, stabilize, and prevent its proteasomal degradation in a murine microglia-like cell line (Kim K. S. et al., 2018). The computer-aided drug design can prevent LPS-induced LRRK2 upregulation and microglia activation in a mouse model of neuroinflammation induced by LPS (Li et al., 2014). The overexpression of human pathogenic LRRK2 mutations exhibits long-term lipopolysaccharide-induced DA neuronal loss

in mice, accompanied by exacerbated neuroinflammation in the brain (Kozina et al., 2018). Overall, LRRK2 is associated with the cellular pathways in microglia (Figure 2), and LRRK2 mutations with increased kinase activity might be one of the possible mechanisms for microglia-exacerbated neuroinflammation in PD.

Alpha-Synuclein

Alpha-synuclein (α -syn), a 140-amino-acid protein, is abundantly expressed at a high level in the brain. Under physiological conditions, the functions of α -syn include the regulation of the dopamine transporter. The expression level of α -syn is regarded as a significant determinant of its neurotoxic potential. In contrast, secreted extracellular α -syn has emerged as an additional important factor in PD's pathological process (Chen Y. et al., 2018). However, dysfunctional modifications of α -syn ultimately cause the pathogenesis of neurodegeneration in PD. In this pathological process, it has been shown that α -syn could trigger inflammation and oxidative stress through the activation of microglia (Harms et al., 2018). When the pathological deposition of α -syn occurs, microglia will migrate to the extracellular α -syn through endocytosis to prevent α -syn accumulation in neurons; however, the extensive uptake of α -syn with microglia could generate glial inclusions and induce inflammation (Vekrellis et al., 2011). Some *in vitro* and *in vivo* studies show that the release of α -syn from neurons can activate microglia through TLRs, followed by the initiation of neuroinflammation and progressive neuronal damage in PD (Zhang et al., 2005; Kim et al., 2013) (Figure 3). Besides, the deposition of α -syn in glial cells induces neuroinflammation, which promotes the degeneration of neurons and aggravates the pathogenesis of PD, and the deposition of α -syn also could further propagate to other glial cells and neurons (Chistiakov and Chistiakov, 2017). Specifically, the over-expression of α -syn could drive microglia into having a reactive phenotype characterized by enhanced levels of cytokine secretion, such as TNF- α and IL-6, as well as nitric oxide (NO), arachidonic acid metabolizing enzymes, and reactive nitrogen species, all superimposed upon impaired phagocytic potential (Rojanathammanee et al., 2011). Fibrillar α -syn, a potent inducer of pro-inflammatory immune, responds to microglia cells and highlights the level of fibrillization of α -syn as a significant feature for its efficient internalization and the activation process of microglia mainly depend on their aggregation state (Hoffmann et al., 2016). In the cerebrospinal fluid and blood of PD patients, researchers have found the aggregated and non-aggregated forms of α -syn. The type of secretion of α -syn into the medium has implied that this form of release from neurons may activate the inflammatory response in a microglial cell line (Alvarez-Erviti et al., 2011). Furthermore, α -syn deficiency promotes neuroinflammation by increasing Th1 cell-mediated immune responses. Endogenous α -syn plays a functional role in immunological processes during early experimental autoimmune encephalomyelitis (EAE) as a new regulator of Th1 responses in neuroinflammation (Ettle et al., 2016).

Dysfunctional modifications of α -syn affecting the activation of microglia are involved in many pathways in PD, and



blocking these pathways can effectively control or attenuate neuroinflammation in PD. For instance, suppressing the Janus kinase/signal transducers and activators of transcription (JAK/STAT) pathway can prevent neuroinflammation and neurodegeneration by inhibiting microglial activation, macrophage, and CD4(+) T-cell infiltration, and the production of pro-inflammatory cytokines/chemokines induced by α -syn (Qin et al., 2016). Furthermore, six monosaccharides (6-mer), a specific inhibitor of the α -syn output, could efficiently modulate neuroinflammation and α -syn expression in neuron-like SH-SY5Y cells by blocking NF- κ B activation (Scuruchi et al., 2016). Meanwhile, α -M inhibits α -syn-induced microglial neuroinflammation and neurotoxicity by targeting nicotinamide adenine dinucleotide phosphate (NADPH) oxidase as a therapeutic possibility in preventing PD progression (Hu et al., 2016). Furthermore, ginsenoside Rg1 could attenuate motor impairment and

neuroinflammation in the MPTP-probenecid-induced PD mouse model via targeting α -syn abnormalities in the substantia nigra (Heng et al., 2016). FK506, as a well-known immunosuppressive drug, could decrease neuroinflammation and DA neurodegeneration, pointing to a causal role of neuroinflammation in an α -syn-based rat model of PD (Van der Perren et al., 2015). The administration of hypoxostoxide could reduce neuroinflammation, neurodegeneration, and α -syn accumulation in a mouse model of PD via modulating the activity of NF- κ B, suggesting that hypoxostoxide may be a potent anti-PD drug (Kim et al., 2015). Curcumin has been found to afford its neuroprotective effect and inhibit α -syn aggregation through the inhibition of oxidative stress generation, replenishing glutathione levels, and preventing glial-associated inflammatory response in an LPS-induced PD model (Sharma et al., 2017; Sharma and Nehru, 2018). Endogenous high-mobility group protein B1, which has been demonstrated to mediate




FIGURE 3 | The signaling pathways and molecular factors involved in neuroinflammation. α -syn together with inflammasome form a network to regulate the activation of microglia. Blocking these signaling pathways and molecular factors can effectively improve apoptosis or the death of dopamine neurons caused by neuroinflammation.

persistent neuroinflammation and consequent progressive neurodegeneration by promoting multiple inflammatory and neurotoxic factors, could promote the autophagic degradation α -syn via the Atg 5-dependent autophagy-initiation pathway in PD (Guan et al., 2018). Immunotherapy targeting TLR2 alleviates α -syn accumulation in neuronal and astroglial cells and attenuates neuroinflammation, neurodegeneration, and behavioral deficits in PD models of synucleinopathy by modulating α -syn transmission and neuroinflammation (Kim C. et al., 2018). Indeed, neuroinflammation might be mediated by the microglial expression of MHC II, a vital regulator of the

immune response, given the central role for microglial MHCII in the activation of both innate and adaptive immune responses to α -syn in PD (Harms et al., 2013).

The PINK1–Parkin Axis Parkin

Parkin is predominantly expressed in the brain. It has been implicated in many biological processes, such as synaptic excitability, inflammation, and immunity. The protein comprises a C-terminal R1-in-between-ring-RING2 motif, an N-terminal

ubiquitin-like (UBL) domain, RING0, RING1 IBR. In them, the Ubl and RING0 domains are unique to Parkin (Arkinson and Walden, 2018). The UBL domain interacts with the R1 environment, which negatively adjusts the activity of E3 ligase and parkin translocation to the mitochondria and parkin-dependent mitophagy. As an E3 ubiquitin-ligating enzyme, Parkin plays a critical role in the cell, which works together with E1 ubiquitin-activating enzymes and E2 ubiquitin-conjugating enzymes in a ubiquitin-proteasome system to ubiquitinate the misfolded or aggregated proteins (van der Merwe et al., 2015). Thus, Parkin can act as an activator to disrupt the autoinhibitory mechanisms and bridge the distance between catalytic sites. Parkin shows conformational changes after point mutation, disrupting the unique autoinhibitory features that release both REP and Ubl domain activity (Tang et al., 2017). Furthermore, the phosphorylation of Parkin leads to decreases in its E3 ubiquitin ligase activity (Aguirre et al., 2018). Besides, Parkin maintains mitochondrial quality control and turnover (McWilliams and Muqit, 2017).

The knockout of Parkin is a crucial way to study the role of Parkin in the pathogenesis of PD (Matheoud et al., 2019). The decrease in Parkin's solubility and stability is associated with the degeneration of substantia nigra neurons in PD (Lonskaya et al., 2013). The MPTP treatment on mice increased the expression of Parkin and neuroinflammation (Mendes et al., 2019). Earlier studies reported the mutant Parkin in *Drosophila* showed age-dependent degeneration of dorsomedial dopaminergic neurons (Cha et al., 2005; Whitworth et al., 2005). Recent studies have shown that Parkin's S-nitrosylation could diminish its protective effects against α -synuclein-mediated neurotoxicity and destroy its ubiquitin ligase activity (Wahabi et al., 2018). It has been reported patients with mutated Parkin have clinical symptoms that are identical to patients with PD (Pickrell and Youle, 2015). Parkin mutation carriers are clinically characterized by slow disease progression and by having an excellent response to levodopa treatment. To date, more than 100 different mutations of Parkin have been reported, and the mutations are largely associated with the pathogenesis of PD (Abbas et al., 1999; Ferreira and Massano, 2017). The overexpression of Parkin could protect against manganese-induced cell death and dopaminergic toxicity (Higashi et al., 2004; Jiang et al., 2004).

Parkin shows a potential role in preventing neuroinflammation from progressing in PD. Mice with Parkin mutations appear to have selective DA neuron degeneration and some motor deficits with intraperitoneal LPS (Vivekanantham et al., 2015). Parkin levels and phenocopy Parkin loss-of-function mutations were found to be decreased in chronic inflammatory conditions, and the expression of TNF, IL-1 β , and iNOS was increased in Parkin-null mice (Tran et al., 2011). In BV2 and primary microglia cell lines, the knockdown of Parkin was found to increase LPS-induced microglial activation by elevating the activity of NF- κ B and JNK, which protected neurons from zVAD-mediated necroptosis (Mouton-Liger et al., 2018; Dionisio et al., 2019). Parkin deficiency could enhance NLRP3 inflammasome signaling by attenuating an A20-dependent negative feedback loop in mice (Mouton-Liger et al., 2018). The overexpression of thioredoxin reductase

(TrxR) 2, a novel mediator of the inflammatory response, could efficiently alleviate inflammation-mediated neuronal death by activating the Akt–Parkin pathway and decreasing oxidative stress (Gao et al., 2019). TNF- α -mediated neuronal inflammation could be attenuated by mitochondrial acid-5 via augmenting the AMPK–Sirt3 pathways and activating Parkin-related mitophagy (Huang et al., 2019).

PTEN-Induced Putative Kinase1 (PINK1)-Parkin Axis

Recent studies have highlighted that mitochondrial dysfunction and DNA abnormalities complicatedly associate with the pathogenesis of PD. PINK1 is a mitochondrial surveillance kinase that contributes to the processes involved in ridding the cell of damaged mitochondria; PINK1 mutations are a common genetic cause of familial PD. The mutations include truncating mutations, point mutations, missense, and deletions. PINK1-associated PD has an earlier age of onset and slower progression (Kawajiri et al., 2011). The examination of humans' brains with PINK1-linked PD shows the pathology of Lewy bodies and neuronal loss in the substantia nigra, which are accompanied by microgliosis and astrocytic gliosis (Steele et al., 2015).

The PINK1 protein sequence contains a predicted C-terminal kinase domain and a mitochondrial targeting sequence of the N-terminus (Pickrell and Youle, 2015). The protein is imported into the mitochondria via the translocase of outer and inner membrane complexes. PINK1 controls mitochondrial quality control by removing the dysfunctional mitochondria. Mitochondrial damage is a significant cause of DA death in PD patients. PINK1 recruits the Parkin protein at the outer mitochondrial membrane while the damage of mitochondria occurred (van der Merwe et al., 2017). The activation of Parkin catalyzes the ubiquitination of outer mitochondrial membrane proteins with ubiquitin which is then degraded by the ubiquitin-proteasome system (Cornelissen et al., 2018). Autophagosomes engulf the dysfunctional organelles, and then lysosomal enzymes typically digest both of them via the process of mitophagy. PINK1/parkin-mediated mitophagy has been found in various neuronal and non-neuronal cells, especially exposing to mitochondrial depolarizing agents (Akabane et al., 2016; Newman and Shadel, 2018). The process prevents the accumulation of products from dysfunctional mitochondria, such as increased ROS and mtDNA damage (Truban et al., 2017).

In a poor state of PINK1, the prevention of accumulated products from dysfunctional mitochondria was obstructed, and mtDNA mutational stress resulted in an inflammatory response and activated the DNA-sensing cGAS–STING pathway, which connected mitoflammation with PD pathology (Sliter et al., 2018). In this study, the researchers also found that the expression of multiple cytokines such as IL-6, -12, and -13; IFN β ; CXCL1; and CCL2 and 4 increased efficiently in *Pink1*^{-/-} and *Parkin*^{-/-} mice. Meanwhile, it could also promote the expression of pro-inflammatory type-I IFN and inflammatory cytokine production and activate NF- κ B signaling (West et al., 2015; West and Shadel, 2017). Furthermore, the mtRNA and their double-stranded mitochondrial RNA activated the inflammasome and the RNA-sensing immune receptor MDA5 (Dhir et al., 2018; Zhong et al., 2018). Moreover, one recent study by Yao et al. (2019a) reported

that hydrogen-rich saline alleviated the inflammatory response and apoptosis via PINK1/Parkin-mediated mitophagy (Figure 4).

Inflammasome

Inflammasomes, the central protein (varies with the type of inflammasome), which on activation recruits the adaptor apoptosis speck-like protein (ASC), is multimeric complexes

consisted of a central protein, an adaptor protein ASC and a caspase-1 protein, forming in response to a variety of physiological and pathogenic stimuli. Inflammasome activation, accrued in both health and disease in the CNS, is an essential component of the innate immune. However, excessive activation of the inflammasome is also a significant driver of autoimmune and metabolic disorders, underlying the importance





FIGURE 4 | PINK1 could phosphorylate Parkin on the mitochondrial surface, resulting in the activation of Parkin. The activated Parkin proteins form phospho-polyubiquitin chains on damaged mitochondria. Finally, the dysfunctional mitochondria are cleared via autophagy. In this way, mitophagy inhibits neuroinflammation in PD and increases microglial phagocytosis. However, the mutation of PINK1 or Parkin would alter the balance of fission to fusion by preventing cells from responding to mitochondrial damage. The expression of ROS and pro-inflammatory factors are increased, which aggravates the development of PD. PD, Parkinson's disease; ROS, reactive oxygen species.

of understanding these physiological and pathological contexts (Sharma and Kanneganti, 2016; Mamik and Power, 2017). Recent work has mainly focused on the existence of inflammasome-mediated inflammatory pathways in CNS disorders. Pattern recognition receptors (PRRs) which are primarily expressed by glial cells, play an integral role in the innate immune response through the recognition of pathogen-specific proteins (PAMPs) and damage-associated proteins (DAMPs) (Singhal et al., 2014). So far, researchers have characterized four different inflammasomes and their activators, including NLRP1, NLRP2, NLRP3, nod-like receptor family CARD domain-containing protein 4 (NLRC4), and absent in melanoma 2

(AIM2) (Mariathasan et al., 2004; Boyden and Dietrich, 2006; Mariathasan et al., 2006; Rathinam et al., 2010; de Rivero Vaccari et al., 2014). Pattern recognition receptors have three distinguishing features: universal expression, fast response, and recognizing many microbes (Zhu et al., 2018). Based on these above features, PRRs efficiently initiate the signaling pathways culminating in the activation of MAPK, NF- κ B, and interferon regulatory factors (IRFs), which control the transcription of genes encoding pro-inflammatory factors (Zhu et al., 2018). In neuroinflammation, inflammasomes can regulate microglial activation and subsequent neuroinflammatory processes in brain pathology (Scholz and Eder, 2017). Otherwise, α -syn enters into



BV2 cells in an endocytosis-dependent manner and subsequently triggers NLRP3 inflammasome activation via inducing lysosomal swelling and increasing cathepsin B release (Zhou et al., 2016).

Meanwhile, it also has been found that inflammasomes cause caspase-1 activation following the stimulation of microglia with lysophosphatidylcholine (LPC), depending on LPS prestimulation, NLRP3, and adaptor ASC, and knockdown of inflammasome NLRC4 inhibits LPC-stimulated caspase-1 activity in microglia (Figure 5) (Scholz and Eder, 2017). Further, inflammasomes are also involved in the inflammatory pathogenesis of PD. For example, β -hydroxy butyrate (BHB), an effective inhibitor of the NLRP3 inflammasome in response to multiple activation stimuli including adenosine triphosphate (ATP), silica, and monosodium urate (MSU) crystals, almost completely blocks all aspects of inflammasome activation and pyroptosis induced by ATP and MSU crystals in PD (Youm et al., 2015; Deora et al., 2017). Crucially, modern studies suggest that the NLRP3 inflammasome could be a major disease-modifying therapeutic target in PD's inflammatory pathogenesis. For instance, miR-7 directly targets NLRP3 expression (besides α -syn) and modulates NLRP3 inflammasome activation to attenuate DA neuronal degeneration accompanied by the amelioration of microglial activation in an MPTP-induced mouse model of PD (Zhou et al., 2016). In the heightened microglial activation response, an exaggerated ROS/c-Abelson murine leukemia viral oncogene homolog (c-Abl)/NLRP3 signaling axis evaluates in LPS-primed rotenone (ROT)-stimulated microglial cells and suggests that targeting c-Abl-regulated NLRP3 inflammasome signaling offers a novel therapeutic strategy for PD treatment (Lawana et al., 2017). Parkin deficiency modulates NLRP3 inflammasome activation by attenuating an A20-dependent negative feedback loop in Parkin's pathogenesis (PARK2)-linked PD, paving the way for the exploration of its potential as a biomarker and treatment target (Mouton-Liger et al., 2018).

CONCLUSION

Chronic inflammation of the CNS is mediated by neuroimmune microglial cells and has been implicated as a pathological contributor to PD. The activation of microglia and DA neuronal damage form a self-propelled degeneration cycle in PD; thus, microglia are more likely to play critical roles in establishing and maintaining inflammatory responses in PD.

REFERENCES

- Abbas, N., Lucking, C. B., Ricard, S., Durr, A., Bonifati, V., De Michele, G., et al. (1999). A wide variety of mutations in the parkin gene are responsible for autosomal recessive parkinsonism in Europe. French Parkinson's disease genetics study group and the European consortium on genetic susceptibility in Parkinson's disease. *Hum. Mol. Genet.* 8, 567–574. doi: 10.1093/hmg/8.4.567
- Aguirre, J. D., Dunkerley, K. M., Lam, R., Rusal, M., and Shaw, G. S. (2018). Impact of altered phosphorylation on loss of function of juvenile Parkinsonism-associated genetic variants of the E3 ligase parkin. *J. Biol. Chem.* 293, 6337–6348. doi: 10.1074/jbc.RA117.000605

Currently, the signaling pathways and molecular factors involved in neuroinflammation have become an important research method to identify PD's pathogenesis. The anti-inflammatory treatment has been found to exert a robust neuroprotective effect in a mouse model of PD. Studies on animal and cell models of PD have shown that dietary supplements containing polyphenolic compounds have beneficial effects and are recommended for treating and preventing inflammation-mediated neurodegeneration of DA neurons (Singh et al., 2020).

Studies have shown that blocking these signaling pathways and molecular factors can effectively improve apoptosis or the death of dopamine neurons caused by neuroinflammation. This paradigm is being shifted from theory to reality as a potential target for developing new drugs for PD. Going forward, focusing on these signaling pathways and molecular factors involved in neuroinflammation would provide a better understanding of the occurrence and development of PD. Ongoing research in this field may open a new door for developing pharmacological strategies toward the prevention and modification of the pathogenesis of PD.

AUTHOR CONTRIBUTIONS

LY and JW performed the majority of the literature search and predominantly contributed to the writing of the article. SK and GL assisted with the literature search. All authors read and approved the final manuscript.

FUNDING

This study was supported by the National Natural Science Foundation of China (82060249), Key project of National Natural Science Foundation of Jiangxi Province (20202ACBL206005), General project of Natural Science Foundation of Jiangxi Province (20192BAB205042), General project of Natural Science Foundation of Jiangxi Province (20202BABL206098), and the major academic and technical leaders training plan of Jiangxi Province-Youth Training Program (20204BCJ23019).

ACKNOWLEDGMENTS

We would like to thank Yuqing Wang for helping us to draw the illustrations.

- Akabane, S., Uno, M., Tani, N., Shimazaki, S., Ebara, N., Kato, H., et al. (2016). PKA regulates PINK1 stability and parkin recruitment to damaged mitochondria through phosphorylation of MIC60. *Mol. Cell* 62, 371–384. doi: 10.1016/j.molcel.2016.03.037
- Alvarez-Erviti, L., Couch, Y., Richardson, J., Cooper, J. M., and Wood, M. J. (2011). Alpha-synuclein release by neurons activates the inflammatory response in a microglial cell line. *Neurosci. Res.* 69, 337–342. doi: 10.1016/j.neures.2010.12.020
- Anitua, E., Pascual, C., érez-Gonzalez, R. P., Orive, G., and Carro, E. (2015). Intranasal PRGF-Endoret enhances neuronal survival and attenuates NF- κ B-dependent inflammation process in a mouse model of Parkinson's disease. *J. Control. Release* 203, 170–180.

- Aoki, E., Yano, R., Yokoyama, H., Kato, H., and Araki, T. (2009). Role of nuclear transcription factor kappa B (NF-kappaB) for MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine)-induced apoptosis in nigral neurons of mice. *Exp. Mol. Pathol.* 86, 57–64. doi: 10.1016/j.yexmp.2008.10.004
- Arkinson, C., and Walden, H. (2018). Parkin function in Parkinson's disease. *Science* 360, 267–268. doi: 10.1126/science.aar6606
- Azizi, S. A., and Azizi, S.-A. (2018). Synucleinopathies in neurodegenerative diseases: accomplices, an inside job and selective vulnerability. *Neurosci. Lett.* 672, 150–152.
- Bassani, T. B., Vital, M. A., and Rauh, L. K. (2015). Neuroinflammation in the pathophysiology of Parkinson's disease and therapeutic evidence of anti-inflammatory drugs. *Arq. Neuropsiquiatr.* 73, 616–623. doi: 10.1590/0004-282X20150057
- Birla, H., Rai, S. N., Singh, S. S., Zahra, W., Rawat, A., Tiwari, N., et al. (2019). *Tinospora cordifolia* Suppresses neuroinflammation in Parkinsonian mouse model. *Neuromol. Med.* 21, 42–53. doi: 10.1007/s12017-018-08521-7
- Boyden, E. D., and Dietrich, W. F. (2006). Nalp1b controls mouse macrophage susceptibility to anthrax lethal toxin. *Nat. Genet.* 38, 240–244. doi: 10.1038/ng1724
- Brockmann, K., Apel, A., Schulte, C., Schneiderhan-Marra, N., Pont-Sunyer, C., Vilas, D., et al. (2016). Inflammatory profile in LRRK2-associated prodromal and clinical PD. *J. Neuroinflammation* 13:122. doi: 10.1186/s12974-016-0588-5
- Brundin, L., Bergkvist, L., and Brundin, P. (2018). Fire prevention in the Parkinson's disease brain. *Nat. Med.* 24, 900–902. doi: 10.1038/s41591-018-0109-4
- Calatayud, C., Carola, G., Consiglio, A., and Raya, A. (2017). Modeling the genetic complexity of Parkinson's disease by targeted genome edition in iPS cells. *Curr. Opin. Genet. Dev.* 46, 123–131.
- Cha, G. H., Kim, S., Park, J., Lee, E., Kim, M., Lee, S. B., et al. (2005). Parkin negatively regulates JNK pathway in the dopaminergic neurons of *Drosophila*. *Proc. Natl. Acad. Sci. U.S.A.* 102, 10345–10350. doi: 10.1073/pnas.0500346102
- Chan, S. L., and Tan, E. K. (2017). Targeting LRRK2 in Parkinson's disease: an update on recent developments. *Expert Opin. Ther. Targets* 21, 601–610.
- Charvin, D., Medori, R., Hauser, R. A., and Rascol, O. (2018). Therapeutic strategies for Parkinson disease: beyond dopaminergic drugs. *Nat. Rev. Drug Discov.* 17, 804–822. doi: 10.1038/nrd.2018.136
- Chen, H., Zhang, S. M., Hernan, M. A., Schwarzschild, M. A., Willett, W. C., Colditz, G. A., et al. (2003). Nonsteroidal anti-inflammatory drugs and the risk of Parkinson disease. *Arch. Neurol.* 60, 1059–1064. doi: 10.1001/archneur.60.8.1059
- Chen, J., Su, P., Luo, W., and Chen, J. (2018). Role of LRRK2 in manganese-induced neuroinflammation and microglial autophagy. *Biochem. Biophys. Res. Commun.* 498, 171–177. doi: 10.1016/j.bbrc.2018.02.007
- Chen, Y., Zhang, N., Ji, D., Hou, Y., Chen, C., Fu, Y., et al. (2018). Dysregulation of bcl-2 enhanced rotenone-induced alpha-synuclein aggregation associated with autophagic pathways. *Neuroreport* 29, 1201–1208. doi: 10.1097/WNR.0000000000001097
- Cheng, C., and Zhu, X. (2019). Cordycepin mitigates MPTP-induced Parkinson's disease through inhibiting TLR/NF-kappaB signaling pathway. *Life Sci.* 223, 120–127. doi: 10.1016/j.lfs.2019.02.037
- Chistiakov, D. A., and Chistiakov, A. A. (2017). alpha-Synuclein-carrying extracellular vesicles in Parkinson's disease: deadly transmitters. *Acta Neurol. Belg.* 117, 43–51. doi: 10.1007/s13760-016-0679-1
- Choi, I., Kim, B., Byun, J. W., Baik, S. H., Huh, Y. H., Kim, J. H., et al. (2015). LRRK2 G2019S mutation attenuates microglial motility by inhibiting focal adhesion kinase. *Nat. Commun.* 6:8255. doi: 10.1038/ncomms9255
- Choubey, V., Safulina, D., Vaarmann, A., Cagalinec, M., Wareski, P., Kuum, M., et al. (2011). Mutant A53T alpha-synuclein induces neuronal death by increasing mitochondrial autophagy. *J. Biol. Chem.* 286, 10814–10824. doi: 10.1074/jbc.M110.132514
- Cornelissen, T., Vilain, S., Vints, K., Gounko, N., Verstreken, P., and Vandenbergh, W. (2018). Deficiency of parkin and PINK1 impairs age-dependent mitophagy in *Drosophila*. *Elife* 7:e35878. doi: 10.7554/eLife.35878
- Cuttillo, G., Simon, D. K., and Eleuteri, S. (2020). VPS35 and the mitochondria: connecting the dots in Parkinson's disease pathophysiology. *Neurobiol. Dis.* 145:105056.
- de Araujo, D. P., Nogueira, P. C. N., Santos, A. D. C., Costa, R. O., de Lucena, J. D., Jataí Gadelha-Filho, C. V., et al. (2018). Viana: *Aspidosperma pyrifolium* Mart: neuroprotective, antioxidant and anti-inflammatory effects in a Parkinson's disease model in rats. *J. Pharm. Pharmacol.* 70, 787–796. doi: 10.1111/jphp.12866
- de Rivero Vaccari, J. P., Dietrich, W. D., and Keane, R. W. (2014). Activation and regulation of cellular inflammasomes: gaps in our knowledge for central nervous system injury. *J. Cereb. Blood Flow Metab.* 34, 369–375. doi: 10.1038/jcbfm.2013.227
- Dehmer, T., Heneka, M. T., Sastre, M., Dichgans, J., and Schulz, J. B. (2004). Protection by pioglitazone in the MPTP model of Parkinson's disease correlates with IkB α induction and block of NF κ B and iNOS activation. *J. Neurochem.* 88, 494–501.
- Deng, H., Fan, K., and Jankovic, J. (2018). The role of TMEM230 gene in Parkinson's disease. *J. Parkinsons Dis.* 8, 469–477.
- Deora, V., Albornoz, E. A., Zhu, K., Woodruff, T. M., and Gordon, R. (2017). The ketone body beta-hydroxybutyrate does not inhibit synuclein mediated inflammasome activation in microglia. *J. Neuroimmune Pharmacol.* 12, 568–574. doi: 10.1007/s11481-017-9754-5
- Dhir, A., Dhir, S., Borowski, L. S., Jimenez, L., Teitell, M., Rotig, A., et al. (2018). Mitochondrial double-stranded RNA triggers antiviral signalling in humans. *Nature* 560, 238–242. doi: 10.1038/s41586-018-0363-0
- Di Maio, R., Hoffman, E. K., Rocha, E. M., Keeney, M. T., Sanders, L. H., De Miranda, B. R., et al. (2018). LRRK2 activation in idiopathic Parkinson's disease. *Sci. Transl. Med.* 10:eaar5429. doi: 10.1126/scitranslmed.aar5429
- Dionisio, P. E. A., Oliveira, S. R., Amaral, J., and Rodrigues, C. M. P. (2019). Loss of microglial parkin inhibits necroptosis and contributes to neuroinflammation. *Mol. Neurobiol.* 56, 2990–3004. doi: 10.1007/s12035-018-1264-9
- Ehgoetz Martens, K. A., and Shine, J. M. (2018). The interactions between non-motor symptoms of Parkinson's disease. *Expert Rev. Neurother.* 18, 457–460. doi: 10.1080/14737175.2018.1472578
- Esposito, E., Di Matteo, V., Benigno, A., Pierucci, M., Crescimanno, G., and Di Giovanni, G. (2007). Non-steroidal anti-inflammatory drugs in Parkinson's disease. *Exp. Neurol.* 205, 295–312. doi: 10.1016/j.expneurol.2007.02.008
- Ettle, B., Kuhbandner, K., Jorg, S., Hoffmann, A., Winkler, J., and Linker, R. A. (2016). alpha-Synuclein deficiency promotes neuroinflammation by increasing Th1 cell-mediated immune responses. *J. Neuroinflammation* 13:201. doi: 10.1186/s12974-016-0694-4
- Fan, F. S. (2020). Ractopamine residue in meat might protect people from Parkinson disease. *Med. Hypotheses* 145:110397.
- Ferreira, M., and Massano, J. (2017). An updated review of Parkinson's disease genetics and clinicopathological correlations. *Acta Neurol. Scand.* 135, 273–284. doi: 10.1111/ane.12616
- Forero, D. A., Trujillo, M. L., and Lopez-Leon, S. (2020). "Genome plasticity and neuropsychiatric disorders," in *Genome Plasticity in Health and Disease*, eds D. A. Forero and G. P. Patrinos (Amsterdam: Elsevier).
- Fu, S. P., Wang, J. F., Xue, W. J., Liu, H. M., Liu, B. R., Zeng, Y. L., et al. (2015). Anti-inflammatory effects of BHBa in both in vivo and in vitro Parkinson's disease models are mediated by GPR109A-dependent mechanisms. *J. Neuroinflammation* 12:9. doi: 10.1186/s12974-014-0230-3
- Gago, M., Machado, A., and Rocha, S. (2020). "Current clinical approaches in neurodegenerative diseases," in *Handbook of Innovations in Central Nervous System Regenerative Medicine*, ed. A. J. Salgado (Amsterdam: Elsevier), 79.
- Gan, P., Xia, Q., Hang, G., Zhou, Y., Qian, X., Wang, X., et al. (2018). Knockdown of cathepsin D protects dopaminergic neurons against neuroinflammation-mediated neurotoxicity through inhibition of NF-kappaB signalling pathway in Parkinson's disease model. *Clin. Exp. Pharmacol. Physiol.* 46, 337–349. doi: 10.1111/1440-1681.13052
- Gao, J., Li, Y., Li, W., and Wang, H. (2019). TrxR2 overexpression alleviates inflammation-mediated neuronal death via reducing the oxidative stress and activating the Akt-Parkin pathway. *Toxicol. Res. (Camb.)* 8, 641–653. doi: 10.1039/c9tx00076c
- Giacoppo, S., Rajan, T. S., De Nicola, G. R., Iori, R., Rollin, P., Bramanti, P., et al. (2017). The isothiocyanate isolated from *Moringa oleifera* shows potent anti-inflammatory activity in the treatment of murine subacute Parkinson's disease. *Rejuvenation Res.* 20, 50–63. doi: 10.1089/rej.2016.1828
- Goes, A. T. R., Jesse, C. R., Antunes, M. S., Lobo Ladd, F. V., Lobo Ladd, A. A. B., Luchese, C., et al. (2018). Protective role of chrysin on 6-hydroxydopamine-induced neurodegeneration a mouse model of Parkinson's disease: involvement

- of neuroinflammation and neurotrophins. *Chem. Biol. Interact.* 279, 111–120. doi: 10.1016/j.cbi.2017.10.019
- Goldwurm, S., Zini, M., Mariani, L., Tesi, S., Miceli, R., Sironi, F., et al. (2007). Evaluation of LRRK2 G2019S penetrance: relevance for genetic counseling in Parkinson disease. *Neurology* 68, 1141–1143. doi: 10.1212/01.wnl.0000254483.19854.ef
- Greggio, E., Bubacco, L., and Russo, I. (2017). Cross-talk between LRRK2 and PKA: implication for Parkinson's disease? *Biochem. Soc. Trans.* 45, 261–267. doi: 10.1042/bst20160396
- Guan, Y., Li, Y., Zhao, G., and Li, Y. (2018). HMGB1 promotes the starvation-induced autophagic degradation of alpha-synuclein in SH-SY5Y cells Atg 5-dependently. *Life Sci.* 202, 1–10. doi: 10.1016/j.lfs.2018.03.031
- Han, Z., Wei, B., Zhang, C., Zhu, H., Tang, L., Lin, R., et al. (2019). PARK16 rs708730 polymorphism decreases Parkinson's disease risk in European ancestry population: a meta-analysis. *J. Geriatr. Med.* 1, 15–22.
- Harms, A. S., Cao, S., Rowse, A. L., Thome, A. D., Li, X., Mangieri, L. R., et al. (2013). MHCII is required for alpha-synuclein-induced activation of microglia, CD4 T cell proliferation, and dopaminergic neurodegeneration. *J. Neurosci.* 33, 9592–9600. doi: 10.1523/JNEUROSCI.5610-12.2013
- Harms, A. S., Thome, A. D., Yan, Z., Schonhoff, A. M., Williams, G. P., Li, X., et al. (2018). Peripheral monocyte entry is required for alpha-Synuclein induced inflammation and Neurodegeneration in a model of Parkinson disease. *Exp. Neurol.* 300, 179–187. doi: 10.1016/j.expneurol.2017.11.010
- Heng, Y., Zhang, Q. S., Mu, Z., Hu, J. F., Yuan, Y. H., and Chen, N. H. (2016). Ginsenoside Rg1 attenuates motor impairment and neuroinflammation in the MPTP-probenecid-induced parkinsonism mouse model by targeting alpha-synuclein abnormalities in the substantia nigra. *Toxicol. Lett.* 243, 7–21. doi: 10.1016/j.toxlet.2015.12.005
- Higashi, Y., Asanuma, M., Miyazaki, I., Hattori, N., Mizuno, Y., and Ogawa, N. (2004). Parkin attenuates manganese-induced dopaminergic cell death. *J. Neurochem.* 89, 1490–1497. doi: 10.1111/j.1471-4159.2004.02445.x
- Hilla, A. M., Diekmann, H., and Fischer, D. (2017). Microglia are irrelevant for neuronal degeneration and axon regeneration after acute injury. *J. Neurosci.* 37, 6113–6124. doi: 10.1523/jneurosci.0584-17.2017
- Ho, D. H., Je, A. R., Lee, H., Son, I., Kweon, H. S., Kim, H. G., et al. (2018). LRRK2 kinase activity induces mitochondrial fission in microglia via Drp1 and modulates neuroinflammation. *Exp. Neurobiol.* 27, 171–180. doi: 10.5607/en.2018.27.3.171
- Hoffmann, A., Ettl, B., Bruno, A., Kulinich, A., Hoffmann, A. C., von Wittgenstein, J., et al. (2016). Alpha-synuclein activates BV2 microglia dependent on its aggregation state. *Biochem. Biophys. Res. Commun.* 479, 881–886. doi: 10.1016/j.bbrc.2016.09.109
- Hu, X. L., Lin, J., Lv, X. Y., Feng, J. H., Zhang, X. Q., Wang, H., et al. (2018). Synthesis and biological evaluation of clovamide analogues as potent anti-neuroinflammatory agents in vitro and in vivo. *Eur. J. Med. Chem.* 151, 261–271. doi: 10.1016/j.ejmech.2018.03.081
- Hu, Z., Wang, W., Ling, J., and Jiang, C. (2016). alpha-mangostin inhibits alpha-Synuclein-induced microglial neuroinflammation and neurotoxicity. *Cell. Mol. Neurobiol.* 36, 811–820. doi: 10.1007/s10571-015-0264-9
- Huang, B., Liu, J., Meng, T., Li, Y., He, D., Ran, X., et al. (2018). Polydatin prevents lipopolysaccharide (LPS)-induced Parkinson's disease via regulation of the AKT/GSK3beta-Nrf2/NF-kappaB signaling axis. *Front. Immunol.* 9, 2527. doi: 10.3389/fimmu.2018.02527
- Huang, D., Liu, M., and Jiang, Y. (2019). Mitochondrial acid-5 attenuates TNF-alpha-mediated neuronal inflammation via activating Parkin-related mitophagy and augmenting the AMPK-Sirt3 pathways. *J. Cell. Physiol.* 234, 22172–22182. doi: 10.1002/jcp.28783
- Ikedo-Matsuo, Y., Miyata, H., Mizoguchi, T., Ohama, E., Naito, Y., Uematsu, S., et al. (2019). Microsomal prostaglandin E synthase-1 is a critical factor in dopaminergic neurodegeneration in Parkinson's disease. *Neurobiol. Dis.* 124, 81–92. doi: 10.1016/j.nbd.2018.11.004
- Jiang, H., Ren, Y., Zhao, J., and Feng, J. (2004). Parkin protects human dopaminergic neuroblastoma cells against dopamine-induced apoptosis. *Hum. Mol. Genet.* 13, 1745–1754. doi: 10.1093/hmg/ddh180
- Kachidian, P., and Gubellini, P. (2021). "Genetic models of Parkinson's disease," in *Clinical Trials In Parkinson's Disease*, ed. S. Perez-Lloret (New York, NY: Springer).
- Kawai, T., and Akira, S. (2007). Signaling to NF-kappaB by Toll-like receptors. *Trends Mol. Med.* 13, 460–469. doi: 10.1016/j.molmed.2007.09.002
- Kawajiri, S., Saiki, S., Sato, S., and Hattori, N. (2011). Genetic mutations and functions of PINK1. *Trends Pharmacol. Sci.* 32, 573–580. doi: 10.1016/j.tips.2011.06.001
- Kim, B., Yang, M. S., Choi, D., Kim, J. H., Kim, H. S., Seol, W., et al. (2012). Impaired inflammatory responses in murine Lrrk2-knockdown brain microglia. *PLoS One* 7:e34693. doi: 10.1371/journal.pone.0034693
- Kim, C., Ho, D.-H., Suk, J.-E., You, S., Michael, S., Kang, J., et al. (2013). Neuron-released oligomeric alpha-synuclein is an endogenous agonist of TLR2 for paracrine activation of microglia. *Nat. Commun.* 4:1562.
- Kim, C., Ojo-Amaize, E., Spencer, B., Rockenstein, E., Mante, M., Desplats, P., et al. (2015). Hypoestoxide reduces neuroinflammation and alpha-synuclein accumulation in a mouse model of Parkinson's disease. *J. Neuroinflammation* 12:236. doi: 10.1186/s12974-015-0455-9
- Kim, C., Spencer, B., Rockenstein, E., Yamakado, H., Mante, M., Adame, A., et al. (2018). Immunotherapy targeting toll-like receptor 2 alleviates neurodegeneration in models of synucleinopathy by modulating alpha-synuclein transmission and neuroinflammation. *Mol. Neurodegener.* 13:43. doi: 10.1186/s13024-018-0276-2
- Kim, K. S., Marcogliese, P. C., Yang, J., Callaghan, S. M., Resende, V., Abdel-Messih, E., et al. (2018). Lrrk2 in Inflammation and D. S. Park: regulation of myeloid cell phagocytosis by LRRK2 via WAVE2 complex stabilization is altered in Parkinson's disease. *Proc. Natl. Acad. Sci. U.S.A.* 115, E5164–E5173. doi: 10.1073/pnas.1718946115
- Kluss, J. H., Mamais, A., and Cookson, M. R. (2019). LRRK2 links genetic and sporadic Parkinson's disease. *Biochem. Soc. Trans.* 47, 651–661. doi: 10.1042/BST20180462
- Koshibu, K., van Asperen, J., Gerets, H., Garcia-Ladona, J., Lorthioir, O., and Courade, J. P. (2015). Alternative to LRRK2-IN-1 for pharmacological studies of Parkinson's disease. *Pharmacology* 96, 240–247. doi: 10.1159/000439136
- Kozina, E., Sadasivan, S., Jiao, Y., Dou, Y., Ma, Z., Tan, H., et al. (2018). Mutant LRRK2 mediates peripheral and central immune responses leading to neurodegeneration in vivo. *Brain* 141, 1753–1769. doi: 10.1093/brain/awy077
- Lawana, V., Singh, N., Sarkar, S., Charli, A., Jin, H., Anantharam, V., et al. (2017). Involvement of c-Abl kinase in microglial activation of NLRP3 inflammasome and impairment in autolysosomal system. *J. Neuroimmune Pharmacol.* 12, 624–660. doi: 10.1007/s11481-017-9746-5
- Lee, B. D., Shin, J. H., VanKampen, J., Petrucelli, L., West, A. B., Ko, H. S., et al. (2010). Inhibitors of leucine-rich repeat kinase-2 protect against models of Parkinson's disease. *Nat. Med.* 16, 998–1000. doi: 10.1038/nm.2199
- Lee, J. A., Kim, J. H., Woo, S. Y., Son, H. J., Han, S. H., Jang, B. K., et al. (2015). A novel compound VSC2 has anti-inflammatory and antioxidant properties in microglia and in Parkinson's disease animal model. *Br. J. Pharmacol.* 172, 1087–1100. doi: 10.1111/bph.12973
- Lee, K. M., Lee, Y., Chun, H. J., Kim, A. H., Kim, J. Y., Lee, J. Y., et al. (2016). Neuroprotective and anti-inflammatory effects of morin in a murine model of Parkinson's disease. *J. Neurosci. Res.* 94, 865–878. doi: 10.1002/jnr.23764
- Leonoudakis, D., Rane, A., Angeli, S., Lithgow, G. J., Andersen, J. K., and Chinta, S. J. (2017). Anti-inflammatory and neuroprotective role of natural product securinine in activated glial cells: implications for Parkinson's disease. *Mediators Inflamm.* 2017:8302636. doi: 10.1155/2017/8302636
- Li, D., Mastaglia, F. L., Fletcher, S., and Wilton, S. D. (2020). Progress in the molecular pathogenesis and nucleic acid therapeutics for Parkinson's disease in the precision medicine era. *Med. Res. Rev.* 40, 2650–2681.
- Li, T., Yang, D., Zhong, S., Thomas, J. M., Xue, F., Liu, J., et al. (2014). Novel LRRK2 GTP-binding inhibitors reduced degeneration in Parkinson's disease cell and mouse models. *Hum. Mol. Genet.* 23, 6212–6222. doi: 10.1093/hmg/ddu341
- Li, Y., Ibañez, D. P., Fan, W., Zhao, P., Chen, S., Abdul, M. M., et al. (2020). Generation of an induced pluripotent stem cell line (GIBHi004-A) from a Parkinson's disease patient with mutant DJ-1/PARK7 (p. L10P). *Stem Cell Res.* 46:101845.
- Lima, F. A. V., Joventino, I. P., Joventino, F. P., de Almeida, A. C., Neves, K. R. T., do Carmo, M. R., et al. (2017). Neuroprotective activities of *Spirulina platensis* in the 6-OHDA Model of Parkinson's disease are related to its anti-inflammatory effects. *Neurochem. Res.* 42, 3390–3400. doi: 10.1007/s11064-017-2379-5

- Liu, Z. H., Lee, J., Krummey, S., Lu, W., Cai, H. B., and Lenardo, M. J. (2011). The kinase LRRK2 is a regulator of the transcription factor NFAT that modulates the severity of inflammatory bowel disease. *Nat. Immunol.* 12, 1063–1070. doi: 10.1038/ni.2113
- Lonskaya, I., Hebron, M. L., Algarzae, N. K., Desforgues, N., and Moussa, C. E. (2013). Decreased parkin solubility is associated with impairment of autophagy in the nigrostriatum of sporadic Parkinson's disease. *Neuroscience* 232, 90–105. doi: 10.1016/j.neuroscience.2012.12.018
- Lu, C., Zhang, J., Shi, X., Miao, S., Bi, L., Zhang, S., et al. (2014). Neuroprotective effects of tetramethylpyrazine against dopaminergic neuron injury in a rat model of Parkinson's disease induced by MPTP. *Int. J. Biol. Sci.* 10, 350–357. doi: 10.7150/ijbs.8366
- Lv, R., Du, L., Liu, X., Zhou, F., Zhang, Z., and Zhang, L. (2019). Rosmarinic acid attenuates inflammatory responses through inhibiting HMGB1/TLR4/NF-kappaB signaling pathway in a mouse model of Parkinson's disease. *Life Sci.* 223, 158–165. doi: 10.1016/j.lfs.2019.03.030
- Lv, Y., Zhang, Z., Hou, L., Zhang, L., Zhang, J., Wang, Y., et al. (2015). Phytic acid attenuates inflammatory responses and the levels of NF-kappaB and p-ERK in MPTP-induced Parkinson's disease model of mice. *Neurosci. Lett.* 597, 132–136. doi: 10.1016/j.neulet.2015.04.040
- Ma, B., Xu, L., Pan, X., Sun, L., Ding, J., Xie, C., et al. (2016). LRRK2 modulates microglial activity through regulation of chemokine (C-X3-C) receptor 1-mediated signalling pathways. *Hum. Mol. Genet.* 25, 3515–3523. doi: 10.1093/hmg/ddw194
- Maekawa, T., Sasaoka, T., Azuma, S., Ichikawa, T., Melrose, H. L., Farrer, M. J., et al. (2016). Leucine-rich repeat kinase 2 (LRRK2) regulates alpha-synuclein clearance in microglia. *BMC Neurosci.* 17:77. doi: 10.1186/s12868-016-0315-2
- Mamik, M. K., and Power, C. (2017). Inflammasomes in neurological diseases: emerging pathogenic and therapeutic concepts. *Brain* 140, 2273–2285. doi: 10.1093/brain/awx133
- Mariathasan, S., Newton, K., Monack, D. M., Vucic, D., French, D. M., Lee, W. P., et al. (2004). Differential activation of the inflammasome by caspase-1 adaptors ASC and Ipaf. *Nature* 430, 213–218. doi: 10.1038/nature02664
- Mariathasan, S., Weiss, D. S., Newton, K., McBride, J., O'Rourke, K., Roose-Girma, M., et al. (2006). Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature* 440, 228–232. doi: 10.1038/nature04515
- Matheoud, D., Cannon, T., Voisin, A., Penttinen, A. M., Ramet, L., Fahmy, A. M., et al. (2019). Intestinal infection triggers Parkinson's disease-like symptoms in Pink1(-/-) mice. *Nature* 571, 565–569. doi: 10.1038/s41586-019-1405-y
- Mattson, M. P., and Camandola, S. (2001). Camandola: NF-kB in neuronal plasticity and neurodegenerative disorders. *J. Clin. Invest.* 107, 247–254.
- Mazo, N. A., Echeverria, V., Cabezas, R., vila-Rodriguez, M. A., Tarasov, V. V., Yarla, N. S., et al. (2017). Medicinal plants as protective strategies against Parkinson's disease. *Curr. Pharm. Des.* 23, 4180–4188. doi: 10.2174/1381612823666170316142803
- McWilliams, T. G., and Muqit, M. M. (2017). PINK1 and Parkin: emerging themes in mitochondrial homeostasis. *Curr. Opin. Cell Biol.* 45, 83–91. doi: 10.1016/j.ceb.2017.03.013
- Mendes, M. O., Rosa, A. I., Carvalho, A. N., Nunes, M. J., Dionisio, P., Rodrigues, E., et al. (2019). Neurotoxic effects of MPTP on mouse cerebral cortex: modulation of neuroinflammation as a neuroprotective strategy. *Mol. Cell. Neurosci.* 96, 1–9. doi: 10.1016/j.mcn.2019.01.003
- Michel, H. E., Tadros, M. G., Esmat, A., Khalifa, A. E., and Abdel-Tawab, A. M. (2017). Tetramethylpyrazine ameliorates rotenone-induced Parkinson's disease in rats: involvement of its anti-inflammatory and anti-apoptotic actions. *Mol. Neurobiol.* 54, 4866–4878. doi: 10.1007/s12035-016-0028-7
- Moehle, M. S., Webber, P. J., Tse, T., Sukar, N., Standaert, D. G., DeSilva, T. M., et al. (2012). LRRK2 inhibition attenuates microglial inflammatory responses. *J. Neurosci.* 32, 1602–1611. doi: 10.1523/JNEUROSCI.5601-11.2012
- Mondal, S., Roy, A., Jana, A., Ghosh, S., Kordower, J. H., and Pahan, K. (2012). Testing NF-kappaB-based therapy in hemiparkinsonian monkeys. *J. Neuroimmune Pharmacol.* 7, 544–556. doi: 10.1007/s11481-012-9377-9
- Morato Torres, C. A., Wassouf, Z., Zafar, F., Sastre, D., Outeiro, T. F., and Schüle, B. (2020). The role of alpha-synuclein and other Parkinson's genes in neurodevelopmental and neurodegenerative disorders. *Int. J. Mol. Sci.* 21:5724.
- Mouton-Liger, F., Rosazza, T., Sepulveda-Diaz, J., Ieang, A., Hassoun, S. M., Claire, E., et al. (2018). Parkin deficiency modulates NLRP3 inflammasome activation by attenuating an A20-dependent negative feedback loop. *Glia* 66, 1736–1751. doi: 10.1002/glia.23337
- Muda, K., Bertinetti, D., Gesellchen, F., Hermann, J. S., von Zweydford, F., Geerlof, A., et al. (2014). Parkinson-related LRRK2 mutation R1441C/G/H impairs PKA phosphorylation of LRRK2 and disrupts its interaction with 14-3-3. *Proc. Natl. Acad. Sci. U.S.A.* 111, E34–E43. doi: 10.1073/pnas.1312701111
- Newman, L. E., and Shadel, G. S. (2018). Pink1/Parkin link inflammation, mitochondrial stress, and neurodegeneration. *J. Cell. Biol.* 217, 3327–3329. doi: 10.1083/jcb.201808118
- Phani, S., Loike, J. D., and Przedborski, S. (2012). Neurodegeneration and inflammation in Parkinson's disease. *Parkinsonism Relat. Disord.* 18(Suppl. 1), S207–S209. doi: 10.1016/S1353-8020(11)70064-5
- Pickrell, A. M., and Youle, R. J. (2015). The roles of PINK1, parkin, and mitochondrial fidelity in Parkinson's disease. *Neuron* 85, 257–273. doi: 10.1016/j.neuron.2014.12.007
- Poly, T. N., Islam, M. M. R., Yang, H. C., and Li, Y. J. (2019). Non-steroidal anti-inflammatory drugs and risk of Parkinson's disease in the elderly population: a meta-analysis. *Eur. J. Clin. Pharmacol.* 75, 99–108. doi: 10.1007/s00228-018-2561-y
- Puccini, J. M., Marker, D. F., Fitzgerald, T., Barbieri, J., Kim, C. S., Miller-Rhodes, P., et al. (2015). Leucine-rich repeat kinase 2 modulates neuroinflammation and neurotoxicity in models of human immunodeficiency virus 1-associated neurocognitive disorders. *J. Neurosci.* 35, 5271–5283. doi: 10.1523/jneurosci.0650-14.2015
- Qin, H., Buckley, J. A., Li, X., Liu, Y., Fox, T. H. III, Meares, G. P., et al. (2016). Inhibition of the JAK/STAT pathway protects against alpha-synuclein-induced neuroinflammation and dopaminergic neurodegeneration. *J. Neurosci.* 36, 5144–5159. doi: 10.1523/JNEUROSCI.4658-15.2016
- Rai, S. N., Birla, H., Singh, S. S., Zahra, W., Patil, R. R., Jadhav, J. P., et al. (2017). Mucuna pruriens protects against MPTP intoxicated Neuroinflammation in Parkinson's disease through NF-kB/PAKT signaling pathways. *Front. Aging Neurosci.* 9:421. doi: 10.3389/fnagi.2017.00421
- Rai, S. N., Zahra, W., Singh, S. S., Birla, H., Keswani, C., Dilmashin, H., et al. (2019). Anti-inflammatory activity of ursolic acid in MPTP-induced Parkinsonism mouse model. *Neurotox. Res.* 36, 452–462. doi: 10.1007/s12640-019-00038-6
- Rathinam, V. A., Jiang, Z., Waggoner, S. N., Sharma, S., Cole, L. E., Waggoner, L., et al. (2010). The AIM2 inflammasome is essential for host defense against cytosolic bacteria and DNA viruses. *Nat. Immunol.* 11, 395–402. doi: 10.1038/ni.1864
- Rey, N. L., George, S., Steiner, J. A., Madaj, Z., Luk, K. C., Trojanowski, J. Q., et al. (2018). Spread of aggregates after olfactory bulb injection of alpha-synuclein fibrils is associated with early neuronal loss and is reduced long term. *Acta Neuropathol.* 135, 65–83. doi: 10.1007/s00401-017-1792-9
- Rojanathamane, L., Murphy, E. J., and Combs, C. K. (2011). Expression of mutant alpha-synuclein modulates microglial phenotype in vitro. *J. Neuroinflammation* 8:44. doi: 10.1186/1742-2094-8-44
- Russo, I., Berti, G., Plotegher, N., Bernardo, G., Filograna, R., Bubacco, L., et al. (2015). Leucine-rich repeat kinase 2 positively regulates inflammation and down-regulates NF-kappaB p50 signaling in cultured microglia cells. *J. Neuroinflammation* 12:230. doi: 10.1186/s12974-015-0449-7
- Russo, I., Bubacco, L., and Greggio, E. (2014). LRRK2 and neuroinflammation: partners in crime in Parkinson's disease? *J. Neuroinflammation* 11:52. doi: 10.1186/1742-2094-11-52
- Saez-Atienzar, S., Bonet-Ponce, L., Blesa, J. R., Romero, F. J., Murphy, M. P., Jordan, J., et al. (2014). The LRRK2 inhibitor GSK2578215A induces protective autophagy in SH-SY5Y cells: involvement of Drp-1-mediated mitochondrial fission and mitochondrial-derived ROS signaling. *Cell Death Dis.* 5:e1368. doi: 10.1038/cddis.2014.320
- Sarrafchi, A., Bahmani, M., Shirzad, H., and Rafeian-Kopaei, M. (2016). Oxidative stress and Parkinson's disease: new hopes in treatment with herbal antioxidants. *Curr. Pharm. Des.* 22, 238–246. doi: 10.2174/1381612822666151112151653
- Schapira, A. H., Chaudhuri, K. R., and Jenner, P. (2017). Non-motor features of Parkinson disease. *Nat. Rev. Neurosci.* 18:435.
- Scholz, H., and Eder, C. (2017). Lysophosphatidylcholine activates caspase-1 in microglia via a novel pathway involving two inflammasomes. *J. Neuroimmunol.* 310, 107–110. doi: 10.1016/j.jneuroim.2017.07.004
- Scuruchi, M., D'Ascola, A., Avenoso, A., Campana, S., Abusamra, Y. A., Spina, E., et al. (2016). 6-Mer hyaluronan oligosaccharides modulate neuroinflammation and alpha-synuclein expression in neuron-Like SH-SY5Y Cells. *J. Cell. Biochem.* 117, 2835–2843. doi: 10.1002/jcb.25595

- Sharma, D., and Kanneganti, T. D. (2016). The cell biology of inflammasomes: Mechanisms of inflammasome activation and regulation. *J. Cell. Biol.* 213, 617–629. doi: 10.1083/jcb.201602089
- Sharma, N., and Nehru, B. (2018). Curcumin affords neuroprotection and inhibits alpha-synuclein aggregation in lipopolysaccharide-induced Parkinson's disease model. *Inflammopharmacology* 26, 349–360. doi: 10.1007/s10787-017-0402-8
- Sharma, N., Sharma, S., and Nehru, B. (2017). Curcumin protects dopaminergic neurons against inflammation-mediated damage and improves motor dysfunction induced by single intranigral lipopolysaccharide injection. *Inflammopharmacology* 25, 351–368. doi: 10.1007/s10787-017-0346-z
- Shih, R. H., Wang, C. Y., and Yang, C. M. (2015). NF-kappaB signaling pathways in neurological inflammation: a mini review. *Front. Mol. Neurosci.* 8:77. doi: 10.3389/fnmol.2015.00077
- Singh, S. S., Rai, S. N., Birla, H., Zahra, W., Kumar, G., Gedda, M. R., et al. (2018). Effect of chlorogenic acid supplementation in MPTP-intoxicated mouse. *Front. Pharmacol.* 9:757. doi: 10.3389/fphar.2018.00757
- Singh, S. S., Rai, S. N., Birla, H., Zahra, W., Rathore, A. S., and Singh, S. P. (2020). NF- κ B-mediated neuroinflammation in Parkinson's disease and potential therapeutic effect of polyphenols. *Neurotox. Res.* 37, 491–507. doi: 10.1007/s12640-019-00147-2
- Singhal, G., Jaehne, E. J., Corrigan, F., Toben, C., and Baune, B. T. (2014). Inflammasomes in neuroinflammation and changes in brain function: a focused review. *Front. Neurosci.* 8:315. doi: 10.3389/fnins.2014.00315
- Sita, G., Hrelia, P., Tarozzi, A., and Morroni, F. (2016). Isothiocyanates are promising compounds against oxidative stress, neuroinflammation and cell death that may benefit neurodegeneration in Parkinson's disease. *Int. J. Mol. Sci.* 17:1454. doi: 10.3390/ijms17091454
- Sliter, D. A., Martinez, J., Hao, L., Chen, X., Sun, N., Fischer, T. D., et al. (2018). Parkin and PINK1 mitigate STING-induced inflammation. *Nature* 561, 258–262. doi: 10.1038/s41586-018-0448-9
- Sokolova, O., and Naumann, M. (2017). NF-kappaB signaling in gastric cancer. *Toxins (Basel)* 9:119. doi: 10.3390/toxins9040119
- Spencer, J. P., Vafeiadou, K., Williams, R. J., and Vauzour, D. (2012). Neuroinflammation: modulation by flavonoids and mechanisms of action. *Mol. Aspects Med.* 33, 83–97. doi: 10.1016/j.mam.2011.10.016
- Steele, J. C., Guella, I., Szu-Tu, C., Lin, M. K., Thompson, C., Evans, D. M., et al. (2015). Defining neurodegeneration on Guam by targeted genomic sequencing. *Ann. Neurol.* 77, 458–468. doi: 10.1002/ana.23436
- Sudira, P. G., Subagya, S., and Sutarni, S. (2018). Aspek genetik dan manifestasi klinis varian young onset Parkinson disease. *Berkala NeuroSains* 17, 119–124.
- Takeda, A., Shinozaki, Y., Kashiwagi, K., Ohno, N., Eto, K., Wake, H., et al. (2018). Microglia mediate non-cell-autonomous cell death of retinal ganglion cells. *Glia* 66, 2366–2384. doi: 10.1002/glia.23475
- Tang, M. Y., Vranas, M., Krahn, A. I., Pundlik, S., Trempe, J. F., and Fon, E. A. (2017). Structure-guided mutagenesis reveals a hierarchical mechanism of Parkin activation. *Nat. Commun.* 8:14697. doi: 10.1038/ncomms14697
- Tentillier, N., Etzerodt, A., Olesen, M. N., Rizalar, F. S., Jacobsen, J., Bender, D., et al. (2016). Anti-inflammatory modulation of microglia via CD163-targeted glucocorticoids protects dopaminergic neurons in the 6-OHDA Parkinson's disease model. *J. Neurosci.* 36, 9375–9390. doi: 10.1523/jneurosci.1636-16.2016
- Tran, T. A., Nguyen, A. D., Chang, J., Goldberg, M. S., Lee, J. K., and Tansey, M. G. (2011). Lipopolysaccharide and tumor necrosis factor regulate Parkin expression via nuclear factor-kappa B. *PLoS One* 6:e23660. doi: 10.1371/journal.pone.0023660
- Truban, D., Hou, X., Caulfield, T. R., Fiesel, F. C., and Springer, W. (2017). PINK1, parkin, and mitochondrial quality control: what can we learn about Parkinson's disease pathobiology? *J. Parkinsons Dis.* 7, 13–29. doi: 10.3233/jpd-160989
- Tufekci, K. U., Meuwissen, R., Genc, S., and Genc, K. (2012). "Inflammation in Parkinson's disease," in *Advances in Protein Chemistry and Structural Biology*, ed. R. Donev (Amsterdam: Elsevier).
- van der Merwe, C., Jalali Sefid Dashti, Z., Christoffels, A., Loos, B., and Bardiën, S. (2015). Evidence for a common biological pathway linking three Parkinson's disease-causing genes: parkin, PINK1 and DJ-1. *Eur. J. Neurosci.* 41, 1113–1125. doi: 10.1111/ejn.12872
- van der Merwe, C., van Dyk, H. C., Engelbrecht, L., van der Westhuizen, F. H., Kinnear, C., Loos, B., et al. (2017). Curcumin rescues a PINK1 knock down SH-SY5Y cellular model of Parkinson's disease from mitochondrial dysfunction and cell death. *Mol. Neurobiol.* 54, 2752–2762. doi: 10.1007/s12035-016-9843-0
- Van der Perren, A., Macchi, F., Toelen, J., Carlon, M. S., Maris, M., de Loor, H., et al. (2015). FK506 reduces neuroinflammation and dopaminergic neurodegeneration in an alpha-synuclein-based rat model for Parkinson's disease. *Neurobiol. Aging* 36, 1559–1568. doi: 10.1016/j.neurobiolaging.2015.01.014
- Vekrellis, K., Xilouri, M., Emmanouilidou, E., Rideout, H. J., and Stefanis, L. (2011). Pathological roles of alpha-synuclein in neurological disorders. *Lancet Neurol.* 10, 1015–1025. doi: 10.1016/S1474-4422(11)70213-7
- Velez-Pardo, C., and Jimenez-Del-Rio, M. (2020). "Oxidative stress signaling and regulated cell death in Parkinson's disease," in *Genetics, Neurology, Behavior, and Diet in Parkinson's Disease*, eds C. R. Martin and V. R. Preedy (Amsterdam: Elsevier).
- Vivekanantham, S., Shah, S., Dewji, R., Dewji, A., Khatri, C., and Ologunde, R. (2015). Neuroinflammation in Parkinson's disease: role in neurodegeneration and tissue repair. *Int. J. Neurosci.* 125, 717–725. doi: 10.3109/00207454.2014.982795
- Wahabi, K., Perwez, A., and Rizvi, M. A. (2018). Parkin in Parkinson's disease and cancer: a double-edged sword. *Mol. Neurobiol.* 55, 6788–6800. doi: 10.1007/s12035-018-0879-1
- Wang, Y. D., Bao, X. Q., Xu, S., Yu, W. W., Cao, S. N., Hu, J. P., et al. (2016). A novel Parkinson's disease drug candidate with potent anti-neuroinflammatory effects through the Src signaling pathway. *J. Med. Chem.* 59, 9062–9079. doi: 10.1021/acs.jmedchem.6b00976
- West, A. P., Khoury-Hanold, W., Staron, M., Tal, M. C., Pineda, C. M., Lang, S. M., et al. (2015). Mitochondrial DNA stress primes the antiviral innate immune response. *Nature* 520, 553–557. doi: 10.1038/nature14156
- West, A. P., and Shadel, G. S. (2017). Mitochondrial DNA in innate immune responses and inflammatory pathology. *Nat. Rev. Immunol.* 17, 363–375. doi: 10.1038/nri.2017.21
- Whitworth, A. J., Theodore, D. A., Greene, J. C., Benes, H., Wes, P. D., and Pallanck, L. J. (2005). Increased glutathione S-transferase activity rescues dopaminergic neuron loss in a *Drosophila* model of Parkinson's disease. *Proc. Natl. Acad. Sci. U.S.A.* 102, 8024–8029. doi: 10.1073/pnas.0501078102
- Yadav, S. K., Rai, S. N., and Singh, S. P. (2017). Mucuna pruriens reduces inducible nitric oxide synthase expression in Parkinsonian mice model. *J. Chem. Neuroanat.* 80, 1–10. doi: 10.1016/j.jchemneu.2016.11.009
- Yan, T., Sun, Y., Gong, G., Li, Y., Fan, K., Wu, B., et al. (2019). The neuroprotective effect of schisandrol A on 6-OHDA-induced PD mice may be related to PI3K/AKT and IKK/IkappaB/NF-kappaB pathway. *Exp. Gerontol.* 128:110743. doi: 10.1016/j.exger.2019.110743
- Yao, L., Chen, H., Wu, Q., and Xie, K. (2019a). Hydrogen-rich saline alleviates inflammation and apoptosis in myocardial I/R injury via PINK-mediated autophagy. *Int. J. Mol. Med.* 44, 1048–1062. doi: 10.3892/ijmm.2019.4264
- Yao, L., Ye, Y., Mao, H., Lu, F., He, X., Lu, G., et al. (2018). MicroRNA-124 regulates the expression of MEKK3 in the inflammatory pathogenesis of Parkinson's disease. *J. Neuroinflammation* 15:13. doi: 10.1186/s12974-018-1053-4
- Yao, L., Zhu, Z., Wu, J., Zhang, Y., Zhang, H., Sun, X., et al. (2019b). MicroRNA-124 regulates the expression of p62/p38 and promotes autophagy in the inflammatory pathogenesis of Parkinson's disease. *FASEB J.* 33, 8648–8665. doi: 10.1096/fj.201900363R
- Yin, J., Valin, K. L., Dixon, M. L., and Leavenworth, J. W. (2017). The role of microglia and macrophages in CNS homeostasis, autoimmunity, and cancer. *J. Immunol. Res.* 2017:5150678. doi: 10.1155/2017/5150678
- Youm, Y. H., Nguyen, K. Y., Grant, R. W., Goldberg, E. L., Bodogai, M., Kim, D., et al. (2015). The ketone metabolite beta-hydroxybutyrate blocks NLRP3 inflammasome-mediated inflammatory disease. *Nat. Med.* 21, 263–269. doi: 10.1038/nm.3804
- Yuan, L., Song, Z., Deng, X., Yang, Z., Yang, Y., Guo, Y., et al. (2017). Genetic analysis of FBXO2, FBXO6, FBXO12, and FBXO41 variants in Han Chinese patients with sporadic Parkinson's disease. *Neurosci. Bull.* 33, 510–514.
- Zahra, W., Rai, S. N., Birla, H., Singh, S. S., Rathore, A. S., Dilmashin, H., et al. (2020). Neuroprotection of rotenone-induced Parkinsonism by ursolic acid in PD Mouse model. *CNS Neurol. Disord. Drug Targets* 19, 527–540. doi: 10.2174/1871527319666200812224457

- Zanon, A. (2020). *Establishment of an Alginate Based 3D Culture System for the Generation of Dopaminergic Neurons*. Mainz: Institute of Molecular Biology.
- Zella, S. M. A., Metzendorf, J., Ciftci, E., Ostendorf, F., Muhlack, S., Gold, R., et al. (2019). Emerging immunotherapies for Parkinson disease. *Neurol. Ther.* 8, 29–44. doi: 10.1007/s40120-018-0122-z
- Zhang, W., Wang, T., Pei, Z., Miller, D. S., Wu, X., Block, M. L., et al. (2005). Aggregated α -synuclein activates microglia: a process leading to disease progression in Parkinson's disease. *FASEB J.* 19, 533–542.
- Zhang, Y., Feng, S., Nie, K., Li, Y., Gao, Y., Gan, R., et al. (2018). TREM2 modulates microglia phenotypes in the neuroinflammation of Parkinson's disease. *Biochem. Biophys. Res. Commun.* 499, 797–802. doi: 10.1016/j.bbrc.2018.03.226
- Zhang, Z. T., Cao, X. B., Xiong, N., Wang, H. C., Huang, J. S., Sun, S. G., et al. (2010). Morin exerts neuroprotective actions in Parkinson disease models in vitro and in vivo. *Acta Pharmacol. Sin.* 31, 900–906. doi: 10.1038/aps.2010.77
- Zhong, H. H., May, M. J., Jimi, E., and Ghosh, S. (2002). The phosphorylation status of nuclear NF-kappa B determines its association with CBP/p300 or HDAC-1. *Mol. Cell* 9, 625–636. doi: 10.1016/S1097-2765(02)00477-X
- Zhong, Z., Liang, S., Sanchez-Lopez, E., He, F., Shalpour, S., Lin, X. J., et al. (2018). New mitochondrial DNA synthesis enables NLRP3 inflammasome activation. *Nature* 560, 198–203. doi: 10.1038/s41586-018-0372-z
- Zhou, Y., Lu, M., Du, R. H., Qiao, C., Jiang, C. Y., Zhang, K. Z., et al. (2016). MicroRNA-7 targets Nod-like receptor protein 3 inflammasome to modulate neuroinflammation in the pathogenesis of Parkinson's disease. *Mol. Neurodegener.* 11:28. doi: 10.1186/s13024-016-0094-3
- Zhu, G., Xu, Y., Cen, X., Nandakumar, K. S., Liu, S., and Cheng, K. (2018). Targeting pattern-recognition receptors to discover new small molecule immune modulators. *Eur. J. Med. Chem.* 144, 82–92. doi: 10.1016/j.ejmech.2017.12.026

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The handling editor is currently organizing a Research Topic with one of the authors GL.

Copyright © 2021 Yao, Wu, Koc and Lu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.