Check for updates

OPEN ACCESS

EDITED BY Valerie Kouskoff, The University of Manchester, United Kingdom

REVIEWED BY Armel Hervé Nwabo Kamdje, University of Garoua, Cameroon Serena Barachini, University of Pisa, Italy

*CORRESPONDENCE

Qi Wang, i wangqi@bjmu.edu.cn Quanjun Wang, i anquan@saifulab.com Sikai Ling, i sikai.ling@bdgene.cn

RECEIVED 08 April 2025 ACCEPTED 28 May 2025 PUBLISHED 16 June 2025

CITATION

Dai X, Li Z, Chen S, Huang Y, Ling S, Wang Q and Wang Q (2025) Preclinical efficacy and safety evaluation of BD211 autologous CD34⁺ hematopoietic stem cell injection for transfusion-dependent β -thalassemia in NCG-X mice. *Front. Cell Dev. Biol.* 13:1607707.

doi: 10.3389/fcell.2025.1607707

COPYRIGHT

© 2025 Dai, Li, Chen, Huang, Ling, Wang and Wang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Preclinical efficacy and safety evaluation of BD211 autologous CD34⁺ hematopoietic stem cell injection for transfusion-dependent β-thalassemia in NCG-X mice

Xuedong Dai^{1,2}, Zike Li², Shuang Chen², Ying Huang², Sikai Ling³*, Quanjun Wang²* and Qi Wang¹*

¹Department of Toxicology, School of Public Health, Peking University, Beijing, China, ²SAFE Pharmaceutical Technology Co., Ltd., Beijing, China, ³BDGENE Therapeutics Co. Ltd., BDGENE Therapeutics, Shanghai, China

Introduction: Autologous CD34⁺ hematopoietic stem cell-based therapies have shown promise in addressing therapeutic needs. However, a comprehensive evaluation of their efficacy and safety is crucial before clinical application. This study aimed to assess the efficacy and safety profile of BD211 autologous CD34⁺ hematopoietic stem cell injection in NCG-X mice.

Methods: NCG-X mice were administered BD211 intravenously at doses of 4.0 $\times 10^5$ and 1.2×10^6 cells per mouse, followed by withdrawal and observation for 13 weeks. Efficacy was evaluated by monitoring the engraftment and differentiation of BD211 into human erythroid cells within the mouse bone marrow and blood. Safety was assessed through clinical observation, pathology, organ weight measurements, and histopathology. Toxicokinetic studies and distribution of BD211 were determined via validated quantitative PCR.

Results: Mortality was observed in all groups of mice with no correlation to dose or BD211. No abnormal effects related to BD211 administration on clinical responses, body weight, or food intake were observed. BD211 successfully engrafted and differentiated into human erythroid cells within the mouse bone marrow and blood.

Conclusion: The no observed adverse effect level of BD211 was established at 1.2×10^6 cells per mouse. BD211 shows potential as a safe therapeutic approach for treating transfusion-dependent thalassemia.

KEYWORDS

transfusion-dependent β -thalassemia, BD211, NCG-X mice, no-observed-adverseeffect level, toxicokinetic, distribution

1 Introduction

Transfusion-dependent β -thalassemia (TDT) is among the most prevalent monogenic disorders globally, with a particularly high incidence in the Eastern Mediterranean, Southeast Asia, and southern China (Weatherall, 2010; Thong and Ngim, 2021). Approximately 60,000 new cases are diagnosed annually. TDT arises from mutations in the β -globin gene (HBB), leading to diminished synthesis (β +) or complete absence (β 0) of functional β-globin (Boulad et al., 2022; Galanello and Origa, 2010). The aberrant production of β -globin disrupts the equilibrium between β -globin and α -globin chains, resulting in the aggregation of surplus unpaired a-globin chains and the formation of precipitates (Li et al., 2024; Chinese Medical Association, 2022). This process damages the red blood cell membrane and induces intravascular hemolysis (Cappellini et al., 2014; Lang and Lang, 2015). For individuals with TDT, conventional treatments encompass lifelong red blood cell transfusions and periodic iron chelation therapy (Shah et al., 2019; Origa, 2017; Cappellini et al., 2011). Although allogeneic hematopoietic stem cell transplantation (HSCT) from human leukocyte antigen (HLA)-matched donors offer a cure for TDT, its application is constrained by the limited availability of suitable donors and the associated risks of transplantation, including graft failure and graft-versus-host disease (Yang et al., 2024; Kleinschmidt et al., 2025).

Advances in scientific technology have facilitated significant progress in gene therapies for TDT (Brendel et al., 2020; Malay et al., 2024). In May 2019, Bluebird Bio's Zynteglo received conditional approval in the European Union, marking it the world's first gene therapy for TDT (Locatelli et al., 2022). Numerous gene therapies are currently undergoing clinical investigation (Segura et al., 2023; Magrin et al., 2022; Wattanapanitch et al., 2018; Zeng et al., 2020). However, $\beta 0/\beta 0$ thalassemia represents the most severe form of TDT and poses a significant challenge for gene therapy. The efficacy of existing gene therapies for β^0/β^0 thalassemia still requires further investigation to be fully validated (Thompson et al., 2018). Against this backdrop, BD211 autologous CD34⁺ hematopoietic stem cell injection (BD211) has emerged as a promising therapeutic approach. BD211 uses an optimized lentiviral vector system to deliver the $\beta^{A-T87Q}\mbox{-globin gene into patient-derived hematopoietic stem cells.}$ The product's distinctive technological advantages include (1) a proprietary insulator design integrated into the lentiviral vector architecture to minimize potential insertional mutagenesis risks and (2) an optimized hemoglobin subunit beta (HBB) gene expression cassette engineered to achieve physiological hemoglobin levels comparable to those observed in healthy individuals. In preliminary studies, BD211 was successfully administered to two patients with severe $\beta 0/\beta 0$ thalassemia. A 2-year follow-up demonstrated its excellent safety and significant efficacy, bringing new hope and promise to the gene therapy of TDT (Li et al., 2024).

In this study, a single-dose toxicity test of BD211 was conducted via intravenous injection in NCG-X mice, followed by a 13week recovery period. The aim was to assess the nature, severity, dose-response and time-effect relationships, and reversibility of potential toxic reactions induced by the test substance. Additionally, this study aimed to identify target organs or tissues of toxicity and conduct concurrent studies on oncogenicity, toxicokinetics, and tissue distribution. This research provides valuable animal experimental data for clinical research and highlights key indicators that must be closely monitored in clinical practice.

2 Materials and methods

2.1 Cells and *in vitro* erythroid differentiation

The BD211 autologous CD34⁺ hematopoietic stem cell injection, non-transduced stem cell, and its cryopreservation solution were sourced from BDGENE Therapeutics Co. Ltd. Human CD34⁺ hematopoietic stem cells were isolated from the peripheral blood of healthy donors following Filgrastim (Recombinant Human Granulocyte Colony-stimulating Factor) mobilization. On the day of use, the BD211 cells and non-transduced stem cells were thawed and maintained on ice until transplantation, with the entire transplantation procedure completed within 4 h after cell thawing. Viability was assessed using the trypan blue exclusion assay, with 78%–90% of the cells demonstrating viability. The hematopoietic differentiation potential and target gene expression of BD211 were confirmed *in vitro* through flow cytometry and PCR analysis (Figure 2).

2.2 Animals

Specific pathogen-free (SPF) grade NCG-X mice, evenly categorized between males and females and aged 7-9 weeks, were procured from Jiangsu Jicui Yaokang Biotechnology Co., Ltd. (GemPharmatech Co., Ltd., Nanjing, China). The mice were housed in individually ventilated cages, with the animal room temperature maintained at 22.0°C-26.0°C and relative humidity controlled at 40%-70%. The central air conditioning system provided ventilation at a rate of \geq 15 times per hour, and the lighting was set to a 12-h light/12-h dark cycle. Cage ventilation was set at 30-80 times per hour. The facilities used in this study have been accredited by the Association for Assessment and Accreditation of Laboratory Animal Care. All experimental animals were bred and used for scientific research purposes. All procedures were reviewed and approved by the Institutional Animal Care and Use Committee of Guoke Saifu Hebei Pharmaceutical Technology Co., Ltd. (approval numbers: IACUC-2023-149 and IACUC-2023-220).

2.3 Experimental design

This study is part of the preclinical safety evaluation of the BD211 autologous CD34⁺ hematopoietic stem cell injection for treating TDT. The study design adhered to ICH guidelines M3 (R2) and relevant technical guidelines (M3R2 I, 2009; National Medical Products Administration, 2017; National Medical Products Administration, 2021).

Overall, 222 SPF-grade NCG-X mice, aged 7–9 weeks, were randomly assigned to four groups, each with an equal number of males and females. The groups included a vehicle control group (n = 40) that received 0.4 mL of Biolife Solutions CS10 cell cryopreservation solution, a mock control group (n = 52) that received 0.4 mL of non-transduced CD34⁺ hematopoietic stem cells $(3 \times 10^{6} \text{ cells/mL})$, and BD211 low-dose (n = 52) and high-dose groups (n = 78) that received 0.13 mL and 0.4 mL of BD211 $(3 \times 10^{6} \text{ cells/mL})$, respectively. The administered doses were $0, 1.2 \times 10^{6}$, 4.0×10^{5} , and 1.2×10^{6} cells per mouse, equivalent to $0, 6.0 \times 10^{7}$, 2.0×10^{7} , and $6.0 \times 10^{7} \text{ cells/kg}$, respectively. Based on the results of previous studies, the low dose was considered the effective dose, while the high dose was anticipated to induce adverse reactions. Mice were administered a single intravenous injection via the tail vein. The first day of administration was designated as Day 1 (D1), and the mice were observed continuously for 13 weeks (D92).

Evaluations were conducted at multiple time points to assess various parameters. Throughout the experimental period, daily observations were made of the general condition of the mice. Body weight and food intake were measured weekly, and ophthalmological examinations were conducted on D0 and D90. On D2 (24 h post-administration), D57, and D92, assessments focused on in vivo erythroid differentiation of stem cells, including the differentiation efficiency of hCD235a⁺ and hCD71⁺ cells, along with the expression levels of the β^{A-T87Q} -globin target gene. Additionally, a more comprehensive series of analyses were performed on D2, D29, D57, and D92, encompassing stem cell stemness through lymphocyte and lymphocyte subset profiling, measurement of reconstitution efficiency in mouse, and cellular differentiation ratios in peripheral blood, spleen, and bone marrow. The analyses also included cytokine level analysis (IFN-y, TNF-a, IL-2, IL-4, IL-6, and IL-10), tissue distribution (BD211 high-dose group only), vector copy number (VCN) quantification, and toxicokinetic parameters, including BD211 levels based on Woodchuck Hepatitis Virus Posttranscriptional Regulatory Element (WPRE) gene concentration and expression analysis of both $\beta\text{-globin}$ and $\beta^{A-T87Q}\text{-globin}$ genes. The solvent and mock control groups were tested only on D2 and D29. On D92, hematological, blood biochemical, gross dissection, organ weighing, bone marrow smear examination, histopathological examination, and oncogenicity assessment were performed (Figure 1).

Manufacturer details for reagents and instruments are listed in Supplementary Tables S5, 6.

2.4 Clinical examination

During the trial period, daily observations were conducted to record the appearance, dosing site, behavior, skin condition, respiration, mouth and nose, glandular secretion, fecal and urinary characteristics, and any instances of death. Body weight and food intake were measured weekly. Ophthalmic examinations were performed the day before administration and on D90, with a necropsy conducted on D92. Hematological analyses (including coagulation), biochemical analyses (including electrolytes), and histopathological examinations were performed. Gross dissections were carried out, and major organs were weighed.

2.5 Detection of VCN

Genomic DNA for VCN analysis was extracted from BD211transduced CD34⁺ cells cultured for 14 days post-transduction,

and whole blood and bone marrow samples were collected from NCG-X mice on D2, D29, D57, and D92 post-administration. Dual fluorescent quantitative PCR was performed using a realtime PCR instrument (Thermo Fisher Scientific, Quant Studio 5) to amplify the viral element WPRE and the human cellular reference gene RNase P using the primers and probe sequences provided in Supplementary Table S4. Thermocycling conditions were as follows: 37°C for 2 min, 95°C for 5 min, followed by 45 cycles of 95°C for 15 s and 60°C for 40 s. Fluorescence signals were monitored during the 60°C phase through the FAM (WPRE) and VIC (RNAseP) channels. WPRE-containing vector RNA undergoes reverse transcription during transduction and becomes integrated into the host genome. It thus serves as a molecular marker for lentiviral VCN quantification via duplex qPCR. Plasmid standards (Lenti-genome-Copies) containing WPRE (vector-specific) and RNAseP (endogenous control) sequences were serially diluted in Yeast tRNA solution with NCG mouse matrix blanks to generate standard curves ($R^2 > 0.99$). Experimental DNA samples were processed identically, with VCN determined by interpolating Ct values into the standard curve, enabling the normalization of vector copies to host genome equivalents. VCN was calculated using the formula: WPRE copy number×2/RNase P copy number.

2.6 Mononuclear cell isolation

The bone marrow cavities were subsequently flushed with DPBS utilizing a 1 mL syringe until the cavities appeared visibly devoid of residual marrow contents. The resultant cell suspension underwent filtration through a 70- μ m cell strainer. A centrifugation process at 330 × g for 5 min at 4°C ensued to pellet the cells. The pelleted cells were subjected to treatment with red cell lysis buffer to eliminate red blood cells, followed by a washing step with DPBS. Subsequent experiments utilized the acquired bone marrow mononuclear cells.

Splenic mononuclear cell isolation involved the mechanical dissociation of spleens on a 70- μ m strainer. The cell suspension was centrifuged at 330 g for 5 min at 4°C, after which red cell lysis and DPBS washing were performed to eradicate erythrocytes and debris.

Peripheral blood mononuclear cell isolation commenced with blood collection via retro-orbital puncture. Dual erythrocyte lysis cycles were implemented: blood samples were centrifuged at 400 gfor 5 min at 4°C. The pellet was resuspended in red cell lysis buffer. This lysis procedure was repeated to ensure thorough removal of red blood cells. Ultimately, cells were washed with DPBS, and the resulting peripheral blood mononuclear cells were collected.

2.7 Flow cytometry analysis

Single-cell suspensions of peripheral blood, spleen, and bone marrow were prepared as described in method 2.6. Cells were adjusted to 1×10^6 cells/mL in DPBS. For staining, $100 \,\mu$ L aliquots were incubated with $100 \,\mu$ L Zombie Aqua Fixable Viability Kit for 15 min at room temperature, protected from light. After DPBS washing ($300 \times g$, 5 min), cells were resuspended in $100 \,\mu$ L DPBS and stained with pre-titrated antibody cocktails (Supplementary Table S5) for 30 min. Unbound antibodies were removed via two sequential DPBS washes ($300 \times g$, 5 min).



Final pellets were resuspended in 300 µL DPBS and stored at 4°C protected from light until acquisition. Representative gating schematics are provided in Supplementary Figures S1–4.

On D2, D57, and D92, femurs, tibias, and sternums were dissected to obtain bone marrow cells. Flow cytometry was used to assess the efficiency of erythroid maturation and differentiation by detecting hCD235a and hCD71 with the following gating strategy: mCD45⁻hCD45⁻hCD235a⁺CD71⁻ (mature erythrocytes), mCD45⁻hCD45⁻hCD71⁺CD235a⁻ (proerythroblasts), and mCD45⁻hCD45⁻hCD235a⁺CD71⁺ (intermediate erythroblasts).

On D2, D29, D57, and D92, peripheral blood, spleens, and bone marrow from femurs, tibias, and sternums were dissected. The analysis of myeloid and lymphoid cell differentiation cell populations was conducted as follows: T cells (hCD45⁺hCD3⁺), CD8⁺ T cells (hCD45⁺hCD3⁺), CD4⁺ T cells (hCD45⁺hCD3⁺), CD4⁺ T cells (hCD45⁺hCD13⁺), B cells (hCD45⁺hCD19⁺), and NK cells (hCD45⁺hCD56⁺). This analysis was conducted to evaluate myeloid and lymphoid cell differentiation. Additionally, hCD45 was measured in peripheral blood to calculate the peripheral blood chimerism rate.

All assays were performed using a DxFLEX cytometer, and data analysis was conducted using CytExpert for DxFLEX.

2.8 Colony forming cell (CFC) assay

Frozen cells were rapidly thawed at 37°C and diluted into 5 mL IMDM medium, followed by centrifugation at 500 g for 5 min. Viable cells were resuspended in IMDM, adjusted to 1.1×10^4 cells/mL using trypan blue exclusion counting, and serially diluted in pre-warmed MethoCult H4435 Enriched Medium at a 1:10 ratio to achieve a final density of 1×10^3 cells/mL. After vigorous vortex mixing (2–5 min) and 10-min deaeration at room temperature,

1 mL aliquots were plated in triplicate onto 6-well plates. Plates were gently tilted to ensure even distribution and incubated for 14 days at 37° C/5% CO₂ under humidified conditions. Colony enumeration was performed manually using inverted microscopy, with morphological criteria distinguishing colony types: burst-forming unit erythroid (BFU-E), colony-forming unit granulocyte (CFU-G), macrophage (CFU-M), granulocyte-macrophage (CFU-GM), and multipotential granulocyte-erythrocyte-macrophagemegakaryocyte (CFU-GEMM) as defined by the manufacturer.

2.9 Cytokine analysis

Serum cytokines were quantified using the V-PLEX Proinflammatory Panel 1 Kit on a QuickPlex SQ 120 platform. Samples (2-fold diluted in Diluent 41) underwent 2 h incubation at 25°C with orbital shaking (600 rpm). Post-washing (PBS-T, 3×), SULFO-TAG-conjugated detection antibodies were added (1 h incubation), followed by Read Buffer T application and electrochemiluminescence signal measurement. Analytes included IFN-y (0.37-1,500 pg/mL), IL-2 (0.37-1,530 pg/mL), IL-6 (0.18-720 pg/mL), IL-4 (0.06–258 pg/mL), IL-10 (0.09–384 pg/mL), and TNF- α (0.09–362 pg/mL). Measurements below detection thresholds were excluded per manufacturer specifications.

2.10 Hematoxylin and eosin (H&E) staining

Tissue sections $(4 \,\mu\text{m})$ were deparaffinized in xylene, rehydrated through graded ethanol, and stained with Mayer's hematoxylin (10 min). After differentiation (1% acid ethanol, 1 s) and bluing (tap water, 7 min), cytoplasmic counterstaining was performed with eosin Y (5 min). Sections were dehydrated, cleared in xylene, and mounted with neutral resin for brightfield microscopy.

2.11 Toxicokinetic and distribution analysis

Mice were euthanized on D2, D29, D57, and D92 following the injection. Blood (with EDTA anticoagulant) and tissues, including bone marrow, spinal cord, brain, lungs, heart, liver, spleen, kidneys, skeletal muscle, joints, stomach, duodenum, jejunum, colon, epididymis, ovaries, or whole blood, were collected from each mouse. Samples were temporarily stored on ice and quickly transferred to conditions below -70°C for tissue distribution analysis. Genomic DNA was extracted using the MagaBio Plus Universal Genomic DNA Purification Kit II. Quantitative PCR was employed to analyze the WPRE gene DNA in mouse tissues to determine the in vivo distribution of BD211. The PCR parameters were set to 37°C for 2 min, 95°C for 5 min, followed by 45 cycles of 95°C for 15 s and 60°C for 40 s. A standard curve was generated by linear regression of the WPRE plasmid Ct and concentration log values. The actual sample Ct values were substituted into the standard curve to calculate the WPRE DNA content in whole blood and tissue samples. The primer sequence information is detailed in Supplementary Table S4.

2.12 Quantification of WT/GAPDH and T87Q/GAPDH mRNA

Total RNA was extracted from tissues using the MagaBio Plus Total RNA Purification Kit II (BSC69L1E/BSC69M1E, BioFlux) and from whole blood and bone marrow using the Whole Blood Total RNA Kit (5201050, Hangzhou Xingjing Biotech Co., Ltd.). Quantitative PCR was used to analyze the human β^{A} -globin, wild-type β^{A} -globin, and GAPDH mRNA in mouse tissues. The parameters were set to 95°C 15 min; 95°C 10 s, 60°C 20 s, 72°C 20 s, and 40 cycles. The Ct values for the samples were obtained using the instrument software. The relative expression levels were calculated as follows:

WT/GAPDH (wild-type β^{A} -globin/GAPDH) = Ct_{WT}/Ct_{GAPDH}×100%;

 $T87Q/GAPDH \quad (\beta^{A-T87Q}\text{-globin/GAPDH}) = Ct_{T87Q}/Ct_{GAPDH} \times 100\%$

2.13 Oncogenicity assessment

All preserved tissues and organs from the vehicle control, mock control, and high-dose BD211 groups underwent microscopic examination for proliferative or neoplastic changes using H&E staining. Should tumors be detected in high-dose animals, corresponding organs from low-dose groups would have been analyzed.

2.14 Statistical analysis

Statistical analyses were performed using SPSS Statistics for Windows (v.23.0, IBM Corp., Armonk, N.Y., United States). Measurement data, including body weight, food intake, hematological indicators, serum biochemical indicators, and organ weight and coefficient, were presented as mean ± standard deviation (x ± SD). One-way analyses of variance (ANOVAs) were used to assess significance. If a statistically significant difference was found among the groups, further pairwise comparisons were conducted. Levene's test was used to determine the homogeneity of variance: if groups were homoscedastic, the least significant difference method was applied, and if heteroscedastic, the Games-Howell test was used for pairwise comparisons. Count data, such as behavioral and pathological incidences, were described as fractions, and non-parametric tests (Kruskal-Wallis test) were employed for significance analysis. When a statistically significant difference was observed among the overall groups, differences between pairwise groups were compared. The P-value of <0.05 was considered statistically significant.

3 Results

3.1 *In vitro* differentiation, β^{A-T87Q} -globin expression, and VCN of BD211

The colony-forming cell assay is an *in vitro* method used to assess the proliferation and differentiation capabilities of hematopoietic stem cells (Kronstein-Wiedemann and Tonn, 2019). Both the mock control and test groups were cultured using semi-solid culture media for 14 days and successfully differentiated into 5 colony types: erythrocytic burst-forming unit, granulocyte colony-forming unit, macrophage colony-forming unit, granulocyte, macrophage, and megakaryocyte colony-forming units (Figures 2A,B).

During the directed differentiation of CD34⁺ hematopoietic stem cells into erythrocytes, the expression of CD71 gradually decreased, while that of CD235a increased. Early erythroid precursors (proerythroblasts) primarily exhibited a CD71⁺CD235a⁻, intermediate erythroid precursors (basophilic, polychromatophilic, and orthochromatic erythroblasts) exhibited a CD71⁺CD235a⁺ and mature erythrocytes (reticulocytes and mature red blood cells) predominantly displayed CD71⁻CD235a⁺ markers. Both mock control cells and four batches of test cells successfully differentiated into erythroid cells *in vitro* (Figure 2C).

Hemoglobin, the primary component of erythrocytes, is composed of three proteins, with hemoglobin A (HbA) being the most abundant, accounting for over 95% of total hemoglobin in the human body. Flow cytometry analysis revealed that both mock control cells and four batches of test cells successfully detected HbA-positive cells *in vitro*, with positivity exceeding 90% (Figure 2D).

The relative expression levels of WT/GAPDH in mock control cells and four batches of BD211 cells significantly increased with prolonged *in vitro* differentiation time. Similarly,



of red blood cells after 14 days of culture of mock cells and BD211 *in vitro*. CD71⁺CD235a⁻: Proerythroblasts; CD71⁺CD235a⁺: Basophilic, Polychromatophilic & Orthochromatic Erythroblasts; CD71⁻CD235a⁺: Reticulocytes & Mature Red Blood Cells. (**D**). Percentage of hemoglobin β^+ cells cultured *in vitro* after 14 days (**E**,**F**). The mRNA expression levels of β^{A} -globin and β^{A-T87O} -globin on days 0, 7, and 14. Wild-type β^{A} -globin/GAPDH (WT/GAPDH) = Ct_{WT}/Ct_{GAPDH}×100%; β^{A-T87O} -globin/GAPDH (T87Q/GAPDH) = Ct_{T87O}/Ct_{GAPDH}×100% (**G**). The VCN value of mock cells and BD211 cultured *in vitro* after 14 days. VCN = WPRE/(Rnase P/2). BFU-E: burst-forming unit erythroid; CFU-G: colony-forming unit granulocyte; CFU-M: colony-forming unit macrophage; CFU-GM: colony-forming unit granulocyte-macrophage; and CFU-GEMM: colony-forming unit multipotential granulocyte-erythrocyte-macrophage-megakaryocyte; VCN: Vector Copy Number.

the relative expression levels of T87Q/GAPDH in the four batches of BD211 cells also significantly increased. T87Q/GAPDH expression was not detected in mock control cells throughout the experiment (Figures 2E,F).

The mock control cells did not exhibit positivity for the BD211 vector, whereas all four batches of test groups showed BD211 vector positivity, with VCN ranging from 2.84 to 3.63 (Figure 2G). The VCN values for all four batches of test groups were within



FIGURE 3

Survival curve, body weight, and food consumption after a single dose of BD211. (A). Survival curve of NCG-X mice from D0 to D92. Body weight of male (B) and female (C) mice from D0 to D89. Food consumption of male (D) and female (E) mice from D0 to D85. Data are expressed as the mean \pm SD and were analyzed using one-way ANOVAs.

the safety range of 1–5 copies/cell in accordance with relevant guidelines (U. S. Food and Drug, 2024).

3.2 Basic physiological and behavioral monitoring in animals

These results demonstrate that BD211 successfully differentiated into various stages of human erythroid cells *in vitro* and expressed β^{A-T87Q} -globin during the directed differentiation process.

During the trial period, mortality or near-death events occurred in all groups, with proportions of 17.50%, 19.23%, 11.54%, and 14.10%,

TABLE 1 Hematology analysis of NCG-X mice treated with BD211 on D92. Blood samples collected in EDTA-K2 were analyzed within	8 h at room
temperature using the Automatic Hematology Analyzer XN-1000V.	

Parameter	Male mice				Female mic	e		
	Control	Mock	Low dose	High dose	Control	Mock	Low dose	High dose
Number of animals	9	11	12	11	8	4	9	8
RBC (10 ¹² /L)	5.520 ± 0.303	3.803 ± 1.134**	5.357 ± 0.316 ^{##}	4.937 ± 0.250****	5.299 ± 0.779	3.588 ± 0.685***	4.972 ± 0.290 ^{###}	4.483 ± 0.420*#
HGB (g/L)	102.9 ± 4.9	72.8 ± 20.0**	101.0 ± 5.0 ^{###}	95.3 ± 3.8**##	100.1 ± 15.0	70.3 ± 12.8***	96.6 ± 6.1 ^{###}	87.9 ± 7.1 ^{##}
HCT (%)	33.46 ± 1.69	24.55 ± 5.93**	32.88 ± 1.41 ^{##}	31.35 ± 1.26** ^{##}	32.65 ± 4.68	24.15 ± 3.57***	31.91 ± 1.99 ^{###}	28.90 ± 2.42 [#]
MCV (fL)	60.63 ± 0.84	65.44 ± 3.63**	61.47 ± 1.72 ^{##}	63.55 ± 1.44***	61.66 ± 0.82	67.83 ± 3.46**	64.17 ± 1.02***#	64.53 ± 1.21***
MCH (Pg)	18.64 ± 0.32	$19.25 \pm 0.63^{*}$	18.86 ± 0.47	$19.31 \pm 0.48^{*}$	18.88 ± 0.50	$19.63 \pm 0.44^{*}$	19.40 ± 0.33	19.63 ± 0.49**
MCHC (g/L)	307.4 ± 2.8	294.5 ± 9.9**	307.2 ± 3.3 ^{##}	303.8 ± 2.9*#	306.3 ± 7.6	289.8 ± 13.7	302.6 ± 4.5	304.0 ± 5.6
#RETIC (10 ⁹ /L)	296.99 ± 39.99	235.12 ± 61.72*	309.48 ± 41.35 ^{##}	289.78 ± 50.59 [#]	317.31 ± 32.81	401.88 ± 122.44	330.76 ± 82.77	297.43 ± 83.70
%RETIC (%)	5.369 ± 0.535	6.368 ± 1.306	5.779 ± 0.689	5.848 ± 0.843	6.089 ± 0.936	11.253 ± 2.926***	6.656 ± 1.586 ^{##}	6.715 ± 2.159 ^{##}
WBC (10 ⁹ /L)	0.988 ± 0.686	0.351 ± 0.142**	0.753 ± 0.564 ^{##}	$0.424 \pm 0.192^*$	0.479 ± 0.134	0.323 ± 0.074	0.509 ± 0.308	0.738 ± 0.994
#NEUT (10 ⁹ /L)	0.584 ± 0.444	0.192 ± 0.087**	0.497 ± 0.384 ^{##}	$0.249 \pm 0.110^{*}$	0.273 ± 0.089	0.200 ± 0.056	0.290 ± 0.190	0.503 ± 0.815
%NEUT (%)	59.09 ± 10.50	54.17 ± 7.60	65.37 ± 6.23 ^{##}	59.11 ± 7.07	57.55 ± 11.90	61.55 ± 4.2	56.71 ± 14.16	58.29 ± 11.71
#LYMPH (10 ⁹ /L)	0.248 ± 0.182	$0.103 \pm 0.037^*$	0.158 ± 0.125	$0.105 \pm 0.052^*$	0.140 ± 0.060	0.078 ± 0.021	0.131 ± 0.090	0.138 ± 0.094
%LYMPH (%)	24.41 ± 5.85	30.08 ± 7.15	20.78 ± 4.51 ^{##}	25.44 ± 6.90	28.70 ± 9.39	24.03 ± 4.27	25.53 ± 10.41	25.79 ± 10.49
#MONO (10 ⁹ /L)	0.153 ± 0.092	0.051 ± 0.021**	0.098 ± 0.062	$0.068 \pm 0.040^{*}$	0.066 ± 0.026	0.045 ± 0.006	0.086 ± 0.052	0.084 ± 0.058
%MONO (%)	16.28 ± 5.57	14.60 ± 1.81	13.85 ± 2.88	15.29 ± 3.43	13.75 ± 3.66	14.43 ± 3.47	17.27 ± 4.34	15.50 ± 4.25
#EOS (10 ⁹ /L)	0.002 ± 0.004	0.005 ± 0.015	0 ± 0	0.001 ± 0.003	0 ± 0	0 ± 0	0.002 ± 0.007	0.013 ± 0.035
%EOS (%)	0.22 ± 0.45	1.15 ± 2.78	0 ± 0	0.16 ± 0.54	0 ± 0	0 ± 0	0.49 ± 1.47	0.39 ± 1.10
#BASO (10 ⁹ /L)	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.001 ± 0.004
%BASO (%)	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0.04 ± 0.11
PLT (10 ⁹ /L)	1,363.4 ± 155.3	938.0 ± 230.0***	1,348.8 ± 274.3 ^{###}	1,179.8 ± 128.4 [#]	1,062.5 ± 402.3	786.5 ± 65.6	1,022.6 ± 90.8 ^{##}	939.8 ± 111.4 [#]

Note: The data were expressed as mean \pm SD. """" or, "**" indicates a statistically significant difference at p < 0.05, p < 0.01 or p < 0.001 when compared to the control group; "#"" ##" or, "###" indicates a statistically significant difference at p < 0.05, p < 0.01 or p < 0.001 when compared to the control group; "#"" ###" indicates a statistically significant difference at p < 0.05, p < 0.01 or p < 0.001 when compared to the mock group; Abbreviation: RBC, red blood cell count; HGB, hemoglobin concentration; HCT, hematocrit; MCV, mean cell volume; MCH, mean cell hemoglobin; MCHC, mean cell hemoglobin concentration; RETIC, reticulocyte; WBC, white blood cell; NEUT, neutrophils; LYMPH, lymphocytes; MONO, monocytes; EOS, eosinophils; BASO, basophils; PLT, platelets.

respectively. No dose-dependent relationship was observed, and these events were not attributed to the test substance (Figure 3A).

Throughout the study, there were no significant differences in body weight gain between the BD211 low- and high-dose groups and the solvent control group for both male and female mice. Body weight increased over time in all groups (Figures 3B,C). Food consumption in BD211 groups fluctuated within normal ranges, with no significant changes noted. A transient increase in food consumption was observed in male mice of the mock control group, which was not considered toxicologically significant (Figures 3D,E).

tome	
ow cy	
by flo	
/zed	
analy	
as, as	
D92	
, and	
D57	
D29,	
D2,	
1 on	
3D21	
/ith E	
ed v	
treat	
nice	
×	
NCG-X	
v of NCG-X	
arrow of NCG-X	
ie marrow of NCG-X	
bone marrow of NCG-X	
the bone marrow of NCG-X	
on in the bone marrow of NCG-X	
tiation in the bone marrow of NCG-X	
erentiation in the bone marrow of NCG-X	
differentiation in the bone marrow of NCG-X	
cell differentiation in the bone marrow of NCG-X	
hoid cell differentiation in the bone marrow of NCG-X	
ymphoid cell differentiation in the bone marrow of NCG-X	
nd lymphoid cell differentiation in the bone marrow of NCG-X	
vid and lymphoid cell differentiation in the bone marrow of NCG-X	
Ayeloid and lymphoid cell differentiation in the bone marrow of NCG-X	
2 Myeloid and lymphoid cell differentiation in the bone marrow of NCG-X	
ABLE 2 Myeloid and lymphoid cell differentiation in the bone marrow of NCG-X	

Frontiers in Cell and Developmental Biology

TABLE 2 Myeloid and lymph	noid cell di	fferentiation in the b				מווט עשב, מז מוומועבר	a by now cyconicaly.		
Parameter		Female mice				Male mice			
		Control	Mock	Low dose	High dose	Control	Mock	Low dose	High dose
	D2	0.003 ± 0.005	0 ± 0	0 ± 0	$0.153 \pm 0.072^{*\#}$	0 ± 0	0.003 ± 0.006	0.023 ± 0.021	0.077 ± 0.080
hCD45 ⁺ hCD3 ⁺ in	D29	0.003 ± 0.005	$3.153 \pm 1.092^{*}$	$0.760 \pm 0.421^{*\#}$	$2.155 \pm 0.896^{*}$	0 ± 0	$1.913 \pm 1.187^{*}$	0.110	$0.772 \pm 0.523^{*}$
hCD45 ⁺ /mCD45 ⁺ (%)	D57	1	5.843 ± 1.698	$2.668 \pm 0.445^{\#}$	4.439 ± 1.695	1	4.363 ± 1.768	1.645 ± 1.082	3.549 ± 2.065
	D92	I	7.348 ± 1.629	$4.278 \pm 1.096^{\#}$	$4.134 \pm 1.324^{\#}$	I	5.953 ± 2.597	1.150 ± 0.290	5.562 ± 3.768
	D2	0 ± 0	$0.033 \pm 0.006^{*}$	0.010 ± 0.010	$0.102 \pm 0.062^{*}$	0.003 ± 0.005	0.013 ± 0.006	0.033 ± 0.035	0.058 ± 0.062
hCD45 ⁺ hCD34 ⁺ in	D29	0.005 ± 0.010	$13.247 \pm 5.610^{*}$	$4.127 \pm 2.134^{*}$	$10.342 \pm 4.119^{*}$	0.003 ± 0.005	$5.983 \pm 3.985^*$	0.470	$3.183 \pm 1.691^{*}$
hCD45 ⁺ /mCD45 ⁺ (%)	D57		23.108 ± 2.932	$13.660 \pm 2.316^{\#}$	19.796 ± 4.029		16.883 ± 3.954	$5.933 \pm 4.311^{\#}$	12.990 ± 6.579
	D92		24.018 ± 7.064	22.425 ± 2.086	20.182 ± 3.358		24.407 ± 5.124	$7.837 \pm 3.650^{\#}$	$16.224 \pm 3.834^{\#}$
	D2	0 ± 0	$0.033 \pm 0.006^{*}$	$0.010 \pm 0.010^{\#}$	$0.182 \pm 0.089^{*\#}$	0 ± 0	0.013 ± 0.006	0.063 ± 0.047	0.112 ± 0.130
hCD45 ⁺ hCD13 ⁺ in	D29	0.005 ± 0.010	$8.830 \pm 3.717^{*}$	$2.087 \pm 0.880^{*\#}$	4.403 ± 1.653 **	0 ± 0	$4.090 \pm 2.869^{*}$	0.320	$1.238 \pm 0.479^{*\#}$
hCD45 ⁺ /mCD45 ⁺ (%)	D57	I	23.595 ± 3.188	6.145 ± 1.241 [#]	16.164 ± 7.083	I	13.643 ± 6.386	$2.500 \pm 1.115^{\#}$	$5.746 \pm 3.250^{\#}$
	D92	1	28.055 ± 6.003	$18.420 \pm 1.660^{\#}$	$15.544 \pm 8.343^{\#}$	I	25.107 ± 12.945	$2.837 \pm 0.448^{\#}$	13.766 ± 6.263
	D2	0.003 ± 0.005	0.003 ± 0.006	0.033 ± 0.058	$0.245 \pm 0.109^{*\#}$	0.008 ± 0.005	0.003 ± 0.006	0.230 ± 0.215	0.305 ± 0.245
hCD45 ⁺ hCD19 ⁺ in	D29	0.003 ± 0.005	$8.770 \pm 4.465^*$	$3.830 \pm 1.537^{*}$	$12.992 \pm 6.008^{*}$	0.003 ± 0.005	$3.670 \pm 2.600^{*}$	0.420	$3.515 \pm 1.676^{*}$
hCD45 ⁺ /mCD45 ⁺ (%)	D57	Ι	48.148 ± 6.956	$31.443 \pm 7.352^{*}$	46.027 ± 7.721		35.840 ± 5.665	12.115 ± 10.960	26.793 ± 13.707
	D92	I	46.065 ± 4.312	56.003 ± 4.712	51.752 ± 9.378		46.690 ± 9.047	24.337 ± 15.415	43.944 ± 11.964
	D2	0 ± 0	0.003 ± 0.006	0.007 ± 0.012	$0.217 \pm 0.158^{*\#}$	0.008 ± 0.005	0.003 ± 0.006	0.077 ± 0.068	0.145 ± 0.182
hCD45 ⁺ hCD56 ⁺ in	D29	0.003 ± 0.005	$0.720 \pm 0.139^{*}$	$0.273 \pm 0.126^{*\#}$	$0.633 \pm 0.203^{*}$	0 ± 0	$0.543 \pm 0.281^{*}$	0.030	$0.268 \pm 0.133^{*}$
hCD45 ⁺ /mCD45 ⁺ (%)	D57		1.728 ± 0.568	$0.653 \pm 0.171^{\#}$	$1.076 \pm 0.211^{\#}$		1.505 ± 0.702	$0.478 \pm 0.438^{\#}$	0.857 ± 0.402
	D92		1.405 ± 0.422	$0.738 \pm 0.139^{\#}$	1.036 ± 0.298		1.933 ± 1.099	0.537 ± 0.234	1.764 ± 1.675
Note: The data were expressed as m compared to the mock group. The <u>p</u> mouse CD45 ⁺ cells.	ean ± SD. """ ercentages in	or"**** indicates a statistics idicate the proportion of c	ally significant difference at just expressing the specified	p < 0.05 and $p < 0.01$ when c antigens. For gating strateg	compared to the control grou y details, refer to the Methon	up; "#»"##" or,"###" indi ds section. h: human; m:	cates a statistically significar mouse; hCD45/mCD45 ⁺ in	nt difference at $p < 0.05$, $p <$ dicates the presence of eithe	0.01 or $p < 0.001$ when r human CD45 ⁺ cells or



Erythroid differentiation *in vivo*, peripheral blood chimerism rate and VCN of NCG-X mice after a single administration of BD211. (**A–C**). The Proportion of red blood cells in the bone marrow of NCG-X mice on D2, D29, D57, and D92 was analyzed by flow cytometry. h: human; m: mouse.*p < 0.05 (one-way ANOVA compared with the Control group); #p < 0.05 (one-way ANOVA compared with the Mock group). (**D**). peripheral blood chimerism rate in blood of NCG-X mice on D2, D29, D57, and D92 analyzed by flow cytometry. hCD45/mCD45⁺ indicates human CD45⁺ cells and/or mouse CD45⁺ cells. (**E,F)**. mRNA expression levels of β^{A} -globin and β^{A-T870} -globin in the bone marrow of NCG-X mice on D2, D29, D57, and D92. (**G**). The VCN value of BD211 cultured in bone marrow and blood of NCG-X mice on D2, D29, D57, and D92.



D57, and D92

3.3 Hematology and biochemical assessment

At the end of the 13-week dosing period (D92), the hematological parameters of the mock control and BD211 groups, including red blood cell indices (RBC, HGB, HCT, MCHC, and

RET), white blood cell indices (WBC, NEUT, LYMPH, MONO), and platelets (PLT), showed a decrease in absolute values compared to the solvent control group, exhibiting a dose-dependent relationship with the human CD34⁺ hematopoietic stem cell dose (Table 1). No biochemically significant changes were observed in the BD211 lowand high-dose groups (Supplementary Table S1).



3.4 Therapeutic efficacy

From D29 onwards, the percentage of hCD3⁺, hCD13⁺, hCD19⁺, hCD34⁺, and hCD56⁺ in the bone marrow of both the mock control and BD211 groups increased in a dose-dependent manner, reaching the highest level by D92 (Table 2). Similarly, starting from D29, the percentage of hCD3⁺, hCD13⁺, and hCD19⁺ in the spleen of the mock control and BD211 groups increased in a dose-dependent manner, reaching the highest level by D92. The percentages of other indicators (hCD4⁺, hCD8⁺, hCD56⁺) did not show significant changes compared to their D2 levels (Table 3).

From D2 to D92, the levels of hCD3⁺, hCD4⁺, hCD8⁺, hCD4⁺/hCD8⁺, hCD56⁺, hCD13⁺, and hCD19⁺ in peripheral blood of the mock control and BD211 groups remained very low level, with no significant increase. This was likely due to the residual immune function of NCG-X mice, as suggested by background information from the animal supplier (unpublished), which indicated that monocytes or macrophages in the tissues could phagocytose and clear the xenogeneic human terminal blood cells released into the peripheral blood after the reconstruction and differentiation of human CD34⁺ hematopoietic stem cells (Supplementary Table S2).

Both the mock control and BD211 groups successfully differentiated into erythroid cells in the bone marrow, with a significant increase in the percentage of hCD71⁺hCD235a⁺ cells, indicating a high proportion of immature erythrocytes, consistent with *in vitro* differentiation results (Figures 4A–C).

Starting from D29, the percentage of hCD45⁺ cells and their proportion in peripheral blood increased in a dose-dependent manner for both the mock control and BD211 groups, reaching the

highest level by D92. On D92, the hCD45⁺ chimerism rates for the mock, BD211 low-dose, and high-dose groups were 73.10%, 43.24%, and 58.85%, respectively (Figure 4D).

On D29, occasional expression of β -globin mRNA and β^{A-T87Q} globin mRNA was observed in the bone marrow in the BD211 highdose group (Figures 5E,F). By D57 and D92, widespread expression of β -globin mRNA was observed in the mock group, whereas β globin mRNA and β^{A-T87Q} -globin mRNA in BD211 groups were detected in the bone marrow (Figures 4E,F).

In the high-dose group, WPRE was detected in the bone marrow of only a few mice (3/12) on D2, with a VCN value of 5.63. From D29 onwards, WPRE was consistently detected in the bone marrow and blood of all mice, with VCN values ranging from 2.07 to 2.84. No significant inter-sample or gender differences in VCN values were observed, and VCN values remained relatively stable throughout the study (Figure 4G).

During the study, human-specific cytokine levels of IL-2, IL-4, IL-6, IL-10, TNF- α , and IFN- γ were mostly below the limit of quantification for all groups at each testing point, failing to meet the requirements for statistical analysis (data not shown).

3.5 Toxicokinetics and distribution

From D2 to D92, the detection rate and genomic concentration of BD211 (representing BD211 and its differentiated nucleated cells in toxicokinetic and tissue distribution studies, with the WPRE gene as the detection marker) increased with the extension of observation time. In the early stages of administration (D2 to

e
È
2
5
0
≥
0
f
\geq
D D
й
S
a
01
as
2
õ
-
2
ar
N
6
Š
5
õ
_
N
0
-
1
õ
-
⇒
>
D.
Ę
g
ž
4
8
-ĕ
~
×
X-2
CG-X r
NCG-X r
NCG-X r
of NCG-X r
n of NCG-X r
en of NCG-X r
leen of NCG-X r
pleen of NCG-X r
spleen of NCG-X r
ie spleen of NCG-X r
the spleen of NCG-X r
n the spleen of NCG-X r
in the spleen of NCG-X r
in in the spleen of NCG-X r
ion in the spleen of NCG-X r
ation in the spleen of NCG-X r
tiation in the spleen of NCG-X r
intiation in the spleen of NCG-X r
rentiation in the spleen of NCG-X r
erentiation in the spleen of NCG-X r
ifferentiation in the spleen of NCG-X r
differentiation in the spleen of NCG-X r
ll differentiation in the spleen of NCG-X r
ell differentiation in the spleen of NCG-X r
cell differentiation in the spleen of NCG-X r
id cell differentiation in the spleen of NCG-X r
oid cell differentiation in the spleen of NCG-X r
shoid cell differentiation in the spleen of NCG-X r
nphoid cell differentiation in the spleen of NCG-X r
mphoid cell differentiation in the spleen of NCG-X r
lymphoid cell differentiation in the spleen of NCG-X r
d lymphoid cell differentiation in the spleen of NCG-X r
ind lymphoid cell differentiation in the spleen of NCG-X r
and lymphoid cell differentiation in the spleen of NCG-X r
id and lymphoid cell differentiation in the spleen of NCG-X r
oid and lymphoid cell differentiation in the spleen of NCG-X r
eloid and lymphoid cell differentiation in the spleen of NCG-X r
lyeloid and lymphoid cell differentiation in the spleen of NCG-X r
Myeloid and lymphoid cell differentiation in the spleen of NCG-X r
Myeloid and lymphoid cell differentiation in the spleen of NCG-X r
3 Myeloid and lymphoid cell differentiation in the spleen of NCG-X r
LE 3 Myeloid and lymphoid cell differentiation in the spleen of NCG-X r
BLE 3 Myeloid and lymphoid cell differentiation in the spleen of NCG-X r
ABLE 3 Myeloid and lymphoid cell differentiation in the spleen of NCG-X r

Frontiers in Cell and Developmental Biology

TABLE 3 Myeloid and lymphoic	I cell differ	entiation in the sple	en of NCG-X mice tre	eated with BD211 on D	12, D29, D57, and D92, i	as analyzed by flow	cytometry.		
Parameter		Female mice				Male mice			
		Control	Mock	Low dose	High dose	Control	Mock	Low dose	High dose
	D2	0.015 ± 0.006	$0.140 \pm 0.036^{*}$	0.053 ± 0.055	$0.025 \pm 0.021^{\#}$	0.025 ± 0.013	$0.127 \pm 0.070^{*}$	$0.003 \pm 0.006^{*\#}$	$0.027 \pm 0.033^{\#}$
hCD45 ⁺ hCD3 ⁺ in	D29	0 ± 0	0.230 ± 0.159	0.053 ± 0.055	0.143 ± 0.119	0 ± 0	0.137 ± 0.142	0.020	0.123 ± 0.102
hCD45 ⁺ /mCD45 ⁺ (%)	D57	I	0.340 ± 0.096	0.153 ± 0.093	0.301 ± 0.119	1	0.688 ± 0.198	$0.095 \pm 0.098^{*}$	$0.250 \pm 0.128^{\#}$
	D92	1	1.015 ± 0.205	0.945 ± 0.747	0.920 ± 0.593	1	0.987 ± 0.127	$0.163 \pm 0.100^{*}$	1.106 ± 0.821
	D2	0.003 ± 0.005	0 ± 0	0.007 ± 0.006	0.000 ± 0.000	0.008 ± 0.005	0.003 ± 0.006	0.003 ± 0.006	0.002 ± 0.004
hCD45 ⁺ hCD3 ⁺ hCD8 ⁺ in	D29	0 ± 0	0±0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0
hCD45 ⁺ /mCD45 ⁺ (%)	D57	I	0±0	0.003 ± 0.005	0.001 ± 0.004	I	0 ± 0	0 ± 0	0.001 ± 0.004
	D92	Ι	0.008 ± 0.005	0.018 ± 0.010	0.004 ± 0.005	Ι	0.010 ± 0.000	0 ± 0	0.006 ± 0.009
	D2	0.008 ± 0.005	0 ± 0	0.007 ± 0.012	0.002 ± 0.004	0 ± 0	0.003 ± 0.006	0 ± 0	0 ± 0
hCD45 ⁺ hCD3 ⁺ hCD4 ⁺ in	D29	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0 ± 0	0	0 ± 0	0 ± 0
hCD45 ⁺ /mCD45 ⁺ (%)	D57	1	0 ± 0	0 ± 0	0 ± 0	1	0.003 ± 0.005	0 ± 0	0 ± 0
	D92	I	0.013 ± 0.013	0.015 ± 0.017	0.018 ± 0.016	I	0.007 ± 0.006	0 ± 0	0.012 ± 0.011
	D2	0.010 ± 0.008	$0.597 \pm 0.155^{*}$	$0.113 \pm 0.023^{*\#}$	$0.367 \pm 0.159^{*}$	0.018 ± 0.013	$0.450 \pm 0.131^{*}$	$0.130 \pm 0.020^{*\#}$	$0.352 \pm 0.140^{*}$
hCD45 ⁺ hCD13 ⁺ in	D29	0 ± 0	$0.790 \pm 0.762^{*}$	$0.157 \pm 0.139^{*}$	$0.440 \pm 0.229^{*}$	0	$0.397 \pm 0.318^{*}$	0.030	$0.237 \pm 0.170^{*}$
hCD45 ⁺ /mCD45 ⁺ (%)	D57	I	1.268 ± 0.342	$0.370 \pm 0.227^{*}$	1.153 ± 0.699	I	2.090 ± 0.652	$0.245 \pm 0.259^{\#}$	$0.654 \pm 0.344^{\#}$
	D92	I	4.163 ± 0.598	3.348 ± 3.166	3.038 ± 2.018	1	3.747 ± 0.129	$0.440 \pm 0.304^{\#}$	4.478 ± 2.903
	D2	0.010 ± 0.000	0.037 ± 0.015	0.050 ± 0.056	0.035 ± 0.029	0.028 ± 0.013	0.040 ± 0.020	0.010 ± 0.017	0.025 ± 0.029
hCD45 ⁺ hCD19 ⁺ in	D29	0 ± 0	$0.517 \pm 0.497^{*}$	$0.217 \pm 0.204^{*}$	$0.420 \pm 0.329^{*}$	0 ± 0	0.160 ± 0.118	0.030	0.205 ± 0.171
hCD45 ⁺ /mCD45 ⁺ (%)	D57	Ι	16.540 ± 1.889	$3.935 \pm 2.035^{*}$	11.033 ± 4.728		16.288 ± 2.760	$2.128 \pm 1.924^{\#}$	$4.241 \pm 2.219^{\#}$
	D92	Ι	39.985 ± 5.676	22.355 ± 14.315	27.784 ± 17.623		30.257 ± 5.020	$3.277 \pm 2.173^{*}$	22.364 ± 14.396
								(Conti	nued on the following page)

TABLE 3 (Continued) Myeloic	and lympl	noid cell differentiati	on in the spleen of NC	CG-X mice treated wit	h BD211 on D2, D29, D	057, and D92, as anal	yzed by flow cytome	try.	
Parameter		Female mice				Male mice			
		Control	Mock	Low dose	High dose	Control	Mock	Low dose	High dose
	D2	0.010 ± 0.008	0.080 ± 0.026	0.057 ± 0.074	0.020 ± 0.021	0.028 ± 0.013	0.053 ± 0.023	$0.003\pm0.006^{*\#}$	0.015 ± 0.023
hCD45 ⁺ hCD56 ⁺ in	D29	0 ∓ 0	$0.167 \pm 0.159^{*}$	$0.067 \pm 0.045^{*}$	$0.092 \pm 0.080^{*}$	0 ± 0	0.077 ± 0.067	0.020	0.077 ± 0.066
hCD45 ⁺ /mCD45 ⁺ (%)	D57	I	0.585 ± 0.159	$0.210 \pm 0.154^{*}$	0.476 ± 0.131		0.795 ± 0.161	$0.115 \pm 0.138^{\#}$	$0.280 \pm 0.154^{*}$
	D92		1.140 ± 0.143	0.798 ± 0.382	0.890 ± 0.524		1.157 ± 0.032	0.193 ± 0.116	1.192 ± 0.730
Note: The data were expressed as me	an ± SD. "*"or'	***** indicates a statistically	significant difference at $p <$	< 0.05 and $p < 0.01$ when co	ompared to the control group	2; "#","##" or,"###" indica	tes a statistically significan	it difference at $p < 0.05$, $p < 0$.01 or $p < 0.001$ when

compared to the mock group. The percentages indicate the proportion of cells expressing the specified antigens. For gating strategy details, refer to the Methods section. h: human; m: mouse; hCD45⁺ indicates the presence of either human CD45⁺ cells or cells mouse CD45⁺ D57), the concentration of BD211 in the blood of high-dose animals was significantly higher than that of the low-dose group. However, as time progressed, by D92, the BD211 concentrations in the blood of both groups tended to converge (Figure 5A). The expression trends of β^{A-T87Q} -globin and β -globin mRNA were essentially consistent with the distribution trend of BD211. Over time, the concentration and detection rate of BD211 in the blood and the detection rate and concentration levels of β^{A-T87Q} -globin and β -globin mRNA exhibited a clear increasing trend within a certain range (Figures 5B,C).

On D2, BD211 was predominantly detected in the bone marrow, lungs, liver, spleen, femur, and stomach, with concentrations ranging from 1.06×10^2 to 4.71×10^3 copies/µg gDNA. Over time, the concentration and distribution range of BD211 gradually increased. By D29, it was distributed in multiple organs, with the highest concentration in the bone marrow, followed by the elbow joint, while concentrations in other tissues were relatively lower. The overall detection concentration range was 1.12×10^2 to 5.34×10^5 copies/µg gDNA, with specific ranges of 2.07×10^4 to 5.34×10^5 copies/µg gDNA in the bone marrow and 3.54×10^3 to 2.12×10^5 copies/µg gDNA in the elbow joint. The distribution trend of BD211 in various tissues and organs at D57 and D92 was consistent with that of D29. The concentration growth trend from D57 to D92 was not significant, suggesting a possible steady state (Figure 5D).

The changes in β^{A-T87Q} -globin mRNA and β -globin mRNA were largely consistent. They were primarily detected in tissues, such as the lungs, elbow joint, bone marrow, and spinal cord. Over time, the range of detected tissues gradually increased until almost all tissues were involved (Figures 5E,F).

3.6 Anatomical and pathological safety evaluation

At D92, gross pathological examination of animals in the BD211 groups revealed no gross pathological changes related to the test substance nor any proliferative or tumorigenic changes associated with BD211.

The spleen weight, as well as the organ-to-body and organto-brain coefficients, were significantly elevated in male animals of the BD211 high-dose and mock control groups (Table 4). In both male and female mice of the BD211 high-dose and mock groups, an increase in the number of bone marrow (sternum) cells, thymic lymphocytes, and cells surrounding the splenic arteries was observed. Additionally, extramedullary hematopoiesis was observed in male animals. In the BD211 low-dose group, both male and female animals exhibited increased bone marrow (sternum) cells, and extramedullary hematopoiesis was noted in male animals (Figure 6). These histopathological changes were due to the engraftment, reconstitution, and differentiation of human CD34⁺ hematopoietic stem cells in NCG-X mice rather than specific changes induced by the test substance BD211.

Bone marrow smears at D92 showed active bone marrow proliferation in both the mock and BD211 groups, with a significant increase in the proportion of lymphocytes and a relative decrease in the number of granulocytes and erythrocytes, consistent with the trends observed in hematological changes (Supplementary Table S3).

Parameter		Male mice				Female mice			
		Control	Mock	Low dose	High dose	Control	Mock	Low dose	High dose
Number of anim	nals	6	11	12	11	8	4	6	8
	Weight (g)	0.125 ± 0.012	$0.138 \pm 0.009^{*}$	$0.127 \pm 0.010^{#\#}$	0.139 ± 0.025	0.117 ± 0.013	0.122 ± 0.004	0.115 ± 0.015	0.116 ± 0.016
Heart	Organ coefficient (%)	0.425 ± 0.033	0.479 ± 0.043	0.462 ± 0.055	0.481 ± 0.076	0.492 ± 0.040	0.487 ± 0.021	0.464 ± 0.061	0.47 ± 0.089
	Organ brain coefficient	27.559 ± 2.222	30.433 ± 2.227	28.091 ± 1.858	30.671 ± 5.540	25.616 ± 2.681	26.653 ± 1.356	25.592 ± 3.165	25.265 ± 3.828
	Weight (g)	1.516 ± 0.126	1.554 ± 0.175	1.395 ± 0.179	1.459 ± 0.166	1.308 ± 0.365	1.309 ± 0.125	1.264 ± 0.056	1.200 ± 0.136
Liver	Organ coefficient (%)	5.163 ± 0.2229	5.367 ± 0.392	5.095 ± 0.908	5.044 ± 0.285	5.521 ± 1.564	5.226 ± 0.366	5.105 ± 0.199	4.890 ± 0.325
	Organ brain coefficient	334.891 ± 24.133	341.672 ± 31.887	309.586 ± 37.999	321.205 ± 33.841	286.589 ± 76.929	285.763 ± 15.815	281.927 ± 17.742	261.124 ± 30.327
	Weight (g)	0.051 ± 0.011	$0.068 \pm 0.013^{**}$	0.058 ± 0.012	$0.070 \pm 0.012^{**}$	0.071 ± 0.058	0.109 ± 0.026	$0.072 \pm 0.015^{\#\#}$	$0.0821 \pm 0.013^{\#}$
Spleen	Organ coefficient (%)	0.174 ± 0.035	$0.235 \pm 0.046^{**}$	0.212 ± 0.048	$0.244 \pm 0.041^{**}$	0.302 ± 0.250	0.436 ± 0.098	$0.290 \pm 0.057^{*}$	$0.334 \pm 0.037^{*}$
	Organ brain coefficient	11.340 ± 2.571	$14.944 \pm 2.889^{*}$	12.888 ± 2.589	15.512 ± 2.764	15.609 ± 12.469	23.815 ± 5.153	$16.051 \pm 3.501^{##}$	$17.863 \pm 2.812^{\#}$
	Weight (g)	0.393 ± 0.051	0.394 ± 0.038	0.372 ± 0.044	0.380 ± 0.032	0.279 ± 0.022	0.275 ± 0.007	0.258 ± 0.023	0.273 ± 0.033
Kidney	Organ coefficient (%)	1.334 ± 0.103	1.361 ± 0.066	1.348 ± 0.074	1.316 ± 0.072	1.175 ± 0.063	1.100 ± 0.040	$1.041 \pm 0.057^{***}$	1.111 ± 0.083
	Organ brain coefficient	86.663 ± 9.620	$86.663 \pm 7,032$	82.619 ± 8.732	83.641 ± 6.449	61.118 ± 4.329	60.164 ± 1.821	57.592 ± 5.603	59.340 ± 7.176
	Weight (g)	0.149 ± 0.040	0.171 ± 0.015	0.164 ± 0.022	0.172 ± 0.010	0.015 ± 0.004	0.018 ± 0.003	0.017 ± 0.003	0.016 ± 0.004
Testis/Ovary	Organ coefficient (%)	0.504 ± 0.130	0.592 ± 0.056	0.595 ± 0.095	0.600 ± 0.065	0.064 ± 0.015	0.071 ± 0.012	0.070 ± 0.011	0.065 ± 0.016
	Organ brain coefficient	32.811 ± 8.677	37.619 ± 3.236	36.254 ± 4.584	37.921 ± 2.342	3.354 ± 0.854	3.872 ± 0.703	3.884 ± 0.791	3.468 ± 1.016
	Weight (g)	0.074 ± 0.012	0.079 ± 0.015	0.079 ± 0.012	0.081 ± 0.010	0.105 ± 0.035	0.119 ± 0.049	0.108 ± 0.056	0.146 ± 0.060
Epididymis/Uterus	Organ coefficient (%)	0.251 ± 0.031	0.271 ± 0.027	0.287 ± 0.036	0.281 ± 0.044	0.445 ± 0.145	0.471 ± 0.181	0.439 ± 0.243	0.589 ± 0.219
	Organ brain coefficient	16.293 ± 2.333	17.288 ± 2.069	17.558 ± 2.538	17.806 ± 2.366	23.113 ± 7.676	25.863 ± 10.226	24.104 ± 12.374	31.826 ± 12.936
	Weight (g)	0.007 ± 0.001	0.007 ± 0.010	0.007 ± 0.002	0.006 ± 0.001	0.009 ± 0.001	0.009 ± 0.001	0.008 ± 0.001	0.008 ± 0.001
Adrenal gland	Organ coefficient (%)	0.020 ± 0.004	0.023 ± 0.005	0.025 ± 0.006	0.022 ± 0.005	0.037 ± 0.003	0.035 ± 0.004	0.033 ± 0.003	0.032 ± 0.003
	Organ brain coefficient	1.329 ± 0.312	1.435 ± 0.314	1.490 ± 0.358	1.359 ± 0.292	1.903 ± 0.222	1.900 ± 0.237	1.795 ± 0.179	1.726 ± 0.205
			-		-			(Continue	ed on the following page)

Parame	eter	Male mice				Female mice			
		Control	Mock	Low dose	High dose	Control	Mock	Low dose	High dose
Numbe	er of animals	6	11	12	11	8	4	6	8
	Weight (g)	0.453 ± 0.021	0.454 ± 0.013	0.450 ± 0.014	0.454 ± 0.007	0.456 ± 0.008	0.457 ± 0.020	0.449 ± 0.020	0.460 ± 0.012
DIAIII	Organ coefficient (%)	1.546 ± 0.083	1.574 ± 0.069	1.642 ± 0.133	1.578 ± 0.092	1.926 ± 0.091	1.828 ± 0.052	1.816 ± 0.124	1.884 ± 0.134
Note: The dat compared to t	a were expressed as mean ± SD. "*"or"* ihe mock group;	*** indicates a statistically :	significant difference at $p < $	c 0.05 and <i>p</i> < 0.01 when co	mpared to the control group	', "#","##" or,"###" indicat	es a statistically significant	t difference at $p < 0.05$, $p < 0$	0.01 or p < 0.001 when

Organ coefficient (%) = g organ weight/g body weight×100%; Organ brain coefficient = g organ weight/g brain weight

Dai et al.

4 Discussion

Zynteglo, the first FDA-approved gene therapy for β -thalassemia (August 2022), employs lentiviral-mediated βA-T87Q-globin gene addition to restore functional β -globin production in hematopoietic stem cells (HSCs) - a mechanism shared by BD211. While clinical trials of Zynteglo demonstrate promising transfusion independence rates, its preclinical characterization in animal models remains unpublished, limiting mechanistic insights into biodistribution and long-term safety. This data gap extends to most HSC-based gene therapies currently in development, which predominantly report clinical rather than preclinical findings. The systematic evaluation of BD211 in NCG-X murine models addresses this critical need, providing empirical evidence of stable transgene integration, erythroid-specific expression, and absence of genotoxicity signals. This preclinical evidence not only supports the IND application of BD211 but also establishes a benchmark for comparative safety assessments of lentiviral therapies, particularly regarding insertional mutagenesis risks in immunodeficient models.

NCG-X mice, derived from NCG severely immunodeficient mice, have been genetically edited to introduce a W41 point mutation in the c-Kit gene (Mcintosh et al., 2015). They exhibit deficiencies in T/B/NK cell immunity and suppressed hematopoietic stem cell function, enabling the reconstitution of higher levels of erythroid cells. This makes them an excellent model for human HSCT and the study of thalassemia (Bao et al., 2021). Under the experimental conditions used in this study, after 8 and 13 weeks of administration, the bone marrow of mice in the mock and BD211 groups showed a significant increase in erythroid cells, primarily characterized by a high proportion of immature erythrocytes (hCD235a⁺hCD71⁺). This indicates a dynamic process of erythroid maturation and differentiation. Additionally, widespread expression of β -globin mRNA and β^{A-T87Q} -globin mRNA was observed in the bone marrow. The cell classification results from the bone marrow and spleen also demonstrated a significant increase in various subtypes of cells derived from human hematopoietic stem cells, including hCD3⁺, hCD13⁺, hCD19⁺, hCD34⁺, and hCD56⁺. These findings suggest that BD211 has successfully engrafted in the mouse bone marrow, reconstituting the human erythroid system within the mice and differentiating stably into various stages of human erythroid cells. Furthermore, during the directed differentiation process, it has expressed human erythroid-specific genes and the target gene of interest.

During the trial period, mortality or near-death events occurred in all groups, with proportions of 17.50%, 19.23%, 11.54%, and 14.10%, respectively, showing no dose-dependent relationship. Most animals with unclear causes of near-death or death exhibited a mild to moderate increase in bone marrow (sternum) cell numbers, accompanied by neutrophilia. Combining the results of bone marrow smears and hematological examinations at D92, we analyzed that the cause of animal deaths may be related to the implantation of human hematopoietic stem cells. While reconstituting the human erythroid system, this implantation could potentially affect the hematopoietic function of the NCG-X mouse's bone marrow stem cells, leading to exacerbated anemia. The animal supplier provided data also indicated that the natural mortality rate of these mice begins to increase at 5 weeks of age, reaching up to 30% by 25 weeks, which is significantly higher than that of ordinary

TABLE 4 (*Continued*) Organ weight and coefficient of NCG-X mice treated with BD211 on D92

NCG mice (Data not published). Moreover, no reports of treatmentrelated deaths have been found in the safety studies of BD211 or other gene therapies for TDT (Mirza et al., 2025; Kwiatkowski et al., 2024). Therefore, the cause of death in these near-death/dead animals is mainly associated with the spontaneous anemia or immunodeficiency of the model mice rather than BD211.

After 13 weeks of administration (D92), hematological indicators in the mock and BD211 groups showed a decrease in absolute values compared to the solvent control group. This included reductions in red blood cell indices (RBC, HGB, HCT, MCHC, and RET), white blood cell indices (WBC, NEUT, LYMPH, MONO), and platelets (PLT). The decrease was dose-dependent with respect to the human CD34⁺ hematopoietic stem cell dose. Bone marrow smears at D92 also indicated active bone marrow proliferation in the mock control and BD211 groups, with a significant increase in the proportion of lymphocytes and a relative decrease in the number of granulocytes and erythrocytes, consistent with hematological changes. However, in our previous assessment of adverse events following BD211 transplantation, an increase in WBC and lymphocyte counts was observed in patients after intravenous infusion, accompanied by the recovery of other subtypes of cells and stabilization within the normal range (Li et al., 2024). The opposite results observed in animals may be due to the engraftment of human CD34⁺ hematopoietic stem cells in NCG-X mice, which can differentiate into various human cell types, leading to a relatively lower proportion of mouse-derived cells. Additionally, NCG-X mice retain some immune function, and their monocytes (macrophages) in the peripheral blood can continuously phagocytose xenogeneic human terminal blood cells that are released into the mouse peripheral blood after the reconstruction and differentiation of human CD34⁺ hematopoietic stem cells. These changes are attributed to the engraftment, reconstitution, and differentiation of human CD34⁺ hematopoietic stem cells in NCG-X mice rather than specific changes induced by the BD211.

The no observed adverse effect level of intravenously administered BD211 in NCG-X mice was 1.2×10^6 cells per mouse. Subsequently, comprehensive safety studies were conducted on BD211, which supported the application for an Investigational New Drug (IND) clinical trial. The application for "BD211 autologous CD34⁺ hematopoietic stem cell injection" was granted implied permission by the China National Drug Administration (acceptance number: CXSL2300710). The indication for this treatment is TDT.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary Material, further inquiries can be directed to the corresponding authors.

Ethics statement

The animal studies were approved by the Institutional Animal Care and Use Committee of Guoke Saifu Hebei Pharmaceutical Technology Co., Ltd. (approval numbers: IACUC-2023-149 and IACUC-2023-220) following the guidelines of the Association for Assessment and Accreditation of Laboratory Animal Care International (AAALAC).

Author contributions

XD: Data curation, Formal Analysis, Investigation, Methodology, Writing – original draft. ZL: Data curation, Formal Analysis, Investigation, Methodology, Writing – review and editing. SC: Conceptualization, Formal Analysis, Methodology, Writing – review and editing. YH: Data curation, Formal Analysis, Investigation, Writing – review and editing. SL: Project administration, Supervision, Writing – review and editing. QuW: Project administration, Supervision, Writing – review and editing. QiW: Project administration, Supervision, Writing – review and editing.

Funding

The author(s) declare that financial support was received for the research and/or publication of this article. This research was sponsored by BDGENE Therapeutics, Shanghai.

Conflict of interest

Authors XD, ZL, SC, YH, and QuW were employed by SAFE Pharmaceutical Technology Co., Ltd. Author SL was employed by BDGENE Therapeutics Co. Ltd.

The remaining author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The authors declare that this study received funding from BDGENE Therapeutics, Shanghai. The funder had the following involvement in the study: the study design, data collection and analysis, and the preparation of the manuscript. Additionally, the funder agreed to the decision to publish the article.

Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: https://www.frontiersin.org/articles/10.3389/fcell.2025. 1607707/full#supplementary-material

Bao, X., Zhang, X., Wang, L., Wang, Z., Huang, J., Zhang, Q., et al. (2021). Epigenetic inactivation of ERF reactivates γ -globin expression in β -thalassemia. *Am. J. Hum. Genet.* 108, 709–721. doi:10.1016/j.ajhg.2021.03.005

Boulad, F., Maggio, A., Wang, X., Moi, P., Acuto, S., Kogel, F., et al. (2022). Lentiviral globin gene therapy with reduced-intensity conditioning in adults with β -thalassemia: a phase 1 trial. *Nat. Med.* 28, 63–70. doi:10.1038/s41591-021-01554-9

Brendel, C., Negre, O., Rothe, M., Guda, S., Parsons, G., Harris, C., et al. (2020). Preclinical evaluation of a novel lentiviral vector driving lineage-specific BCL11A knockdown for sickle cell gene therapy. *Mol. Ther. Methods Clin. Dev.* 17, 589–600. doi:10.1016/j.omtm.2020.03.015

Cappellini, M. D., Bejaoui, M., Agaoglu, L., Canatan, D., Capra, M., Cohen, A., et al. (2011). Iron chelation with deferasirox in adult and pediatric patients with thalassemia major: efficacy and safety during 5 years' follow-up. *Blood* 118, 884–893. doi:10.1182/blood-2010-11-316646

Cappellini, M. D., Cohen, A., Porter, J., Taher, A., and Viprakasit, V. (2014). Guidelines for the Management of Transfusion Dependent Thalassaemia (TDT) [Internet], 3rd edition. Nicosia, Cyprus: Thalassaemia International Federation.

Chinese Medical Association (2022). Chinese guideline for diagnosis and treatment of transfusion dependent β -thalassemiA 2022. Zhonghua Xue Ye Xue Za Zhi 43, 889–896. doi:10.3760/cma.j.issn.0253-2727.2022.11.002

U. S. Food and Drug (2024). *Considerations for the development of chimeric antigen receptor (CAR) T cell products.*

Galanello, R., and Origa, R. (2010). Beta-thalassemia. Orphanet J. Rare Dis. 5, 11. doi:10.1186/1750-1172-5-11

Kleinschmidt, K., Penkivech, G., Troeger, A., Foell, J., Hanafee-Alali, T., Leszczak, S., et al. (2025). αß T-cell depleted haploidentical stem cell transplantation for pediatric and young adult patients with transfusion-dependent thalassemia. *Bone Marrow Transpl.* 60, 682–689. doi:10.1038/s41409-025-02546-w

Kronstein-Wiedemann, R., and Tonn, T. (2019). Colony Formation: an assay of hematopoietic progenitor cells. *Methods Mol. Biol.* 2017, 29–40. doi:10.1007/978-1-4939-9574-5_3

Kwiatkowski, J. L., Walters, M. C., Hongeng, S., Yannaki, E., Kulozik, A. E., Kunz, J. B., et al. (2024). Betibeglogene autotemcel gene therapy in patients with transfusion-dependent, severe genotype β -thalassaemia (HGB-212): a non-randomised, multicentre, single-arm, open-label, single-dose, phase 3 trial. Lancet 404, 2175–2186. doi:10.1016/S0140-6736(24)01884-1

Lang, E., and Lang, F. (2015). Mechanisms and pathophysiological significance of eryptosis, the suicidal erythrocyte death. *Semin. Cell Dev. Biol.* 39, 35-42. doi:10.1016/j.semcdb.2015.01.009

Li, S., Ling, S., Wang, D., Wang, X., Hao, F., Yin, L., et al. (2024). Modified lentiviral globin gene therapy for pediatric $\beta(0)/\beta(0)$ transfusion-dependent β -thalassemia: a single-center, single-arm pilot trial. *Cell Stem Cell* 31, 961–73.e8. doi:10.1016/j.stem.2024.04.021

Locatelli, F., Thompson, A. A., Kwiatkowski, J. L., Porter, J. B., Thrasher, A. J., Hongeng, S., et al. (2022). Betibeglogene autotemcel gene therapy for non- $\beta(0)/\beta(0)$ genotype β -thalassemia. *N. Engl. J. Med.* 386, 415–427. doi:10.1056/NEJMoa2113206

M3(R2) I (2009). Guidelines for non-clinical safety studies to support the clinical trial and marketing of drugs. FDA-2008-D-0470.

Magrin, E., Semeraro, M., Hebert, N., Joseph, L., Magnani, A., Chalumeau, A., et al. (2022). Long-term outcomes of lentiviral gene therapy for the β -hemoglobinopathies: the HGB-205 trial. *Nat. Med.* 28, 81–88. doi:10.1038/s41591-021-01650-w

Malay, J., Salama, R. A. A., Alam Qureshi, G. S., Ammar, ARAA, Janardhan, G., Safdar, M., et al. (2024). Gene therapy: a revolutionary step in treating thalassemia. *Hematol. Rep.* 16, 656–668. doi:10.3390/hematolrep16040064

Mcintosh, B. E., Brown, M. E., Duffin, B. M., Maufort, J. P., Vereide, D. T., Slukvin, I. I., et al. (2015). Nonirradiated NOD,B6.SCID Il2rγ-/- Kit(W41/W41) (NBSGW) mice support multilineage engraftment of human hematopoietic cells. *Stem Cell Rep.* 4, 171–180. doi:10.1016/j.stemcr.2014.12.005

Mirza, A., Ritsert, M. L., Tao, G., Thakar, H., Lobitz, S., Heine, S., et al. (2025). Gene therapy in transfusion-dependent non- $\beta 0/\beta 0$ genotype β -thalassemia: first real-world experience of beti-cel. *Blood Adv.* 9, 29–38. doi:10.1182/bloodadvances.2024014104

National Medical Products Administration (2017). *Technical guidelines for the research and evaluation of cell therapy products (trial)*. China: National medical products administration.

National Medical Products Administration (2021). Technical guidelines for nonclinical research of genetically modified cell therapy products (trial). China: National Medical Products Administration.

Origa, R. (2017). β-Thalassemia. Genet. Med. 19, 609–619. doi:10.1038/gim.2016.173

Segura, E. E. R., Ayoub, P. G., Hart, K. L., and Kohn, D. B. (2023). Gene therapy for β -hemoglobinopathies: from discovery to clinical trials. *Viruses* 15, 713. doi:10.3390/v15030713

Shah, F. T., Sayani, F., Trompeter, S., Drasar, E., and Piga, A. (2019). Challenges of blood transfusions in β -thalassemia. Blood Rev. 37, 100588. doi:10.1016/j.blre.2019.100588

Thompson, A. A., Walters, M. C., Kwiatkowski, J., Rasko, J. E. J., Ribeil, J. A., Hongeng, S., et al. (2018). Gene therapy in patients with transfusion-dependent β -thalassemia. *N. Engl. J. Med.* 378, 1479–1493. doi:10.1056/NEJMoa1705342

Thong, M. K., and Ngim, C. F. (2021). β -Thalassemias. N. Engl. J. Med. 384, 2165. doi:10.1056/NEJMc2105064

Wattanapanitch, M., Damkham, N., Potirat, P., Trakarnsanga, K., Janan, M., U-Pratya, Y., et al. (2018). One-step genetic correction of hemoglobin E/beta-thalassemia patient-derived iPSCs by the CRISPR/Cas9 system. *Stem Cell Res. Ther.* 9, 46. doi:10.1186/s13287-018-0779-3

Weatherall, D. J. (2010). The inherited diseases of hemoglobin are an emerging global health burden. *Blood* 115, 4331–4336. doi:10.1182/blood-2010-01-251348

Yang, H., Li, X., Que, L., Canatan, D., Capra, M., Cohen, A., et al. (2024). A novel conditioning regimen with pre-transplantation immunosuppression reduces the complication rates in hematopoietic stem cell transplantation in transfusion-dependent β -thalassemia. *Stem Cell Res. Ther.* 15, 495. doi:10.1186/s13287-024-04103-6

Zeng, J., Wu, Y., Ren, C., Bonanno, J., Shen, A. H., Shea, D., et al. (2020). Therapeutic base editing of human hematopoietic stem cells. *Nat. Med.* 26, 535–541. doi:10.1038/s41591-020-0790-y