## Check for updates

### **OPEN ACCESS**

APPROVED BY Frontiers Editorial Office, Frontiers Media SA, Switzerland

\*CORRESPONDENCE María Ángeles Muñoz-Fernández, mmunoz.hgugm@gmail.com

RECEIVED 27 June 2025 ACCEPTED 30 June 2025 PUBLISHED 11 July 2025

### CITATION

Magro-Lopez E, Vazquez-Alejo E, Espinar-Buitrago MdlS and Muñoz-Fernández MÁ (2025) Correction: Optimizing Nodal, Wnt and BMP signaling pathways for robust and efficient differentiation of human induced pluripotent stem cells to intermediate mesoderm cells. *Front. Cell Dev. Biol.* 13:1655115. doi: 10.3389/fcell.2025.1655115

### COPYRIGHT

© 2025 Magro-Lopez, Vazquez-Alejo, Espinar-Buitrago and Muñoz-Fernández. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Correction: Optimizing Nodal, Wnt and BMP signaling pathways for robust and efficient differentiation of human induced pluripotent stem cells to intermediate mesoderm cells

Esmeralda Magro-Lopez<sup>1,2,3</sup>, Elena Vazquez-Alejo<sup>1,2,3</sup>, María de la Sierra Espinar-Buitrago<sup>1,2,3</sup> and María Ángeles Muñoz-Fernández<sup>1,2,3</sup>\*

<sup>1</sup>Molecular Immuno-Biology Laboratory, Immunology Section, Hospital General Universitario Gregorio Marañón (HGUGM), Madrid, Spain, <sup>2</sup>Instituto de Investigación Sanitaria Gregorio Marañón (IiSGM), Madrid, Spain, <sup>3</sup>Biomedical Research Networking Center in Bioengineering, Biomaterials and Nanomedicine (CIBER-BBN), Instituto de Salud Carlos III (ISCIII), Madrid, Spain

### KEYWORDS

hiPSC, mesoderm progenitors, intermediate mesoderm, Nodal, Wnt, BMPs, 3D organoids, urogenital organoids

### A Correction on:

Optimizing Nodal, Wnt and BMP signaling pathways for robust and efficient differentiation of human induced pluripotent stem cells to intermediate mesoderm cells

by Magro-Lopez E, Vazquez-Alejo E, Espinar-Buitrago MdlS and Muñoz-Fernández MÁ (2024). Front. Cell Dev. Biol. 12:1395723. doi: 10.3389/fcell.2024.1395723

In the published article, there was an omission in the Funding statement.

The co-funders Ministerio de Ciencia e Innovación and Unión Europea-European Regional Development Fund were erroneously omitted.

## Funding

The author(s) declare that financial support was received for the research and/or publication of this article. This work has been supported with resources from ISCIII (grant number PI23/01137) and co-funded by the European Union, and by CIBER - Consorcio Centro de Investigación Biomédica en Red- (CIBER-BBN) (CB22/01/00041), Instituto de Salud Carlos III, Ministerio de Ciencia e Innovación and Unión Europea – European Regional Development Fund. EM-L was supported by the Instituto de Investigación Sanitaria Gregorio Marañón under grant agreement "2022-II-POSTDOC-01"

through the Intramural Program." EV-A was supported by CIBER -BBN, ISCIII (CB22/01/00041), Instituto de Salud Carlos III, Ministerio de Ciencia e Innovación and Unión Europea – European Regional Development Fund. MSE-B was supported with resources from ISCIII (grant number PI19/01638) and co-funded by the European Union.

The original article has been updated.

# Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.