



Corrigendum: Association of Hantavirus Infections and Leptospirosis With the Occurrence of Chronic Kidney Disease of Uncertain Etiology in the North Central Province of Sri Lanka: A Prospective Study With Patients and Healthy Persons

OPEN ACCESS

Approved by:

Frontiers Editorial Office,
Frontiers Media SA, Switzerland

***Correspondence:**

N. P. Sunil-Chandra
sunilchandra@kln.ac.lk

Specialty section:

This article was submitted to
Virus and Host,
a section of the journal
*Frontiers in Cellular
and Infection Microbiology*

Received: 20 November 2020

Accepted: 25 November 2020

Published: 17 December 2020

Citation:

Sunil-Chandra NP, Jayaweera JAAS, Kumbukgolla W and Jayasundara MVML (2020) Corrigendum: Association of Hantavirus Infections and Leptospirosis With the Occurrence of Chronic Kidney Disease of Uncertain Etiology in the North Central Province of Sri Lanka: A Prospective Study With Patients and Healthy Persons. *Front. Cell. Infect. Microbiol.* 10:631515. doi: 10.3389/fcimb.2020.631515

N. P. Sunil-Chandra^{1*}, J. A. A. S. Jayaweera², W. Kumbukgolla²
and M. V. M. L. Jayasundara¹

¹ Department of Medical Microbiology, Faculty of Medicine, University of Kelaniya, Kelaniya, Sri Lanka, ² Faculty of Medicine & Allied Sciences, Rajarata University of Sri Lanka, Anuradhapura, Sri Lanka

Keywords: hantaviruses, leptospira, sero-prevalence, chronic kidney disease, CKDu

A Corrigendum on

Association of Hantavirus Infections and Leptospirosis With the Occurrence of Chronic Kidney Disease of Uncertain Etiology in the North Central Province of Sri Lanka: A Prospective Study With Patients and Healthy Persons

by Sunil-Chandra NP, Jayaweera JAAS, Kumbukgolla W and Jayasundara MVML (2020). *Front. Cell. Infect. Microbiol.* 10:556737. doi: 10.3389/fcimb.2020.556737

In the original article, there was an error in the ethical clearance number. The correct number for ethical clearance is ERC/2012/33.

A correction has been made to **Materials and Methods, Paragraph 3:**

According to CKDu case definition, an individual identified with an albumin-to-creatinine ratio (ACR) $\geq 30 \text{ mg/g}$ urine creatinine during the initial visit and at a follow-up visit, a normal glycosylated hemoglobin ($\text{HbA}_{1c} < 6.5\%$), not on treatment for diabetes, no elevated blood pressure, and no past history of kidney disease or snake bite were included. Patients with other known causes of chronic kidney disease (CKD) and those who did not consent were excluded from the study (16,13). Studies involving human patients/participants were reviewed and approved by the Research, Ethical Review and Higher degree

committee of the faculty of Medicine and Allied Sciences, Rajarata University, Sri Lanka. Patients/participants provided their written informed consent to participate in this study (ERC/2012/33).

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2020 Sunil-Chandra, Jayaweera, Kumbukgolla and Jayasundara. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.