



# Corrigendum: Activation of transient receptor potential vanilloid 4 increases NMDA-activated current in hippocampal pyramidal neurons

Lin Li<sup>1,2</sup>, Weijun Qu<sup>1</sup>, Libin Zhou<sup>1</sup>, Zihong Lu<sup>1</sup>, Pinghui Jie<sup>1</sup>, Lei Chen<sup>1,\*</sup> and Ling Chen<sup>1,2\*</sup>

<sup>1</sup> Department of Physiology, Nanjing Medical University, Nanjing, China

<sup>2</sup> State Key Laboratory of Reproductive Medicine, Department of Histology and Embryology, Nanjing Medical University, Nanjing, China

\*Correspondence: chenl@njmu.edu.cn

**Edited and reviewed by:**

Dieter Wicher, Max Planck Institute for Chemical Ecology, Germany

**Keywords:** TRPV4, NMDA receptor, NR2B subunit, phosphorylation, excitotoxicity, cerebral ischemia

## A corrigendum on

### Activation of transient receptor potential vanilloid 4 increases NMDA-activated current in hippocampal pyramidal neurons

by Li, L., Qu, W., Zhou, L., Lu, Z., Jie, P., Chen, L., et al. (2013). *Front. Cell. Neurosci.* 7:17. doi: 10.3389/fncel.2013.00017

The dosage of HC-067047 used in the article by Li et al. (2013) should be “10 μM/2 μl/mouse.” In addition, in appendix, the perfusion solution for recording EPSCs should be composed

of (in mM): NaCl 126, CaCl<sub>2</sub> 1, KCl 2.5, MgCl<sub>2</sub> 1, NaHCO<sub>3</sub> 26, KH<sub>2</sub>PO<sub>4</sub> 1.25, D-glucose 20 and bicuculline 0.01.

## REFERENCES

- Li, L., Qu, W., Zhou, L., Lu, Z., Jie, P., Chen, L., et al. (2013). Activation of transient receptor potential vanilloid 4 increases NMDA-activated current in hippocampal pyramidal neurons. *Front. Cell. Neurosci.* 7:17. doi: 10.3389/fncel.2013.00017

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 25 April 2014; accepted: 26 April 2014;  
published online: 19 May 2014.

Citation: Li L, Qu W, Zhou L, Lu Z, Jie P, Chen L and Chen L (2014) Corrigendum: Activation of transient receptor potential vanilloid 4 increases NMDA-activated current in hippocampal pyramidal neurons. *Front. Cell. Neurosci.* 8:130. doi: 10.3389/fncel.2014.00130

This article was submitted to the journal *Frontiers in Cellular Neuroscience*.

Copyright © 2014 Li, Qu, Zhou, Lu, Jie, Chen and Chen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.