



Presynaptic mechanisms of neuronal plasticity and their role in epilepsy

Jochen C. Meier^{1*}, Marcus Semtner¹, Aline Winkelmann¹ and Jakob Wolfart²

¹ RNA Editing and Hyperexcitability Disorders, Max Delbrück Centre for Molecular Medicine, Neurosciences, Berlin, Germany

² Oscar Langendorff Institute of Physiology, University of Rostock, Rostock, Germany

Edited by:

Jaichandar Subramanian,
Massachusetts Institute of
Technology, USA

Reviewed by:

Detlev Boison, Legacy Research
Institute, USA

Rory McQuiston, Virginia
Commonwealth University, USA

***Correspondence:**

Jochen C. Meier, RNA Editing and
Hyperexcitability Disorders, Max
Delbrück Centre for Molecular
Medicine, Neurosciences, Building
C31.2, Robert-Rössle-Straße 10,
13125 Berlin, Germany
e-mail: jochen.meier@
mdc-berlin.de

Synaptic communication requires constant adjustments of pre- and postsynaptic efficacies. In addition to synaptic long term plasticity, the presynaptic machinery underlies homeostatic regulations which prevent out of range transmitter release. In this minireview we will discuss the relevance of selected presynaptic mechanisms to epilepsy including voltage- and ligand-gated ion channels as well as cannabinoid and adenosine receptor signaling.

Keywords: epilepsy, axon, RNA editing, potassium channels, glycine receptor, homeostatic regulation, neuropsychiatric disorders, hippocampus

INTRODUCTION

Many studies highlighted the importance of homeostasis in neuronal signaling either within neurons, then called “intrinsic plasticity” or between neurons, referred to as “synaptic scaling” (Davis and Bezprozvanny, 2001; Eichler and Meier, 2008; Turrigiano, 2011). Classic synaptic plasticity in its non-homeostatic “Hebbian” form and pathological disturbances need counterbalancing homeostatic scaling mechanisms (Abbott and Nelson, 2000). The latter was mainly regarded from the postsynaptic perspective (Thiagarajan et al., 2005; Groth et al., 2011). However, in addition to presynaptic expression of synaptic long term plasticity (Nicoll and Schmitz, 2005), slow homeostatic regulations occur in the presynapse, e.g., in form of chronic receptor or ion channel modulations.

PRESYNAPTIC ION CHANNEL AND GLYCINE RECEPTOR PLASTICITY

Because transmitter release is controlled by action potential-(AP)-triggered calcium influx in the synaptic terminal, regulation of ion channels which shape the axonal AP and terminal depolarization is an effective mechanism of presynaptic plasticity. With this definition, the AP initiation zone (AIZ) could be viewed as part of the presynaptic equipment. A striking form of homeostatic plasticity has been documented for the AIZ: this entire subcellular structure including voltage-gated Na^+ (Na_v) and K^+ (K_v) channels can be shifted along the axon (Figure 1), thereby counteracting hyperexcitation by increasing thresholds for AP generation (Grubb and Burrone, 2010). Such axonal remodeling may be facilitated via ion channel trafficking

regulated by alternative splicing, as shown for “shaw-related” $\text{Kv}3$ channels (Gu et al., 2012). Other important constituents of presynaptic control are “shaker-related” $\text{Kv}1$ channels (Wang et al., 1994). Their role is well demonstrated for the large glutamatergic mossy fiber boutons of dentate granule cells, which impinge on hippocampal CA3 pyramidal cells (Geiger and Jonas, 2000; Bischofberger et al., 2006). Here, K_v channels could gain further importance during temporal lobe epilepsy (TLE), when seizures invade the hippocampus and feedforward inhibition of CA3 pyramidal cells via interneurons is compromised (Lawrence and McBain, 2003). Indeed, seizures trigger a transcriptional upregulation of $\text{K}_v1.1$ channels in granule cells, thereby delaying their AP responses considerably, as recently shown in a TLE mouse model (Kirchheim et al., 2013). Consistent with the view that $\text{K}_v1.1$ is a promising antiepileptic target, $\text{K}_v1.1$ knockout mice develop epilepsy (Wenzel et al., 2007) and lentiviral over-expression of $\text{K}_v1.1$ ameliorates seizures in an animal model of neocortical epilepsy (Wykes et al., 2012). The interaction of activity-dependent downscaling and potentiation of presynaptic excitability may involve the adenylyl cyclase pathway (Nicoll and Schmitz, 2005) but it is still unclear how exactly these seemingly opposed mechanisms interact in the same presynaptic compartment.

Epilepsy often comes with cognitive dysfunction and neuropsychiatric comorbidities (García-Morales et al., 2008). We discovered a molecule which in this regard may have an important impact: an RNA variant of the neurotransmitter receptor for glycine (GlyR). The GlyRs are subject to increased RNA editing in resected hippocampi of TLE patients (Eichler et al., 2008)




FIGURE 1 | Scheme depicting strategic molecules relevant for homeostasis of presynaptic function and epilepsy discussed in this minireview. In glutamatergic neurons, gain-of-function of molecules colored red/orange will increase network excitability, whereas those colored green will

decrease it. The opposite holds true if the changes occur in GABAergic neurons. Abbreviations: Na_V , Ca_V , K_V , voltage-gated sodium, calcium and potassium channels; AIZ, action potential (AP) initiation zone; GlyR, glycine receptors; A1, adenosine receptor; CB₁, cannabinoid receptor 1.

which profoundly influences biophysical receptor properties. The reason is an amino acid substitution in the ligand binding domain leading to gain-of-function receptors with increased neurotransmitter affinity (Meier et al., 2005; Eichler et al., 2008; Legendre et al., 2009) and spontaneous channel activity (Kletke et al., 2013; Winkelmann et al., 2014). In addition, RNA splicing governs presynaptic GlyR expression (Winkelmann et al., 2014), and in hippocampal neurons, the lack of the GlyR β subunit (Weltzien et al., 2012) which governs postsynaptic receptor clustering (Meyer et al., 1995; Meier et al., 2000, 2001; Eichler et al., 2009; Förster et al., 2010; Kowalczyk et al., 2013), certainly facilitates GlyR expression and function at presynapses (Figure 1). Presynaptic GlyRs are tightly packed (~ 200 receptor channels in a cluster with ~ 100 nm radius; Notelaers et al., 2012, 2014a,b), which implies that a single presynaptic cluster from the spontaneously active GlyR RNA variant will have a considerable functional impact on synaptic neurotransmitter release, even if the contribution of the glycinergic system to this brain region appears limited (Zeilhofer et al., 2005). Consistent with the excitatory nature of presynaptic chloride channels and the well documented presynaptic GlyR expression in the hippocampus (Kubota et al., 2010; Ruiz et al., 2010; Waseem and Fedorovich, 2010; Winkelmann et al., 2014), we found that the spontaneously active GlyR RNA variant actually increased presynaptic excitability and the functional impact of glutamatergic neurons or parvalbumin-positive interneurons *in vivo* and, depending on the type of neuron, triggered cognitive dysfunction or anxiety in our mouse model of epilepsy (Winkelmann et al., 2014). In agreement with the proposed critical role of presynaptic GlyRs in the regulation of neural network excitability, application of a low, non-receptor-saturating, glycine concentration ($10 \mu\text{M}$) to corticohippocampal slice preparations was sufficient to enhance epileptiform activity induced by block of K_V channels (Chen et al., 2014).

RETROGRADE AUTOCRINE AND PARACRINE SIGNALING

Although the idea of cannabis as a potential antiepileptic drug is ancient, it remained elusive how it could work reliably (Adams and Martin, 1996; Miller, 2013). Recent discoveries on endogenous cannabinoid receptors (CB), of which particularly CB₁ is widely expressed in presynaptic terminals of excitatory and inhibitory neurons (Figure 1), could lead to a better

understanding of CB mechanisms in epilepsy (Alger, 2004; Katona and Freund, 2008; Hill et al., 2012). In GABAergic neurons, activation of CB₁, e.g., via neuronal activity-dependent retrograde post-to-presynapse release of CBs anandamide or 2-AG has been shown to decrease synaptic GABA release, a mechanism termed depolarization-induced suppression of inhibition (DSI; Ohno-Shosaku et al., 2001; Wilson et al., 2001). Consistently, elevated CB₁ presence observed in epilepsy models and TLE patients (Goffin et al., 2011; Karlóca et al., 2011; Bojnik et al., 2012) has been interpreted as proconvulsive (Chen et al., 2003, 2007). On the other hand, CB₁ is also expressed on glutamatergic terminals, where its activation reduces glutamate release (Domenici et al., 2006; Kawamura et al., 2006). Furthermore, CB₁ activation increases inward rectifier K⁺ (Kir) currents (Mackie et al., 1995; Chemin et al., 2001) mediated via postsynaptic channels which are also upregulated in TLE (Young et al., 2009; Stegen et al., 2012). In summary, while elevation of CB₁ at GABAergic synapses and reduction at glutamatergic synapses likely constitute endogenous adaptations to epilepsy, exogenous CB₁ overexpression and activation in principal neurons, possibly via receptors physiologically rarely activated, could effectively protect against seizures (Blair et al., 2006; Guggenhuber et al., 2010; Hofmann and Frazier, 2013).

Adenosine triphosphate (ATP) is released from astrocytes and can enhance neuronal excitability through its direct action onto purinergic receptors. However, ATP can also exert indirect effects upon its enzymatic conversion to adenosine and signaling through adenosine A1 receptors, which reduces synaptic glutamate release (Nicoll and Schmitz, 2005; Boison, 2013; Dias et al., 2013). Therefore, adenosine signaling is another mechanism of presynaptic homeostasis with recognized relevance to epilepsy; while too much adenosine clearance via gliosis-enhanced adenosine kinase activity is a proconvulsive factor, adenosine augmentation in the epileptic focus represents a powerful anticonvulsive principle (Boison, 2012).

PERSPECTIVE

In agreement with the proposed critical role of presynaptic compartments in the regulation of neural network homeostasis, diverse pharmacological agents with a presynaptic mode of action were reported to be effective in the treatment of epilepsy. In

particular drugs which provide rapid adaptation against excessive excitation, e.g., via use-dependent inhibition of Na_v or Ca_v channels (phenytoin, carbamazepine, lamotrigine, topiramate, and levetiracetam), likely act primarily in axons (Stefani et al., 1996; Wu et al., 1998; Catterall, 1999; Vogl et al., 2012). Interestingly, these drugs also have effects on psychiatric symptoms (Barbosa et al., 2003; Lexi-Comp, 2009; Andrus and Gilbert, 2010) indicating common underlying mechanisms of cognitive dysfunction and psychiatric symptoms of epilepsy. With more research on neuron type-specific roles in behavior (Lovett-Barron et al., 2014; Winkelmann et al., 2014), new antiepileptic strategies could ground on these insights and specifically target presynaptic molecules in the affected cell types.

REFERENCES

- Abbott, L. F., and Nelson, S. B. (2000). Synaptic plasticity: taming the beast. *Nat. Neurosci.* 3(Suppl), 1178–1183. doi: 10.1038/81453
- Adams, I. B., and Martin, B. R. (1996). Cannabis: pharmacology and toxicology in animals and humans. *Addiction* 91, 1585–1614. doi: 10.1046/j.1360-0443.1996.91115852.x
- Alger, B. E. (2004). Endocannabinoids and their implications for epilepsy. *Epilepsy Curr.* 4, 169–173. doi: 10.1111/j.1535-7597.2004.04501.x
- Andrus, M. R., and Gilbert, E. (2010). Treatment of civilian and combat-related posttraumatic stress disorder with topiramate. *Ann. Pharmacother.* 44, 1810–1816. doi: 10.1345/aph.1P163
- Barbosa, L., Berk, M., and Vorster, M. (2003). A double-blind, randomized, placebo-controlled trial of augmentation with lamotrigine or placebo in patients concomitantly treated with fluoxetine for resistant major depressive episodes. *J. Clin. Psychiatry* 64, 403–407. doi: 10.4088/jcp.v64n0407
- Bischofberger, J., Engel, D., Frotscher, M., and Jonas, P. (2006). Timing and efficacy of transmitter release at mossy fiber synapses in the hippocampal network. *Pflügers Arch.* 453, 361–372. doi: 10.1007/s00424-006-0093-2
- Blair, R. E., Deshpande, L. S., Sombati, S., Falenski, K. W., Martin, B. R., and DeLorenzo, R. J. (2006). Activation of the cannabinoid type-1 receptor mediates the anticonvulsant properties of cannabinoids in the hippocampal neuronal culture models of acquired epilepsy and status epilepticus. *J. Pharmacol. Exp. Ther.* 317, 1072–1078. doi: 10.1124/jpet.105.100354
- Boison, D. (2012). Adenosine dysfunction in epilepsy. *Glia* 60, 1234–1243. doi: 10.1002/glia.22285
- Boison, D. (2013). Adenosine kinase: exploitation for therapeutic gain. *Pharmacol. Rev.* 65, 906–943. doi: 10.1124/pr.112.006361
- Bojniki, E., Turunc, E., Armanag, G., Kanit, L., Benyhe, S., Yalcin, A., et al. (2012). Changes in the cannabinoid (CB1) receptor expression level and G-protein activation in kainic acid induced seizures. *Epilepsy Res.* 99, 64–68. doi: 10.1016/j.eplepsyres.2011.10.020
- Catterall, W. A. (1999). Molecular properties of brain sodium channels: an important target for anticonvulsant drugs. *Adv. Neurol.* 79, 441–456.
- Chemin, J., Monteil, A., Perez-Reyes, E., Nargeot, J., and Lory, P. (2001). Direct inhibition of T-type calcium channels by the endogenous cannabinoid anandamide. *EMBO J.* 20, 7033–7040. doi: 10.1093/emboj/20.24.7033
- Chen, K., Neu, A., Howard, A. L., Foldy, C., Echegoyen, J., Hilgenberg, L., et al. (2007). Prevention of plasticity of endocannabinoid signaling inhibits persistent limbic hyperexcitability caused by developmental seizures. *J. Neurosci.* 27, 46–58. doi: 10.1523/jneurosci.3966-06.2007
- Chen, R., Okabe, A., Sun, H., Sharopov, S., Hangau-Opatz, I. L., Kolbaev, S. N., et al. (2014). Activation of glycine receptors modulates spontaneous epileptiform activity in the immature rat hippocampus. *J. Physiol.* 592(Pt. 10), 2153–2168. doi: 10.1113/jphysiol.2014.271700
- Chen, K., Ratzliff, A., Hilgenberg, L., Gulyas, A., Freund, T. F., Smith, M., et al. (2003). Long-term plasticity of endocannabinoid signaling induced by developmental febrile seizures. *Neuron* 39, 599–611. doi: 10.1016/s0896-6273(03)00499-9
- Davis, G. W., and Bezprozvanny, I. (2001). Maintaining the stability of neural function: a homeostatic hypothesis. *Annu. Rev. Physiol.* 63, 847–869. doi: 10.1146/annurev.physiol.63.1.847
- Dias, R. B., Rombo, D. M., Ribeiro, J. A., Henley, J. M., and Sebastiao, A. M. (2013). Adenosine: setting the stage for plasticity. *Trends Neurosci.* 36, 248–257. doi: 10.1016/j.tins.2012.12.003
- Domenici, M. R., Azad, S. C., Marsicano, G., Schierloh, A., Wotjak, C. T., Dodt, H. U., et al. (2006). Cannabinoid receptor type 1 located on presynaptic terminals of principal neurons in the forebrain controls glutamatergic synaptic transmission. *J. Neurosci.* 26, 5794–5799. doi: 10.1523/jneurosci.0372-06.2006
- Eichler, S. A., Förster, B., Smolinsky, B., Jüttner, R., Lehmann, T. N., Fahling, M., et al. (2009). Splice-specific roles of glycine receptor $\alpha 3$ in the hippocampus. *Eur. J. Neurosci.* 30, 1077–1091. doi: 10.1111/j.1460-9568.2009.06903.x
- Eichler, S. A., Kirischuk, S., Jüttner, R., Schäfermeier, P. K., Legendre, P., Lehmann, T. N., et al. (2014). Glycinergic tonic inhibition of hippocampal neurons with depolarising GABAergic transmission elicits histopathological signs of temporal lobe epilepsy. *J. Cell. Mol. Med.* 12, 2848–2866. doi: 10.1111/j.1582-4934.2008.00357.x
- Eichler, S. A., and Meier, J. C. (2008). E-I balance and human diseases - from molecules to networking. *Front. Mol. Neurosci.* 1:2. doi: 10.3389/neuro.02.002.2008
- Förster, B., Belaidi, A. A., Jüttner, R., Bernert, C., Tsokos, M., Lehmann, T. N., et al. (2010). Irregular RNA splicing curtails postsynaptic gephyrin in the cornu ammonis of patients with epilepsy. *Brain* 133, 3778–3794. doi: 10.1093/brain/awq298
- Garcia-Morales, I., de la Pena, M. P., and Kanner, A. M. (2008). Psychiatric comorbidities in epilepsy: identification and treatment. *Neurologist* 14, S15–S25. doi: 10.1097/01.nrl.0000340788.07672.51
- Geiger, J. R., and Jonas, P. (2000). Dynamic control of presynaptic $\text{Ca}^{(2+)}$ inflow by fast-inactivating $\text{K}(+)$ channels in hippocampal mossy fiber boutons. *Neuron* 28, 927–939. doi: 10.1016/s0896-6273(00)00164-1
- Goffin, K., Van, P. W., and Van, L. K. (2011). In vivo activation of endocannabinoid system in temporal lobe epilepsy with hippocampal sclerosis. *Brain* 134, 1033–1040. doi: 10.1093/brain/awq385
- Groth, R. D., Lindskog, M., Thiagarajan, T. C., Li, L., and Tsien, R. W. (2011). Beta $\text{Ca}^{2+}/\text{CaM}$ -dependent kinase type II triggers upregulation of GluA1 to coordinate adaptation to synaptic inactivity in hippocampal neurons. *Proc. Natl. Acad. Sci. U.S.A.* 108, 828–833. doi: 10.1073/pnas.1018022108
- Grubb, M. S., and Burrone, J. (2010). Activity-dependent relocation of the axon initial segment fine-tunes neuronal excitability. *Nature* 465, 1070–1074. doi: 10.1038/nature09160
- Gu, Y., Barry, J., McDougel, R., Terman, D., and Gu, C. (2012). Alternative splicing regulates kv3.1 polarized targeting to adjust maximal spiking frequency. *J. Biol. Chem.* 287, 1755–1769. doi: 10.1074/jbc.m11.299305
- Guggenhuber, S., Monory, K., Lutz, B., and Klugmann, M. (2010). AAV vector-mediated overexpression of CB1 cannabinoid receptor in pyramidal neurons of the hippocampus protects against seizure-induced excitotoxicity. *PLoS One* 5:e15707. doi: 10.1371/journal.pone.0015707
- Hill, A. J., Williams, C. M., Whalley, B. J., and Stephens, G. J. (2012). Phytocannabinoids as novel therapeutic agents in CNS disorders. *Pharmacol. Ther.* 133, 79–97. doi: 10.1016/j.pharmthera.2011.09.002
- Hofmann, M. E., and Frazier, C. J. (2013). Marijuana, endocannabinoids and epilepsy: potential and challenges for improved therapeutic intervention. *Exp. Neurol.* 244, 43–50. doi: 10.1016/j.expneurol.2011.11.047
- Karlócaí, M. R., Toth, K., Watanabe, M., Ledent, C., Juhasz, G., Freund, T. F., et al. (2011). Redistribution of CB1 cannabinoid receptors in the acute and chronic phases of pilocarpine-induced epilepsy. *PLoS One* 6:e27196. doi: 10.1371/journal.pone.0027196
- Katona, I., and Freund, T. F. (2008). Endocannabinoid signaling as a synaptic circuit breaker in neurological disease. *Nat. Med.* 14, 923–930. doi: 10.1038/nfm.1869
- Kawamura, Y., Fukaya, M., Maejima, T., Yoshida, T., Miura, E., Watanabe, M., et al. (2006). The CB1 cannabinoid receptor is the major cannabinoid receptor at excitatory presynaptic sites in the hippocampus and cerebellum. *J. Neurosci.* 26, 2991–3001. doi: 10.1523/jneurosci.4872-05.2006
- Kirchheim, F., Tinnes, S., Haas, C. A., Stegen, M., and Wolfart, J. (2013). Regulation of action potential delays via voltage-gated potassium Kv1.1 channels in dentate granule cells during hippocampal epilepsy. *Front. Cell. Neurosci.* 7:248. doi: 10.3389/fncel.2013.00248
- Kletke, O., Sergeeva, O. A., Lorenz, P., Oberland, S., Meier, J. C., Hatt, H., et al. (2013). New insights in endogenous modulation of ligand-gated ion channels:

- Histamine is an inverse agonist at strychnine sensitive glycine receptors. *Eur. J. Pharmacol.* 710, 59–66. doi: 10.1016/j.ejphar.2013.04.002
- Kowalczyk, S., Winkelmann, A., Smolinsky, B., Förster, B., Neundorf, I., Schwarz, G., et al. (2013). Direct binding of GABA(A) receptor $\beta 2$ and $\beta 3$ subunits to gephyrin. *Eur. J. Neurosci.* 37, 544–554. doi: 10.1111/ejn.12078
- Kubota, H., Alle, H., Betz, H., and Geiger, J. R. (2010). Presynaptic glycine receptors on hippocampal mossy fibers. *Biochem. Biophys. Res. Commun.* 393, 587–591. doi: 10.1016/j.bbrc.2010.02.019
- Lawrence, J. J., and McBain, C. J. (2003). Interneuron diversity series: containing the detonation—feedforward inhibition in the CA3 hippocampus. *Trends Neurosci.* 26, 631–640. doi: 10.1016/j.tins.2003.09.007
- Legendre, P., Förster, B., Jüttner, R., and Meier, J. C. (2009). Glycine receptors caught between genome and proteome—functional implications of RNA editing and splicing. *Front. Mol. Neurosci.* 2:23. doi: 10.3389/neuro.02.023.2009
- Lexi-Comp. (2009). “Carbamazepine,” in *The Merck Manual*, ed R. S. Porter (Whitehouse Station, NJ, U.S.A: Merck Sharp and Dohme Corp., a subsidiary of Merck and Co., Inc.).
- Lovett-Barron, M., Kaifosh, P., Kheirbek, M. A., Danielson, N., Zaremba, J. D., Reardon, T. R., et al. (2014). Dendritic inhibition in the hippocampus supports fear learning. *Science* 343, 857–863. doi: 10.1126/science.1247485
- Mackie, K., Lai, Y., Westenbroek, R., and Mitchell, R. (1995). Cannabinoids activate an inwardly rectifying potassium conductance and inhibit Q-type calcium currents in AtT20 cells transfected with rat brain cannabinoid receptor. *J. Neurosci.* 15, 6552–6561.
- Meier, J. C., Henneberger, C., Melnick, I., Racca, C., Harvey, R. J., Heinemann, U., et al. (2005). RNA editing produces glycine receptor $\alpha 3\beta 1\gamma 5\delta 1$ resulting in high agonist potency. *Nat. Neurosci.* 8, 736–744. doi: 10.1038/nn1467
- Meier, J., Meunier-Durmort, C., Forest, C., Triller, A., and Vannier, C. (2000). Formation of glycine receptor clusters and their accumulation at synapses. *J. Cell Sci.* 113, 2783–2795.
- Meier, J., Vannier, C., Serge, A., Triller, A., and Choquet, D. (2001). Fast and reversible trapping of surface glycine receptors by gephyrin. *Nat. Neurosci.* 4, 253–260. doi: 10.1038/85099
- Meyer, G., Kirsch, J., Betz, H., and Langosch, D. (1995). Identification of a gephyrin binding motif on the glycine receptor beta subunit. *Neuron* 15, 563–572. doi: 10.1016/0896-6273(95)90145-0
- Miller, J. W. (2013). Slim evidence for cannabinoids for epilepsy. *Epilepsy Curr.* 13, 81–82. doi: 10.5698/1535-7597-13.2.81
- Nicoll, R. A., and Schmitz, D. (2005). Synaptic plasticity at hippocampal mossy fibre synapses. *Nat. Rev. Neurosci.* 6, 863–876. doi: 10.1038/nrn1786
- Notelaers, K., Smisdom, N., Rocha, S., Janssen, D., Meier, J. C., Rigo, J. M., et al. (2012). Ensemble and single particle fluorimetric techniques in concerted action to study the diffusion and aggregation of the glycine receptor alpha3 isoforms in the cell plasma membrane. *Biochim. Biophys. Acta* 1818, 3131–3140. doi: 10.1016/j.bbamem.2012.08.010
- Notelaers, K., Rocha, S., Paesen, R., Smisdom, N., De, C. B., Meier, J. C., et al. (2014a). Analysis of alpha3 GlyR single particle tracking in the cell membrane. *Biochim. Biophys. Acta* 1843, 544–553. doi: 10.1016/j.bbamcr.2013.11.019
- Notelaers, K., Rocha, S., Paesen, R., Swinnen, N., Vangindertael, J., Meier, J. C., et al. (2014b). Membrane distribution of the glycine receptor alpha3 studied by optical super-resolution microscopy. *Histochem. Cell Biol.* doi: 10.1007/s00418-014-1197-y. [Epub ahead of print].
- Ohno-Shosaku, T., Maejima, T., and Kano, M. (2001). Endogenous cannabinoids mediate retrograde signals from depolarized postsynaptic neurons to presynaptic terminals. *Neuron* 29, 729–738. doi: 10.1016/s0896-6273(01)00247-1
- Ruiz, A., Campanac, E., Scott, R. S., Rusakov, D. A., and Kullmann, D. M. (2010). Presynaptic GABA(A) receptors enhance transmission and LTP induction at hippocampal mossy fiber synapses. *Nat. Neurosci.* 13, 431–438. doi: 10.1038/nn.2512
- Stefani, A., Spadoni, F., Siniscalchi, A., and Bernardi, G. (1996). Lamotrigine inhibits Ca²⁺ currents in cortical neurons: functional implications. *Eur. J. Pharmacol.* 307, 113–116. doi: 10.1016/0014-2999(96)00265-8
- Stegen, M., Kirchheim, F., Hanuschkin, A., Staszewski, O., Veh, R. W., and Wolfart, J. (2012). Adaptive intrinsic plasticity in human dentate gyrus granule cells during temporal lobe epilepsy. *Cereb. Cortex* 22, 2087–2101. doi: 10.1093/cercor/bhr294
- Thiagarajan, T. C., Lindskog, M., and Tsien, R. W. (2005). Adaptation to synaptic inactivity in hippocampal neurons. *Neuron* 47, 725–737. doi: 10.1016/j.neuron.2005.06.037
- Turrigiano, G. (2011). Too many cooks? Intrinsic and synaptic homeostatic mechanisms in cortical circuit refinement. *Annu. Rev. Neurosci.* 34, 89–103. doi: 10.1146/annurev-neuro-060909-153238
- Vogl, C., Mochida, S., Wolff, C., Whalley, B. J., and Stephens, G. J. (2012). The synaptic vesicle glycoprotein 2A ligand levetiracetam inhibits presynaptic Ca²⁺ channels through an intracellular pathway. *Mol. Pharmacol.* 82, 199–208. doi: 10.1124/mol.111.076687
- Wang, H., Kunkel, D. D., Schwartzkroin, P. A., and Tempel, B. L. (1994). Localization of Kv1.1 and Kv1.2, two K channel proteins, to synaptic terminals, somata and dendrites in the mouse brain. *J. Neurosci.* 14, 4588–4599.
- Waseem, T. V., and Fedorovich, S. V. (2010). Presynaptic glycine receptors influence plasma membrane potential and glutamate release. *Neurochem. Res.* 35, 1188–1195. doi: 10.1007/s11064-010-0174-7
- Weltzien, F., Puller, C., O'Sullivan, G. A., Paarmann, I., and Betz, H. (2012). Distribution of the glycine receptor beta-subunit in the mouse CNS as revealed by a novel monoclonal antibody. *J. Comp. Neurol.* 520, 3962–3981. doi: 10.1002/cne.23139
- Wenzel, H. J., Vacher, H., Clark, E., Trimmer, J. S., Lee, A. L., Sapolsky, R. M., et al. (2007). Structural consequences of Kcnal1 gene deletion and transfer in the mouse hippocampus. *Epilepsia* 48, 2023–2046. doi: 10.1111/j.1528-1167.2007.01189.x
- Wilson, R. I., Kunos, G., and Nicoll, R. A. (2001). Presynaptic specificity of endocannabinoid signaling in the hippocampus. *Neuron* 31, 453–462. doi: 10.1016/s0896-6273(01)00372-5
- Winkelmann, A., Maggio, N., Eller, J., Caliskan, G., Semtner, M., Häussler, U., et al. (2014). Changes in neural network homeostasis trigger neuropsychiatric symptoms. *J. Clin. Invest.* 124, 696–711. doi: 10.1172/jci.71472
- Wu, S. P., Tsai, J. J., and Gean, P. W. (1998). Frequency-dependent inhibition of neuronal activity by topiramate in rat hippocampal slices. *Br. J. Pharmacol.* 125, 826–832. doi: 10.1038/sj.bjp.0702096
- Wykes, R. C., Heeroma, J. H., Mantoan, L., Zheng, K., MacDonald, D. C., Deisseroth, K., et al. (2012). Optogenetic and potassium channel gene therapy in a rodent model of focal neocortical epilepsy. *Sci. Transl. Med.* 4:161ra152. doi: 10.1126/scitranslmed.3004190
- Young, C. C., Stegen, M., Bernard, R., Muller, M., Bischofberger, J., Veh, R. W., et al. (2009). Upregulation of inward rectifier K⁺ (Kir2) channels in dentate gyrus granule cells in temporal lobe epilepsy. *J. Physiol.* 587, 4213–4233. doi: 10.1113/jphysiol.2009.170746
- Zeilhofer, H. U., Studler, B., Arabadzisz, D., Schweizer, C., Ahmadi, S., Layh, B., et al. (2005). Glycinergic neurons expressing enhanced green fluorescent protein in bacterial artificial chromosome transgenic mice. *J. Comp. Neurol.* 482, 123–141. doi: 10.1002/cne.20349

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 22 April 2014; accepted: 29 May 2014; published online: 17 June 2014.

Citation: Meier JC, Semtner M, Winkelmann A and Wolfart J (2014) Presynaptic mechanisms of neuronal plasticity and their role in epilepsy. *Front. Cell. Neurosci.* 8:164. doi: 10.3389/fncel.2014.00164

This article was submitted to the journal *Frontiers in Cellular Neuroscience*.

Copyright © 2014 Meier, Semtner, Winkelmann and Wolfart. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.