



Applications of Mass Spectrometry in the Onset of Amyloid Fibril Formation: Focus on the Analysis of Early-Stage Oligomers

Jiaojiao Hu and Qiuling Zheng*

Key Laboratory of Drug Metabolism and Pharmacokinetics, State Key Laboratory of Natural Medicines, Department of Pharmaceutical Analysis, School of Pharmacy, China Pharmaceutical University, Nanjing, China

Amyloid fibril formation is a hallmark of diverse neurodegenerative and metabolic diseases, such as Alzheimer's disease (AD), Parkinson's disease (PD), and type 2 diabetes mellitus (T2DM). Conventional diagnosis is based on the appearance of fibrils or plaques, while neglects the role of early-stage oligomers in the disease progression. Recent studies have uncovered that it is the early-stage oligomer, rather than the mature fibril, that greatly contributes cytotoxicity. The formation of oligomers involves complicate structural conversions and it is essential to investigate their conformational changes for a better understanding of aggregation mechanism. The coexistence of soluble early-stage oligomers, intermediates, and pre-fibril species makes it difficult to be differentiated by morphological methods, and only average structural information is provided as they lack the ability of separation. Therefore, mass spectrometry (MS) becomes an alternative technique that presents new and complementary insights into the onset of amyloid fibrils. This review highlights the hotspots and important achievements by MS in the field of amyloid formation mechanism, including the direct detection and differentiation of soluble oligomers (native MS), unambiguous identification of interacted sites involved in the onset of aggregation [hydrogen/deuterium exchange (HDX) and chemical cross-linking (CX)], and conformational switch that leads to fibrilization [collision cross section (CCS) regularity by ion mobility (IM)].

Keywords: mass spectrometry, amyloid fibril, aggregation mechanism, early-stage oligomers, conformational analysis

INTRODUCTION

Amyloid fibril formation is a hallmark of numerous neurodegenerative and metabolic diseases, such as Alzheimer's disease (AD), Parkinson's disease (PD), and type 2 diabetes mellitus (T2DM) (Chiti and Dobson, 2017). *In vitro* studies suggest that most amyloid fibril formation follows nucleated polymerization mechanism, involving lag phase and growth phase before mature fibril formed. It has long been recognized that mature fibril is the leading cause of related diseases and techniques are mainly applied for diagnosis of its appearance.

With the development of electron microscopy, such as transmission electron microscope (TEM) and atomic force microscope (AFM), amyloid fibrils ranging from 6 to 130 Å in width and 1,000 to 16,000 Å in length could be detected (Sipe and Cohen, 2000). Dyeing methods, including thioflavin

OPEN ACCESS

Edited by:

Shusheng Zhang,
Linyi University, China

Reviewed by:

Lingxin Chen,
Yantai Institute of Coastal Zone
Research (CAS), China
Xia Guan,
Louisiana State University,
United States

*Correspondence:

Qiuling Zheng
qiuling_zheng@cpu.edu.cn

Specialty section:

This article was submitted to
Analytical Chemistry,
a section of the journal
Frontiers in Chemistry

Received: 22 October 2019

Accepted: 30 March 2020

Published: 05 May 2020

Citation:

Hu J and Zheng Q (2020) Applications of Mass Spectrometry in the Onset of Amyloid Fibril Formation: Focus on the Analysis of Early-Stage Oligomers. *Front. Chem.* 8:324. doi: 10.3389/fchem.2020.00324

T (ThT) and Congo red, enable the measurement of β -sheet in solution and are applied for kinetic monitoring of aggregation (Munishkina and Fink, 2007). Besides, solid-state nuclear magnetic resonance (ss-NMR) offers molecular level information of amyloid fibrils with its superiority of interaction distance determination, placing constraints on the backbone and side-chain torsion angles, as well as tertiary and quaternary contact identification (Tycko, 2011). X-ray diffraction (XRD) is another high-resolution technique which allows the investigation of molecular structure at the atomic level by obtaining folding information through reflection or transmission diffraction (Nelson et al., 2005).

Recent studies uncover that early-stage oligomer, rather than mature fibril, is the one that greatly contributes to cytotoxicity (Stefani, 2012). However, the physicochemical properties of early-stage oligomer are significantly different from that of fibrils, and no fibril formation makes it difficult to be monitored by the above-mentioned morphological techniques. Though dyeing methods support the solution-based detection, early-stage oligomers with no sufficient β -sheet still could not be sensed. What's more, the coexistence of oligomers, intermediates, and pre-fibril species during early stage yields a heterogeneous solution state and increases detection complexity. Therefore, sensitive and high spatial resolution methods are urgently required.

Mass spectrometry (MS) is a powerful analytical technique that attracts increasing attention in bioanalytical fields. Molecules are converted to gaseous ions in the ion source and separated according to their mass-to-charge ratios (m/z) in the mass analyzer. Besides measuring the m/z , structural information is provided by tandem MS through dissociation (Domon and Aebersold, 2006). Therefore, MS enables the identification of oligomerization number (the number of monomers that consist

of the oligomer, denoted as n) according to molecular weight measurement. Different conformations of the same oligomer are discovered upon gas-phase separation, which are derived from formation pathways. On the basis, MS has emerged as a highly sensitive technique for amyloid aggregation studies and presents new and complementary insights into the onset of amyloid fibrils, including oligomer characterization, aggregation mechanism investigation, and inhibitor screening. This review surveys the combination of MS with selected analytical techniques and their advanced achievements in aggregation studies (selected studies are listed in **Table 1**).

MASS SPECTROMETRY

Native MS

Electrospray ionization (ESI) is one of the soft ionization methods, which has been widely used for protein ionization. During ESI, protein molecules are transferred from solution to the gas phase with the assistance of high voltage (2.5–3.0 kV) and nebulization gas (e.g., N_2). Denaturant, such as organic solvent, is usually added to enhance the MS signal and achieve high detection sensitivity. However, it is not compatible with holding non-covalent bindings within oligomers, and it will cause dissociation during ionization. Therefore, native MS is an alternative ionization method that enables the direct detection of oligomers (Mitra, 2019). The mild ionization process and the usage of volatile ammonium-based buffer (e.g., ammonium acetate) instead of denaturant not only maintain non-covalent bindings within oligomers but also hold interactions between ligands and oligomers (Bereszczak et al., 2012). For instance, Collingwood and coworkers successfully investigated complexes formed between amyloid protein and nine metal ions by native MS. On the basis, intact oligomer or complex ions are subjected to tandem MS and resulted fragments reveal the information

TABLE 1 | Representative studies done by mass spectrometry (MS)-based techniques for early-stage oligomer analysis.

Protein/peptide	MS-based techniques	Achievements	References
A β _{1–42}	Native top-down	Interaction between A β _{1–42} and metal ion	Lermyte et al., 2019
NNQQNY VEALYL SSTNVG	ESI-IM-MS	Mechanism of peptide assembly	Bleholder et al., 2011
hIAPP A β _{1–40}	ESI-IM-MS	Mechanisms and binding patterns of small molecule inhibitors	Young et al., 2015; Hoffmann et al., 2017
PrP	HDX-MS	Amyloid fibril formation mechanism	Singh and Udgaonkar, 2013
β 2m	HDX-MS	Mechanism of Cu(II) induced amyloid formation	Borotto et al., 2017
A β _{1–42}	FPOP-MS	Conformational change of the aggregation process	Li et al., 2016
Sup 35	CX-MS	Mechanism of different folding patterns	Wong and King, 2015
α -crystallin	CX-MS	Mechanism of α A66–80 peptide induced aggregation	Kannan et al., 2013

A_{1–42}: Amyloid β _{1–42}; hIAPP: human islet amyloid polypeptide; A β _{1–40}: Amyloid β _{1–40}; PrP: prion protein; β 2m: β 2-microglobulin; ESI: electrospray ionization; IM: ion mobility; HDX: hydrogen/deuterium exchange; FPOP: fast photochemical oxidation of proteins; CX: Chemical cross-linking.

of binding/interacting sites (top-down strategy) (Tipton et al., 2011; Gregorich and Ge, 2014). Two histidine residues near the N-terminus (His 6, His 13) of A β _{1–42} were identified as major binding regions by observation of corresponding fragment ions upon electron capture dissociation (ECD), infrared multiphoton dissociation (IRMPD), and collision-induced dissociation (CID) (Lermyte et al., 2019). Native MS serves as a promising method for the detection of intact assemblies. In most cases, the corresponding oligomeric states can be established according to their *m/z* and isotopic distributions. However, there are still cases that species are overlapped and could not be well differentiated: (1) the overlapped species have significant intensity difference; (2) the overlapped species have high charge states so that the isotopic distributions could not be well resolved; or (3) a single oligomer contains different conformations. Therefore, further separation is required.


Ion Mobility MS (IM-MS)

As there are still cases that oligomers overlapped at the same *m/z* or a single oligomer has different conformations, multidimensional separation is essential for further exploration. IM is such a designed technique that enables gas-phase separation of ions based on their mobility in a buffer gas. Gas-phase ions generated from the ion source are subsequently pushed into an IM cell which is filled with a buffer gas (e.g., He). The gas-phase separation depends on their charges, sizes, and shapes, and leads to drift time difference (Kanu et al., 2008). The obtained drift time is then converted to collision cross section (CCS) value to deduce stoichiometry information. Therefore, oligomer ions sharing with the same *m/z*, but with different charges or conformations, could be separated by IM. For instance, MS detection of human islet amyloid polypeptide (hIAPP) showed a doubly charged monomer (denoted a monomer²⁺) predominantly as well as minor dimer³⁺ and trimer⁵⁺, which were identified according to their *m/z* and isotope distributions. Upon IMS separation, co-populated species sharing the same *m/z* were differentiated according to their CCS values, such as dimer³⁺ overlapped with tetramer⁶⁺. Moreover, two CCS values of monomer differed by 15% were resolved, referring to compact and expanded conformers (Young et al., 2014). CCS is applied for aggregation mechanism studies by coupling with morphological techniques. The observation of the β -sheet transformation of an aggregating peptide was first achieved by combining with AFM. In this work, based on the CCS values of peptides NNQQNY, VEALYL, SSTNVG, and YGGFL, two assembly pathways of oligomers, including isotopic growth and fibril assembly, were revealed: $\sigma = \sigma_{mon}n^{\frac{2}{3}}$ (isotopic growth, σ : the CCS values of oligomer, σ_{mon} : the CCS value of monomer, and n : the oligomeric state); $\sigma = \alpha n + \kappa$ (fibril assembly, α : describes the fibril shell area per monomer unite, and κ : means the fibril base given by the fibril diameter) (Bleiholder et al., 2011). Bowers and coworkers also correlated the conformational difference of early-stage oligomer with the structure and morphology of mature fibril by taking six mutants of sup35 (Figure 1) and five mutants of [Leu-5]-Enkephalin as examples (Do et al., 2013, 2014). IM separation connected with infrared spectroscopy (IR) analysis by Pagel and coworkers offered secondary structural analysis of gas-phase ions. This combination overcame the defect

that only average structural information could be provided by IR, while no secondary structural information provided by MS. Oligomer ions generated from peptide VEALYL and its variants were first separated by IM and subjected to secondary structural analysis, upon which a positive correlation was revealed between CCS that deviates from isotopic growth and the presence of β -sheets (Seo et al., 2017). The combination of IM-MS and IR also revealed that the hexapeptide NFGAIL followed a nucleation-dependent aggregation mechanism at neutral pH (Hoffmann et al., 2018). Besides the investigation of amyloid aggregation, Ashcroft and coworkers performed a high throughput screening of small molecule inhibitors by using IM-MS. Various modes of inhibition were identified, including interaction with different species (monomer or oligomer); through different binding patterns (specific, nonspecific, or colloidal), or different effects on aggregation (monomer binding, binding with different conformations of monomer, or disassembly of oligomers). A novel inhibitor (a nonobvious structural mimetic of chloronaphthoquinone-tryptophan) of hIAPP aggregation was discovered accordingly by screening a library of small molecules (Young et al., 2015; Hoffmann et al., 2017). Therefore, IM-MS serves as a promising method in the analysis of amyloid proteins by revealing conformational information of monomers, differentiating oligomeric states of copopulated species, as well as mechanism study of inhibitor binding (Woods et al., 2013).

Hydrogen/Deuterium Exchange MS (HDX-MS)

Upon IM, co-populated species could be separated in the gas phase, especially the differentiation of conformational intermediates during amyloid formation. However, it lacks the ability to investigate which region is involved in fibril formation. Therefore, HDX is employed regarding its capability of monitoring protein folding/unfolding dynamics and locating sites or regions that are involved in conformational changes. For HDX, proteins are incubated with D₂O and amide hydrogens of protein backbone would be replaced by the deuterium in solution. The HDX rate greatly depends on the compactness of the protein as solvent-accessible portions exchange rapidly compared with portions that are buried in the center of the structure (Koneremann et al., 2011). For amyloid protein analysis, the formed fibrils were incubated with a deuterated buffer to initiate the exchange and subsequently dissociated to monomer prior to proteolysis and MS analysis (Kheterpal and Wetzel, 2006). Upon HDX, Udgaonkar and coworkers successfully demonstrated the conformation conversion reaction from the cellular prion protein (PrP^C) to the aggregated form (PrP^{Sc}). Their work suggested that PrP^{Sc} underwent two-step conversions before fibril formation, including the formation of the β -sheet core in region 159–225, and the structure loss in the α 1 region for further monomer addition to pre-fibril. The attachment of monomer to fibril induced two-step conformational changes, including helix 1 unfolding only after helices 2 and 3 transformed into β -sheet (Singh and Udgaonkar, 2013). With the combination of covalent labeling-MS using dimethyl(2-hydroxy-5-nitrobenzyl)sulfonium bromide, Vachet and coworkers discovered that only Cu(II)



binding induced the *cis-trans* isomerization of the His31-Pro32 amide bond, and increased dynamics in the A, B, D, and E β -strands of β -2-microglobulin, which contributed to the formation of an amyloid-competent dimer interface (Borotto et al., 2017). However, the composition of aggregation converts between monomers and oligomers dynamically, resulting in the coexistence of various intermediates that cannot be captured and resolved by conventional HDX. Thus, kinetic pulse labeling HDX strategy was introduced owing to the capability of the characterization of short-lived aggregates (Carulla et al., 2010). Upon this method, temperature, agitation, and metal ion effects on the A β aggregation process were investigated by Gross and coworkers. Their work showed that the center region of

A β was first to aggregate, and soluble intermediates of lag phase were affected by Cu (II) and temperature (Zhang et al., 2013). These unique advantages make HDX-MS an indispensable weapon on amyloid protein studies, and there are a variety of works done by HDX-MS including coupled with top-down strategy (Pan et al., 2012), monitoring the oligomeric stability of insulin analogs (Nakazawa et al., 2012), aggregation kinetic study of α -synuclein (Illes-Toth et al., 2018), side-chain level oligomer-specific interaction of A β (Przygonska et al., 2018), oligomer structure of α -synuclein (Paslawski et al., 2014), and conformational dynamics of β 2m (Hodkinson et al., 2012), α -synuclein (Stephens et al., 2018), and tau (Huang et al., 2018). Similarly, there is another chemical labeled method named

as fast photochemical oxidation of proteins (FPOP), and it was based on the *in-situ* modification by hydroxyl radicals generated by the pulsed KrF laser (248 nm). With the advantage of fast sampling timescale ($\sim\mu\text{s}$) and native or near-native state detection of proteins, FPOP combined with MS can offer high special resolution structural information of proteins or protein complexes (Li et al., 2018). For example, Gross and coworkers applied FPOP-MS to monitor the conformational change and solvent accessibility at various stages in the $\text{A}\beta_{1-42}$ aggregation process, and their work showed that the middle domain of $\text{A}\beta_{1-42}$ played a major role in aggregation, whereas the N-terminus remained most of its solvent-accessibility during aggregation, and the hydrophobic C-terminus was involved to an intermediate extent (Li et al., 2016).

Chemical Cross-Linking MS (CX-MS)

HDX-MS is able to define residue or region involved in conformational changes during the aggregation process, yet, tedious sample preparation and back-exchange issue somehow limit its applications. CX appears as an alternative method to monitor protein 3D structure and protein-protein interactions. CX involves bifunctional groups containing a cross-linker that has reactivity toward specific amino acids. The two functional groups are linked by a spacer with defined length so that the side chain of amino acids within this distance could be permanently linked and detected by MS after enzymatic digestion (Sinz, 2018). As CX-MS is compatible with physiological pH, it has been widely used for structural elucidation of aggregates, including the structure of distinct tau monomers (inert and seed-competent) (Mirbaha et al., 2018), β 2-microglobulin (Hall et al., 2016), tau (Mirbaha et al., 2015), and prion protein oligomers (Onisko et al., 2005). King and coworkers have applied a photo-reactive cross-linker (p-benzoyl-L-phenylalanine) to probe amino acid proximities of Sup35 prion strains using two mutants. The detection of different cross-link products between two mutants, including intra- and inter-peptide cross-link products between amino acid residues 3/28 and 32/55, clearly indicated different folding patterns (Wong and King, 2015). Additionally, by applying the CX-MS, the interactions between hetero-protein and their effects on protein aggregation can also be obtained. For instance, homobifunctional cross-linker N-hydroxy succinimide esters BS2G and its isotope labeled form were used by Sharma and coworkers to find the specific low molecular weight peptide which induced α -crystallin fibrillization. Results showed that α A66-80 peptide bound to multiple sites in α -crystallin and was close enough to be fixed by BS2G, including the chaperone site, C-terminal extension, and subunit interaction sites. The identification of interactions between α A66-80 peptide and α -crystallin gave a better understanding of protein aggregation mechanism in cataract formation, which would have an impact on therapeutic strategies (Kannan et al., 2013). Another work was done to monitor interactions between β -amyloid with transthyretin monomers and tetramers, which was later confirmed to be biologically relevant. Upon the analysis of cross-linked $\text{A}\beta$ and TTR, the modified Lys-15, Lys-76, and Lys-9 of TTR suggested that A strand and EF helix were the major binding sites involving in $\text{A}\beta$ and TTR association, which was further confirmed to

be significant for biological study of the two proteins (Du and Murphy, 2010).

COMPLEMENTARY APPROACHES

There are also complementary approaches that are necessary to study the early stage of amyloid progression. Matrix-assisted laser desorption ionization-MS (MALDI-MS) enables direct detection of oligomers (Severinovskaya et al., 2014). According to the oligomer signals obtained from wild-type $\text{A}\beta_{1-40}$ and random sequence $\text{A}\beta_{1-40}$ variant, Yeung and coworkers were able to differentiate the peaks of $\text{A}\beta$ oligomers generated by MALDI-MS, which came from $\text{A}\beta$ complexes assembled in solution rather than random gas-phase aggregation during ionization (Wang et al., 2018). Additionally, MALDI-MS was applied to the inhibition studies. For example, with the detection and characterization of the early stage and nucleating species of $\text{A}\beta$ (25–35) aggregation process by MALDI-MS, the myricetin was confirmed to inhibit the aggregation of $\text{A}\beta$ (25–35) by delaying the transition from short oligomers to more organized soluble intermediates (Fiori et al., 2013). ^{19}F NMR is one of the alternative methods for oligomeric species detection as the chemical shift of the ^{19}F nucleus is sensitive to small changes in chemical environment and its detection would not be disturbed by background signals. Taking this advantage, Marsh et al. successfully captured multiple types of oligomers coexisted during the lag phase of aggregation by a ^{19}F -labeled $\text{A}\beta_{1-40}$. The combination with circular dichroism (CD) and AFM helped obtain more information about secondary structure and confirmed that it was a heterogeneous mixture of oligomers that existed during the lag phase (Suzuki et al., 2013). Additionally, Raman spectroscopy is another method that can provide the secondary structural information of oligomers, which is more sensitive to α -helical signal. Raman microscopy can be applied to the detection of the whole aggregation process starting with soluble species to insoluble mature fibril. Moreover, the combination of Raman microscopy with AFM further explores the changes in secondary structure and correlates it to morphology changes of aggregates. Anderson and coworkers investigated the aggregation process of α -synuclein by the Raman microscopy and AFM, and their work showed that in the aggregation progress, the content of β -sheet increases, while α -helix and disordered secondary structure decreases. The unchanged vibrational spectra during the fibrillization process suggested that a cooperative conformational change would probably contribute to the kinetic control of fibrillization (Apetri et al., 2006). What's more, various fluorescence probes are developed for the identification of prefibril species, but most of them are specially designed for certain amyloid proteins, such as the N-aryl amino naphthalene sulfonate and its analog are designed for the detection of α -synuclein aggregates (Celej et al., 2008). Recently, Carver and coworkers reported a generally applied fluorescent probe bis (triphenylphosphonium) tetraphenylethene (TPE-TPP), which would emit fluorescence upon the interaction with hydrophobic residues of aggregates. Thus, TPE-TPP dye is more sensitive than ThT dye in the detection of early-stage oligomers as they

contain a significant number of exposed hydrophobic residues while lacking sufficient β -sheets (Kumar et al., 2017).

CONCLUSION AND FUTURE POTENTIALS OF MS IN AMYLOID RESEARCH

This review describes the analysis and mechanism studies of amyloid formation by MS coupled with other analytical methodologies, which is complementary to that of biological and morphological methods. Conventional methods offer average structural information by regarding them as an entirety. Early-stage aggregation is actually a mixture of oligomers, which could be clearly differentiated by MS according to their molecular weights. In addition, oligomer ions are further separated in the gas phase by IM so that various conformations are discovered. MS coupling with HDX and CX also enables insight into conformational changes and unambiguous identification of interacted sites involved in the onset of aggregation. However,

limitations still exist: (i) current studies are done *in vitro* and results obtained may be different between simulated and real biological systems; (ii) how the mechanism obtained from the simulated system could be verified in real biological systems; and (iii) there are more factors that affect aggregation process during *in vivo* studies which increases the detection complexity.

AUTHOR CONTRIBUTIONS

JH did literature research and created the draft. QZ reviewed and revised the manuscript. All authors read and approved the manuscript.

FUNDING

This work was supported by National Natural Science Foundation of China (81703471), the Natural Science Foundation of Jiangsu Province (BK20170740).

REFERENCES

- Apetri, M. M., Maiti, N. C., Zagorski, M. G., Carey, P. R., and Anderson, V. E. (2006). Secondary structure of α -synuclein oligomers: characterization by Raman and atomic force microscopy. *J. Mol. Biol.* 355, 63–71. doi: 10.1016/j.jmb.2005.10.071
- Bereszczak, J. Z., Barbu, I. M., Tan, M., Xia, M., Jiang, X., van Duijn, E., et al. (2012). Structure, stability and dynamics of norovirus P domain derived protein complexes studied by native mass spectrometry. *J. Struct. Biol.* 177, 273–282. doi: 10.1016/j.jsb.2012.01.005
- Bleiholder, C., Dupuis, N. F., Wyttenbach, T., and Bowers, M. T. (2011). Ion mobility-mass spectrometry reveals a conformational conversion from random assembly to beta-sheet in amyloid fibril formation. *Nat. Chem.* 3, 172–177. doi: 10.1038/nchem.945
- Borotto, N. B., Zhang, Z., Dong, J., Burant, B., and Vachet, R. W. (2017). Increased β -sheet dynamics and D-E loop repositioning are necessary for Cu(II)-induced amyloid formation by β -2-microglobulin. *Biochemistry* 56, 1095–1104. doi: 10.1021/acs.biochem.6b01198
- Carulla, N., Zhou, M., Giralt, E., Robinson, C. V., and Dobson, C. M. (2010). Structure and intermolecular dynamics of aggregates populated during amyloid fibril formation studied by hydrogen/deuterium exchange. *Acc. Chem. Res.* 43, 1072–1079. doi: 10.1021/ar9002784
- Celej, M. S., Jares-Erijman, E. A., and Jovin, T. M. (2008). Fluorescent N-arylaminoanthralene sulfonate probes for amyloid aggregation of alpha-synuclein. *Biophys. J.* 94, 4867–4879. doi: 10.1529/biophysj.107.125211
- Chiti, F., and Dobson, C. M. (2017). Protein misfolding, amyloid formation, and human disease: a summary of progress over the last decade. *Annu. Rev. Biochem.* 86, 27–68. doi: 10.1146/annurev-biochem-061516-045115
- Do, T. D., Economou, N. J., Lapointe, N. E., Kincannon, W. M., Bleiholder, C., Feinstein, S. C., et al. (2013). Factors that drive peptide assembly and fibril formation: experimental and theoretical analysis of Sup35 NNQNY mutants. *J. Phys. Chem. B* 117, 8436–8446. doi: 10.1021/jp4046287
- Do, T. D., Lapointe, N. E., Sangwan, S., Teplow, D. B., Feinstein, S. C., Sawaya, M. R., et al. (2014). Factors that drive peptide assembly from native to amyloid structures: experimental and theoretical analysis of [leu-5]-enkephalin mutants. *J. Phys. Chem. B* 118, 7247–7256. doi: 10.1021/jp502473s
- Domon, B., and Aebersold, R. (2006). Review - mass spectrometry and protein analysis. *Science* 312, 212–217. doi: 10.1126/science.1124619
- Du, J., and Murphy, R. M. (2010). Characterization of the interaction of β -amyloid with transthyretin monomers and tetramers. *Biochemistry* 49, 8276–8289. doi: 10.1021/bi101280t
- Fiori, J., Naldi, M., and Andrisano, V. (2013). Mass spectrometry as an efficient tool for the characterization of amyloid beta peptide 25-35 self-assembly species in aggregation and inhibition studies. *Eur. J. Mass Spectrom* 19, 483–490. doi: 10.1255/ejms.1255
- Gregorich, Z. R., and Ge, Y. (2014). Top-down proteomics in health and disease: challenges and opportunities. *Proteomics* 14, 1195–1210. doi: 10.1002/pmic.201300432
- Hall, Z., Schmidt, C., and Politis, A. (2016). Uncovering the early assembly mechanism for amyloidogenic β 2-microglobulin using crosslinking and native mass spectrometry. *J. Biol. Chem.* 291, 4626–4637. doi: 10.1074/jbc.M115.691063
- Hodkinson, J. P., Radford, S. E., and Ashcroft, A. E. (2012). The role of conformational flexibility in beta2-microglobulin amyloid fibril formation at neutral pH. *Rapid. Commun. Mass Spectrom* 26, 1783–1792. doi: 10.1002/rcm.6282
- Hoffmann, W., Folmert, K., Moschner, J., Huang, X., Von Berlepsch, H., Koksche, B., et al. (2018). NFGAIL amyloid oligomers: the onset of beta-sheet formation and the mechanism for fibril formation. *J. Am. Chem. Soc.* 140, 244–249. doi: 10.1021/jacs.7b09510
- Hoffmann, W., von Helden, G., and Pagel, K. (2017). Ion mobility-mass spectrometry and orthogonal gas-phase techniques to study amyloid formation and inhibition. *Curr. Opin. Struct. Biol.* 46, 7–15. doi: 10.1016/j.sbi.2017.03.002
- Huang, R. Y., Iacob, R. E., Sankaranarayanan, S., Yang, L., Ahljianian, M., Tao, L., et al. (2018). Probing conformational dynamics of tau protein by hydrogen/deuterium exchange mass spectrometry. *J. Am. Soc. Mass Spectrom* 29, 174–182. doi: 10.1007/s13361-017-1815-8
- Illes-Toth, E., Rempel, D. L., and Gross, M. L. (2018). Pulsed hydrogen-deuterium exchange illuminates the aggregation kinetics of α -synuclein, the causative agent for parkinson's disease. *ACS Chem. Neurosci.* 9, 1469–1476. doi: 10.1021/acschemneuro.8b00052
- Kannan, R., Santhoshkumar, P., Mooney, B. P., and Sharma, K. K. (2013). The α A66-80 peptide interacts with soluble α -crystallin and induces its aggregation and precipitation: a contribution to age-related cataract formation. *Biochemistry* 52, 3638–3650. doi: 10.1021/bi301662w
- Kanu, A. B., Dwivedi, P., Tam, M., Matz, L., and Hill, H. H. Jr. (2008). Ion mobility-mass spectrometry. *J. Mass Spectrom* 43, 1–22. doi: 10.1002/jms.1383
- Khetarpal, I., and Wetzel, R. (2006). Hydrogen/deuterium exchange mass spectrometry—a window into amyloid structure. *Acc. Chem. Res.* 39, 584–593. doi: 10.1021/ar050057w
- Konermann, L., Pan, J., and Liu, Y. H. (2011). Hydrogen exchange mass spectrometry for studying protein structure and dynamics. *Chem. Soc. Rev.* 40, 1224–1234. doi: 10.1039/C0CS00113A
- Kumar, M., Hong, Y., Thorn, D. C., Ecroyd, H., and Carver, J. A. (2017). Monitoring early-stage protein aggregation by an

- aggregation-induced emission fluorogen. *Anal. Chem.* 89, 9322–9329. doi: 10.1021/acs.analchem.7b02090
- Lermyte, F., Everett, J., Lam, Y. P. Y., Wootton, C. A., Brooks, J., Barrow, M. P., et al. (2019). Metal ion binding to the amyloid beta monomer studied by native top-down ftrc mass spectrometry. *J. Am. Soc. Mass Spectrom* 30, 2123–2134. doi: 10.1007/s13361-019-02283-7
- Li, K. S., Rempel, D. L., and Gross, M. L. (2016). Conformational-sensitive fast photochemical oxidation of proteins and mass spectrometry characterize amyloid beta 1-42 aggregation. *J. Am. Chem. Soc.* 138, 12090–12098. doi: 10.1021/jacs.6b07543
- Li, K. S., Shi, L., and Gross, M. L. (2018). Mass spectrometry-based fast photochemical oxidation of proteins (fpop) for higher order structure characterization. *Acc. Chem. Res.* 51, 736–744. doi: 10.1021/acs.accounts.7b00593
- Mirbaha, H., Chen, D., Morazova, O. A., Ruff, K. M., Sharma, A. M., Liu, X., et al. (2018). Inert and seed-competent tau monomers suggest structural origins of aggregation. *eLife* 7:e36584. doi: 10.7554/eLife.36584
- Mirbaha, H., Holmes, B. B., Sanders, D. W., Bieschke, J., and Diamond, M. I. (2015). Tau Trimers are the minimal propagation unit spontaneously internalized to seed intracellular aggregation. *J. Biol. Chem.* 290, 14893–14903. doi: 10.1074/jbc.M115.652693
- Mitra, G. (2019). Application of native mass spectrometry in studying intrinsically disordered proteins: a special focus on neurodegenerative diseases. *Biochim. Biophys. Acta Proteins Proteomics* 1867:140260. doi: 10.1016/j.bbapap.2019.07.013
- Munishkina, L. A., and Fink, A. L. (2007). Fluorescence as a method to reveal structures and membrane-interactions of amyloidogenic proteins. *Biochim. Biophys. Acta, Biomembr.* 1768, 1862–1885. doi: 10.1016/j.bbamem.2007.03.015
- Nakazawa, S., Hashii, N., Harazono, A., and Kawasaki, N. (2012). Analysis of oligomeric stability of insulin analogs using hydrogen/deuterium exchange mass spectrometry. *Anal. Biochem.* 420, 61–67. doi: 10.1016/j.ab.2011.09.002
- Nelson, R., Sawaya, M. R., Balbirnie, M., Madsen, A. O., Riekel, C., Grothe, R., et al. (2005). Structure of the cross-beta spine of amyloid-like fibrils. *Nature* 435, 773–778. doi: 10.1038/nature03680
- Onisko, B., Guitian Fernandez, E., Louro Freire, M., Schwarz, A., Baier, M., Camina, F., et al. (2005). Probing PrPsc structure using chemical cross-linking and mass spectrometry: evidence of the proximity of Gly90 amino termini in the PrP 27-30 aggregate. *Biochemistry* 44, 10100–10109. doi: 10.1021/bi0501582
- Pan, J., Han, J., Borchers, C. H., and Konermann, L. (2012). Structure and dynamics of small soluble A β (1-40) oligomers studied by top-down hydrogen exchange mass spectrometry. *Biochemistry* 51, 3694–3703. doi: 10.1021/bi3002049
- Paslawski, W., Mysling, S., Thomsen, K., Jorgensen, T. J., and Otzen, D. E. (2014). Co-existence of two different alpha-synuclein oligomers with different core structures determined by hydrogen/deuterium exchange mass spectrometry. *Angew. Chem. Int. Ed.* 53, 7560–7563. doi: 10.1002/anie.201400491
- Przygonska, K., Poznanski, J., Mistarz, U. H., Rand, K. D., and Dadlez, M. (2018). Side-chain moieties from the N-terminal region of A β are involved in an oligomer stabilizing network of interactions. *PLoS ONE* 13:e0201761. doi: 10.1371/journal.pone.0201761
- Seo, J., Hoffmann, W., Warnke, S., Huang, X., Gewinner, S., Schollkopf, W., et al. (2017). An infrared spectroscopy approach to follow beta-sheet formation in peptide amyloid assemblies. *Nat. Chem.* 9, 39–44. doi: 10.1038/nchem.2615
- Severinovskaya, O. V., Kovalska, V. B., Losytsky, M. Y., Cherepanov, V. V., Subramaniam, V., and Yarmoluk, S. M. (2014). Application of MALDI-TOF mass spectrometry for study on fibrillar and oligomeric aggregates of alpha-synuclein. *Biopolym. Cell* 30, 190–196. doi: 10.7124/bc.000895
- Singh, J., and Udgaonkar, J. B. (2013). Dissection of conformational conversion events during prion amyloid fibril formation using hydrogen exchange and mass spectrometry. *J. Mol. Biol.* 425, 3510–3521. doi: 10.1016/j.jmb.2013.06.009
- Sinz, A. (2018). Cross-linking/mass spectrometry for studying protein structures and protein-protein interactions: where are we now and where should we go from here? *Angew. Chem. Int. Ed.* 57, 6390–6396. doi: 10.1002/anie.201709559
- Sipe, J. D., and Cohen, A. S. (2000). Review: history of the amyloid fibril. *J. Struct. Biol.* 130, 88–98. doi: 10.1006/jsbi.2000.4221
- Stefani, M. (2012). Structural features and cytotoxicity of amyloid oligomers: implications in Alzheimer's disease and other diseases with amyloid deposits. *Prog. Neurobiol.* 99, 226–245. doi: 10.1016/j.pneurobio.2012.03.002
- Stephens, A. D., Nespovitaya, N., Zacharopoulou, M., Kaminski, C. F., Phillips, J. J., and Schierle, G. S. K. (2018). Different structural conformers of monomeric alpha-synuclein identified after lyophilizing and freezing. *Anal. Chem.* 90, 6975–6983. doi: 10.1021/acs.analchem.8b01264
- Suzuki, Y., Brender, J. R., Soper, M. T., Krishnamoorthy, J., Zhou, Y., Ruotolo, B. T., et al. (2013). Resolution of oligomeric species during the aggregation of A β 1-40 using 19F NMR. *Biochemistry* 52, 1903–1912. doi: 10.1021/bi400027y
- Tipton, J. D., Tran, J. C., Catherman, A. D., Ahlf, D. R., Durbin, K. R., and Kelleher, N. L. (2011). Analysis of intact protein isoforms by mass spectrometry. *J. Biol. Chem.* 286, 25451–25458. doi: 10.1074/jbc.R111.239442
- Tycko, R. (2011). Solid-state NMR studies of amyloid fibril structure. *Annu. Rev. Phys. Chem.* 62, 279–299. doi: 10.1146/annurev-physchem-032210-103539
- Wang, J. S., Whitehead, S. N., and Yeung, K. K. (2018). Detection of Amyloid Beta (Abeta) oligomeric composition using matrix-assisted laser desorption ionization mass spectrometry (MALDI MS). *J. Am. Soc. Mass Spectrom* 29, 786–795. doi: 10.1007/s13361-018-1896-z
- Wong, S.-H., and King, C.-Y. (2015). Amino acid proximities in two sup35 prion strains revealed by chemical cross-linking. *J. Biol. Chem.* 290, 25062–25071. doi: 10.1074/jbc.M115.676379
- Woods, L. A., Radford, S. E., and Ashcroft, A. E. (2013). Advances in ion mobility spectrometry-mass spectrometry reveal key insights into amyloid assembly. *Biochim. Biophys. Acta Proteins Proteomics* 1834, 1257–1268. doi: 10.1016/j.bbapap.2012.10.002
- Young, L. M., Cao, P., Raleigh, D. P., Ashcroft, A. E., and Radford, S. E. (2014). Ion mobility spectrometry-mass spectrometry defines the oligomeric intermediates in amylin amyloid formation and the mode of action of inhibitors. *J. Am. Chem. Soc.* 136, 660–670. doi: 10.1021/ja406831n
- Young, L. M., Saunders, J. C., Mahood, R. A., Revill, C. H., Foster, R. J., Tu, L. H., et al. (2015). Screening and classifying small-molecule inhibitors of amyloid formation using ion mobility spectrometry-mass spectrometry. *Nat. Chem.* 7, 73–81. doi: 10.1038/nchem.2129
- Zhang, Y., Rempel, D. L., Zhang, J., Sharma, A. K., Mirica, L. M., and Gross, M. L. (2013). Pulsed hydrogen-deuterium exchange mass spectrometry probes conformational changes in amyloid beta (A beta) peptide aggregation. *Proc. Natl. Acad. Sci. U.S.A.* 110, 14604–14609. doi: 10.1073/pnas.1309175110

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Hu and Zheng. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.