



## OPEN ACCESS

## APPROVED BY

Frontiers in Ecology And Evolution  
Editorial Office,  
Frontiers Media SA, Switzerland

## \*CORRESPONDENCE

Florian P. Schiestl  
[florian.schiestl@systbot.uzh.ch](mailto:florian.schiestl@systbot.uzh.ch)

RECEIVED 09 October 2023

ACCEPTED 16 October 2023

PUBLISHED 27 October 2023

## CITATION

Huber FK and Schiestl FP (2023) Corrigendum: Scent releasing silicone septa: a versatile method for bioassays with volatiles. *Front. Ecol. Evol.* 11:1310147. doi: 10.3389/fevo.2023.1310147

## COPYRIGHT

© 2023 Huber and Schiestl. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Corrigendum: Scent releasing silicone septa: a versatile method for bioassays with volatiles

Franz K. Huber and Florian P. Schiestl\*

Institute of Systematic and Evolutionary Botany, University of Zurich, Zurich, Switzerland

## KEYWORDS

floral scent, pollination, volatile, chemical ecology, bioassay, herbivore

## A corrigendum on

Scent releasing silicone septa: a versatile method for bioassays with volatiles

by Huber FK and Schiestl FP (2022). *Front. Ecol. Evol.* 10:958982.  
doi: 10.3389/fevo.2022.958982

In the published article, there was an error in the unit used for soaking concentration: we wrote “mg ml<sup>-1</sup>” where we meant “μg ml<sup>-1</sup>”. The text has been updated in the following locations.

### Materials and methods, paragraph 1:

For soaking septa, each compound was prepared in four concentrations: 1, 10, 100, and 1,000 mg ml<sup>-1</sup>, in dichloromethane (highperformance liquid chromatography (HPLC) grade) as solvent.

has been updated to read:

For soaking septa, each compound was prepared in four concentrations: 1, 10, 100, and 1,000 μg ml<sup>-1</sup>, in dichloromethane (highperformance liquid chromatography (HPLC) grade) as solvent.

### Results, paragraph 1:

For example, for benzaldehyde, coefficients of variation in scent emission across time points were 0.31, 0.08, 0.10, and 0.16 for soaking concentrations 1, 10, 100, and 1,000 mg ml<sup>-1</sup>, respectively (Supplementary Table S2).


has been updated to read:

For example, for benzaldehyde, coefficients of variation in scent emission across time points were 0.31, 0.08, 0.10, and 0.16 for soaking concentrations 1, 10, 100, and 1,000 μg ml<sup>-1</sup>, respectively (Supplementary Table S2).

The same error occurred in Figure 2. The corrected figure and its caption appear below.

The same error occurred in Supplementary Tables S1, S2, S3, S4. The corrected tables are now published with the original article.

The authors apologize for these errors and state that they do not change the scientific conclusions of the article in any way. The original article has been updated.



**FIGURE 2**  
Volatile emission rate from silicone septa for benzaldehyde. Five trials were done for each soaking concentration and time points after drying.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.