



Corrigendum: ACE Gene Variants Rise the Risk of Severe COVID-19 in Patients With Hypertension, Dyslipidemia or Diabetes: A Spanish Pilot Study

María Íñiguez¹, Patricia Pérez-Matute¹, Pablo Viloslada-Blanco¹, Emma Recio-Fernandez¹, Diana Ezquierro-Pérez², Jorge Alba², M. Lourdes Ferreira-Laso³ and José A. Oteo^{1,2*}

¹ Infectious Diseases, Microbiota and Metabolism Unit, Infectious Diseases Department, Center for Biomedical Research of La Rioja (CIBIR), Logroño, Spain, ² Infectious Diseases Department, Hospital Universitario San Pedro, Logroño, Spain,

³ Department of Anesthesiology and Postoperative Care, Hospital Universitario San Pedro, Logroño, Spain

OPEN ACCESS

Edited and reviewed by:

Hamad Ali,
Kuwait University, Kuwait

*Correspondence:

José A. Oteo
jaoteo@riojasalud.es

Specialty section:

This article was submitted to
Clinical Diabetes,
a section of the journal
Frontiers in Endocrinology

Received: 06 September 2021

Accepted: 08 September 2021

Published: 20 September 2021

Citation:

Íñiguez M, Pérez-Matute P,
Viloslada-Blanco P,

Recio-Fernandez E, Ezquierro-Pérez D,

Alba J, Ferreira-Laso ML and Oteo JA

(2021) Corrigendum: ACE Gene

Variants Rise the Risk of Severe

COVID-19 in Patients With

Hypertension, Dyslipidemia or

Diabetes: A Spanish Pilot Study.

Front. Endocrinol. 12:771445.

doi: 10.3389/fendo.2021.771445

Keywords: COVID-19, angiotensin converting enzyme, polymorphisms, hypertension, dyslipidemia, diabetes

A Corrigendum on

ACE Gene Variants Rise the Risk of Severe COVID-19 in Patients With Hypertension, Dyslipidemia or Diabetes: A Spanish Pilot Study

By Íñiguez M, Pérez-Matute P, Viloslada-Blanco P, Recio-Fernandez E, Ezquierro-Pérez D, Alba J, Ferreira-Laso ML and Oteo JA (2021). *Front. Endocrinol.* 12:688071. doi: 10.3389/fendo.2021.688071.

ERROR IN FIGURE/TABLE

In the original article, there was a mistake in **Table 2** as published. Alleles of rs4343 polymorphism were incorrectly written. The corrected **Table 2** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Íñiguez, Pérez-Matute, Viloslada-Blanco, Recio-Fernandez, Ezquierro-Pérez, Alba, Ferreira-Laso and Oteo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

TABLE 2 | Genotype frequencies of rs4341 and rs4343 polymorphisms in COVID-19 patients according to its hypertensive, dyslipidemic or diabetic status.

rs4341				
Hypertension	G/G	G/C	C/C	P-value
Hospital ward	6 (20%)	16 (53.3%)	8 (26.7%)	0.031*
ICU	6 (50%)	5 (41.7%)	1 (8.3%)	0.03\$
Dyslipidemia				
	G/G	G/C	C/C	P-value
Hospital ward	2 (15.4%)	6 (46.1%)	5 (38.5%)	0.043*
ICU	4 (57.1%)	3 (42.9%)	0 (0%)	
Diabetes Mellitus				
	G/G	G/C	C/C	P-value
Hospital ward	1 (8.3%)	7 (58.3%)	4 (33.3%)	0.033*
ICU	3 (100%)	0 (0%)	0 (0%)	0.009\$

rs4343				
Hypertension	G/G	G/A	A/A	P-value
Hospital ward	15 (16.6%)	17 (56.7%)	8 (26.7%)	0.03%
ICU	5 (41.7%)	6 (50%)	1 (8.3%)	
Dyslipidemia				
	G/G	G/A	A/A	P-value
Hospital ward	2 (15.4%)	6 (46.1%)	5 (38.5%)	0.043#
ICU	4 (57.1%)	3 (42.9%)	0 (0%)	
Diabetes Mellitus				
	G/G	G/A	A/A	P-value
Hospital ward	1 (8.3%)	7 (58.3%)	4 (33.3%)	0.033#
ICU	3 (100%)	0 (0%)	0 (0%)	0.009%

Data are presented as number of patients and percentages.

*p-value for group difference between GG and GC and CC.

\$p-value for group difference between GG+GC and CC.

#p-value for group difference between GG+GA and AA.

%p-value for group difference between GG+GA and AA.

Significant differences appear in bold case.

All data were adjusted for age, sex, BMI and number of comorbidities.