



The Impact of Hormonal Contraceptive Use on Serotonergic Neurotransmission and Antidepressant Treatment Response: Results From the NeuroPharm 1 Study

Søren Vinther Larsen^{1,2}, Brice Ozenne^{1,3}, Kristin Köhler-Forsberg^{1,2,4}, Asbjørn Seenithamby Poulsen¹, Vibeke Høyrup Dam¹, Claus Svarer¹, Gitte Moos Knudsen^{1,2}, Martin Balslev Jørgensen⁴ and Vibe Gedso Frokjaer^{1,2,4*}

OPEN ACCESS

Edited by:

Birgit Derntl,
University of Tübingen, Germany

Reviewed by:

Neven Henigsberg,
University of Zagreb, Croatia
Jeffrey Miller,
Columbia University, United States

*Correspondence:

Vibe Gedso Frokjaer
vibe.frokjaer@nru.dk

Specialty section:

This article was submitted to
Neuroendocrine Science,
a section of the journal
Frontiers in Endocrinology

Received: 21 October 2021

Accepted: 31 January 2022

Published: 11 March 2022

Citation:

Larsen SV, Ozenne B, Köhler-Forsberg K, Poulsen AS, Dam VH, Svarer C, Knudsen GM, Jørgensen MB and Frokjaer VG (2022) The Impact of Hormonal Contraceptive Use on Serotonergic Neurotransmission and Antidepressant Treatment Response: Results From the NeuroPharm 1 Study. *Front. Endocrinol.* 13:799675. doi: 10.3389/fendo.2022.799675

¹ Neurobiology Research Unit, Copenhagen University Hospital Rigshospitalet, Copenhagen, Denmark, ² Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark, ³ Department of Public Health, Section of Biostatistics, University of Copenhagen, Copenhagen, Denmark, ⁴ Psychiatric Center Copenhagen, Mental Health Services in the Capital Region of Denmark, Copenhagen, Denmark

Background: Hormonal contraceptive (HC) use has been associated with an increased risk of developing a depressive episode. This might be related to HC's effect on the serotonergic brain system as suggested by recent cross-sectional data from our group, which show that healthy oral contraceptive (OC) users relative to non-users have lower cerebral serotonin 4 receptor (5-HT₄R) levels. Here, we determine if cerebral 5-HT₄R binding differs between HC non-users, OC users, and hormonal intrauterine device (HIUD) users among women with an untreated depressive episode. Also, we test if antidepressant drug treatment response and its association with pre-treatment 5-HT₄R binding depends on HC status.

Methods: [¹¹C]-SB207145 Positron Emission Tomography imaging data from the NeuroPharm-NP1 Study (NCT02869035) were available from 59 depressed premenopausal women, of which 26 used OCs and 10 used HIUDs. The participants were treated with escitalopram. Treatment response was measured as the relative change in the Hamilton Depression Rating Scale 6 items (rΔHAMD₆) from baseline to week eight. Latent variable models were used to evaluate the association between global 5-HT₄R binding and OC and HIUD use as well as rΔHAMD₆.

Results: We found no evidence of a difference in global 5-HT₄R binding between depressed HC users and non-users (p≥0.51). A significant crossover interaction (p=0.02) was observed between non-users and OC users in the association between baseline global 5-HT₄R binding and week eight rΔHAMD₆; OC users had 3-4% lower binding compared to non-users for every 10% percent less improvement in HAMD₆. Within the groups, we observed a trend towards a positive association in non-users

($p_{\text{adj}}=0.10$) and a negative association in OC users ($p_{\text{adj}}=0.07$). We found no strong evidence of a difference in treatment response between the groups ($p=0.13$).

Conclusions: We found no difference in 5-HT4R binding between HC users vs. non-users in depressed women, however, it seemed that 5-HT4R settings differed qualitatively in their relation to antidepressant drug treatment response between OC users and non-users. From this we speculate that depressed OC users constitutes a special serotonin subtype of depression, which might have implications for antidepressant drug treatment response.

Keywords: hormonal contraception, oral contraception, hormonal intrauterine device, [^{11}C]SB207145, serotonin, major depressive disorder, serotonin 4 receptor, sex steroid hormones

INTRODUCTION

Hormonal contraception (HC) is used by millions of women worldwide to avoid pregnancies as well as for other indications such as dysmenorrhea, menorrhagia, and acne (1, 2). In spite of HC being used for more than 60 years it is still debated whether HC use causes mood deterioration and development of depressive episodes (3). Recent findings from large epidemiological studies have suggested that starting on HC is associated with an increased risk of being diagnosed with Major Depressive Disorder (MDD), being prescribed antidepressants (4, 5), and even attempting or completing suicide (6–8). Notably, the risk is higher when HC is initiated in adolescence (9, 10) and moreover, HC use in adolescence is associated with lasting vulnerability for MDD in adulthood (11), which may relate to HC being introduced in a critical stage of brain development (12). Even though HCs are widely used, we still have limited understanding on how HC affects the brain and which implications it may have for mental health (13).

HCs exist in different types in terms of hormonal content and route of administration such as oral contraceptives (OCs), hormonal intrauterine devices (HIUDs), vaginal rings, injections and subdermal implants (14). OCs and HIUDs are the types most widely used (1). HCs contain synthetic female sex hormones either in form of a progestin alone (HIUDs and progesterone-only pills) or in a combination with an estrogen (combined OCs), most frequently ethinylestradiol (14). The synthetic steroids in combined OCs and in high-dose progesterone-only pills act by suppressing the hypothalamic-pituitary-gonadal hormonal axis resulting in suppression of the endogenous hormone production, disrupted follicular maturation, and inhibition of ovulation (15). HIUDs and low-dose progesterone-only pills on the other hand only inhibit ovulation in 60–85% of the time and even as low as 15% of the time after one year of HIUD use. Instead, their primary mechanism of action is induction of local inflammation in the cervical mucous to prevent access of sperm and by thinning of the uterus lining to prevent implantation of a fertilized egg (16, 17). Further, the synthetic steroids in HC induce an increase in the sex hormone binding globulin level further lowering the bioavailable fraction of sex hormones (18). These profound changes in the sex hormone milieu may shape brain biology both in terms of brain structure (19) and function (20). Recently,

we have shown that OC use appears to affect the brain's serotonin signaling system (21), which is a key system for maintaining mental health and is involved in the pathophysiology of MDD and the treatment thereof (22). We used the molecular imaging technique ([^{11}C]SB207145-radioligand positron emission tomography (PET)) to quantify the postsynaptic serotonin 4 receptors (5-HT4R). The 5-HT4Rs are sensitive to chronic synaptic serotonin manipulation such that they are inversely correlated to serotonergic tone (23–25) making it an interesting tool to study MDD pathophysiology. In a population of healthy women, we found that women using OCs had 9–12% lower 5-HT4R level globally in the brain compared to non-users (21). In comparison, our group also found 7–8% lower 5-HT4R global binding in unmedicated depressed individuals compared to healthy controls, and intriguingly, this gap was only evident in those who remitted after eight weeks of antidepressant treatment with a selective reuptake inhibitor (SSRI), indicating that those responding to the treatment may have a serotonergic subtype of MDD (26). The SSRI works by targeting the serotonin system, which induces an additional downregulation of 5-HT4R levels in neostriatum, a phenomenon confirmed in both a depressed and healthy cohort (23, 26). From this, we speculate whether the mechanisms causing the lower 5-HT4R binding in OC users and depressed individuals are similar and if it has any implications for the SSRI treatment response in women with a depressive episode.

We sought to investigate this by determining 1) if the 5-HT4R binding differs between depressed women who do not use HC vs. those using OCs and HIUDs, 2) if antidepressant drug treatment response is affected by HC use, 3) if an association between pre-treatment 5-HT4R brain binding and antidepressant drug treatment response depends on HC status, and 4) if neostriatal 5-HT4R levels are sensitive to SSRI treatment in HC users. We hypothesize that if the effects from OC and MDD are not additive, we will not be able to detect a difference in baseline 5-HT4R binding. However, when we account for treatment response, and thus may account for the serotonergic subtype of MDD, we will be able to detect a negative main effect of OC use on 5-HT4R binding. In line with that, we hypothesize that an association between baseline binding and treatment response will depend on OC use, such that an association is found in non-users but not in OC users, as they may constitute a more homogenous group in terms of 5-HT4R downregulation. We further

hypothesize that the effect of HIUD use on 5-HT4R binding will be in the same direction as of OC use, but with smaller effect size, due to the smaller degree of synthetic steroid exposure and larger degree of preserved ovulations (i.e., preserved hormonal cycle).

METHODS

We included patient data from the NeuroPharm-NP1 clinical trial (27), which aimed to predict MDD treatment outcome with the use of potential biomarkers including 5-HT4R PET brain scans. The study was approved by the Committees on Health Research Ethics in the Capital Region of Denmark (H-15017713), the Danish Data Protection Agency (04711/RH-2016-163), and the Danish Medicines Agency (NeuroPharm-NP1, EudraCT-number 2016-001626-34) and was pre-registered on clinicaltrials.gov (NCT02869035). A detailed description of the study design and the study elements can be found in the clinical trial protocol (27). The present study used information on HC use and baseline 5-HT4R PET imaging data together with clinical outcome measures after eight weeks of antidepressant drug treatment. It represents initially unplanned analyses motivated by the observed difference in 5-HT4R binding in healthy women who use OC vs. those who do not use OC (21). The treatment started at a daily dose of 10 mg of the SSRI, escitalopram, and was increased to a daily dose of maximally 20 mg, depending on treatment response and adverse reactions. In case of non-response at week four, patients were switched to duloxetine, a serotonin-norepinephrine reuptake inhibitor (SNRI). Compliance was confirmed by serum drug levels after eight weeks of treatment. A subset of the study population was rescanned after eight weeks of treatment to map the change in 5-HT4R level. Allocation to rescan happened continuously until allotted rescans were completed.

Study Population

The study population consists of all the depressed premenopausal women (defined as < 50 years of age) from the NeuroPharm-NP1 study with available baseline 5-HT4R PET data and information on HC use. One patient was excluded from the analyses on 5-HT4R imaging data due to an interrupted PET scan (n=59 for the first analysis). Eight patients dropped out during follow-up, six OC users and two non-users (n=52 for the second and n=51 for the third analysis). Twenty-six of the remaining women were rescanned at follow-up; 15 were non-users, six were OC users and five were HIUD users (n=26 for the fourth analysis). The women were suffering from an unmedicated moderate to severe (> 17 points on the Hamilton Depression Rating Scale 17 items (HAMD₁₇)) depressive episode confirmed by a psychiatrist at inclusion. The current depressive episode was single or recurrent and did not involve acute severe suicidal ideation or psychosis. They had no prior or present history of other major psychiatric disorders confirmed by use of the diagnostic tool, the Mini International Neuropsychiatric Interview version 6.0 (28), or any other severe somatic illness confirmed by a basic somatic screening. Level of education was

calculated as the number of completed school years (7-12 years) summed with a Likert score indexing highest completed or commenced degree ranging from one (no vocational degree) to five (>four years of higher learning at university level). Information about relationship status was acquired *via* interview. Blood tests to assess plasma estradiol and progesterone levels were taken at the baseline PET scan as part of standard biochemical screening.

Hormonal Contraception

Information about HC use was acquired *via* face-to-face interview at the time of the PET scan and *via* a written questionnaire which included a question on the specific HC type. The HC type was divided into HIUD and OC. OCs included different generations of combined OCs (n=20) and progestogen-only pills (n=6), as specified in **Table S1**.

Clinical Outcome Measure

The clinical outcome measure was the relative change in Hamilton Depression Rating Scale 6 items (rΔHAMD₆) in percentage from baseline to week eight. In the larger NeuroPharm-NP1 study, the categorical treatment response categories, remitters vs. non-responders, were used as the primary clinical outcome and rΔHAMD₆ was included as the secondary outcome (27). For this study, which only includes the women, we only applied rΔHAMD₆ to avoid underpowered statistics.

Imaging

PET acquisition and quantification is detailed in the trial protocol (27) and briefly summarized here: A high-resolution research tomography Siemens PET scanner (CTI/Siemens, Knoxville, TN, USA) (256 × 256 × 207 voxels; 1.22 × 1.22 × 1.22 mm) was used for acquiring PET scans. Each scan was obtained from a 120 min dynamic PET acquisition immediately after a 20 second intravenous bolus injection of the [¹¹C] SB207145 tracer ligand. Motion correction was performed using the AIR 5.2.5 software (29). High-resolution structural T1-weighted magnetic resonance images were acquired on a Siemens 3-Tesla Prisma scanner with a 64-channel head coil. The images were segmented into cerebrospinal fluid, white- and gray matter and were co-registered with PET images using the Statistical Parametric Mapping software (SPM8, The Wellcome Centre for Human Neuroimaging, UCL, London, UK). Regions were automatically delineated from the MR image *via* the user-independent algorithm in the Pvelab software (30). Regions of interest (ROIs) were neostriatum, hippocampus and neocortex, as these were lower in 5-HT4R binding in the depressed cohort in the NeuroPharm-NP1 Study (26), and as these regions represent low, intermediate, and high expression levels of brain 5-HT4Rs (31) and lastly, as these regions have previously shown lower 5-HT4R levels in OC users (21). As previous studies only found an effect of SSRI treatment on 5-HT4R binding in neostriatum, we only used this region to test whether this was also true in the HC users. Regional non-displaceable binding potentials (BP_{ND}) were quantified using the simplified reference tissue model with cerebellum as reference tissue (32).

Statistics

For the descriptive statistics, we compared between the HC non-users, the OC users, and the HIUD users, and *p*-values were computed with Fisher's exact test for the categorical outcomes, Kruskal-Wallis Rank Sum Test for the continuous and the discrete variables and when relevant, Dunn's test was used for *post hoc* analyses where multiple comparisons were corrected for with the Bonferroni-Holm method. For plasma progesterone and estradiol, we used Gehan test (33) due to censored values (18 estradiol- and 35 progesterone samples were below the detection limit of 0.09 nM and 0.6 nM, respectively).

To evaluate if HC use was associated with global 5-HT4R brain binding (aim 1), we used a linear latent variable model (LVM) where the effects of HIUD- and OC use were mediated through a shared latent variable (hereafter phrased as the global LV) across brain regions of low (neocortex), intermediate (hippocampus) and high (neostriatum) 5-HT4R binding (31). By introducing the global LV, we were able to account for the large inter-correlation in 5-HT4R binding between brain regions. The binding in each region was adjusted independently, i.e., not through the global LV, for age, BMI, 5-HTTLPR genotype (LALA or non-LALA), and injected [11C]SB207145 mass per kg bodyweight as these are considered to influence 5-HT4R PET measurements (34–36). 5-HT4R BP_{ND} values were log-transformed prior to modeling and the regional estimates are expressed as a percent difference in 5-HT4R binding in the OC users *vs.* the non-users and in the HIUD users *vs.* the non-users. To test if the treatment response differed between the groups (aim 2), we used the Kruskal-Wallis Rank Sum Test. For reporting the effect size, we used the Mann-Whitney parameter (37), which gives the probability that a randomly selected individual from one group had a worse treatment response compared to one from another group. An estimate of 0.5 indicates no difference in treatment response. To evaluate if an association between baseline 5-HT4R BP_{ND} and week eight rΔHAMD₆ depends on OC/HIUD use (aim 3), we extended our LVM by including an interaction term between rΔHAMD₆ and the HC group to be mediated through the global LV. The estimates related to rΔHAMD₆ are expressed as the percent change in 5-HT4R BP_{ND} per 10% change in HAMD₆ (r₁₀ΔHAMD₆). To evaluate the effect of eight weeks of SSRI/SNRI treatment on neostriatal 5-HT4R binding across groups (aim 4), we used linear regression adjusting for the difference in injected [11C]SB207145 mass per kg bodyweight between baseline and week eight scans. Due to the smaller re-test samples within the OC and the HIUD group, the *p*-values were also computed with a permutation test with 10,000 permutations.

Diagnostic tools were used to assess the adequacy of the models' assumptions. We used chi-squared tests to evaluate specification of the covariance structure for each LVM with chi-square < 0.05 indicating suboptimal specification [section 6.2.4 in (38)]. Missing data in the analyses regarding treatment response were handled using complete case analysis. Two-sided statistical tests were used and *p* < 0.05 was considered statistically significant. Confidence intervals and *p*-values computed for the regional effects were adjusted for three comparisons using the

single-step Dunnett's procedure (39). Statistical analyses were performed in R (40) and LVMs were estimated with the lava package (38).

RESULTS

Study Population Profile

The clinical profile and PET parameters of the study population at baseline are shown in **Table 1**. The non-users, the HIUD users, and the OC users were similar in terms of proportion suffering from first depressive episode and in clinician rated depressive symptoms (HAMD_{6/17}) at baseline. The depressive symptoms scored within a moderate-to-severe depressive episode as restricted by the inclusion criteria. The educational level tended to differ between the groups (*p*=0.06) such that the HC users tended to have lower education. As expected, the OC users had lower plasma estradiol levels (median [Q1, Q3]: 0.09 nM [0.09, 0.18]) compared to the non-users (median [Q1, Q3]: 0.30 nM [0.16, 0.64]), *p*_{adj}=0.0002, and the HIUD users (median [Q1, Q3]: 0.23 [0.17, 0.26]), *p*_{adj}=0.01. The OC users had lower plasma progesterone levels but only compared to the non-users (*p*_{adj}=0.04). The HIUD users did not seem to differ from the non-users in plasma hormone levels.

Hormonal Contraception and Serotonin 4 Receptor Binding

In support of the LVM structure, we found significant region-specific bindings loading onto the global LV for all regions (*p*<0.001). From the chi-squared test we found no evidence for a lack of fit (*p*_{adj}>0.79), so no additional covariance was added to the model. There was no evidence for an association between 5-HT4R binding and OC use (*p*=0.51) or HIUD use (*p*=0.73) (**Figure 1**). The corresponding non-significant regional estimates for OC use varied between -2.8% and -1.9%, and for HIUD use between -2.1% and -1.4%. No gross deviation from the normality assumption was observed.

Antidepressant Treatment Response

Week eight depression profiles are shown in **Table 2**. We found no evidence of a difference in proportion of patients who switched to duloxetine after week four (*p*=0.34) or in week eight treatment response (rΔHAMD₆) between the groups (*p*=0.13). The estimated probability of finding a smaller reduction in HAMD₆ in an OC and an HIUD user was 66% (95% CI: [0.48; 0.80], *p*=0.08) and 68% (95% CI: [0.46; 0.83], *p*=0.11), respectively, compared to in a non-user (**Figure 2**), and notably, this was at a non-significant level as the confidence intervals included 0.5 (*p*-values were not corrected for multiple comparisons). The OC users seemed to require higher doses of escitalopram compared to the non-users (*p*_{adj}=0.04) and the HIUD users (*p*_{adj}=0.01); 77.8% of the OC users were treated with the highest recommended daily dose of 20 mg compared to 35.0% and 22.2% of the non-users and the HIUD users, respectively. Correspondingly, the OC users also had a higher plasma concentration of escitalopram at week eight compared to the HIUD users (*p*_{adj}=0.03), and at a trend level compared to the non-users (*p*_{adj}=0.06). The dropouts' depression profiles, in

TABLE 1 | Clinical profile and PET parameters at baseline.

	Non-users (n=23)	HIUD users (n=11)	OC users (n=26)		
	n (%)	n (%)	n (%)	p-value^a	n
First MDD episode	10 (43.5%)	5 (45.5%)	8 (30.8%)	0.61	60
5-HTTLPR L _A L _A genotype	8 (34.8%)	1 (9.1%)	7 (26.9%)	0.32	60
In relationship	7 (30.4%)	6 (54.5%)	9 (34.6%)	0.47	60
	Median [Q1, Q3]	Median [Q1, Q3]	Median [Q1, Q3]	p-value^b	n
Age	26.2 [22.9, 30.3]	23.7 [22.5, 24.4]	23.3 [21.6, 25.7]	0.25	60
BMI [kg/m ²]	23.9 [20.5, 31.2]	22.1 [19.2, 23.9]	21.8 [20.0, 24.7]	0.31	60
Educational level	16.0 [16.0, 17.0]	15.0 [12.5, 16.5]	14.5 [13.0, 16.0]	0.06	54
HAMD ₆	12.0 [11.5, 13.5]	13.0 [12.0, 13.5]	12.5 [12.0, 13.0]	0.87	60
HAMD ₁₇	22.0 [21.0, 24.0]	25.0 [21.0, 27.0]	23.0 [20.2, 25.0]	0.58	60
P-estradiol [nM]	0.30 [0.16, 0.64]	0.23 [0.17, 0.26]	0.09 [0.09, 0.18]	0.20 ^c	60
				0.0002 ^d	
				0.01 ^e	
P-progesterone [nM]	0.90 [0.60, 15.00]	0.60 [0.60, 3.85]	0.60 [0.60, 0.67]	0.59 ^f	60
				0.04 ^g	
				0.59 ^c	
Injected dose [MBq]	600.2 [573.6, 604.2]	591.3 [526.8, 602.4]	602.5 [589.8, 605.2]	0.21	59
Injected tracer mass/kg [μg/kg]	8.6x10 ⁻³ [6.5x10 ⁻³ , 1.3x10 ⁻²]	8.6x10 ⁻³ [6.0x10 ⁻³ , 1.7x10 ⁻²]	9.2x10 ⁻³ [6.4x10 ⁻³ , 1.2x10 ⁻²]	0.91	59
Cerebellum, area under curve [kBq/ml]	10380.8 [9105.6, 12660.3]	10755.5 [9055.6, 11657.2]	11118.2 [9058.6, 13413.4]	0.47	59

HIUD, hormonal intrauterine device; OC, oral contraceptive; MDD, Major Depressive Disorder; BMI, Body Mass Index; HAMD_{6/17}, Hamilton Depression Rating Scale 17 or 6 items; MDI, Major Depressive Inventory; ^ap-values are computed with Fisher's exact test. ^bp-values are computed with Kruskal-Wallis Rank Sum Test except for P-estradiol and P-progesterone, for which Gehan test was used due to censored values and they were corrected for three comparisons with the Bonferroni-Holm method. ^cNon-users vs. HIUD users. ^dNon-users vs. OC users. ^eHIUD users vs. OC users.




FIGURE 1 | The estimated latent variable model for the effect of oral contraceptive (OC) and hormonal intrauterine device (HIUD) use on baseline 5-HT4R BP_{ND} in women with an untreated depressive episode. γ is the effect on the global latent variable interpreted as global (log-transformed) 5-HT4R BP_{ND} effects. λ is the loading on each region. The boxes beneath the loadings indicate the percentage difference in 5-HT4R binding for each brain region in OC- and HIUD users compared to non-users. Regional bindings were adjusted for Age, BMI, 5-HTTLPR genotype and injected tracer mass per kg bodyweight (not shown). P-values and confidence intervals are adjusted for 3 comparisons by use of the Dunnett's test.

terms of baseline HAMD_{6/17} as well as relative change in HAMD₆ in percentage from baseline to week one, two, four and eight and reason for dropout, are presented in **Table S2**.

Pre-treatment Serotonin 4 Receptor Binding and Antidepressant Treatment Response

Again, the LVM was supported by the region-specific bindings loading onto the global LV ($p < 0.001$). The chi-squared test showed no evidence of misspecification of the covariance structure ($p_{adj} > 0.41$). The estimated associations between

baseline global 5-HT4R binding (represented as the global LV) and treatment responses across each group are shown in **Figure 3** with a trend towards a positive slope in the non-users (1.33×10^{-2} ($p = 0.10$)), a trend towards a negative slope in the OC users (-1.27×10^{-2} ($p = 0.07$)), and a non-significant negative slope in the HIUD users (-3.34×10^{-4} ($p = 0.75$)). The corresponding correlation coefficients are 0.39 for the non-users, -0.46 for the OC users, and -0.12 for the HIUD users. In the OC users, the slope differed by -2.59×10^{-2} ($p = 0.02$) from the non-users. In **Figure 4**, the LVM is summarized in three layers (A, B and C). The first (A) shows the trend towards a positive association

TABLE 2 | Clinical depression profile at week eight.

	Non-users (n=21)	HIUD users (n=11)	OC users (n=20)	p-value ^α	n
Switchers to duloxetine	n (%) 1 (4.8%)	n (%) 2 (18.2)	n (%) 2 (10)	0.34	52
rΔHAMD ₆ [%]	Median [Q1, Q3] -63.6 [-78.6, -46.2]	Median [Q1, Q3] -50.0 [-64.1, -23.7]	Median [Q1, Q3] -45.3 [-64.9, -15.1]	p-value ^β 0.13	52
P-escitalopram [nM]	68.1 [43.3, 102.8]	42.5 [36.0, 108.5]	105.7 [74.1, 139.8]	0.40 ^γ	47
				0.06 ^δ	
				0.03 ^ε	
Escitalopram dose:	n (%)	n (%)	n (%)	p-value ^α	n
5 mg	0 (0.0)	1 (11.1)	0 (0.0)	0.26 ^γ	47
10 mg	2 (10.0)	2 (22.2)	1 (5.6)	0.04 ^δ	
15 mg	11 (55.0)	4 (44.4)	3 (16.7)	0.01 ^ε	
20 mg	7 (35.0)	2 (22.2)	14 (77.8)		

HIUD, hormonal intrauterine device; OC, oral contraceptive; rΔHAMD₆, relative change in Hamilton Depression Rating Scale 6 items from baseline. ^αp-value is computed with Fischer's exact test. ^βp-values are computed with Kruskal-Wallis Rank Sum Test and, for post hoc analyses, with Dunn's test corrected for three comparisons with the Bonferroni-Holm method. ^γNon-users vs. HIUD users. ^δNon-users vs. OC users. ^εHIUD users vs. OC users.




FIGURE 2 | Antidepressant drug treatment response at week eight across hormonal contraceptive user status. Difference in relative change in Hamilton Depression Rating Scale 6 items (HAMD₆) from baseline between the HC non-users, the hormonal intrauterine device (HIUD) users and the oral contraceptive (OC) users. A larger negative change corresponds to a better improvement of depressive symptoms. P-values are computed with Mann-Whitney tests with no correction for multiple comparisons.




FIGURE 3 | Estimated global 5-HT4R latent variable summarizing the association between baseline global 5-HT4R brain binding and week eight antidepressant drug treatment response across the groups. The slopes are the association between baseline (un-medicated) 5-HT4R binding and change in Hamilton Depression Rating Scale 6 items (HAMD₆) in each group, respectively. A negative change in HAMD₆ mirrors an improvement of depressive symptoms. The slopes are 1.33×10^{-2} ($p_{\text{adj}}=0.10$) for the non-users, -3.34×10^{-3} ($p_{\text{adj}}=0.75$) for the hormonal intrauterine device (HIUD) users, and -1.27×10^{-2} ($p_{\text{adj}}=0.07$) for the oral contraceptive (OC) users.

between the 5-HT4R binding and $r_{10}\Delta\text{HAMD}_6$ mediated through the global LV in the non-users ($p=0.10$). The second layer (B) shows how this association differs in the HC users from the non-users. The regional estimates show that per 10% less improvement in HAMD₆, the OC users had 2.6–4.0% ($p_{\text{adj}} \leq 0.03$) lower binding at baseline compared to the non-users. The third layer (C) shows the estimated effect of HC use on 5-HT4R binding if no treatment response was achieved at week eight, i.e.,

the estimated HC effect when accounting for the confounding factor of a suggested serotonergic subtype of MDD. However, as we have only few observations at this part of the response scale, these estimates are calculated based on the model and less on actual observed differences. At the regional level, the percent effect of OC use varied between -24.2% and -16.3% ($p_{\text{adj}} \leq 0.01$). No gross deviation from the normality assumption was observed for any of the LVs.




FIGURE 4 | The estimated latent variable model for the association between baseline 5-HT4R binding and week eight antidepressant treatment response as a function of hormonal contraceptive status shown in three layers (A–C). (A) The association between baseline 5-HT4R binding and treatment response in non-users. (B) The difference in the association between baseline 5-HT4R binding and treatment response between the non-users and the oral contraceptive (OC) and the hormonal intrauterine device (HIUD) users, respectively. (C) The estimated effect of OC and HIUD use on 5-HT4R binding when no change in Hamilton Depression Rating Scale 6 items (HAMD6). γ is the estimates of the association with the global latent variable interpreted as global (log-transformed) 5-HT4R BP_{ND} estimates. λ is the loading on each region. The boxes beneath the loadings indicate the percentage difference in 5-HT4R binding for each brain region. Regional bindings were adjusted for Age, BMI, 5-HTTLPR genotype and injected tracer mass per kg bodyweight (not shown). P-values and confidence intervals are adjusted for 3 comparisons. $r_{10}\Delta\text{HAMD}_6$: 10% relative change in Hamilton Depression Rating Scale 6 items.


Antidepressants' Effect on Serotonin 4 Receptor Binding

In **Figure 5**, we show the effects of eight weeks of SSRI/SNRI treatment on the 5-HT4R BP_{ND} across the groups. A reduction in 5-HT4R BP_{ND} appeared to be present in all groups with an estimated reduction in the non-users of 7.0% (95% CI: [-12.1; -1.6], $p=0.02$). Due to the small sample sizes in the HIUD- and the OC group, the effect sizes are less reliable, and the p-values are also computed after 10,000 permutations. It revealed that, if the null hypothesis is true (i.e., no change in 5-HT4R BP_{ND} after

eight weeks of SSRI treatment), the observed lower 5-HT4R BP_{ND} would be found by chance in 9–10% of the cases, hence, a reduction in 5-HT4R binding is only found at a trend level in the HIUD and the OC users.

DISCUSSION

In this study, we found no evidence of a difference in 5-HT4R binding in unmedicated depressed women who use OCs or



HIUDs relative to non-users. We found a trend towards an association between baseline global 5-HT4R BP_{ND} and treatment response in the non-users and the OC users and these were in opposite directions with a significant crossover interaction, such that the OC users had 3–4% lower binding compared to the non-users for every 10% percent less improvement in $HAMD_6$ at week eight. Based on our model, the main effect of OC use on regional 5-HT4R binding at zero improvement in depressive symptoms was estimated to be between -16 and -24%. In addition, we observed no strong evidence of a difference in treatment response between the groups, although, the estimated probability of finding a smaller reduction in $HAMD_6$ in an OC and an HIUD user was numerically higher compared to in a non-user. Last, we found that as in the non-users, both the HIUD users and the OC users seemed to react to SSRI treatment in terms of neostriatal 5-HT4R downregulation, however, only at a trend level.

Hormonal Contraceptive Use and Brain Serotonin

In a healthy population we found OC use to be associated with 9–12% lower global 5-HT4R binding (21), which was not seen in this depressed cohort, however, the depressed cohort is also 7–8% lower in binding compared to a healthy population (26). Hence, the OC effect on 5-HT4R binding may be obscured by the effects of the MDD as expected, and thus we see no indication of the effects of OC and MDD being additive. Comparable to OC use, we found a numerically lower, but still statistically non-significant effect of HIUD use. In the previous study on the healthy population (21), we were underpowered to investigate an effect of HIUD use. Thus, brain signatures of HIUD use remain

to be studied since HIUD use is also associated with an increased risk of developing MDD (4, 5).

Whereas we found no difference in 5-HT4R binding level at baseline between the groups, we observed a crossover interaction between the OC users and the non-users in the association between baseline 5-HT4R BP_{ND} and treatment response, but notably, the association within each group was only borderline significant, which may be due to power issues. As we were not expecting to see a negative trend in the association in the OC users, this highlights the question whether the 5-HT4R setting may differ qualitatively, i.e., whether the mechanisms behind a lower 5-HT4R level differ. As shown in a mixed male-female cohort of MDD patients, only those responding to treatment had a lower baseline (unmedicated) 5-HT4R level compared to healthy controls, so it has been speculated whether they have a serotonergic subtype of MDD, in which the lower binding is due to a compensatory upregulation of the serotonergic tonus as an attempt to maintain euthymia (26). In contrast, we speculate whether all depressed OC users belong to a special serotonergic subgroup with varying degree of serotonergic involvement. We suggest that the mechanism affecting the 5-HT4R level in OC users may be driven by a hormone-dependent decrease in gene expression levels due to a suppressed hormone state in OC users, which align with previous observations in healthy OC users (21). This also aligns with preclinical research showing that estradiol supports serotonergic signaling in terms of increased capacity of serotonin synthesis and reuptake (41, 42), reduced capacity of serotonin degradation (41), increased neural firing (43) and increased serotonin receptor availability (44), which mainly happen *via* gene expression through estrogen receptor alpha and -beta (45). Also notably, estradiol increases 5-HT4R mRNA expression in the anterior pituitary cells in rats (46). We speculate that the more the 5-HT4R expression level as well as other parts of the serotonin signaling are compromised by the suppressed hormone state in OC users, the worse the treatment response, and in contrast, only the non-users with a serotonergic component show a better response to an SSRI targeting this system. If this theory is applied on our results and we compare the 5-HT4R levels at $r\Delta HAMD_6$ equal to 0% (i.e., the results from the third layer of the LVM), we compare the effect of OC use with a population with a non-serotonergic subtype of MDD, which reveals an OC effect equal to about -20%. However, since this result is based on our model with rather few observations in this range of the treatment response, it should be interpreted with some caution.

The association between baseline binding and treatment response did not differ between the HIUD users and the non-users, however, the estimate was in the same direction as in the OC users. It is possible that, if replicated in larger sample sizes, this reflects an effect of HIUD use on serotonin brain signaling.

Hormonal Contraceptive Use and Antidepressant Treatment Response

The estimated probability of finding a smaller reduction in $HAMD_6$ in an OC and an HIUD user compared to a non-user was 66% and 68%, respectively, but the confidence intervals included 50%, so we have insufficient evidence to conclude that

differences in the treatment response exist. This might be a question of power as the confidence intervals were wide, e.g., the OC users *vs.* the non-users 95% confidence spanned from 48% to 80%. Larger samples would help narrow the confidence intervals to help us discriminate between differences in treatment response *vs.* no or very small differences. The median $r\Delta\text{HAMD}_6$ was, at the non-significant level, about 18 percent point and 14 percent point lower in the OC and the HIUD users, respectively, relative to the non-users. This is equivalent to about two points based on median HAMD_6 scores at baseline. Treatment response could be affected by educational level (47) as the non-users tended to have higher educational level, however, this could also be related to the HC users not having started their final degree, yet, as the median age in the HC users was about 23 *vs.* 26 in non-users. Adjusting for educational level did not have any notable impact on the result (not shown). Limited and insufficient clinical evidence exists addressing whether HC use affects SSRI treatment response. The STAR*D study reported a trend towards better remission rates to citalopram treatment in 226 HC users compared to 670 HC non-users, however, this effect was not robust to adjustment for potential confounders (48). Another study, based on 17 double-blind, placebo-controlled clinical trials with 1698 women, found no difference in treatment response between the OC users and the non-users measured by change in HAMD_{17} (49). It has been highlighted that the studies are limited by lacking information on the types of HCs used, for not including or missing response rates across the trials, and for including studies with variable SSRI dosages (50). The latter could be relevant as higher dosage was required in the OC users, which could make up for less response in OC users relative to non-users. Correspondingly, serum levels of escitalopram seemed to be higher in the OC users, which might also be a result of an OC-induced inhibition of the cytochrome P450 (CYP) hepatic enzymes, CYP2C19 and CYP3A4 (51), which makes up about 70% of the escitalopram metabolism (52).

Preclinical evidence points towards a modulatory role of estradiol on antidepressant drug effects in rats (53–56). One study found that exogenous administration of 17 beta-estradiol as well as ethinylestradiol in ovariectomized female rats facilitates the antidepressant-like effects of fluoxetine and desipramine in a forced swim test (53). In support of that, other studies found that deficiency of brain estrogen levels attenuated sertraline- (54) and duloxetine (55)-induced antidepressive behaviors in mice and it correlated with serotonin turnover in hippocampus and prefrontal cortex (54, 55). In contrast, another study found that acute administration of estrogen with progesterone blocked the antidepressant-like effect of desipramine, but only after acute, not chronic administration (56).

Suppressed levels of endogenous estradiol thus might affect SSRI treatment response and possibly this involves the 5-HT4R. A difference in the 5-HT4R setting could potentially affect the treatment effects as it plays a key role in serotonergic neuronal firing from the dorsal raphe nuclei with projections to most parts of the brain (57). Thus, a lower 5-HT4R agonism capacity could make the serotonergic brain function less adaptable to

environmental demands. This aligns with rodent data showing that 5-HT4R partial agonists reduce stress-induced antidepressive behavior (58), have fast acting antidepressant-like effects (59), and that the antidepressant-like effects of SSRIs depends on 5-HT4R activation (60). Whereas our data highlight that treatment response depends on the initial 5-HT4R setting in an OC-dependent manner, we also see a trend towards the expected SSRI-effect on the neostriatal 5-HT4R level after eight weeks of treatment, supporting that the 5-HT4R setting in HC users is sensitive to SSRIs.

CONCLUSION AND PERSPECTIVES

In a depressed cohort, we found no evidence of a difference in the 5-HT4R level between the non-users and the HC users, but the associations between 5-HT4R binding and treatment response were in opposite directions in the OC users and the non-users; the lower the global 5-HT4R binding before treatment start the worse the treatment response in the OC users in contrast to better treatment response in the non-users. We found no strong evidence of a difference in treatment response between the groups. From this study, we speculate that depressed OC users constitutes a special serotonin subtype of MDD which might have implications for SSRI treatment. We will in a planned longitudinal study address the causal relationship between OC use and lower 5-HT4R binding and future and properly-designed studies must address if OC use affects treatment response in depressed women and whether they perhaps could benefit from tapering off OC use. Also, the effects of HIUD use on brain serotonin signatures remains to be investigated in future studies.

Methodological Considerations

When interpreting the results from this study, some limitations should be considered; 1) the sample size is small, so interpretation should be made with caution, especially with regard to HIUD use and when comparing treatment responses, 2) the OC group was pooled from users of different types of OCs (combined OCs and progesterone-only pills) with different hormone content, which may blur the results, 3) we lacked information on previous HC use among the non-users and information on starting day and pill-free days in the HC users, which could have helped us understand the potential role of HC use in their depressive episode, 4) we had 8 dropouts during follow-up which were mainly OC users. However, based on their last response rate (**Table S2**), they seem more or less equally distributed between those showing some degree of response and those who do not, so it appears not to introduce a selection bias.

DATA AVAILABILITY STATEMENT

The data analyzed in this study is subject to the following licenses/restrictions: For accessing the dataset, a Cimb

database application is required. Requests to access these datasets should be directed to the corresponding author (vibe.frokjaer@nru.dk) or access can be requested through the procedures outlined here: <https://cimbi.dk/index.php/documents/category/3-cimbi-database>.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by The Committees on Health Research Ethics in the Capital Region of Denmark (H-15017713). The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

SL and VF contributed to study conceptualization, data collection, analysis and interpretation of results, participated in drafting the manuscript, and approved the final version prior to submission. BO contributed to analysis and interpretation of data, revised the manuscript critically for important intellectual content, and approved the final version prior to submission. KK-F, AP, VD, and CS contributed to data collection and the acquisition of data, revised the manuscript critically for important intellectual content, and approved the final version prior to submission. GK and MJ contributed to study conceptualization and interpretation of data, revised the manuscript critically for important intellectual content, and approved the final version prior to submission. All authors contributed to the article and approved the submitted version.

REFERENCES

- United Nations. *Contraceptive Use by Method 2019*. UN: Data Booklet (2019). Available at: <https://www.un-ilibrary.org/content/books/9789210046527>.
- Brynhildsen J. Combined Hormonal Contraceptives: Prescribing Patterns, Compliance, and Benefits Versus Risks. *Ther Adv Drug Saf* (2014) 5(5):201–13. doi: 10.1177/2042098614548857
- Schaffir J, Worly BL, Gur TL. Combined Hormonal Contraception and its Effects on Mood: A Critical Review. *Eur J Contracept Reprod Health Care* (2016) 21(5):347–55. doi: 10.1080/13625187.2016.1217327
- Skovlund CW, Mørch LS, Kessing LV, Lidegaard O. Association of Hormonal Contraception With Depression. *JAMA Psychiatry* (2016) 73(11):1154–62. doi: 10.1001/jamapsychiatry.2016.2387
- Zettermark S, Vicente RP, Merlo J. Hormonal Contraception Increases the Risk of Psychotropic Drug Use in Adolescent Girls But Not in Adults: A Pharmacoepidemiological Study on 800 000 Swedish Women. *PLoS One* (2018) 13(3):e0194773. doi: 10.1371/journal.pone.0194773
- Skovlund CW, Mørch LS, Kessing LV, Lange T, Lidegaard J. Association of Hormonal Contraception With Suicide Attempts and Suicides. *Am J Psychiatry* (2018) 175(4):336–42. doi: 10.1176/appi.ajp.2017.17060616
- Jung SJ, Cho SMJ, Kim HC. Association of Oral Contraceptive Use With Suicidal Behavior Among Representative Korean Population: Results From Korea National Health and Nutrition Examination Survey (2007–2016). *J Affect Disord* (2019) 243:8–15. doi: 10.1016/j.jad.2018.09.004
- Edwards AC, Lönn SL, Crump C, Mościcki EK, Sundquist J, Kendler KS, et al. Oral Contraceptive Use and Risk of Suicidal Behavior Among Young Women. *Psychol Med* (2020) 1–8. doi: 10.1017/S0033291720003475
- Wirhn AB, Foldemo A, Josefsson A, Lindberg M. Use of Hormonal Contraceptives in Relation to Antidepressant Therapy: A Nationwide Population-Based Study. *Eur J Contracept Reprod Health Care* (2010) 15(1):41–7. doi: 10.3109/13625181003587004
- De Wit AE, Booij SH, Giltay EJ, Joffe H, Schoevers RA, Oldehinkel AJ. Association of Use of Oral Contraceptives With Depressive Symptoms Among Adolescents and Young Women. *JAMA Psychiatry* (2020) 77(1):52–9. doi: 10.1001/jamapsychiatry.2019.2838
- Anderl C, Li G, Chen FS. Oral Contraceptive Use in Adolescence Predicts Lasting Vulnerability to Depression in Adulthood. *J Child Psychol Psychiatry Allied Discip* (2020) 61(2):148–56. doi: 10.1111/jcpp.13115
- Cahill L. How Does Hormonal Contraception Affect the Developing Human Adolescent Brain? *Curr Opin Behav Sci* (2018) 23:131–5. doi: 10.1016/j.cobeha.2018.06.015
- Lewis CA, Kimmig ACS, Zsido RG, Jank A, Derntl B, Sacher J. Effects of Hormonal Contraceptives on Mood: A Focus on Emotion Recognition and Reactivity, Reward Processing, and Stress Response. *Curr Psychiatry Rep* (2019) 21(11):115. doi: 10.1007/s11920-019-1095-z
- Britton LE, Alspaugh A, Greene MZ, McLemore MR. CE: An Evidence-Based Update on Contraception. *Am J Nurs* (2020) 120(2):22–33. doi: 10.1097/01.NAJ.0000654304.29632.a7
- Rivera R, Yacobson I, Grimes D. The Mechanism of Action of Hormonal Contraceptives and Intrauterine Contraceptive Devices. *Am J Obstet Gynecol* (1999) 181(5 Pt 1):1263–9. doi: 10.1016/S0002-9378(99)70120-1
- Kailasam C, Cahill D. Review of the Safety, Efficacy and Patient Acceptability of the Levonorgestrel-Releasing Intrauterine System. *Patient Prefer Adherence* (2008) 2:293–301. doi: 10.2147/PPA.S3464

FUNDING

The study was funded by Independent Research Fund Denmark (grantID: 0134-00278B & DFF-6120-00038), the Research Fund of the Mental Health Services - Capital Region of Denmark, the Innovation Fund Denmark, H. Lundbeck A/S (grantID: 5189-00087A), the Research Council of Rigshospitalet, the Augustinus Foundation (grantID: 16-0058), and Savværksejer Jeppe Juhl og hustru Ovita Juhls Mindelegat. None of the funders had any influence on study design, data collection, analysis, and result interpretation.

ACKNOWLEDGMENTS

We gratefully acknowledge investigators from the NeuroPharm-NP1 study as well as collaborating general practitioners and the Center for Referral and Diagnostics, Mental Health Services, Capital Region of Copenhagen for helping with recruitment of patients. We wish to thank Lone Freyr, Anders Jørgensen, Peter Jensen, Dorthe Givard, Birgit Tang, Arafat Nasser, Ida Marie Brandt, Elizabeth Landman, Anna Maria Florescu, Agnete Dyssegaard, Jacob Madsen, Sune Keller, Svitlana Olsen, and Gerda Thomsen and technical staff at the Dept. of Nuclear Medicine, Rigshospitalet for superb technical assistance.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fendo.2022.799675/full#supplementary-material>

17. Duijkers IJM, Heger-Mahn D, Drouin D, Colli E, Skouby S. Maintenance of Ovulation Inhibition With a New Progestogen-Only Pill Containing Drospirenone After Scheduled 24-H Delays in Pill Intake. *Contraception* (2016) 93(4):303–9. doi: 10.1016/j.contraception.2015.12.007
18. Panzer C, Wise S, Fantini G, Kang D, Munarriz R, Guay A, et al. Impact of Oral Contraceptives on Sex Hormone-Binding Globulin and Androgen Levels: A Retrospective Study in Women With Sexual Dysfunction. *J Sex Med* (2006) 3(1):104–13. doi: 10.1111/j.1743-6109.2005.00198.x
19. Rehbein E, Hornung J, Sundström Poromaa I, Derntl B. Shaping of the Female Human Brain by Sex Hormones: A Review. *Neuroendocrinology* (2021) 111(3):183–206. doi: 10.1159/000507083
20. Toffoletto S, Lanzemberger R, Gingnell M, Sundström-Poromaa I, Comasco E. Emotional and Cognitive Functional Imaging of Estrogen and Progesterone Effects in the Female Human Brain: A Systematic Review. *Psychoneuroendocrinology* (2014) 50:28–52. doi: 10.1016/j.psyneuen.2014.07.025
21. Larsen SV, Köhler-Forsberg K, Dam VH, Poulsen AS, Svarer C, Jensen PS, et al. Oral Contraceptives and the Serotonin 4 Receptor: A Molecular Brain Imaging Study in Healthy Women. *Acta Psychiatr Scand* (2020) 142(4):294–306. doi: 10.1111/acps.13211
22. Lin SH, Lee LT, Yang YK. Serotonin and Mental Disorders: A Concise Review on Molecular Neuroimaging Evidence. *Clin Psychopharmacol Neurosci* (2014) 12(3):196–202. doi: 10.9758/cpn.2014.12.3.196
23. Haahr ME, Fisher PM, Jensen CG, Frøkjær VG, McMahon B, Madsen K, et al. Central 5-HT₄ Receptor Binding as Biomarker of Serotonergic Tonus in Humans: A [¹¹C]SB207145 PET Study. *Mol Psychiatry* (2014) 19(4):427–32. doi: 10.1038/mp.2013.147
24. Licht CL, Marcussen AB, Wegener G, Overstreet DH, Aznar S, Knudsen GM. The Brain 5-HT₄ Receptor Binding is Down-Regulated in the Flinders Sensitive Line Depression Model and in Response to Paroxetine Administration. *J Neurochem* (2009) 109(5):1363–74. doi: 10.1111/j.1471-4159.2009.06050.x
25. Vidal R, Valdizán EM, Mostany R, Pazos A, Castro E. Long-Term Treatment With Fluoxetine Induces Desensitization of 5-HT₄ Receptor-Dependent Signalling and Functionality in Rat Brain. *J Neurochem* (2009) 110(3):28–52. doi: 10.1111/j.1471-4159.2009.06210.x
26. Köhler-Forsberg K, Ozenne B, Landman EB, Larsen SV, Poulsen AS, Dam VH, et al. Evidence for a Serotonergic Subtype of Major Depressive Disorder: A NeuroPharm-1 Study. *medRxiv* (2021) 2021.06.17.21258740. doi: 10.1101/2021.06.17.21258740
27. Köhler-Forsberg K, Jørgensen A, Dam VH, Stenbæk DS, Fisher PM, Ip CT, et al. Predicting Treatment Outcome in Major Depressive Disorder Using Serotonin 4 Receptor PET Brain Imaging, Functional MRI, Cognitive-, EEG-Based, and Peripheral Biomarkers: A NeuroPharm Open Label Clinical Trial Protocol. *Front Psychiatry* (2020) 11:641. doi: 10.3389/fpsy.2020.00641
28. Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): The Development and Validation of a Structured Diagnostic Psychiatric Interview for DSM-IV and ICD-10. *J Clin Psychiatry* (1998) 59(SUPPL. 20):22–3. doi: 10.1037/t18597-000
29. Woods RP, Cherry SR, Mazziotta JC. Rapid Automated Algorithm for Aligning and Reslicing Pet Images. *J Comput Assist Tomogr* (1992) 16(4):620–33. doi: 10.1097/00004728-199207000-00024
30. Svarer C, Madsen K, Hasselbalch SG, Pinborg LH, Haugbøl S, Frøkjær VG, et al. MR-Based Automatic Delineation of Volumes of Interest in Human Brain PET Images Using Probability Maps. *Neuroimage* (2005) 24(4):969–79. doi: 10.1016/j.neuroimage.2004.10.017
31. Beliveau V, Ganz M, Feng L, Ozenne B, Højgaard L, Fisher PM, et al. A High-Resolution In Vivo Atlas of the Human Brain's Serotonin System. *J Neurosci* (2017) 37(1):120–8. doi: 10.1523/JNEUROSCI.2830-16.2016
32. Marner L, Gillings N, Comley RA, Baaré WFC, Rabiner EA, Wilson AA, et al. Kinetic Modeling Of [¹¹C]-SB207145 Binding to 5-HT₄ Receptors in the Human Brain In Vivo. *J Nucl Med* (2009) 50(6):900–8. doi: 10.2967/jnumed.108.058552
33. Gehan EA. A Generalized Two-Sample Wilcoxon Test for Doubly Censored Data. *Biometrika* (1965) 52(3):650–3. doi: 10.2307/2333721
34. Madsen K, Haahr MT, Marner L, Keller SH, Baaré WF, Svarer C, et al. Age and Sex Effects on 5-HT₄ Receptors in the Human Brain: A [¹¹C]SB207145 PET Study. *J Cereb Blood Flow Metab* (2011) 31(6):1475–81. doi: 10.1038/jcbfm.2011.11
35. Haahr ME, Rasmussen PM, Madsen K, Marner L, Ratner C, Gillings N, et al. Obesity Is Associated With High Serotonin 4 Receptor Availability in the Brain Reward Circuitry. *Neuroimage* (2012) 61(4):884–8. doi: 10.1016/j.neuroimage.2012.03.050
36. Fisher PM, Holst KK, Adamsen D, Klein AB, Frøkjær VG, Jensen PS, et al. BDNF Val66met and 5-HTTLPR Polymorphisms Predict a Human In Vivo Marker for Brain Serotonin Levels. *Hum Brain Mapp* (2015) 36(1):313–23. doi: 10.1002/hbm.22630
37. Fay MP, Malinovsky Y. Confidence Intervals of the Mann-Whitney Parameter That are Compatible With the Wilcoxon-Mann-Whitney Test. *Stat Med* (2018) 37(27):3991–4006. doi: 10.1002/sim.7890
38. Holst KK, Budtz-Jørgensen E. Linear Latent Variable Models: The Lava-Package. *Comput Stat* (2013) 28(4):1385–452. doi: 10.1007/s00180-012-0344-y
39. Dmitrienko A, D'Agostino R. Traditional Multiplicity Adjustment Methods in Clinical Trials. *Stat Med* (2013) 32(29):5172–218. doi: 10.1002/sim.5990
40. Core Computing Team R. R: A Language and Environment for Statistical Computing. *R Found Stat Comput Vienna Austria* (2017) 0. Available at: <https://www.r-project.org>
41. Smith LJ, Henderson JA, Abell CW, Bethea CL. Effects of Ovarian Steroids and Raloxifene on Proteins That Synthesize, Transport, and Degrade Serotonin in the Raphe Region of Macaques. *Neuropsychopharmacology* (2004) 29(11):2035–45. doi: 10.1038/sj.npp.1300510
42. Sánchez MG, Morissette M, Di Paolo T. Oestradiol Modulation of Serotonin Reuptake Transporter and Serotonin Metabolism in the Brain of Monkeys. *J Neuroendocrinol* (2013) 25(6):560–9. doi: 10.1111/jne.12034
43. Robichaud M, Debonnel G. Oestrogen and Testosterone Modulate the Firing Activity of Dorsal Raphe Nucleus Serotonergic Neurons in Both Male and Female Rats. *J Neuroendocrinol* (2005) 17(3):179–85. doi: 10.1111/j.1365-2826.2005.01292.x
44. Kugaya A, Epperson CN, Zoghbi S, Van Dyck CH, Hou Y, Fujita M, et al. Increase in Prefrontal Cortex Serotonin_{2a} Receptors Following Estrogen Treatment in Postmenopausal Women. *Am J Psychiatry* (2003) 160(8):1522–4. doi: 10.1176/appi.ajp.160.8.1522
45. Hernández-Hernández OT, Martínez-Mota L, Herrera-Pérez JJ, Jiménez-Rubio G. Role of Estradiol in the Expression of Genes Involved in Serotonin Neurotransmission: Implications for Female Depression. *Curr Neuropharmacol* (2018) 17(5):459–71. doi: 10.2174/1570159X16666180628165107
46. Papageorgiou A, Deneff C. Estradiol Induces Expression of 5-Hydroxytryptamine (5-HT)₄, 5-HT₅, and 5-HT₆ Receptor Messenger Ribonucleic Acid in Rat Anterior Pituitary Cell Aggregates and Allows Prolactin Release via the 5-HT₄ Receptor. *Endocrinology* (2007) 148(3):1384–95. doi: 10.1210/en.2006-1198
47. Perlman K, Benrimoh D, Israel S, Rollins C, Brown E, Tunteng JF, et al. A Systematic Meta-Review of Predictors of Antidepressant Treatment Outcome in Major Depressive Disorder. *J Affect Disord* (2019) 243:503–15. doi: 10.1016/j.jad.2018.09.067
48. Kornstein SG, Toups M, Rush AJ, Wisniewski SR, Thase ME, Luther J, et al. Do Menopausal Status and Use of Hormone Therapy Affect Antidepressant Treatment Response? Findings From the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) Study. *J Women's Heal* (2013) 22(2):121–31. doi: 10.1089/jwh.2012.3479
49. Koke SC, Brown EB, Miner CM. Safety and Efficacy of Fluoxetine in Patients Who Receive Oral Contraceptive Therapy. *Am J Obstet Gynecol* (2002) 187(3):551–5. doi: 10.1067/mob.2002.124939
50. Berry-Bibee EN, Kim MJ, Simmons KB, Tepper NK, Riley HEM, Pagano HP, et al. Drug Interactions Between Hormonal Contraceptives and Psychotropic Drugs: A Systematic Review. *Contraception* (2016) 94(49):650–67. doi: 10.1016/j.contraception.2016.07.011
51. Damoiseau VA, Proost JH, Jiawan VCR, Melgert BN. Sex Differences in the Pharmacokinetics of Antidepressants: Influence of Female Sex Hormones and Oral Contraceptives. *Clin Pharmacokinet* (2014) 53(6):509–19. doi: 10.1007/s40262-014-0145-2
52. Von Moltke LL, Greenblatt DJ, Giancarlo GM, Granda BW, Harmatz JS, Shader RI. Escitalopram (S-Citalopram) and its Metabolites In Vitro: Cytochromes Mediating Biotransformation, Inhibitory Effects, and Comparison to R-Citalopram. *Drug Metab Dispos* (2001) 29(8):1102–9.

53. Estrada-Camarena E, Fernández-Guasti A, López-Rubalcava C. Interaction Between Estrogens and Antidepressants in the Forced Swimming Test in Rats. *Psychopharmacol (Berl)* (2004) 173(1):139–45. doi: 10.1007/s00213-003-1707-4
54. Ma L, Xu Y, Zhou J, Li Y, Zhang X, Jiang W, et al. Brain Estrogen Alters the Effects of the Antidepressant Sertraline in Middle-Aged Female and Male Mice. *Mol Cell Endocrinol* (2020) 516:110947. doi: 10.1016/j.mce.2020.110947
55. Xu Y, Ma L, Jiang W, Li Y, Wang G, Li R. Study of Sex Differences in Duloxetine Efficacy for Depression in Transgenic Mouse Models. *Front Cell Neurosci* (2017) 11:344. doi: 10.3389/fncel.2017.00344
56. Shah A, Frazer A. Influence of Acute or Chronic Administration of Ovarian Hormones on the Effects of Desipramine in the Forced Swim Test in Female Rats. *Psychopharmacol (Berl)* (2014) 231(18):3685–94. doi: 10.1007/s00213-014-3510-9
57. Lucas G, Compan V, Charnay Y, Neve RL, Nestler EJ, Bockaert J, et al. Frontocortical 5-HT₄ Receptors Exert Positive Feedback on Serotonergic Activity: Viral Transfections, Subacute and Chronic Treatments With 5-HT₄ Agonists. *Biol Psychiatry* (2005) 57(8):918–25. doi: 10.1016/j.biopsych.2004.12.023
58. Chen BK, Mendez-David I, Luna VM, Faye C, Gardier AM, David DJ, et al. Prophylactic Efficacy of 5-HT₄R Agonists Against Stress. *Neuropsychopharmacology* (2020) 45(3):542–52. doi: 10.1038/s41386-019-0540-3
59. Lucas G, Rymar VV, Du J, Mnie-Filali O, Bisgaard C, Manta S, et al. Serotonin (4) (5-HT₄) Receptor Agonists are Putative Antidepressants With a Rapid Onset of Action. *Neuron* (2007) 55(5):712–25. doi: 10.1016/j.neuron.2007.07.041
60. Mendez-David I, David DJ, Darcet F, Wu MV, Kerdine-Römer S, Gardier AM, et al. Rapid Anxiolytic Effects of a 5-HT₄ Receptor Agonist are Mediated by a Neurogenesis-Independent Mechanism. *Neuropsychopharmacology* (2014) 39(6):1366–78. doi: 10.1038/npp.2013.332

Conflict of Interest: VF and GK declare to have received honorarium for advisory meetings at Sage Therapeutics, GK at Sanos, and VF for a symposium lecture at Lundbeck Pharma A/S. MJ has given talks sponsored by Boehringer Ingelheim and H. Lundbeck.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Larsen, Ozanne, Köhler-Forsberg, Poulsen, Dam, Svarer, Knudsen, Jørgensen and Frokjaer. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.