



Molecular Mechanisms of AMH Signaling

James A. Howard¹, Kaitlin N. Hart¹ and Thomas B. Thompson^{2*}

¹ Department of Pharmacology & Systems Physiology, University of Cincinnati, Cincinnati, OH, United States, ² Department of Molecular Genetics, Biochemistry, & Microbiology, University of Cincinnati, Cincinnati, OH, United States

Anti-Müllerian Hormone (AMH) is a secreted glycoprotein hormone with critical roles in reproductive development and regulation. Its chemical and mechanistic similarities to members of the Transforming Growth Factor β (TGF- β) family have led to its placement within this signaling family. As a member of the TGF- β family, AMH exists as a noncovalent complex of a large N-terminal prodomain and smaller C-terminal mature signaling domain. To produce a signal, the mature domain will bind to the extracellular domains of two type I and two type II receptors which results in an intracellular SMAD signal. Interestingly, as will be discussed in this review, AMH possesses several unique characteristics which set it apart from other ligands within the TGF- β family. In particular, AMH has a dedicated type II receptor, Anti-Müllerian Hormone Receptor Type II (AMHR2), making this interaction intriguing mechanistically as well as therapeutically. Further, the prodomain of AMH has remained largely uncharacterized, despite being the largest prodomain within the family. Recent advancements in the field have provided valuable insight into the molecular mechanisms of AMH signaling, however there are still many areas of AMH signaling not understood. Herein, we will discuss what is known about the biochemistry of AMH and AMHR2, focusing on recent advances in understanding the unique characteristics of AMH signaling and the molecular mechanisms of receptor engagement.

Keywords: anti-müllerian hormone (AMH), anti-müllerian hormone receptor type II (AMHR2), transforming growth factor- β (TGF- β), bone morphogenetic protein (BMP), activin, persistent müllerian duct syndrome (PMDS), cell signaling, prodomain

OPEN ACCESS

Edited by:

Rodolfo A Rey,
Hospital de Niños Ricardo Gutiérrez,
Argentina

Reviewed by:

Richard Cate,
Boston University, United States

*Correspondence:

Thomas B. Thompson
Tom.Thompson@uc.edu

Specialty section:

This article was submitted to
Reproduction,
a section of the journal
Frontiers in Endocrinology

Received: 25 April 2022

Accepted: 20 May 2022

Published: 22 June 2022

Citation:

Howard JA, Hart KN and
Thompson TB (2022) Molecular
Mechanisms of AMH Signaling.
Front. Endocrinol. 13:927824.
doi: 10.3389/fendo.2022.927824

INTRODUCTION

Anti-Müllerian Hormone (AMH), also known as Müllerian Inhibiting Substance (MIS), was first described by Alfred Jost in 1946 as a secreted testicular factor which drove the regression of the Müllerian ducts in the male fetus (1). Importantly, dysregulation of this mechanism presents clinically as Persistent Müllerian Duct Syndrome (PMDS), in which loss of function mutations in AMH or its signaling receptor, Anti-Müllerian Hormone Receptor Type II (AMHR2), lead to persistence of Müllerian duct derivatives – uterus, fallopian tubes, cervix, and upper vagina – in males (2, 3). In women, AMH is a negative regulator of folliculogenesis and dysregulation of the signaling pathway has been implicated in two leading causes of female infertility: Polycystic Ovary Syndrome (PCOS) and Primary Ovarian Insufficiency (POI) (4). Since its initial description, more recent characterization of this hormone has provided foundational insights into our current understanding of the structure and function of AMH and its signaling pathway.

AMH is a glycoprotein hormone (5) which shares structural and mechanistic homology with signaling proteins of the Transforming Growth Factor β (TGF- β) family (6). This family consists of over 30 secreted signaling ligands that have essential functions for many processes regulating cell homeostasis and human development, including reproductive development (7). These ligands are synthesized from a precursor consisting of a large N-terminal prodomain and smaller C-terminal mature signaling domain (**Figure 1A**). Folding, dimerization, and secretion are regulated by prodomains, which are cleaved from the smaller signaling domain and, in most cases, remain noncovalently associated (**Figure 1B**) (8). Ligands signal by binding to the extracellular domain (ECD) of two type I and two type II serine/threonine receptor kinases. This complex brings the intracellular kinase domain (ICD) of the constitutively active type II receptor in close enough proximity to phosphorylate the GS domain of the type I receptor ICD, relieving inhibition and activating Smad transcription factors (9) (**Figure 1C**). Signaling within the TGF- β family is limited to specific combinations of the seven type I receptors, Activin-like kinases 1-7 (ALK1-7), and five type II receptors, ActRIIA, ActRIIB, BMPR2, T β R2, and AMHR2 (7). It has been shown that AMH will mainly signal using ALK2 (10, 11) or ALK3 (12–14), type I receptors used by the bone morphogenetic protein (BMP) branch of TGF- β ligands, and activation of BMP R-Smads 1, 5, and 9 as well as activation of BMP reporter genes (15, 16). The other BMP type I receptor, ALK6, has a stimulatory or inhibitory effect depending on the tissue type (17, 18). AMHR2 is unique within the TGF- β family as it is the only receptor specific for a single ligand (19). In this review, we will summarize the current biochemical understanding of AMH as a TGF- β ligand from secretion to signal, with a focus on recent efforts to characterize the binding of AMH to AMHR2 and the looming gaps the field must overcome in order to better understand this important biological pathway.

PROCESSING AND REGULATION OF AMH & AMHR2


AMH was first identified as a TGF- β ligand by sequence similarity of its C-terminal mature signaling domain with Activins and TGF- β s (20) and the proteolytic processing of this domain (6). The full open reading frame of human AMH (UniProtKB P03971-1) consists of a signal sequence (SS) (residues 1-24), prodomain (residues 25-451), and mature domain (residues 452-560) (**Figure 1A**) (6). Human AMH is processed canonically; mammalian proprotein convertases (PCs), such as furin, will cleave the proprotein downstream of an R-X-X-R motif at amino acid position 448-451 to generate the 109 amino acid mature domain (21–24). Similar to other family members, PC cleavage separates the N-terminal prodomain from the C-terminal mature domain, which allows for assembly into a noncovalent complex (**Figure 1B**) (6, 25–27). Only the cleaved, processed, dimer form can properly bind its receptors and induce downstream signaling (28), but evidence of

mixed circulating species of AMH suggests a regulatory role of this processing (29). Both the processed and unprocessed species are found in the serum and follicular fluid (25, 30) in varying ratios depending on age (27), sex (27), and disease state (31, 32). Interestingly, alternative cleavage products resulting from serine proteinase activity within the prodomain (**Figure 1A**) have been described during purification (6, 25, 33–35), however their biological relevance remains unknown.

The processing of AMHR2 (UniProtKB Q16671-1) is less characterized than its ligand. While there has been robust study of type II receptor regulation within the TGF- β family *via* mechanisms of internalization (36–40), homo- and heteromeric complex formation (41–43), and glycosylation (44, 45), for AMHR2, the understanding of regulation is currently limited to biosynthetic processing and surface presentation alone (46, 47). Unlike other type II receptors, it has been suggested that functional presentation of AMHR2 at the membrane is negatively regulated by cleavage or by disulfide-linked oligomerization of the extracellular domain, leading to increased retention within the ER (46). Additionally, those functional receptors that are presented appear to organize in clusters of homo-oligomers, resulting in a lack of lateral mobility (46). In mammals, receptor splice variants have been identified that result in the deletion of either amino acids 377-471 (Amhr2 Δ 9/10) within the kinase domain, or 17-77 (Amhr2 Δ 2) within the extracellular domain (47–49) (**Figure 1A**). Although their mRNA expression level in the testes and brain is 5% or less of the normal receptor, the existence of these variants raises interesting questions about their regulatory function in the AMH signaling pathway (47). Thus, continued investigation of the functional consequences of these or other splice variants is necessary to understand their potential impact on signaling. Lastly, Unlike other TGF- β ligands, investigations into mechanisms of extracellular antagonism of AMH have not been definitive (50–52). Nevertheless, regulation of ligands by protein antagonists represents a significant feature of TGF- β ligands, and the lack of known AMH-binding proteins is either a missing piece of the known mechanism or an interesting aberration from other family ligands.

THE ROLE OF THE AMH PRODOMAIN

It is widely accepted that the prodomains of TGF- β ligands are required for proper folding and dimerization of the mature signaling ligand (53–55). While most prodomains are similar in size, an indication of secondary structure elements, there are exceptions. For example, GDF15 maintains the smallest prodomain of 18.5 kDa whereas AMH has evolved the largest of the prodomains at 45 kDa. Furthermore, unlike the ligands which typically have a conserved pattern of cysteines, the prodomains exhibit significant differences in the number and placement of cysteines, indicating structural divergence (7). For TGF- β 1-3 the prodomains from two different chains are joined by a disulfide bond (56). The intermolecular disulfide bond increases the affinity of the prodomains for the mature ligand through avidity effects. Similarly, AMH also exhibits an



interchain disulfide bond which likely increases its affinity for the mature ligand (6).

For some ligands, such as the TGF- β s, GDF8, and GDF11 the prodomain maintains the ligand in a latent state, and activation occurs through proteolysis (57–59) of the prodomain or an integrin-mediated stretching mechanism (56), both of which liberate the bound ligand from the prodomain. For AMH, BMPs, and other activin ligands, the prodomain does not render the ligand latent and the ligand is either thought to signal in the presence of the prodomain or that the prodomain is readily displaced by binding the signaling receptors. For AMH, the prodomain has been shown to allosterically regulate AMH binding to AMHR2 without inhibiting signal (28, 60). This mechanism is similar to the non-latent BMP7 pro-complex, however unlike AMH, the BMP7 prodomain has a weakly competitive interaction with the BMP type II receptors and unchanged type I receptor interactions (61). Further parallels might be drawn from the crystal structure of the BMP9 pro-complex bound by ALK1, which shows that the type I receptor can associate without displacing the prodomain (62), but this remains untested for AMH. Unlike most BMPs, the AMH prodomain has a 10–100 fold higher affinity for the mature ligand ($K_d = 0.4$ pM) (60). Despite this high-affinity interaction, bivalent binding to AMHR2 presented on a surface is able to disrupt interactions and attenuate binding 1000-fold (60). Displacement appears to be dependent on the avidity as neither monovalent binding nor soluble receptor are able to induce prodomain displacement. Thus, while certain ligands have high affinity for their prodomain and confer latency, many BMP ligands have lower affinity for their prodomains and are more readily displaced by receptor binding (8, 58). AMH appears somewhat unique in that it maintains a very high affinity for the ligand, but the prodomain can also be displaced by cell surface receptors. The high affinity of the prodomain of AMH suggests that the prodomain is likely to play an important role in either protecting AMH or facilitating signaling.

As mentioned, the prodomain seems to be an additional and principal factor of regulation within the signaling pathway. The prodomain is required for proper folding, homo- or heterodimerization, and secretion (7, 63–65), and the presence of PMDS mutations within the prodomain support this mechanism for AMH (2). In the serum, there is no unbound mature AMH ligand (25), suggesting a role for the prodomain in shuttling the mature domain to nearby and distant targets. The endocrine character of AMH is a robust research area, as we have yet to fully comprehend the breadth of extragonadal signaling targets (4, 18, 66, 67). For other ligands, the pro-complex also functions as a shield from extracellular antagonists (23, 33, 64, 68, 69). The interface between BMP antagonist Crossveinless 2 and BMP2 is analogous to the interface between mature BMP9 and the BMP9 prodomain (7), so the large AMH prodomain might function to protect AMH from interactions with a milieu of extracellular matrix (ECM) components. On the other hand, prodomains seem to be important for targeting the mature ligand to the cell surface through interactions with heparin (8), fibrillin (70), and other components of the ECM (7). However, unlike many BMPs, AMH does not have

large positively-charged patches of amino acids which would limit its interactions with heparin; it instead has a significant hydrophobic character (71). We do know, however, that the prodomain is necessary for activity in tissue-based assays (72) but dispensable for cell-based assays (47, 52, 73). This suggests that the prodomain likely does not play a major role in the signaling mechanism but might play a larger role in the availability of the ligand by mediating ECM interactions or conferring protection from degradation or antagonism.

The prodomain itself may also be subject to regulatory mechanisms common to the TGF- β family. In this vein, the previously mentioned alternate cleavage sites of AMH (6, 74) (**Figure 1A**) might have some bearing on the activity of the noncovalent complex. Conformational changes (7), alternative cleavage (57, 75, 76), or other uncharacterized modifications (77) have been shown to prime the noncovalent complex for receptor interactions. Furthermore, there are 2 N-linked glycosylation and several possible O-linked glycosylation sites predicted within the AMH prodomain (78) comprising 13.5% of the complex mass (71). Differential modification of glycosylation may impact protein-protein interactions or cleavage, as observed in *Drosophila* with the ortholog of BMP7, *Gbb* (76). Largely, we lack understanding of the regulatory role of the AMH prodomain beyond its absolute necessity for secretion and activity in the body. Whether the AMH prodomain, which is the family's largest and most divergent, has additional function beyond increasing the availability of the AMH signaling ligand is not known. While recent advances in modeling using AlphaFold can help to visualize structure and domain architecture of the AMH prodomain (79), the lack of structural definition of this region and low homology cause difficulty in effectively modeling the AMH prodomain (62, 63, 80–82). As such, structural and biochemical characterization of the prodomain structure and its interfacing interactions with the mature ligand and receptors will help ascertain its function.

STRUCTURAL DEFINITION OF AMH AND AMHR2

The TGF- β family is part of the cystine knot growth factor (CKGF) superfamily (7) which have a conserved fold and sequence. The overall shape can be described as an opposite-facing left and right hands in a Vulcan salute joined at the palm (**Figure 2A**). This creates a concave pocket between the wrist helix of one chain and the fingertips of the opposing chain, to which the type I receptor is recruited, and a convex surface on a single chain at the “knuckle” region, to which the type II receptor binds for Activins and BMPs (84). The extracellular ligand-binding domain of the type II receptors adopt a three-finger toxin fold, which has also classically been described with a hand-like morphology, consisting of three anti-parallel beta strand fingers and a central palm region (**Figure 2B**). These features are also conserved for AMH and AMHR2 as shown in the recently solved structure of the extracellular complex (85).

In general, ligands have evolved two central binding modes to interact specifically with their type II receptors (**Figure 2C**).

Ligands of the Activin and BMP class bind at the convex, knuckle surface of the ligand fingers, while the TGF- β class ligands bind the fingertips (84). For AMH, the binding mode was unknown until the recently-solved crystal structure of AMH bound to AMHR2 (PDB:7L0J) (85). This structure provided a critical piece for understanding ligand-receptor interactions and disease-causing mutations (**Figures 2D–E**), revealing that while similar to the general binding mode of BMP and Activins, AMH utilizes a modified mode of type II receptor binding (85).

The receptor binding interaction of AMH and AMHR2 is unique within the family. While TGF- β class ligands bind T β R2 using finger 1 of the receptor and the fingertips of the ligand, Activin and BMP class ligands bind ActRIIA and ActRIIB at the palm of the receptor and the knuckles of the ligand (86–91) (**Figure 2C**). Like Activins and BMPs, AMH binds AMHR2 using the palm of the receptor and the knuckles of the ligand, however this interface is shifted towards the fingertips by about 7.5 Å. Additionally, fingers 1 and 3 of AMHR2 wrap around the ligand making unique contacts with sections of AMH not observed with Activins and BMPs, especially within fingers 3 and 4 of AMH and the connecting loop (85). While the mature ligand is similar to each of the other three classes (the root mean square deviation of the C α positions is below 2 Å for BMP2, GDF11, and TGF- β 1) the AMH ligand adopts a flat character of the fingers akin to TGF- β s.

The structure of AMH bound to AMHR2 highlighted structural differences in each that are likely responsible for specificity. Of note, AMH has a truncated finger 1/2 loop relative to other ligands that facilitates the wrap-around mechanism of AMHR2 (85). More significant variance is observed on the receptor side with differences between AMHR2 and other type II receptors undoubtedly contributing to specificity. Most notably is the conformation of finger 1 of AMHR2 which is extended compared to other receptors and forms a favorable interaction with AMH. While the number of disulfide bonds are similar, a shift of one cysteine (Cys60) results in unique structural character of AMHR2. The altered location of the disulfide bond brings together the finger 2/3 loop and finger 3 to create unique surface for AMH binding (85). These distinct conformational features of AMHR2 promote the selectivity of AMH binding and signaling.

Where we still lack critical information is in our understanding of the nature of interactions between AMH and its type I receptors. The affinity of AMH for its type II receptor has been shown to resemble TGF- β s or Activins, while affinity for the type I receptor, though not yet directly tested, is assumed to follow the same low affinity archetype as the above ligand classes (28, 73, 85). What is known is that AMH lacks two conserved tryptophan residues present in the type I receptor binding site of BMPs. Importantly, these residues have been shown to be necessary for proper signaling in members of the BMP class (92). In fact, the entire type I binding interface is dissimilar to that of BMPs (86, 93, 94) and contains more polar and charged residues, yet AMH will signal using the same type I receptors – ALK2, ALK3, ALK6 – as BMPs. It will be interesting to determine how AMH

accommodates for its binding and specificity of type I receptors with these differences. It is possible that type I receptor binding is shifted relative to BMPs and might even potentially interact with the type II receptor in a cooperative mechanism similar to TGF- β , however, this has yet to be explored.

DISCUSSION

Recent studies have revealed a wealth of information about the molecular mechanisms of AMH signaling, but the field has a long way to go towards a full understanding of the intricacies of this




FIGURE 2 | Structural features of AMH and AMHR2. Diagram of the structure of mature AMH (A) and AMHR2 ECD (B). (C) Comparison of the binding modes of each ligand class to a type II receptor. (D) Mapping of PMDS mutations (83), indicated by red spheres, to the binary complex. (E) Sequence of AMH and AMHR2 annotated with secondary structure features and highlighted in red with the above PMDS mutations.

unique pathway. The biochemistry of AMH is certainly less characterized than its TGF- β family counterparts. Knowledge of these structures and their interactions can help explain the expanding genetic information linked to human diseases, such as PMDS and PCOS. For example, several mutations have been identified in the prodomain, however, we lack the structural information needed to better understand how these mutations impact AMH function.

The interaction between mature AMH and the extracellular domain of AMHR2 is perhaps the most actionable piece of recent data characterizing AMH. The crystal structure demonstrates unique features that set AMH apart from other ligand classes, as well as an atomic-level explanation for PMDS mutants which map to the interface (**Figures 2D–E**). This story is, of course, incomplete without a structure of mature AMH bound by a type I receptor. While the mutual exclusivity of the AMH and AMHR2 interaction is an interesting feature within the TGF- β family, equally interesting is how the intracellular kinase domain of AMHR2 can employ type I receptors shared with BMPs yet propagate an AMH-specific signal. Intracellular interactions remain something of a black box for the fields of both TGF- β and AMH biology.

Looking ahead, further structural studies of the AMH ligand and receptor are warranted; these studies must be supported by stronger assay development. Most importantly, the field should address discrepancies between *in vitro* and *in vivo* studies, especially concerning the prodomain. It has been known for some time that the AMH prodomain is required for biological function. This feature is reflected in tissue-based assays but not in cell-based assays where the mature ligand will suffice. Better care

should be taken to include the prodomain, when possible, to better replicate the biological context of AMH and allay concerns about differences between these two assay systems. Additionally, the ability to distinguish between the transcriptional outcome of AMH and BMP signaling would be a great and powerful tool for probing the mechanisms of the signaling pathway at every level. Research into this area might also help to answer a major question of AMH signal in general: is there a signaling cascade unique to AMH, or does AMH modulate a pre-existing BMP signal to generate unique outcomes? Although the research areas in need of attention are difficult, fresh data and new techniques have done wonders to answer critical questions and spark novel hypotheses about how this pathway truly functions.

AUTHOR CONTRIBUTIONS

JH and KH developed the concept. JH wrote the manuscript. KH and TT edited and revised the manuscript for important intellectual content. All authors read and approved the submitted version.

FUNDING

This work was funded by an R01 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development, R01HD105818.

REFERENCES

- Jost A. The Age Factor in the Castration of Male Rabbit Fetuses. *Proc Soc Exp Biol Med* (1947) 66:302. doi: 10.3181/00379727-66-16071
- Belville C, Van Vlijmen H, Ehrenfels C, Pepinsky B, Rezaie AR, Picard J-Y, et al. Mutations of the Anti-Müllerian Hormone Gene in Patients With Persistent Müllerian Duct Syndrome: Biosynthesis, Secretion, and Processing of the Abnormal Proteins and Analysis Using a Three-Dimensional Model. *Mol Endocrinol* (2004) 18:708–21. doi: 10.1210/me.2003-0358
- Josso N, Belville C, di Clemente N, Picard J-Y. AMH and AMH Receptor Defects in Persistent Müllerian Duct Syndrome. *Hum Reprod Update* (2005) 11:351–6. doi: 10.1093/humupd/dmi014
- di Clemente N, Racine C, Pierre A, Taieb J. Anti-Müllerian Hormone in Female Reproduction. *Endocr Rev* (2021) 42:753–82. doi: 10.1210/endo/rev/bnab012
- Picard JY, Tran D, Josso N. Biosynthesis of Labeled Anti-Müllerian Hormone by Fetal Testes: Evidence for the Glycoprotein Nature of the Hormone and for its Disulfide-Bonded Structure. *Mol Cell Endocrinol* (1978) 12:17–30. doi: 10.1016/0303-7207(78)90098-9
- Pepinsky RB, Sinclair LK, Chow EP, Mattaliano RJ, Manganaro TF, Donahoe PK, et al. Proteolytic Processing of Müllerian Inhibiting Substance Produces a Transforming Growth Factor-Beta-Like Fragment. *J Biol Chem* (1988) 263:18961–4. doi: 10.1016/S0021-9258(18)37375-7
- Hinck AP, Mueller TD, Springer TA. Structural Biology and Evolution of the TGF- β Family. *Cold Spring Harb Perspect Biol* (2016) 8:a022103. doi: 10.1101/cshperspect.a022103
- Sengle G, Ono RN, Sasaki T, Sakai LY. Prodomains of Transforming Growth Factor Beta (TGFbeta) Superfamily Members Specify Different Functions: Extracellular Matrix Interactions and Growth Factor Bioavailability. *J Biol Chem* (2011) 286:5087–99. doi: 10.1074/jbc.M110.188615
- Wrana JL, Attisano L, Wieser R, Ventura F, Massagué J. Mechanism of Activation of the TGF-Beta Receptor. *Nature* (1994) 370:341–7. doi: 10.1038/370341a0
- Visser JA, Olaso R, Verhoef-Post M, Kramer P, Themmen AP, Ingraham HA. The Serine/Threonine Transmembrane Receptor ALK2 Mediates Müllerian Inhibiting Substance Signaling. *Mol Endocrinol* (2001) 15:936–45. doi: 10.1210/mend.15.6.0645
- Clarke TR, Hoshiya Y, Yi SE, Liu X, Lyons KM, Donahoe PK. Müllerian Inhibiting Substance Signaling Uses a Bone Morphogenetic Protein (BMP)-Like Pathway Mediated by ALK2 and Induces SMAD6 Expression. *Mol Endocrinol* (2001) 15:946–59. doi: 10.1210/mend.15.6.0664
- Jamin SP, Arango NA, Mishina Y, Hanks MC, Behringer RR. Requirement of Bmpr1a for Müllerian Duct Regression During Male Sexual Development. *Nat Genet* (2002) 32:408–10. doi: 10.1038/ng1003
- Sèdes L, Leclerc A, Moindjie H, Cate RL, Picard J-Y, di Clemente N, et al. Anti-Müllerian Hormone Recruits BMPR-IA in Immature Granulosa Cells. *PLoS One* (2013) 8:e81551. doi: 10.1371/journal.pone.0081551
- Gouédard L, Chen YG, Thevenet L, Racine C, Borie S, Lamarre I, et al. Engagement of Bone Morphogenetic Protein Type IB Receptor and Smad1 Signaling by Anti-Müllerian Hormone and its Type II Receptor. *J Biol Chem* (2000) 275:27973–8. doi: 10.1074/jbc.M002704200
- di Clemente N, Josso N, Gouédard L, Belville C. Components of the Anti-Müllerian Hormone Signaling Pathway in Gonads. *Mol Cell Endocrinol* (2003) 211:9–14. doi: 10.1016/j.mce.2003.09.005
- Orvis GD, Jamin SP, Kwan KM, Mishina Y, Kaartinen VM, Huang S, et al. Functional Redundancy of TGF-Beta Family Type I Receptors and Receptor-Smads in Mediating Anti-Müllerian Hormone-Induced Müllerian Duct Regression in the Mouse. *Biol Reprod* (2008) 78:994–1001. doi: 10.1095/biolreprod.107.066605

17. Belville C, Jamin SP, Picard J-Y, Josso N, di Clemente N. Role of Type I Receptors for Anti-Müllerian Hormone in the SMAT-1 Sertoli Cell Line. *Oncogene* (2005) 24:4984–92. doi: 10.1038/sj.onc.1208686
18. Silva MSB, Giacobini P. New Insights Into Anti-Müllerian Hormone Role in the Hypothalamic-Pituitary-Gonadal Axis and Neuroendocrine Development. *Cell Mol Life Sci* (2021) 78:1–16. doi: 10.1007/s00018-020-03576-x
19. Baarends WM, van Helmond MJ, Post M, van der Schoot PJ, Hoogerbrugge JW, de Winter JP, et al. A Novel Member of the Transmembrane Serine/Threonine Kinase Receptor Family is Specifically Expressed in the Gonads and in Mesenchymal Cells Adjacent to the Müllerian Duct. *Development* (1994) 120:189–97. doi: 10.1242/dev.120.1.189
20. Cate RL, Mattaliano RJ, Hession C, Tizard R, Farber NM, Cheung A, et al. Isolation of the Bovine and Human Genes for Müllerian Inhibiting Substance and Expression of the Human Gene in Animal Cells. *Cell* (1986) 45:685–98. doi: 10.1016/0092-8674(86)90783-x
21. Dubois CM, Laprise MH, Blanchette F, Gentry LE, Leduc R. Processing of Transforming Growth Factor Beta 1 Precursor by Human Furin Convertase. *J Biol Chem* (1995) 270:10618–24. doi: 10.1074/jbc.270.18.10618
22. Constam DB, Robertson EJ. Regulation of Bone Morphogenetic Protein Activity by Pro Domains and Proprotein Convertases. *J Cell Biol* (1999) 144:139–49. doi: 10.1083/jcb.144.1.139
23. Constam DB. Regulation of Tg β 1 and Related Signals by Precursor Processing. *Semin Cell Dev Biol* (2014) 32:85–97. doi: 10.1016/j.semdb.2014.01.008
24. Nachtigal MW, Ingraham HA. Bioactivation of Müllerian Inhibiting Substance During Gonadal Development by a Kex2/Subtilisin-Like Endoprotease. *Proc Natl Acad Sci U.S.A.* (1996) 93:7711–6. doi: 10.1073/pnas.93.15.7711
25. Pankhurst MW, McLennan IS. Human Blood Contains Both the Uncleaved Precursor of Anti-Müllerian Hormone and a Complex of the NH₂- and COOH-Terminal Peptides. *Am J Physiol Endocrinol Metab* (2013) 305:E1241–1247. doi: 10.1152/ajpendo.00395.2013
26. McLennan IS, Pankhurst MW. Anti-Müllerian Hormone is a Gonadal Cytokine With Two Circulating Forms and Cryptic Actions. *J Endocrinol* (2015) 226:R45–57. doi: 10.1530/JOE-15-0206
27. Pankhurst MW, Chong YH, McLennan IS. Relative Levels of the Proprotein and Cleavage-Activated Form of Circulating Human Anti-Müllerian Hormone are Sexually Dimorphic and Variable During the Life Cycle. *Physiol Rep* (2016) 4:e12783. doi: 10.14814/phy2.12783
28. di Clemente N, Jamin SP, Lugovskoy A, Carmillo P, Ehrenfels C, Picard J-Y, et al. Processing of Anti-Müllerian Hormone Regulates Receptor Activation by a Mechanism Distinct From TGF- β . *Mol Endocrinol* (2010) 24:2193–206. doi: 10.1210/me.2010-0273
29. Pankhurst MW, Leathart B-LA, Batchelor NJ, McLennan IS. The Anti-Müllerian Hormone Precursor (proAMH) Is Not Converted to the Receptor-Competent Form (AMHN,C) in the Circulating Blood of Mice. *Endocrinology* (2016) 157:1622–9. doi: 10.1210/en.2015-1834
30. Campbell BK, Clinton M, Webb R. The Role of Anti-Müllerian Hormone (AMH) During Follicle Development in a Monovulatory Species (Sheep). *Endocrinology* (2012) 153:4533–43. doi: 10.1210/en.2012-1158
31. Peigné M, Pigny P, Pankhurst MW, Drumez E, Loyens A, Dewailly D, et al. The Proportion of Cleaved Anti-Müllerian Hormone is Higher in Serum But Not Follicular Fluid of Obese Women Independently of Polycystic Ovary Syndrome. *Reprod BioMed Online* (2020) 41:1112–21. doi: 10.1016/j.rbmo.2020.07.020
32. Pankhurst MW, Shorakae S, Rodgers RJ, Teede HJ, Moran LJ. Efficacy of Predictive Models for Polycystic Ovary Syndrome Using Serum Levels of Two Antimüllerian Hormone Isoforms (proAMH and AMHN,C). *Fertil Steril* (2017) 108:851–857.e2. doi: 10.1016/j.fertnstert.2017.08.012
33. Akiyama T, Marqués G, Wharton KA. A Large Bioactive BMP Ligand With Distinct Signaling Properties is Produced by Alternative Proconvertase Processing. *Sci Signal* (2012) 5:ra28. doi: 10.1126/scisignal.2002549
34. Ragin RC, Donahoe PK, Kenneally MK, Ahmad MF, MacLaughlin DT. Human Müllerian Inhibiting Substance: Enhanced Purification Imparts Biochemical Stability and Restores Antiproliferative Effects. *Protein Expr Purif* (1992) 3:236–45. doi: 10.1016/1046-5928(92)90020-w
35. Almeida J, Ball BA, Conley AJ, Place NJ, Liu IKM, Scholtz EL, et al. Biological and Clinical Significance of Anti-Müllerian Hormone Determination in Blood Serum of the Mare. *Theriogenology* (2011) 76:1393–403. doi: 10.1016/j.theriogenology.2011.06.008
36. Chen Y-G. Endocytic Regulation of TGF- β Signaling. *Cell Res* (2009) 19:58–70. doi: 10.1038/cr.2008.315
37. Chen C-L, Hou W-H, Liu I-H, Hsiao G, Huang SS, Huang JS. Inhibitors of Clathrin-Dependent Endocytosis Enhance Tg β 1 Signaling and Responses. *J Cell Sci* (2009) 122:1863–71. doi: 10.1242/jcs.038729
38. Di Guglielmo GM, Le Roy C, Goodfellow AF, Wrana JL. Distinct Endocytic Pathways Regulate TGF- β Receptor Signalling and Turnover. *Nat Cell Biol* (2003) 5:410–21. doi: 10.1038/ncb975
39. Hartung A, Bitton-Worms K, Rechtmann MM, Wenzel V, Boergermann JH, Hassel S, et al. Different Routes of Bone Morphogenetic Protein (BMP) Receptor Endocytosis Influence BMP Signaling. *Mol Cell Biol* (2006) 26(20):7791–805. doi: 10.1128/MCB.00022-06
40. Yao D, Ehrlich M, Henis YI, Leaf EB. Transforming Growth Factor- β Receptors Interact With AP2 by Direct Binding to β 2 Subunit. *MBoC* (2002) 13:4001–12. doi: 10.1091/mbc.02-07-0104
41. Ehrlich M, Horbelt D, Marom B, Knaus P, Henis YI. Homomeric and Heteromeric Complexes Among TGF- β and BMP Receptors and Their Roles in Signaling. *Cell Signalling* (2011) 23:1424–32. doi: 10.1016/j.celsig.2011.04.004
42. Ehrlich M, Gutman O, Knaus P, Henis YI. Oligomeric Interactions of TGF- β and BMP Receptors. *FEBS Lett* (2012) 586:1885–96. doi: 10.1016/j.febslet.2012.01.040
43. Nohe A, Hassel S, Ehrlich M, Neubauer F, Sebald W, Henis YI, et al. The Mode of Bone Morphogenetic Protein (BMP) Receptor Oligomerization Determines Different BMP-2 Signaling Pathways*. *J Biol Chem* (2002) 277:5330–8. doi: 10.1074/jbc.M102750200
44. Kim Y-W, Park J, Lee H-J, Lee S-Y, Kim S-J. TGF- β Sensitivity is Determined by N-Linked Glycosylation of the Type II TGF- β Receptor. *Biochem J* (2012) 445:403–11. doi: 10.1042/BJ20111923
45. Partridge EA, Le Roy C, Di Guglielmo GM, Pawling J, Cheung P, Granovsky M, et al. Regulation of Cytokine Receptors by Golgi N-Glycan Processing and Endocytosis. *Science* (2004) 306:120–4. doi: 10.1126/science.1102109
46. Hirschhorn T, di Clemente N, Amsalem AR, Pepinsky RB, Picard J-Y, Smorodinsky NI, et al. Constitutive Negative Regulation in the Processing of the Anti-Müllerian Hormone Receptor II. *J Cell Sci* (2015) 128:1352–64. doi: 10.1242/jcs.160143
47. Imhoff FM, Yang D, Mathew SF, Clarkson AN, Kawagishi Y, Tate WP, et al. The Type 2 Anti-Müllerian Hormone Receptor has Splice Variants That are Dominant-Negative Inhibitors. *FEBS Lett* (2013) 587:1749–53. doi: 10.1016/j.febslet.2013.04.014
48. Faure E, Gouédard L, Imbeaud S, Cate R, Picard J-Y, Josso N, et al. Mutant Isoforms of the Anti-Müllerian Hormone Type II Receptor Are Not Expressed at the Cell Membrane*. *J Biol Chem* (1996) 271:30571–5. doi: 10.1074/jbc.271.48.30571
49. di Clemente N, Wilson C, Faure E, Boussin L, Carmillo P, Tizard R, et al. Cloning, Expression, and Alternative Splicing of the Receptor for Anti-Müllerian Hormone. *Mol Endocrinol* (1994) 8:1006–20. doi: 10.1210/mend.8.8.7997230
50. Gipson GR, Goebel EJ, Hart KN, Kappes EC, Kattamuri C, McCoy JC, et al. Structural Perspective of BMP Ligands and Signaling. *Bone* (2020) 140:115549. doi: 10.1016/j.bone.2020.115549
51. Nilsson EE, Larsen G, Skinner MK. Roles of Gremlin 1 and Gremlin 2 in Regulating Ovarian Primordial to Primary Follicle Transition. *Reproduction* (2014) 147:865–74. doi: 10.1530/REP-14-0005
52. Kawagishi Y, Pankhurst MW, Nakatani Y, McLennan IS. Anti-Müllerian Hormone Signaling is Influenced by Follistatin 288, But Not 14 Other Transforming Growth Factor Beta Superfamily Regulators. *Mol Reprod Dev* (2017) 84:626–37. doi: 10.1002/mrd.22828
53. Gentry LE, Nash BW. The Pro Domain of Pre-Pro-Transforming Growth Factor Beta 1 When Independently Expressed Is a Functional Binding Protein for the Mature Growth Factor. *Biochemistry* (1990) 29:6851–7. doi: 10.1021/bi00481a014

54. Gray AM, Mason AJ. Requirement for Activin A and Transforming Growth Factor- β 1 Pro-Regions in Homodimer Assembly. *Science* (1990) 247:1328–30. doi: 10.1126/science.2315700
55. Walton KL, Makanji Y, Wilce MC, Chan KL, Robertson DM, Harrison CA. A Common Biosynthetic Pathway Governs the Dimerization and Secretion of Inhibin and Related Transforming Growth Factor β (TGF β) Ligands. *J Biol Chem* (2009) 284:9311–20. doi: 10.1074/jbc.M808763200
56. Shi M, Zhu J, Wang R, Chen X, Mi L, Walz T, et al. Latent TGF- β Structure and Activation. *Nature* (2011) 474:343–9. doi: 10.1038/nature10152
57. McCoy JC, Goebel EJ, Thompson TB. Characterization of Tolloid-Mediated Cleavage of the GDF8 Procomplex. *Biochem J* (2021) 478(9):1733–47. doi: 10.1042/BCJ20210054
58. Wolfman NM, McPherron AC, Pappano WN, Davies MV, Song K, Tomkinson KN, et al. Activation of Latent Myostatin by the BMP-1/Tolloid Family of Metalloproteinases. *Proc Natl Acad Sci U.S.A.* (2003) 100:15842–6. doi: 10.1073/pnas.2534946100
59. Walker RG, Poggioli T, Katsimpardi L, Buchanan SM, Oh J, Wattruss S, et al. Biochemistry and Biology of GDF11 and Myostatin: Similarities, Differences, and Questions for Future Investigation. *Circ Res* (2016) 118:1125–41. doi: 10.1161/CIRCRESAHA.116.308391
60. Cate RL, di Clemente N, Racine C, Groome NP, Pepinsky RB, Whitty A. The Anti-Müllerian Hormone Prodomain Is Displaced From the Hormone/Prodomain Complex Upon Bivalent Binding to the Hormone Receptor. *J Biol Chem* (2022) 298:101429. doi: 10.1016/j.jbc.2021.101429
61. Sengle G, Ono RN, Lyons KM, Bächinger HP, Sakai LY. A New Model for Growth Factor Activation: Type II Receptors Compete With the Prodomain for BMP-7. *J Mol Biol* (2008) 381:1025–39. doi: 10.1016/j.jmb.2008.06.074
62. Salmon RM, Guo J, Wood JH, Tong Z, Beech JS, Lawera A, et al. Molecular Basis of ALK1-Mediated Signalling by BMP9/BMP10 and Their Prodomain-Bound Forms. *Nat Commun* (2020) 11:1621. doi: 10.1038/s41467-020-15425-3
63. Zhao B, Xu S, Dong X, Lu C, Springer TA. Prodomain-Growth Factor Swapping in the Structure of Pro-TGF- β 1. *J Biol Chem* (2018) 293:1579–89. doi: 10.1074/jbc.M117.809657
64. Harrison CA, Al-Musawi SL, Walton KL. Prodomains Regulate the Synthesis, Extracellular Localisation and Activity of TGF- β Superfamily Ligands. *Growth Factors* (2011) 29:174–86. doi: 10.3109/08977194.2011.608666
65. Neugebauer JM, Kwon S, Kim H-S, Donley N, Tilak A, Sopory S, et al. The Prodomain of BMP4 is Necessary and Sufficient to Generate Stable BMP4/7 Heterodimers With Enhanced Bioactivity. *in vivo Proc Natl Acad Sci U.S.A.* (2015) 112:E2307–2316. doi: 10.1073/pnas.1501449112
66. Malone SA, Papadakis GE, Messina A, Mimouni NEH, Trova S, Imbernon M, et al. Defective AMH Signaling Disrupts GnRH Neuron Development and Function and Contributes to Hypogonadotropic Hypogonadism. *Elife* (2019) 8:e47198. doi: 10.7554/eLife.47198
67. Tata B, El Houda Mimouni N, Barbotin A-L, Malone SA, Loyens A, Pigny P, et al. Elevated Prenatal Anti-Müllerian Hormone Reprograms the Fetus and Induces Polycystic Ovary Syndrome in Adulthood. *Nat Med* (2018) 24:834–46. doi: 10.1038/s41591-018-0035-5
68. Cui Y, Hackenmiller R, Berg L, Jean F, Nakayama T, Thomas G, et al. The Activity and Signaling Range of Mature BMP-4 Is Regulated by Sequential Cleavage at Two Sites Within the Prodomain of the Precursor. *Genes Dev* (2001) 15:2797–802. doi: 10.1101/gad.940001
69. Ashe HL, Levine M. Local Inhibition and Long-Range Enhancement of Dpp Signal Transduction by Sog. *Nature* (1999) 398:427–31. doi: 10.1038/18892
70. Gregory KE, Ono RN, Charbonneau NL, Kuo C-L, Keene DR, Bächinger HP, et al. The Prodomain of BMP-7 Targets the BMP-7 Complex to the Extracellular Matrix. *J Biol Chem* (2005) 280:27970–80. doi: 10.1074/jbc.M504270200
71. Picard JY, Goulut C, Bourrillon R, Josso N. Biochemical Analysis of Bovine Testicular Anti-Müllerian Hormone. *FEBS Lett* (1986) 195:73–6. doi: 10.1016/0014-5793(86)80133-8
72. Wilson CA, di Clemente N, Ehrenfels C, Pepinsky RB, Josso N, Vigier B, et al. Müllerian Inhibiting Substance Requires its N-Terminal Domain for Maintenance of Biological Activity, A Novel Finding Within the Transforming Growth Factor- β Superfamily. *Mol Endocrinol* (1993) 7:247–57. doi: 10.1210/mend.7.2.8469238
73. Hart KN, Pépin D, Czepnik M, Donahoe PK, Thompson TB. Mutational Analysis of the Putative Anti-Müllerian Hormone (AMH) Binding Interface on its Type II Receptor, Amhr2. *Endocrinology* (2020) 161:bqaa066. doi: 10.1210/endo/bqaa066
74. Picard JY, Josso N. Purification of Testicular Anti-Müllerian Hormone Allowing Direct Visualization of the Pure Glycoprotein and Determination of Yield and Purification Factor. *Mol Cell Endocrinol* (1984) 34:23–9. doi: 10.1016/0303-7207(84)90155-2
75. Tilak A, Nelsen SM, Kim H-S, Donley N, McKnight A, Lee H, et al. Simultaneous Rather Than Ordered Cleavage of Two Sites Within the BMP4 Prodomain Leads to Loss of Ligand in Mice. *Development* (2014) 141:3062–71. doi: 10.1242/dev.110130
76. Anderson EN, Wharton KA. Alternative Cleavage of the Bone Morphogenetic Protein (BMP), Gbb, Produces Ligands With Distinct Developmental Functions and Receptor Preferences. *J Biol Chem* (2017) 292:19160–78. doi: 10.1074/jbc.M117.793513
77. Pierre A, Racine C, Rey RA, Fanchin R, Taieb J, Cohen-Tannoudji J, et al. Most Cleaved Anti-Müllerian Hormone Binds Its Receptor in Human Follicular Fluid But Little Is Competent in Serum. *J Clin Endocrinol Metab* (2016) 101:4618–27. doi: 10.1210/jc.2016-1742
78. The UniProt Consortium. UniProt: The Universal Protein Knowledgebase in 2021. *Nucleic Acids Res* (2021) 49:D480–9. doi: 10.1093/nar/gkaa1100
79. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly Accurate Protein Structure Prediction With AlphaFold. *Nature* (2021) 596:583–9. doi: 10.1038/s41586-021-03819-2
80. Mi L-Z, Brown CT, Gao Y, Tian Y, Le VQ, Walz T, et al. Structure of Bone Morphogenetic Protein 9 Procomplex. *Proc Natl Acad Sci U.S.A.* (2015) 112:3710–5. doi: 10.1073/pnas.1501303112
81. Wang X, Fischer G, Hyvönen M. Structure and Activation of Pro-Activin a. *Nat Commun* (2016) 7:12052. doi: 10.1038/ncomms12052
82. Cotton TR, Fischer G, Wang X, McCoy JC, Czepnik M, Thompson TB, et al. Structure of the Human Myostatin Precursor and Determinants of Growth Factor Latency. *EMBO J* (2018) 37:367–83. doi: 10.15252/embj.201797883
83. Picard J-Y, Cate RL, Racine C, Josso N. The Persistent Müllerian Duct Syndrome: An Update Based Upon a Personal Experience of 157 Cases. *Sex Dev* (2017) 11:109–25. doi: 10.1159/000475516
84. Goebel EJ, Hart KN, McCoy JC, Thompson TB. Structural Biology of the Tgf β Family. *Exp Biol Med (Maywood)* (2019) 244:1530–46. doi: 10.1177/1535370219880894
85. Hart KN, Stocker WA, Nagykeri NG, Walton KL, Harrison CA, Donahoe PK, et al. Structure of AMH Bound to AMHR2 Provides Insight Into a Unique Signaling Pair in the TGF- β Family. *PNAS* (2021) 118:1–10. doi: 10.1073/pnas.2104809118
86. Allendorph GP, Vale WW, Choe S. Structure of the Ternary Signaling Complex of a TGF-Beta Superfamily Member. *Proc Natl Acad Sci U.S.A.* (2006) 103:7643–8. doi: 10.1073/pnas.0602558103
87. Groppa J, Hinck CS, Samavarchi-Tehrani P, Zubieta C, Schuermann JP, Taylor AB, et al. Cooperative Assembly of TGF-Beta Superfamily Signaling Complexes Is Mediated by Two Disparate Mechanisms and Distinct Modes of Receptor Binding. *Mol Cell* (2008) 29:157–68. doi: 10.1016/j.molcel.2007.11.039
88. Weber D, Kotsch A, Nickel J, Harth S, Seher A, Mueller U, et al. A Silent H-Bond can be Mutationally Activated for High-Affinity Interaction of BMP-2 and Activin Type IIB Receptor. *BMC Struct Biol* (2007) 7:6. doi: 10.1186/1472-6807-7-6
89. Radaev S, Zou Z, Huang T, Lafer EM, Hinck AP, Sun PD. Ternary Complex of Transforming Growth Factor-Beta1 Reveals Isoform-Specific Ligand Recognition and Receptor Recruitment in the Superfamily. *J Biol Chem* (2010) 285:14806–14. doi: 10.1074/jbc.M109.079921
90. Townson SA, Martinez-Hackert E, Greppi C, Lowden P, Sako D, Liu J, et al. Specificity and Structure of a High Affinity Activin Receptor-Like Kinase 1 (ALK1) Signaling Complex. *J Biol Chem* (2012) 287:27313–25. doi: 10.1074/jbc.M112.377960
91. Goebel EJ, Corpina RA, Hinck CS, Czepnik M, Castonguay R, Grenha R, et al. Structural Characterization of an Activin Class Ternary Receptor Complex Reveals a Third Paradigm for Receptor Specificity. *Proc Natl Acad Sci U.S.A.* (2019) 116:15505–13. doi: 10.1073/pnas.1906253116

92. Klammert U, Mueller TD, Hellmann TV, Wuerzler KK, Kotsch A, Schliermann A, et al. GDF-5 can Act as a Context-Dependent BMP-2 Antagonist. *BMC Biol* (2015) 13:77. doi: 10.1186/s12915-015-0183-8
93. Keller S, Nickel J, Zhang J-L, Sebald W, Mueller TD. Molecular Recognition of BMP-2 and BMP Receptor IA. *Nat Struct Mol Biol* (2004) 11:481–8. doi: 10.1038/nsmb756
94. Kotsch A, Nickel J, Seher A, Sebald W, Müller TD. Crystal Structure Analysis Reveals a Spring-Loaded Latch as Molecular Mechanism for GDF-5-Type I Receptor Specificity. *EMBO J* (2009) 28:937–47. doi: 10.1038/emboj.2009.37

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Howard, Hart and Thompson. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.