



OPEN ACCESS

EDITED BY

Sheerazed Boulkroun,
Institut National de la Santé et de la
Recherche Médicale (INSERM),
France

REVIEWED BY

Letizia Canu,
University of Florence, Italy
Clemens Kratochwil,
Heidelberg University Hospital,
Germany

*CORRESPONDENCE

Stefan Prado-Wohlwend
prado_ste@gva.es

SPECIALTY SECTION

This article was submitted to
Cancer Endocrinology,
a section of the journal
Frontiers in Endocrinology

RECEIVED 30 May 2022

ACCEPTED 30 September 2022

PUBLISHED 20 October 2022

CITATION

Prado-Wohlwend S,
del Olmo-García MI,
Bello-Arques P and Merino-Torres JF
(2022) Response to targeted
radionuclide therapy with [¹³¹I]MIBG
AND [¹⁷⁷Lu]Lu-DOTA-TATE according
to adrenal vs. extra-adrenal primary
location in metastatic paragangliomas
and pheochromocytomas: A
systematic review.
Front. Endocrinol. 13:957172.
doi: 10.3389/fendo.2022.957172

COPYRIGHT

© 2022 Prado-Wohlwend,
del Olmo-García, Bello-Arques and
Merino-Torres. This is an open-access
article distributed under the terms of
the [Creative Commons Attribution
License \(CC BY\)](#). The use, distribution
or reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Response to targeted radionuclide therapy with [¹³¹I]MIBG AND [¹⁷⁷Lu]Lu-DOTA-TATE according to adrenal vs. extra-adrenal primary location in metastatic paragangliomas and pheochromocytomas: A systematic review

Stefan Prado-Wohlwend^{1*}, María Isabel del Olmo-García²,
Pilar Bello-Arques¹ and Juan Francisco Merino-Torres^{2,3}

¹Nuclear Medicine Department, University and Polytechnic Hospital La Fe, Valencia, Spain,

²Endocrinology and Nutrition Department, University and Polytechnic Hospital La Fe, Valencia, Spain, ³Medicine Department, Universitat de València, Valencia, Spain

Purpose: Targeted radionuclide therapy (TRT) with [¹³¹I]MIBG and [¹⁷⁷Lu]Lu-DOTA-TATE is an alternative treatment to the classic schemes in slow progressive metastatic/inoperable paraganglioma (PGL) and pheochromocytoma (PHEO). There is no consensus on which treatment to administer and/or the best sequence in patients who are candidates for both therapies. To clarify these questions, this systematic review assesses the prognostic value of [¹³¹I]MIBG and [¹⁷⁷Lu]Lu-DOTA-TATE (PRRT-Lu) treatments in terms of progression-free survival (PFS) both globally and considering the primary location.

Methods: This review was developed according to the PRISMA Statement with 27 final studies (608 patients). Patient characteristics, treatment procedure, and follow-up criteria were evaluated. In addition, a Bayesian linear regression model weighted according to its sample size and an alternative model, which also included an interaction between the treatment and the proportion of PHEOs, were carried out, adjusted by a Student's *t* distribution.

Results: In linear regression models, [¹³¹I]MIBG overall PFS was, on average, 10 months lower when compared with PRRT-Lu. When considering the interaction between treatment responses and the proportion of PHEOs, PRRT-Lu showed remarkably better results in adrenal location. The PFS of PRRT-Lu was longer when the ratio of PHEOs increased, with a decrease in [¹³¹I]MIBG PFS by 1.9 months for each 10% increase in the proportion of PHEOs in the sample.

Conclusion: Methodology, procedure, and PFS from the different studies are quite heterogeneous. PRRT-Lu showed better results globally and specifically in PHEOs. This fact opens the window to prospective trials comparing or sequencing [¹³¹I]MIBG and PRRT-Lu.

KEYWORDS

paraganglioma, 131I-MIBG therapy, PRRT (peptide receptor radionuclide therapy), theragnostic agents, pheochromocytoma

Introduction

Paragangliomas (PGLs) and pheochromocytomas (PHEOs), altogether PGGLs, are uncommon neuroendocrine chromaffin tumors with an incidence of about 0.6 case per 100,000 persons per year. PHEOs are more common (80%–85%) and arise from the adrenal medulla, while PGLs are less frequent extra-adrenal tumors (15%–20%) derived from sympathetic or parasympathetic ganglia (1).

PGGLs are the most heritable of all human tumors, but paradoxically there is a wide genetic heterogeneity, with almost 32 genes described until the present and related to this disease (2). At least one-third of these patients carry a germline mutation, sorted in three different clusters: pseudohypoxia (cluster 1), kinase signaling (cluster 2), and Wnt signaling (cluster 3). The cluster will determine the primary locations, special molecular features, risk of metastatic disease, and avidity for nuclear medicine radiotracers (3–5).

It is difficult to predict the individual tumoral behavior of PGGLs. Half of them are slow-growing, but the other half behave aggressively, that the term benign or malignant PGGL has been replaced by a risk stratification approach (6).

In limited focal disease, the surgical treatment is the only curative approach. In metastatic disease, the overall 5-year survival is 60%–70%, and the decision as to whether systemic treatment is appropriate is based on multiple factors. The “wait-and-see” strategy may be an option in those patients without any evidence of disease progression, but in symptomatic, progressive, or high-tumor-burden disease, a systemic treatment should be administered (7).

Among the systemic choices, chemotherapy with cyclophosphamide-vincristine-dacarbazine and temozolomide is the classic primary option in patients with quick progression. Targeted therapies such as anti-angiogenic tyrosine-kinase inhibitors, everolimus, and alpha-interferon have been tested in different disease stages and progression indexes, but many questions still remain unsolved.

Among the few therapeutic options, targeted radionuclide therapies (TRTs) are promising and generally well-tolerated treatments that can reach an adequate symptomatic control with little interference in a patient’s life. There are two options within TRT: ¹³¹I-metaiodobenzylguanidine ([¹³¹I]MIBG) and [¹⁷⁷Lu]Lu-DOTA-TATE, also called peptide receptor radionuclide therapy (PRRT-Lu). Both treatments are alternatives to the classic schemes, at least in slow progressive patients, and in many cases, radiopharmaceuticals and even a sequential treatment with them may be appropriate in the same patient. The first step to plan a TRT is to evaluate the therapeutic targets with specific nuclear medicine scans: [¹²³I]MIBG to select the therapy with [¹³¹I]MIBG and somatostatin receptor (SSTR) scans (SSTR SPECT/CT and [⁶⁸Ga]Ga-DOTA-TOC PET/CT) to select PRRT. If the tumor expresses only one target, the decision is easy. However, overexpression of both targets has been described in up to 50% of patients, and it is in those cases where a question arises: which treatment is the most appropriate and/or in what sequence? (8, 9). In other cases, the tumors do not adequately express any of the specific targets and will not be candidates for TRT. The characteristics of the TRT modalities are summarized in [Table 1](#).

Within the PRRT modality, there are two radiopharmaceuticals: [¹⁷⁷Lu]Lu-DOTA-TATE (PRRT-Lu) and [⁹⁰Y]Y-DOTA-TOC (PRRT-Y). The most commonly used agent in PGGLs is PRRT-Lu, and therefore in this review we will focus on it. PRRT-Y has a better profile for larger lesions, while PRRT-Lu has more favorable characteristics to treat small-volume metastatic lesions, such as those present in PGGLs. PRRT-Lu was approved in the treatment of well-differentiated metastatic/inoperable gastroenteropancreatic neuroendocrine tumors, but in metastatic/inoperable PGGL, its use is limited to clinical trials or compassionate use in many countries, so that the published data are limited to small uncontrolled retrospective studies (10, 11).

There are two sorts of [¹³¹I]MIBG radiopharmaceuticals: low-specific-activity or conventional [¹³¹I]MIBG (LSA [¹³¹I]MIBG), which is available worldwide and has been used for

TABLE 1 TRT modality characteristics.

| Targeted radiolabeled therapy | ¹³¹ I-MIBG (Low-specific-activity) | ¹⁷⁷ Lu-DOTATATE |
|-------------------------------|---|---|
| Target | Norepinephrine transporter (principally) | Somatostatin receptors (especially subtype 2) |
| Phenotypic imaging scans | ¹²³ I-MIBG | ¹¹¹ In-Pentetreotide/ ^{99m} Tc-HYNIC-TOC SPECT/CT, ⁶⁸ Ga-DOTATOC/NOC/TATE PET/CT |
| Average beta emission | 0.192 MeV | 0.134 MeV |
| Mean tissue range | 0.5mm | 0.23 mm |
| Storage | Neuroendocrine vesicles | Lysosomes |
| FDA approval | Metastatic/inoperable PGGL | Well-differentiated metastatic/inoperable GEP-NETs |

decades, and high-specific-activity [¹³¹I]MIBG (HAS [¹³¹I]MIBG) approved by the FDA in 2018 but not yet available in many countries. In LSA [¹³¹I]MIBG, which is used up to date, more than 99% of MIBG molecules are not labeled with ¹³¹I (cold MIBG). In HAS [¹³¹I]MIBG, the labeling process is 100–200 times more efficient, reducing the side effects and the competitiveness with the cold MIBG and reaching probably better results than LSA [¹³¹I]MIBG, but several studies will be needed to support it when use of this new therapeutic agent becomes widespread (12).

There is not yet any consensus or therapeutic guide on which TRT radiopharmaceutical should be used, and their place in the therapeutic algorithm is unclear. The ESMO-EURACAN clinical guidelines for metastatic PGGLs published in 2020 recommended an individualized management approach in case of disease progression, including TRT among chemotherapy, local therapies, or additional treatments (13).

The aim of this systematic review in metastatic PGGLs is to assess patient characteristics, treatment procedure, follow-up, and differences in the global response and as a function of the primary location of [¹³¹I]MIBG and [¹⁷⁷Lu]Lu-DOTA-TATE in terms of progression-free survival (PFS). These results could facilitate the selection of the best option in those patients who can be treated with both therapies or in whom a sequential treatment is proposed.

Material and methods


This systematic review was developed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement and conducted using MEDLINE (accessed from PubMed), Google scholar, and ClinicalTrials.gov. The search strategy was based on the Population, Intervention, Comparator, Outcome (PICO) framework and was designed to find studies and reviews including a combination of medical subject headings (MeSH) and non-MeSH keywords related to treatment

with radiopharmaceuticals ([¹⁷⁷Lu]Lu-DOTA-TATE and [¹³¹I]MIBG) in patients with metastatic paraganglioma and/or pheochromocytoma: “Peptide Receptor Radionuclide Therapy” or “¹³¹I-metaiodobenzylguanidine” and “metastatic paraganglioma and/or pheochromocytoma”. Papers referring exclusively to HAS [¹³¹I]MIBG or PRRT-Y were excluded. The total number of final studies included in the qualitative synthesis was 27 (Figure 1).

Patient characteristics, treatment procedure, and administered activity were reviewed. Response to PRRT-Lu and [¹³¹I]MIBG was assessed through PFS. PFS was assessed from the time of treatment performed on disease progression, both overall and based on the primary location of the tumor. The evaluation criteria mentioned in the reports were biochemical and radiological according to Response Evaluation Criteria in Solid Tumors (RECIST) and through clinical follow-up.

Statistical analysis

Data were summarized using mean (standard deviation) and median (first, third quartile) in the case of numerical variables, whereas relative and absolute frequencies were applied in the case of categorical variables. A descriptive study of patient variables and a calculation of the unweighted PFS means were performed. To assess differences in PFS between both treatments, a Bayesian linear regression model was adjusted where each study was weighted according to its sample size, including treatment, age, year of the study, and proportion of PHEOs as explicative variables. Additionally, an alternative model was adjusted where the interaction between the treatment and the proportion of PHEOs was also included. The model was fitted using a Student's *t* distribution for likelihood instead of Gaussian to deal with outliers in the data and to make the model more robust to them. In both models, 95% credibility intervals were estimated for each parameter. The effect of the different explicative variables was also represented in conditional effect plots. All statistical analyses were performed using R (version 4.0.3) and R package brms (version 2.14.4).



Results

This systematic review included 27 studies, 4 of them prospective and 23 retrospective (Table 2). Nine studies were carried out with PRRT-Lu, 17 with [¹³¹I]MIBG, and one study included both modalities.

Only four reports provided PFS based on the adrenal or extra-adrenal location. In relation to the line in the treatment scheme, few articles specified it. [¹³¹I]MIBG was preferably administered as the first systemic line (73.4%), whereas PRRT-Lu was administered first line in 39.7% and second line in 38.5%. Most studies did not report the genetic syndromes of the patients (especially in [¹³¹I]MIBG), so this variable could not be assessed. Regarding the response evaluation criteria, 10 articles evaluated biochemical criteria, 21 RECIST, and 10 clinical follow-up.

A total of 1248 patients were reported in the series, of which 608 were PGGL (264 PHEO and 316 PGL, and in the remaining 28 patients, the primary location was not specified) (Table 3). The largest number of patients was treated with [¹³¹I]MIBG (76.6%). In the PRRT-Lu group, most of the patients were PGL (83.3%), whereas in the [¹³¹I]MIBG group, the numbers of PGL and PHEO were similar (46.9 vs. 53%). The vast majority of PHEOs were treated with [¹³¹I]MIBG (92.4%) (Figure 2).

The PFS estimates from the reviewed studies show a very large spread (12–46 months), and there is also no clear pattern regarding years or treatment modality (Figure 3). The unweighted mean of PFS was 25.43 months for [¹³¹I]MIBG and 29.55 for PRRT-Lu, with no statistical significance ($p: 0.787$) (95% CI [-5.9, 14.1]). The PFS mean based on the primary location could not be calculated due to the lack of specific data in the reports.

In the regression model weighted for each report according to the corresponding sample sizes, [¹³¹I]MIBG showed lower PFS when compared with PRRT-Lu. Specifically, the [¹³¹I]MIBG yielded, on average, 10 months lower PFS than PRRT-Lu (95% CI [-11.7, -8.5]) (Table 4).

As only four reports provided PFS based on the primary location and the previous statistical model did not consider that the treatments do not affect adrenal and extra-adrenal primary location equally, we adjusted a new Bayesian linear regression model considering a possible interaction between treatment responses and the proportion of PHEOs. This new model revealed very different results, and the difference between [¹³¹I]MIBG and PRRT-Lu was dependent on the proportion of PHEOs in each study. PRRT-Lu increased the PFS in PHEOs comparatively with [¹³¹I]MIBG. [¹³¹I]MIBG showed larger PFS

TABLE 2 Studies included in this systematic review.

| TYPE | Year of publication | TRT | Total patients | Number of PGGL patients | Primary location | PFS (months) | PFS (PGL) | PFS (PHEO) | Reference |
|---------------|---------------------|----------------|----------------|-------------------------|-------------------|--------------|-------------|------------|--|
| Retrospective | 2017 | 177Lu-DOTATATE | 20 | 20 | PHEO 8 PGL 12 | 39 | ns* | ns | (Kong et al., 2017) (14) |
| Retrospective | 2019 | 177Lu-DOTATATE | 30 | 30 | PHEO 3 PGL 27 | 30 | 31 | 10 | (Zandee et al., 2019) (15) |
| Retrospective | 2006 | 177Lu-DOTATATE | 22 | 12 | PHEO 1 PGL 12 | 22 | ns | ns | (Van Essen et al., 2006) (16) |
| Retrospective | 2016 | 177Lu-DOTATATE | 5 | 5 | PHEO 0 PGL 5 | 17 | ns | ns | (Pinato et al., 2016) (17) |
| Prospective | 2017 | 177Lu-DOTATATE | 143 | 5 | PHEO 1 PGL 4 | ns | ns | ns | Hamditibar (et al. 2017) (18) |
| Prospective | 2018 | 177Lu-DOTATATE | 200 | 5 | PHEO 2 PGL 3 | 14 | ns | ns | (Garske-Roman et al., 2018) (19) |
| Retrospective | 2018 | 177Lu-DOTATATE | 186 | 12 | ns | 31,4 | ns | ns | (Demirci et al., 2018) (20) |
| Retrospective | 2019 | 177Lu-DOTATATE | 25 | 25 | PHEO 0 PGL 25 | 32 | ns | ns | (Yadav et al., 2019) (21) |
| Retrospective | 2017 | 177Lu-DOTATATE | 22 | 2 | PHEO 0 PGL 2 | 38,5 | 38,5 | ns | (Nastos et al., 2017) ** (22) |
| Retrospective | 2020 | 177Lu-DOTATATE | 15 | 15 | PHEO 5 PGL 10 | Not reached | Not reached | 14 | (Jaiswal et al., 2017) (23) |
| Retrospective | 2017 | 131I-MIBG | 22 | 13 | PHEO 6 PGL 7 | 20,6 | 14,4 | ns | (Nastos et al., 2017) ** |
| Retrospective | 2001 | 131I-MIBG | 6 | 6 | PHEO 5 PGL 1 | 12 | ns | ns | (Hartley et al., 2001) (24) |
| Retrospective | 2001 | 131I-MIBG | 37 | 15 | PHEO 8 PGL 7 | ns | ns | ns | (Mukherjee et al., 2001) (25) |
| Retrospective | 2003 | 131I-MIBG | 25 | 6 | PHEO 0 PGL 6 | 25,8 | ns | ns | (Bomanji et al., 2003) (26) |
| Retrospective | 2003 | 131I-MIBG | 33 | 33 | PHEO 22 PGL 11 | ns | ns | ns | (Safford et al., 2003) (27) |
| Retrospective | 2007 | 131I-MIBG | 19 | 19 | PHEO 13 PGL 6 | 28,5 | ns | ns | (Gedik et al., 2008) (28) |
| Prospective | 2009 | 131I-MIBG | 50 | 50 | PHEO 15 PGL 35 | ns | ns | ns | (Gonias et al., 2009) (29) |
| Retrospective | 2010 | 131I-MIBG | 10 | 10 | PHEO 7 PGL 3 | 17,5 | ns | ns | (Shilkrut et al., 2010) (30) |
| Retrospective | 2011 | 131I-MIBG | 32 | 12 | PHEO 8 PGL 4 | 29 | ns | ns | (Rachh et al., 2011) (31) |
| Retrospective | 2011 | 131I-MIBG | 16 | 6 | PHEO 6 PGL 0 | 48 | ns | 48 | (Szalat et al., 2011) (32) |
| Retrospective | 2012 | 131I-MIBG | 17 | 5 | PHEO 0 PGL 5 | 11,26 | ns | ns | (Fishbein et al., 2012) (33) |
| Retrospective | 2014 | 131I-MIBG | 70 | 48 | PHEO 37 PGL 11 | ns | ns | ns | (Yoshinaga et al., 2014) (34) |
| Prospective | 2006 | 131I-MIBG | 30 | 30 | PHEO 11 PGL 19 | ns | ns | ns | (Fitzgerald et al., 2006) (35) |
| Retrospective | 2013 | 131I-MIBG | 26 | 26 | PHEO 8 PGL 18 | ns | ns | ns | (Wakabayashi et al.2019) (36) |
| Retrospective | 2010 | 131I-MIBG | 12 | 12 | PHEO 4 PGL 8 | 30 | ns | ns | Castellani group 1 before 2001. Low activity (Castellani et al., 2010) (37) |
| Retrospective | 2010 | 131I-MIBG | 16 | 16 | PHEO 11 PGL 5 | 24,92 | ns | ns | Castellani group 2 after 2001. Intermediate activity (Castellani et al., 2010) |

(Continued)

TABLE 2 Continued

| TYPE | Year of publication | TRT | Total patients | Number of PGGL patients | Primary location | PFS (months) | PFS (PGL) | PFS (PHEO) | Reference |
|---------------|---------------------|-----------|----------------|-------------------------|-------------------|--------------|-----------|------------|--------------------------------|
| Retrospective | 2003 | 131I-MIBG | 12 | 12 | PHEO 6 PGL 6 | 32,25 | ns | ns | (Rose et al., 2003) (38) |
| Retrospective | 2015 | 131I-MIBG | 22 | 22 | PHEO 10 PGL 12 | ns | ns | ns | (Rutherford et al., 2015) (39) |
| Retrospective | 2020 | 131I-MIBG | 125 | 125 | PHEO 73 PGL 52 | ns | ns | ns | (Thorpe et al., 2020) (40) |

*Nonspecified. **Study including patients treated with both therapeutic modalities.

TABLE 3 Patient descriptive variables according to the treatment modality. In some variables, the data specified in the papers are scarce.

| Treatment | Total patients PGGL | Primary PGL | Primary PHEO | Mean age (years) | Women/Men | First-line therapy | Second-line therapy | Third-line therapy | Fourth-line therapy and successive | Cumulative activity range (GBq) | Follow up (months) |
|-----------------------|---------------------|----------------|----------------|-------------------|-----------|--------------------|---------------------|--------------------|------------------------------------|---------------------------------|--------------------|
| ¹³¹ I-MIBG | 466 (76.6%) | 216 (68.3%) | 244 (92.2%) | 42.8 (32.5-52) | 205/261 | 36 | 12 | 1 | – | 7.4–39.4 | 28.6 |
| PRRT-Lu | 142 (23.3%) | 100 (31.6%) | 20 (7.5%) | 43.7 (30.9–51) | 52/90 | 33 | 32 | 11 | 7 | 22–29.6 | 31.4 |

than PRRT-Lu when the proportion of PHEOs was low, and PFS was significantly longer with PRRT-Lu when the proportion of PHEOs increased (Table 5). Specifically, with no PHEOs, the mean difference was 5.18 months (95% CI [4.16, 6.06]) in favor of [¹³¹I]MIBG, and it decreased by 1.9 months (95% CI [-2.01, -1.78]) for each 10% increase in the proportion of PHEOs in the sample.

To aid in the interpretation of the interaction model, we also estimated a conditional effects plot, describing the relationship between both treatments, the proportion of PHEOs in the sample, and the estimated PFS (Figure 4).

Discussion

PGGLs have heterogeneous presentation and prognosis. This fact, added to the lack of current guidelines and controlled studies, leads physicians to a pragmatic personalized approach based on the experts' consensus and a benefits–risk balance.

In the [¹³¹I]MIBG series, there is large heterogeneity in the published average cumulative dose (7.4–39.4 GBq), the number of administered cycles (1–12), and PFS (12–32.2 months).




FIGURE 2

Distribution of the administered treatments according to the adrenal or extra-adrenal location of the tumor.




FIGURE 3

PFS estimates from the different studies between 2001 and 2020 with ^{177}Lu Lu-DOTA-TATE (red) and ^{131}I MIBG (blue). Dot sizes are proportional to the sample size of the study.

Most of the ^{131}I MIBG series are retrospective case reports, and there is only one prospective study reported by Gonias et al. (29). This one involved 50 patients with administered activities ranging from 18.2 to 118 GBq, reaching responses in 22% of the cases but at the expense of significant toxicities.

In ^{131}I MIBG, the balance between the benefits and comorbidities of a high-activity treatment remains unclear. Castellani et al. (37) compared two groups of patients using an intermediate vs. low cumulative dose (39.4 vs. 24.1 GBq) and concluded that the most important result using higher activity without reaching myeloablative levels was the shortened time to achieve a significant response. Low-dose treatment results reported by Shilkrut et al. (30) (one to four cycles) with 11.6 GBq obtained a PFS of 17.5 months, and those reported by Rachh et al. (31) (one to five cycles) with 11.8 GBq reached a PFS of 29 months. At the other end, Rose et al. (38) using high therapeutic doses (29.6 GBq) observed a 32.2-month PFS.

Studies with ^{131}I MIBG that specifically classify their results according to adrenal or extra-adrenal primary location are

lacking. Safford et al. (27) observed a trend to decreased survival in those patients with metastatic PGL vs. PHEO (1.8 vs. 4.7 years, respectively), whereas they did not get a statistically significant difference.

As opposed to ^{131}I MIBG, PRRT-Lu published series are quite homogeneous in the cumulative dose that ranges between 22 and 29.6 GBq. However, we found more variability in PFS, ranging between 14 and 39 months. There are only three prospective studies, two of them with five PGGLs and the third one with 20 PGGLs published by Kong et al. (14) Hamiditabar et al. (18), and Garske-RomáVerify that all the equations and special characters are displayed correctly.n et al. (19), respectively. Among the few cases in which the papers specified the therapeutic line, and in contrast to ^{131}I MIBG, most of the PPRT-Lu treatments were not administered as the first line, probably because PRRT-Lu is a more recent treatment, and it is limited to compassionate use in many countries.

The largest PRRT-Lu series, involving 30 patients and published by Zandee et al. (15), observed an overall PFS of 30 months and divided the results according to the adrenal vs.

TABLE 4 Regression model comparing PFS between ^{131}I MIBG and ^{177}Lu Lu-DOTA-TATE treatments.

| Variables | Estimate | Std error | Lower 95 | Upper 95 |
|----------------|----------|-----------|----------|----------|
| Age | -0.327 | 0.031 | -0.39 | -0.27 |
| Year | -0.37 | 0.051 | -0.473 | -0.272 |
| MIBG treatment | -10.066 | 0.824 | -11.722 | -8.504 |
| Prop PHEO | -2.699 | 1.625 | -5.496 | 0.793 |

The variables under study were age, year of publication, ^{131}I MIBG treatment, and proportion of PHEOs.

TABLE 5 Regression model comparing PFS between [¹³¹I]MIBG and [¹⁷⁷Lu]Lu-DOTA-TATE considering that the treatments do not affect equally the adrenal and extra-adrenal primary location.

| Variables | Estimate | Std Error | Lower 95 | Upper 95 |
|--------------------------|----------|-----------|----------|----------|
| Age | -0.447 | 0.008 | -0.463 | -0.429 |
| Year | -0.285 | 0.012 | -0.311 | -0.266 |
| MIBG treatment | 5.181 | 0.484 | 4.157 | 6.063 |
| Prop PHEO | 14.009 | 0.559 | 12.824 | 15.018 |
| MIBG treatment/prop PHEO | -19.046 | 0.575 | -20.126 | -17.845 |

The variables under study were age, year of publication, treatment with [¹³¹I]MIBG, proportion of PHEO, and proportion of PHEO treated with [¹³¹I]MIBG.

extra-adrenal location. They obtained a 30-month PFS in 10 patients with parasympathetic PGL, 13-month PFS in the sympathetic PGLs, and 8–10–14 months in the three PHEO patients.

When considering TRT, the extension and intensity uptake of the targets should be evaluated both on [¹²³I]MIBG and SSTR scans and compared with the whole tumor burden noted on CT/MRI and/or ¹⁸F-FDG PET/CT. Less than 50% of metastatic PGLs display [¹²³I]MIBG avidity. On the other hand, SSTRs are overexpressed in more than 80% of neuroendocrine tumors. In many cases, differences between them are obvious (normally due to the lack of uptake in one of them). But if both scans reveal a similar uptake and extent of disease, the choice becomes complicated. There is only one study comparing the therapeutic outcomes of both TRT modalities published by Nastos et al. (22). They used [¹³¹I]MIBG or a combination of [¹³¹I]MIBG and PRRT (most PRRT-Y) and observed that patients treated with PRRT had an increased PFS and response to treatment compared with [¹³¹I]MIBG ($p < 0.05$) but with no significant difference in overall

survival. When comparing only PGL patients, Nastos et al. (22) observed that response to treatment, overall survival, and PFS were significantly higher in the PRRT group. In this study, the effectiveness of PRRT in PHEOs could not be assessed because, of the seven patients with this primary location, only one was treated with PRRT.

Combining both [¹³¹I]MIBG and PRRT could be a future alternative especially in those patients with aggressive disease. In this line, Bushnell et al. (41) published a phase 1 clinical trial combining treatment with PRRT-Y and [¹³¹I]MIBG with pretreatment tumor dosimetry in three patients with neuroendocrine tumors, with an adequate safety margin (41).

There are no large prospective studies comparing [¹³¹I]MIBG and PRRT-Lu, if both are available and similarly suitable. In the Jha et al. report (8), a group of international experts facing this obstacle recommends considering, in addition to nuclear medicine scans, the patient's profile and the tumor characteristics and prognostic features to guide the decision. If mismatched uptakes are found in nuclear medicine scans, these




FIGURE 4

Conditional effects plot depicting the interaction between treatment and proportion of PHEOs in the sample regarding the estimated PFS in months.

authors suggest a combined sequence of both TRTs to cover all the lesions. Unfortunately, there are still not enough prognostic data available, based on these variables, to support which TRT to choose (Figure 5).


We found several limitations in this review. Firstly, the samples of both treatments were not homogeneous, and, in 28 patients, the primary location of the tumor was not identified, so the means obtained could not be fully compared. In most of the reports, PGGLs were a small number of patients included in larger neuroendocrine tumor series. The majority of the studies did not provide PFS according to the location, and for this reason, linear regression statistical models had to be carried out. Within PHEOs, only 20 patients were treated with PRRT-Lu. However, they were extracted from five different reports, so it is unlikely that the results were due to the bias of a single study. In neuroendocrine tumors, a correlation has been observed between the high uptake of lesions in SSTR scans and the density of SSTR2 expression in biopsies. A higher SSTR density may imply a better response to PRRT, so patients who strongly express this target are preferentially selected for PRRT rather than for ^{131}I MIBG. Given that patients with high SSTR expression are expected to have a better response to PRRT, this fact may imply a patient selection bias due to falsely improving PRRT results (42). Additionally, in the ^{131}I MIBG group, we observed significant heterogeneity in the administered cumulative activities, although it showed a low impact on the PFS, and the reviewed reports did not distinguish whether all

patients were candidates for both treatments. Most of the reviewed articles did not report to which cluster the genetic syndromes belonged, so a correlation could not be done between response to treatment and genetic status.

Despite these limitations, we observed PFS differences globally and according to the primary location and the therapeutic modality administered. The unweighted means and the statistical model weighted to represent the sample size showed greater PFS in the PRRT-Lu group. The model weighted to the proportion of PHEOs showed that the adrenal location responded better to treatment with PRRT-Lu. In patients expressing both targets, this fact could support the selection of PRRT-Lu as a first option, especially in PHEOs, and if a TRT sequence with both therapies is considered, the first one to be administered.

Conclusion

Methodology, procedure, and PFS from the different reviewed studies about TRT in metastatic/inoperable PGGLs are quite heterogeneous. PRRT-Lu seems to have overall better results, particularly in PHEOs, but this question should probably be reassessed shortly with the arrival of HAS ^{131}I MIBG. Although the reports present several limitations, our results make it clear that tumor primary location is a factor to consider in therapy planning.



These results should open the window to prospective trials comparing or sequencing [¹³¹I]MIBG and ¹⁷⁷Lu-DOTATATE. Studies aimed at the evaluation of both treatments are necessary, with special attention to the primary location and the line of therapy, reaching similar administered cumulative activities and correlating them with the genetic cluster within each group.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary Material. Further inquiries can be directed to the corresponding author.

Author contributions

SP-W and MO-G reviewed the articles and conducted the study. PB-A and JM-T formally and scientifically reviewed the manuscript. All authors contributed to the article and approved the submitted version.

References

1. Neumann HPH, Young WF, Eng C. Pheochromocytoma and paraganglioma. *New N Engl J Med* (2019) 381(6):552–65. doi: 10.1056/NEJMra1806651
2. Cascón A, Remacha L, Calsina B, Robledo M. Pheochromocytomas and paragangliomas: Bypassing cellular respiration. *Cancers* (2019) 16(5):683. doi: 10.3390/cancers11050683
3. Taieb D, Hicks RJ, Hindie E, Guillet BA, Avram AM, Ghedini P, et al. European Association of nuclear medicine practice Guideline/Society of nuclear medicine and molecular imaging procedure standard 2019 for radionuclide imaging of phaeochromocytoma and paraganglioma. *Eur J Nucl Med Mol Imaging* (2019) 46(10):2112–37. doi: 10.1007/s00259-019-04398-1
4. Nölting S, Ullrich M, Pietzsch J, Ziegler C, Eisenhofer G, Grossman A, et al. Current management of Pheochromocytoma/Paraganglioma: A guide for the practicing clinician in the era of precision medicine. *Cancers* (2019) 11(10):1505. doi: 10.3390/cancers11101505
5. Crona J, Taieb D, Pacak K. New perspectives on pheochromocytoma and paraganglioma: Toward a molecular classification. *Endocr Rev* (2017) 38(6):489–515. doi: 10.1210/er.2017-00062
6. Turchini J, Cheung VKY, Tischler AS, De Krijger RR, Gill AJ. Pathology and genetics of phaeochromocytoma and paraganglioma. *Histopathology* (2018) 72(1):97–105. doi: 10.1111/his.13402
7. De Filipo G, Maggi M, Mannelli M, Canu L. Management and outcome of metastatic pheochromocytomas/paragangliomas: an overview. *J Endocrinol Invest* (2021) 44(1):15–25. doi: 10.1007/s40618-020-01344-z
8. Jha A, Taieb D, Carrasquillo JA, Pryma DA, Patel M, Millo C, et al. High-specific-activity [¹³¹I]MIBG vs [¹⁷⁷Lu]LU-DOTA-TATE targeted radionuclide therapy for metastatic pheochromocytoma and paraganglioma. *Clin Cancer Res* (2021) 27:2989–95. doi: 10.1158/1078-0432.CCR-20-3703
9. Prado-Wohlwend S, Del Olmo-García MI, Bello-Arques P, Merino-Torres JF. [¹⁷⁷Lu]Lu-DOTA-TATE and [¹³¹I]MIBG phenotypic imaging-based therapy in Metastatic/Inoperable pheochromocytomas and paragangliomas: Comparative results in a single center. *Front Endocrinol* (2022) 7:778322(13). doi: 10.3389/fendo.2022.778322
10. Hirmas N, Jadaan R, Al-Ibraheem A. Peptide receptor radionuclide therapy and the treatment of gastroentero-pancreatic neuroendocrine tumors: Current findings and future perspectives. *Nucl Med Mol Imaging* (2018) 52(3):190–9. doi: 10.1007/s13139-018-0517-x
11. Basu S, Parghane RV, Kamaldeep, Chakrabarty S. Peptide receptor radionuclide therapy of neuroendocrine tumors. *Semin Nucl Med* (2020) 50(5):447–64. doi: 10.1053/j.semnuclmed.2020.05.004
12. Pryma DA, Chin BB, Noto RB, Dillon JS, Perkins S, Solnes L, et al. Efficacy and safety of high-Specific-Activity ¹³¹I-MIBG therapy in patients with advanced pheochromocytoma or paraganglioma. *J Nucl Med* (2019) 60(5):623–30. doi: 10.2967/jnumed.118.217463
13. Fassnacht M, Assie G, Baudin E, Eisenhofer G, de la Fouchardiere C, Haak HR. Adrenocortical carcinomas and malignant phaeochromocytomas: ESMOeURACAN clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* (2020) 31(11):1476–90. doi: 10.1016/j.annonc.2020.08.2099
14. Kong G, Grozinsky-Glasberg S, Hofman MS, Callahan J, Meirovitz A, Maimon O, et al. Efficacy of peptide receptor radionuclide therapy for functional metastatic paraganglioma and pheochromocytoma. *J Clin Endocrinol Metab* (2017) 102(9):3278–87. doi: 10.1210/jc.2017-00816
15. Zandee WT, Feelders RA, Smit Duijzenkunst DA, Hofland J, Metselaar RM, Oldenburg RA, et al. Treatment of inoperable or metastatic paragangliomas and pheochromocytomas with peptide receptor radionuclide therapy using ¹⁷⁷Lu-DOTATATE. *Eur J Endocrinol* (2019) 181(1):45–53. doi: 10.1530/EJE-18-0901
16. Van Essen M, Krenning EP, Kooij PP, Bakker WH, Feelders RA, de Herder WW, et al. Effects of therapy with [¹⁷⁷Lu-DOTA0, Tyr3]Octreotate in patients with paraganglioma, meningioma, small cell lung carcinoma, and melanoma. *J Nucl Med* (2006) 47(10):1599–606.
17. Pinato DJ, Black JRM, Ramaswami R, Tan TM, Adjogatse D, Sharma RR. Peptide receptor radionuclide therapy for metastatic paragangliomas. *Med Oncol* (2016) 33(5):47. doi: 10.1007/s12032-016-0737-9
18. Hamiditabar M, Ali M, Roys J, Wolin EM, O'Dorisio TM, Ranganathan D, et al. Peptide receptor radionuclide therapy with ¹⁷⁷Lu-octreotate in patients with somatostatin receptor expressing neuroendocrine tumors: Six years' assessment. *Clin Nucl Med* (2017) 42(6):436–43. doi: 10.1097/RLU.0000000000001629
19. Garske-Román U, Sandström M, Fröss Baron K, Lundin L, Hellman P, Welin S, et al. Prospective observational study of ¹⁷⁷Lu-DOTA-octreotate therapy in 200 patients with advanced metastasized neuroendocrine tumours (NETs): feasibility and impact of a dosimetry-guided study protocol on outcome and toxicity. *Eur J Nucl Med Mol Imaging* (2018) 45(6):970–88. doi: 10.1007/s00259-018-3945-z
20. Demirci E, Kabasakal L, Toklu T, Ocak M, Şahi O, Alan-Selcuk N, et al. ¹⁷⁷Lu-DOTATATE therapy in patients with neuroendocrine tumours including

Funding

Funding for conducting the review has been received from the La Fe Hospital Research Foundation.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- high-grade (WHO G3) neuroendocrine tumours: response to treatment and long-term survival update. *Nucl Med Commun* (2018) 39(8):789–96. doi: 10.1097/MNM.0000000000000874
21. Yadav M, Ballal S, Bal C. Concomitant [177Lu]LU-DOTA-TATE and capecitabine therapy in malignant paragangliomas. *EJNMMI* (2019) 9(1):1–10. doi: 10.1186/s13550-019-0484-y
22. Nastos K, Cheung VTF, Toumpanakis C, Navalkisoor S, Quigley A, Caplin M, et al. Peptide receptor radionuclide treatment and (131I)-MIBG in the management of patients with metastatic/progressive pheochromocytomas and paragangliomas. *J Surg Oncol* (2017) 115(4):425. doi: 10.1002/jso.24553
23. Jaiswal SK, Sarathi V, Memon SS, Garg R, Malhotra G, Verma P, et al. 177Lu-DOTATATE therapy in metastatic/inoperable pheochromocytoma-paraganglioma. *Endocr Connect* (2020) 9(9):864–73. doi: 10.1530/EC-20-0292
24. Hartley A, Spooner D, Brunt AM. Management of malignant pheochromocytoma: A retrospective review of the use of MIBG and chemotherapy in the West Midlands. *Clin Oncol* (2001) 13(5):361–6. doi: 10.1053/clon.2001.9290
25. Mukherjee JJ, Kaltsas GA, Islam N, Plowman PN, Foley R, Hikmat J, et al. Treatment of metastatic carcinoid tumours, pheochromocytoma, paraganglioma and medullary carcinoma of the thyroid with (131I)-meta-iodobenzylguanidine [(131I)-mIBG]. *Clin Endocrinol* (2001) 55(1):47–60. doi: 10.1046/j.1365-2265.2001.01309.x
26. Bomanji JB, Wong W, Gaze MN, Cassoni A, Waddington W, Solano J, et al. Treatment of neuroendocrine tumours in adults with 131I-MIBG therapy. *Clin Oncol* (2003) 15(4):193–8. doi: 10.1016/s0936-6555(02)00273-x
27. Safford SD, Coleman RE, Gockerman JP, Moore J, Feldman JM, Leight GS, et al. Iodine-131 metaiodobenzylguanidine is an effective treatment for malignant pheochromocytoma and paraganglioma. *Surgery* (2003) 134(6):956–62; discussion 962–3. doi: 10.1016/s0039-6060(03)00426-4
28. Gedik GK, Hoefnagel CA, Bais E, Valdes Olmos RA. [sup.131I]-MIBG therapy in metastatic pheochromocytoma and paraganglioma. *Eur J Nucl Med Mol Imaging* (2008) 35(4):725–33. doi: 10.1007/s00259-007-0652-6
29. Gonias S, Goldsby R, Matthay KK, Hawkins R, Price D, Huberty J, et al. Phase II study of high-dose [131I]Metaiodobenzylguanidine therapy for patients with metastatic pheochromocytoma and paraganglioma. *J Clin Oncol* (2009) 27(25):4162–8. doi: 10.1200/JCO.2008.21.3496
30. Shilkrut M, Bar-Deroma R, Bar-Sela G, Berniger A, Kuten A. Low-dose iodine-131 metaiodobenzylguanidine therapy for patients with malignant pheochromocytoma and paraganglioma single center experience. *Am J Clin Oncol* (2010) 33(1):79–82. doi: 10.1097/COC.0b013e31819e2c28
31. Rachh S, Abhyankar S, Basu S. [131I]Metaiodobenzylguanidine therapy in neural crest tumors: varying outcome in different histopathologies. *Nucl Med Commun* (2011) 32(12):1201–10. doi: 10.1097/MNM.0b013e32834bad97
32. Szalat A, Fraenkel M, Doviner V, Salmon A, Gross D. Malignant pheochromocytoma: predictive factors of malignancy and clinical course in 16 patients at a single tertiary medical center. *Endocrine* (2011) 39(2):160–6. doi: 10.1007/s12020-010-9422-5
33. Fishbein L, Bonner L, Torigian DA, Nathanson KL, Cohen DL, Pryma D, et al. External beam radiation therapy (EBRT) for patients with malignant pheochromocytoma and non-head and -neck paraganglioma: Combination with 131I-MIBG. *Horm Metab Res* (2012) 44(5):405–10. doi: 10.1055/s-0032-1308992
34. Yoshinaga K, Oriuchi N, Wakabayashi H, Tomiyama Y, Jinguji M, Higuchi T, et al. Effects and safety of ¹³¹I-metaiodobenzylguanidine (MIBG) radiotherapy in malignant neuroendocrine tumors: results from a multicenter observational registry. *Endocr J* (2014) 61(12):1171–80. doi: 10.1507/endocrj.EJ14-0211
35. Fitzgerald PA, Goldsby RE, Huberty JP, Price DC, Hawkins RA, Veatch JJ, et al. Malignant pheochromocytomas and paragangliomas: a phase II study of therapy with high-dose 131I-metaiodobenzylguanidine (131I-MIBG). *Ann N Y Acad Sci* (2006) 1073:465–90. doi: 10.1196/annals.1353.050
36. Wakabayashi H, Taki J, Inaki A, Nakamura A, Kayano D, Fukuoka M, et al. Prognostic values of initial responses to low-dose 131I-MIBG therapy in patients with malignant pheochromocytoma and paraganglioma. *Ann Nucl Med* (2013) 27(9):839–46. doi: 10.1007/s12149-013-0755-z
37. Castellani MR, Seghezzi S, Chiesa C, Aliberti GL, Maccauro M, Seregni E, et al. (131I)-MIBG treatment of pheochromocytoma: low versus intermediate activity regimens of therapy. *Q J Nucl Med Mol Imaging* (2010) 54(1):100–13.
38. Rose B, Matthay KK, Price D, Huberty J, Klencke B, Norton JA, et al. High-dose 131I-metaiodobenzylguanidine therapy for 12 patients with malignant pheochromocytoma. *Cancer* (2003) 98(2):239–48. doi: 10.1002/cncr.11518
39. Rutherford MA, Rankin AJ, Yates TM, Mark PB, Perry CG, Reed NS, et al. Management of metastatic pheochromocytoma and paraganglioma: use of iodine-131-meta-iodobenzylguanidine therapy in a tertiary referral centre. *QJM* (2015) 108(5):361–8. doi: 10.1093/qjmed/hcu208
40. Thorpe MP, Kane A, Zhu J, Morse MA, Wong T, Borges-Neto S. Long-term outcomes of 125 patients with metastatic pheochromocytoma or paraganglioma treated with 131 I-MIBG. *J Clin Endocrinol Metab* (2020) 105(3):dgz074. doi: 10.1210/clinem/dgz074
41. Bushnell DL, Bodeker KL, O'dorisio TM, Madsen MT, Menda, Graves SA, et al. Addition of 131I MIBG to PRRT (90Y DOTATOC) for personalized treatment of selected patients with neuroendocrine tumors. *J Nucl Med* (2021) 62(9):1274–7. doi: 10.2967/jnumed.120.254987
42. Kratochwil C, Stefanova M, Mavriopoulou E, Holland-Letz T, Dimitrakopoulou-Strauss A, Afshar-Oromieh A, et al. SUV Of [68Ga] DOTATOC-PET/CT predicts response probability of PRRT in neuroendocrine tumors. *Mol Imaging Biol* (2014) 17(3):313–8. doi: 10.1007/s11307-014-0795-3