

OPEN ACCESS

EDITED AND REVIEWED BY
Roberto Scicali,
University of Catania, Italy

*CORRESPONDENCE

Ping Shuai

✉ shuaiping@med.uestc.edu.cn

Junlin Zhang

✉ zhangjunlin_hx@163.com

Zhengwei Wan

✉ 18715799366@163.com

[†]These authors share first authorship

RECEIVED 16 May 2024

ACCEPTED 21 May 2024

PUBLISHED 27 May 2024

CITATION

Hou Q, Zhang H, Zhang R, Li B, Li L, Li D, Wang X, Liu Y, Wan Z, Zhang J and Shuai P (2024) Corrigendum: Relationship between the longitudinal trajectory of the triglyceride-glucose index and the development of CKD: an 8-year retrospective longitudinal cohort study.

Front. Endocrinol. 15:1433515.

doi: 10.3389/fendo.2024.1433515

COPYRIGHT

© 2024 Hou, Zhang, Zhang, Li, Li, Li, Wang, Liu, Wan, Zhang and Shuai. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Relationship between the longitudinal trajectory of the triglyceride-glucose index and the development of CKD: an 8-year retrospective longitudinal cohort study

Qinchuan Hou^{1,2†}, Huiwang Zhang^{1,2†}, Rui Zhang^{1,2†}, Binghong Li^{1,3†}, Lei Li^{1,3}, Dongyu Li^{1,3}, Xian Wang¹, Yuping Liu¹, Zhengwei Wan^{1*}, Junlin Zhang^{1*} and Ping Shuai^{1,2,3*} 

¹Health Management Center & Health Management Research Institute, Sichuan Provincial People's Hospital, Chengdu, China, ²School of Public Health, Southwest Medical University, Luzhou, China, ³School of Medicine, University of Electronic Science and Technology of China, Chengdu, China

KEYWORDS

triglyceride-glucose index, change trajectory, chronic kidney disease, health check-up population, cohort study

A Corrigendum on

Relationship between the longitudinal trajectory of the triglyceride-glucose index and the development of CKD: an 8-year retrospective longitudinal cohort study

By Hou Q, Zhang H, Zhang R, Li B, Li L, Li D, Wang X, Liu Y, Wan Z, Zhang J and Shuai P (2024). *Front. Endocrinol.* 15:1376166. doi: 10.3389/fendo.2024.1376166

In the published article, there was an error in [Figure 4](#) as published. The red color indicates the high-level group and the green color indicates the medium-level stable group, which is now labeled the other way round. The corrected [Figure 4](#) and its caption appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.




FIGURE 4
The TyG index trajectory groups in the 8-year follow-up. TyG, the triglyceride-glucose index.