

OPEN ACCESS

EDITED AND REVIEWED BY
Rossella Cannarella,
Cleveland Clinic, United States

*CORRESPONDENCE Iva Arato ☑ iva.arato@libero.it

[†]These authors have contributed equally to this work and share last authorship

RECEIVED 08 May 2025 ACCEPTED 30 May 2025 PUBLISHED 19 June 2025

CITATION

Arato I, Giovagnoli S, Di Michele A, Bellucci C, Lilli C, Aglietti MC, Bartolini D, Gambelunghe A, Muzi G, Calvitti M, Eugeni E, Gaggia F, Baroni T, Mancuso F and Luca G (2025) Corrigendum: Nickel oxide nanoparticles exposure as a risk factor for male infertility: "In vitro" effects on porcine pre-pubertal Sertoli cells. Front. Endocrinol. 16:1625122. doi: 10.3389/fendo.2025.1625122

COPYRIGHT

© 2025 Arato, Giovagnoli, Di Michele, Bellucci, Lilli, Aglietti, Bartolini, Gambelunghe, Muzi, Calvitti, Eugeni, Gaggia, Baroni, Mancuso and Luca. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC By). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Nickel oxide nanoparticles exposure as a risk factor for male infertility: "In vitro" effects on porcine pre-pubertal Sertoli cells

Iva Arato^{1*}, Stefano Giovagnoli², Alessandro Di Michele³, Catia Bellucci¹, Cinzia Lilli¹, Maria Chiara Aglietti¹, Desirée Bartolini¹, Angela Gambelunghe¹, Giacomo Muzi¹, Mario Calvitti¹, Elena Eugeni¹, Francesco Gaggia⁴, Tiziano Baroni¹, Francesca Mancuso^{1†} and Giovanni Luca^{1,5,6†}

¹Department of Medicine and Surgery, University of Perugia, Perugia, Italy, ²Department of Pharmaceutical Sciences, University of Perugia, Perugia, Italy, ³Department of Physics and Geology, University of Perugia, Perugia, Italy, ⁴Internal Medicine Endocrine and Metabolic Sciences Unit, Santa Maria della Misericordia Hospital of Perugia, Perugia, Italy, ⁵International Biotechnological Center for Endocrine, Metabolic and Embryo-Reproductive Translational Research (CIRTEMER), Department of Medicine and Surgery, University of Perugia, Perugia, Italy, ⁶Division of Medical Andrology and Endocrinology of Reproduction, Saint Mary Hospital, Terni, Italy

KEYWORDS

Sertoli cells, nickel oxide nanoparticles, ROS, comet, MAPK pathways

A Corrigendum on

Nickel oxide nanoparticles exposure as a risk factor for male infertility: "In vitro" effects on porcine pre-pubertal Sertoli cells

By Arato I, Giovagnoli S, Di Michele A, Bellucci C, Lilli C, Aglietti MC, Bartolini D, Gambelunghe A, Muzi G, Calvitti M, Eugeni E, Gaggia F, Baroni T, Mancuso F and Luca G (2023). Front. Endocrinol. 14:1063916. doi: 10.3389/fendo.2023.1063916

In the published article, there was an error in Figure 6 panel A as published. Duplicate image for four loading controls that were used by our group one year before in Figure 5A of Mancuso et al., 2022, where the conditions are different (TiO2-NPs). The corrected Figure 6 panel A and its caption "Caspase-3 Evaluation by WB analysis. (A) Immunoblots of caspase-3 p35, p19, and p17 in SCs at 24h and 1, 2, and 3 weeks of incubation with NiO-NPs at 1 and 5mg/ml" appear below.


The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Arato et al. 10.3389/fendo.2025.1625122

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.



Caspase-3 Evaluation by WB analysis. **(A)** Immunoblots of caspase-3 p35, p19, and p17 in SCs at 24h and 1, 2, and 3 weeks of incubation with NiO-NPs at 1 and 5mg/ml. Densitometric analysis of the protein bands of caspase-3 p35 **(B)**, p19 **(C)**, and p17 **(D)** in SCs at 24 h and 1, 2, and 3 weeks of incubation with NiO-NPs 1 and 5 μ g/ml. Data represent the mean \pm SEM (**p < 0.001 vs. 0 NiO-NPs of three independent experiments, each performed in triplicate).