



OPEN ACCESS

EDITED AND REVIEWED BY
Jules Zhang-Yin,
Vivalia, Belgium

*CORRESPONDENCE

Isabella Chiardi
✉ isabellachiardi@gmail.com
Pierpaolo Trimboli
✉ pierpaolo.trimboli@usi.ch

RECEIVED 11 July 2025
ACCEPTED 31 July 2025
PUBLISHED 12 August 2025

CITATION

Chiardi I, Makovac P, Leoncini A, Forte F, Rotondi M and Trimboli P (2025) Correction: Ultrasound detection of normal parathyroid glands: detection rate, topographic anatomy and the role of underlying thyroid disease. *Front. Endocrinol.* 16:1664368. doi: 10.3389/fendo.2025.1664368

COPYRIGHT

© 2025 Chiardi, Makovac, Leoncini, Forte, Rotondi and Trimboli. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Correction: Ultrasound detection of normal parathyroid glands: detection rate, topographic anatomy and the role of underlying thyroid disease

Isabella Chiardi ^{1,2*}, Petra Makovac ³, Andrea Leoncini ⁴, Flavio Forte ⁵, Mario Rotondi ^{1,6} and Pierpaolo Trimboli ^{2,7*}

¹Department of Internal Medicine and Therapeutics, University of Pavia, Pavia, Italy, ²Thyroid Unit, Clinic for Endocrinology and Diabetology, Ente Ospedaliero Cantonale (EOC), Bellinzona, Switzerland, ³Service of Surgery, Ospedale Regionale di Mendrisio, Ente Ospedaliero Cantonale (EOC), Mendrisio, Switzerland, ⁴Clinic for Radiology, Imaging Institute of Southern Switzerland, Ente Ospedaliero Cantonale (EOC), Bellinzona, Switzerland, ⁵Urologic Division, Vannini Hospital, Rome, Italy, ⁶Istituti Clinici Scientifici Maugeri IRCCS, Unit of Endocrinology and Metabolism, Laboratory for Endocrine Disruptors, Pavia, Italy, ⁷Faculty of Biomedical Sciences, Università Della Svizzera Italiana (USI), Lugano, Switzerland

KEYWORDS

US, normal parathyroids, parathyroid detection, parathyroid imaging, neck ultrasound

A Correction on

Ultrasound detection of normal parathyroid glands: detection rate, topographic anatomy, and the role of underlying thyroid disease

By Chiardi I, Makovac P, Leoncini A, Forte F, Rotondi M and Trimboli P (2025) *Front. Endocrinol.* 16:1595940. doi: 10.3389/fendo.2025.1595940

In the published article, there was an error in the **Discussion** section regarding the citation of the article by Marchand et al. Our article incorrectly referred to “64 intra-thyroid parathyroids” instead of “infra-thyroid.” Accordingly, the authors have removed this citation from the relevant passage (while keeping it elsewhere as it is cited multiple times), since the citation was no longer appropriate.

A correction has been made to the section **Discussion**, Paragraph 2:

“In our study, PTGs were visible in 51 out of 113 patients, representing a detection rate of 45%. This is lower than detection rates reported in some other studies (2, 3, 6, 7), which may be partially explained by our decision to exclude intra-thyroid PTGs to minimize bias. This exclusion was necessary because differentiating a PTG from thyroid tissue in this region is nearly impossible and would require FNA for confirmation. Table 3 provides a comparison of various studies in literature (2, 3, 6–9), highlighting differences in PTG detection rates, gland sizes, thyroid disorders of patients, and PTG locations.”

The original version of this article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.