



# Changes in the Microbial Community Structure and Soil Chemical Properties of Vertisols Under Different Cropping Systems in Northern China

Xiuli Song<sup>1,2†</sup>, Bo Tao<sup>1\*</sup>, Jing Guo<sup>1†</sup>, Jingjing Li<sup>1†</sup> and Guofeng Chen<sup>2</sup>

<sup>1</sup> Agricultural College, Northeast Agricultural University, Harbin, China, <sup>2</sup> Qiqihar Branch of Heilongjiang Academy of Agricultural Sciences, Qiqihar, China

## OPEN ACCESS

### Edited by:

Sudhakar Srivastava,  
Banaras Hindu University, India

### Reviewed by:

Saurabh Yadav,  
Hemwati Nandan Bahuguna Garhwal  
University, India  
Arnab Majumdar,  
Indian Institute of Science Education  
and Research Kolkata, India

### \*Correspondence:

Bo Tao  
botaol@163.com

<sup>†</sup>These authors have contributed  
equally to this work

### Specialty section:

This article was submitted to  
Agroecology and Ecosystem Services,  
a section of the journal  
Frontiers in Environmental Science

**Received:** 02 May 2018

**Accepted:** 24 October 2018

**Published:** 13 November 2018

### Citation:

Song X, Tao B, Guo J, Li J and  
Chen G (2018) Changes in the  
Microbial Community Structure and  
Soil Chemical Properties of Vertisols  
Under Different Cropping Systems in  
Northern China.  
*Front. Environ. Sci.* 6:132.  
doi: 10.3389/fenvs.2018.00132

The predominant cropping system and management practices play an important role in soil physico-chemical properties and microbiome composition and diversity. This study analyzed the changes in soil fertility and the microbial community in four soybean-based cropping systems over 12 years. Amplification subsequencing techniques were used to compare soil community structures among the systems and identify significant positive and negative fertility factors. Soybean cropping favored the accumulation of OM and the available N, K DTPA Fe, Mn, Zn and Cu contents in soil but not fixed available P. The WS and CS cropping systems were conducive to the fixed available P, but they consumed OM, DTPA K, and Zn. The SC exhibited the lowest soil bacterial and archaeal abundance and diversity but high fungal abundance and diversity. The dominant Proteobacteria in the SC were significantly positively correlated with soil DTPA Fe and Mn. The dominant Actinobacteria were positively correlated with available P, DTPA Cu and Mn. The FS cropping system contained 764 unique bacterial species, 724 unique fungal species and the highest relative abundance of Protista. The FS had high microbial diversity, with high relative abundances of Bacteroidetes and Zygomycota and a significantly lower relative abundance of Actinobacteria. The Bacteroidetes were significantly correlated with available N, OM and DTPA Fe and negatively correlated with available P and DTPA Cu. Zygomycota was negatively correlated with available P and DTPA Cu. In the CS and WS, the soil bacterial abundance and diversity were moderate. The dominant Acidobacteria was significantly negatively correlated with soil DTPA, Fe and Mn. The CS exhibited the lowest fungal abundance and diversity. Furthermore, the relative abundance of Ascomycota was significantly improved in the WS and significantly positively correlated with available P and DTPA Cu. Decreases in the available P, K, DTPA Cu, and Mn of Vertisols greatly affect microbial community structure, and these nutrients regulate bacterial and fungal abundance and diversity. Compared to the SC, the FS, WS, and CS had more balanced soil fertility and microbial stability, but diverse cropping systems are most conducive to soil productivity. These findings are of great relevance for protecting the ecological environment.

**Keywords:** microbial community, microbial diversity, soybean, high-throughput sequencing, 16S rRNA gene, soil

## INTRODUCTION

The microbial community in soil drives important nutrient cycling processes that can influence the soil nutritional profile, crop productivity, and environmental sustainability. Agricultural management affects the chemical, physical, and biological processes in the soil and triggers many changes in the microbial community structure (Fierer et al., 2012; Zhang et al., 2016). The abnormal accumulation or excessive consumption of soil nutrients can result in the rapid multiplication of pathogenic microorganisms, imbalances in the soil microbial population structure, and decreased crop yield and quality (Xing et al., 2011).

Continuous monocultures reduce soil microbial biodiversity and alter the microbial community structure, which decreases its buffering capability to tolerate biotic and abiotic stress, thus increasing the incidence of disease and subsequently affecting plant health (Wang et al., 2011; Mo et al., 2016; Xiong et al., 2016). Continuous cropping could reduce the quantity of bacteria in the soil and increase the relative proportion of fungi. Soil fungal diversity was found to be a good and sensitive indicator of soil fertility (Hobbs et al., 2008; Fierer et al., 2012). Reduced tillage is a sustainable plant management system that can lead to increased surface microbial activity and biomass and protect soil, water, and air quality as well as biodiversity (Tian et al., 2015; Wang et al., 2015; Benitez et al., 2017). Anaerobic bacteria, such as Clostridia and other dominant anaerobic methane-producing archaea, were found in only no-till systems (Luo et al., 2016). However, crop rotation can also modify soil mass and fertility factors and thus influence microbial community structure (Benitez et al., 2017). Microbial diversity is reportedly higher in plots in which oat and maize are grown (Fierer et al., 2012). The rotation of corn and soybeans was found to have a great effect on the microbial community structure of rhizosphere soil microbial groups, with greater impacts being observed on fungi than on bacteria (Benitez et al., 2017). In a plant system containing soy, combining the appropriate crop management techniques is necessary for sustainable production (Fierer et al., 2009). Proteobacteria, Acidobacteria, and Actinobacteria were the main phyla in soybean continuous cropping (SC), soybean-corn, and corn-soybean rotation systems in all soil samples across succession. With the accumulation of soil organic matter (OM) and nutrients, the relative abundance of Actinobacteria decreased, while the relative abundance of *Proteobacteria* increased (Zeng et al., 2017).

Lu et al. (2011) found that increasing the crop rotation of legumes could release nutrients more quickly, leading to higher nitrogen (N) and potassium (K) contents and lower available phosphorus (P) content. Fierer et al. (2009) found that the P, total N, and mineral N contents were significantly improved on the soil surface by no-tillage management systems relative to traditional farming. Rotation could also increase the amount of OM and N in soil, and increasing N enriches the bacterial microorganism composition. In previous studies, microbial communities were significantly correlated with the OM and available N, and external carbon (C), and N additions to a system significantly influenced the microbial community structure (Fierer et al., 2012). However, research on the interactions

among different cropping systems, soil fertility and microbial communities, especially on the effects of mineral elements on microbial communities in soil, is rare. Thus, the potential biological control of crop plants is of great relevance for reducing the amount of fertilizer used (Cai et al., 2015).

High-throughput sequencing provides a useful means to understand the soil microbial community structure and microbial diversity. The microbial community structure is influenced by the physical and chemical properties of the soil (Xun et al., 2015; Cai et al., 2017) as well as by the species and even the genotype of the host plant (Ofek et al., 2014). Heilongjiang Province is the main soybean production area in northern China. From 1999 to 2016, the average soybean production accounted for 36.21% of the national average, while Nenjiang is the main production area in Heilongjiang Province. Therefore, in this study, high-throughput sequencing (i.e., Illumina MiSeq) was conducted to analyze the microorganism community structure and diversity as well as the interactions of the community with soil nutritional profiling under different cropping systems, i.e., the fallow-soybean cropping system (FS), corn-soybean rotation cropping system (CS), wheat-soybean rotation cropping system (WS), and soybean continuous cropping system (SC), across multiple years. The objectives were to (1) identify the appropriate cropping system for maintaining soil fertility, (2) determine the appropriate agricultural cropping system for protecting the soil microbial community, and (3) identify how to enhance the microbial community structure according to the nutritional preferences under different cropping systems. The results of this work will provide a foundation for the regulation of soil nutritional profiling and microorganism community structure, guide cropping system decisions, and protect soil ecology.

## MATERIALS AND METHODS

### Description of the Experimental Field

This study was based on soil samples collected during a 12-year experiment conducted in the experimental ground of the 5th squadron of the fourth field of the Nenjiang River in northern China, known as China's soybean country. The soil type belongs to the Vertisol class according to the American soil classification system (Soil Survey Staff, 2014). This site has a temperate monsoon climate and the following geographical coordinates: 125°41' E longitude and 49° N latitude. The local annual mean temperature is 0.8 to 1.4°C, and the average rainfall is 480 to 512 mm. The winter is cold, long, and dry, with a minimum temperature of -47.3 to -43.7°C. The summer is hot and rainy, with a maximum temperature of 33.9 to 37.4°C. The soil is classified as black, and the region is the main production area of wheat and soybean in China.

### Experimental Design and Treatments

Soybean was planted in experimental plots before March 2005, and rotation trials began in April 2005. Four cropping systems were arranged in a randomized block design: fallow-soybean cropping system (FS), corn-soybean rotation cropping system (CS), wheat-soybean rotation cropping system (WS),

and soybean continuous cropping system (SC). Each year, the dosage of ammonium phosphate was  $349.5 \text{ kg ha}^{-1}$ , the dosage of urea was  $199.5 \text{ kg ha}^{-1}$ , and the dosage of K phosphate was  $199.5 \text{ kg ha}^{-1}$ . The cropping lasted for 12 years. Soil samples were collected after the cropping of soybeans in 2016.

## Soil Sampling

Soil samples were collected at seven different sampling points per plot. The soil cores were obtained using a 5-cm-diameter soil auger that collected soil between depths of 0 and 20 cm. The cores were removed by hand and separated into two parts. One part was air dried and sieved through a 2-mm sieve for soil fertility analysis, while the second part was stored at  $-80^\circ\text{C}$  for DNA extraction.

## Chemical Analyses

Soil samples were dried at room temperature and ground into pieces (that passed through a 0.2-mm sieve). The soil micronutrients Fe, Mn, Zn, and Cu were extracted with diethylenetriaminepentaacetic acid (DTPA) followed by determination by atomic absorption spectrophotometry (Lindsay and Norvell, 1978). Available boron (B) was determined by K imine colorimetry (Malekani and Cresser, 1998), and the pH was measured at a soil-to-water ratio of 2.5:1. Soil organic C was determined following wet digestion (Walkley and Black, 1934), and the values were multiplied by a factor of 1.724 to obtain OM values. The basic N solution was determined using the diffusion method (Bremner and Keeney, 1966), and available P was determined using the  $0.5 \text{ mol. L}^{-1} \text{ NaHCO}_3$  leaching-molybdenum antimony colorimetric method (Föhse et al., 1991). Available K was measured with the  $1 \text{ mol. L}^{-1} \text{ NH}_4\text{OAc}$  extraction-flame photometric method (Petersen and Corey, 1966).

## DNA Extraction, Library Construction, and Sequencing

Total DNA was extracted using the MoBio PowerSoil DNA Isolation Kit (MoBio, Carlsbad, CA, USA) according to the recommended protocol. The concentration and purity of the DNA were determined using a NanoDrop spectrophotometer, and the DNA integrity was determined using 0.8% agarose gel electrophoresis. The primers 515F (5'-GTGCCAGCMGCCGCGG-3') and 907R (5'-CCGTCAATTCMTTTRAGTTT-3') were used to amplify the V4-V5 hypervariable regions of the 16S rRNA gene. The primers ITS1F (5'-CTTGGTCATTTAGAGGAAGTAA-3') and ITS1R (5'-GCTGCGTTCTTCATCGATGC-3') were used to amplify the fungal internal transcribed spacer (ITS1) region. A parallel amplification of *E. coli*/yeast genomic DNA served as the positive control, and sterile water was used as the negative control. Each sample was independently amplified three times. Finally, the PCR products were evaluated by agarose gel electrophoresis, and PCR products from the same sample were pooled. The pooled PCR product was used as a template, and index PCR was performed using index primers to add the Illumina index to the library. The amplification products were evaluated using gel electrophoresis and purified using the Agencourt AMPure

XP Kit (Beckman Coulter, CA, USA). The purified products were indexed in the 16S V4-V5 or ITS1 library. The library quality was assessed on the Qubit@2.0 Fluorometer (Thermo Scientific) and Agilent Bioanalyzer 2100 systems. Finally, the pooled library was sequenced on an Illumina MiSeq 250 Sequencer, generating  $2 \times 250 \text{ bp}$  paired-end reads (Zhu et al., 2016).

## Statistical and Bioinformatics Analyses

Fastx was used to remove bases with end masses lower than Q15; the paired sequences obtained by double-end sequencing were merged into a longer sequence by FLASH in an overlapping relationship; and the obtained splicing sequence (merge) was obtained. Cutadapt software was used to delete the miscellaneous sequence at the end of the short sequence, and the resulting sequence was filtered with USEARCH software to obtain the valid sequence (Primer Filter). The singleton sequence was removed by USEARCH software, and the reference standard was compared to remove the chimeric sequence. The UPARSE clustering method was performed to merge the quality-controlled sequences (97-degree similarity between sequences) into an operational taxonomic unit (OTU) cell for subsequent OTU classification annotations (Edgar, 2013). Mothur software was used to compare the RDP, and the UNITE database was used for taxonomic analysis (Caporaso et al., 2010). All sequences were deposited in the NCBI Sequence Read Archive (SRA) database (accession numbers SRP155104), and the quality of the raw data sequencing is shown in **Supplementary Table 1**. Microbial community composition and species diversity analyses were performed using Lefse in R (R Development Core Team, 2008) and with Mothur software, and regression analyses of the relationships between microorganisms and soil physico-chemical properties were performed using R (Tian et al., 2015). Analyses of variance (ANOVAs) and Duncan's multiple range tests were performed with SPSS 17.0 to test the significance level ( $p < 0.05$ ) of the differences among the treatments.

## RESULTS

### Soil Properties

Analysis of the collected soil samples showed that the four cropping systems (i.e., FS, CS, WS, and SC) had significantly varied fertility characteristics ( $p < 0.05$ ; **Table 1**). The soils of the tillage modes in SC and FS exhibited higher contents of OM, available N, and K and were thus not conducive to the accumulation of available soil P. These results indicated that soybean cultivation resulted in the accumulation of available OM, N, K, Zn, Fe, and Mn in the soil, which resulted in the overconsumption of P. FS tillage reduced the available soil Cu content and the accumulation of Mn, resulting in the loss of available P. The WS cropping system improved the soil available P and the Cu content, but it consumed greater quantities of OM, available N and Fe. The contents of available K and Zn were the lowest in CS.

**TABLE 1** | Soil physico-chemical properties and microbial diversity by cropping system.

Soil properties	FS	CS	WS	SC	
Organic Matter (g/kg)		6.00 ± 0.84 <sup>ab</sup>	5.41 ± 0.14 <sup>bc</sup>	4.60 ± 0.19 <sup>c</sup>	7.40 ± 0.36 <sup>a</sup>
Available N (ppm)		364 ± 2.08 <sup>ab</sup>	311 ± 7.54 <sup>b</sup>	248 ± 8.39 <sup>c</sup>	387 ± 37.54 <sup>a</sup>
Available P (ppm)		12.33 ± 0.88 <sup>d</sup>	38.33 ± 1.20 <sup>b</sup>	47.33 ± 0.88 <sup>a</sup>	25.00 ± 0.58 <sup>c</sup>
Available K (ppm)		228 ± 0.58 <sup>b</sup>	152 ± 11.32 <sup>c</sup>	231 ± 2.52 <sup>b</sup>	323 ± 1.76 <sup>a</sup>
pH		6.13 ± 0.05 <sup>b</sup>	6.25 ± 0.01 <sup>a</sup>	6.19 ± 0.04 <sup>ab</sup>	6.24 ± 0.01 <sup>ab</sup>
DTPA Cu (ppm)		0.65 ± 0.02 <sup>c</sup>	1.33 ± 0.04 <sup>b</sup>	1.84 ± 0.04 <sup>a</sup>	1.29 ± 0.01 <sup>b</sup>
DTPA Zn (ppm)		0.46 ± 0.09 <sup>b</sup>	0.14 ± 0.04 <sup>c</sup>	0.50 ± 0.03 <sup>b</sup>	0.69 ± 0.02 <sup>a</sup>
DTPA Fe (ppm)		244 ± 4.02 <sup>b</sup>	208 ± 9.10 <sup>c</sup>	196 ± 3.47 <sup>c</sup>	304 ± 0.69 <sup>a</sup>
DTPA Mn (ppm)		10.91 ± 0.23 <sup>d</sup>	13.95 ± 0.72 <sup>c</sup>	16.48 ± 0.54 <sup>b</sup>	32.54 ± 0.19 <sup>a</sup>
DTPA B (ppm)		0.30 ± 0.00 <sup>a</sup>	1.02 ± 0.64 <sup>a</sup>	0.96 ± 0.11 <sup>a</sup>	0.70 ± 0.27 <sup>a</sup>
Bacteria	Chao1	5161 ± 204 <sup>a</sup>	4415 ± 148 <sup>b</sup>	4371 ± 46 <sup>b</sup>	4315 ± 159 <sup>b</sup>
	ACE	5112 ± 132 <sup>a</sup>	4358 ± 143 <sup>b</sup>	4362 ± 38 <sup>b</sup>	4307 ± 135 <sup>b</sup>
	Shannon	6.71 ± 0.08 <sup>a</sup>	6.49 ± 0.08 <sup>ab</sup>	6.53 ± 0.03 <sup>ab</sup>	6.42 ± 0.06 <sup>b</sup>
Fungi	Chao1	1222 ± 69 <sup>a</sup>	884 ± 51 <sup>b</sup>	906 ± 69 <sup>b</sup>	1256 ± 21 <sup>a</sup>
	ACE	1202 ± 64 <sup>a</sup>	873 ± 59 <sup>b</sup>	901 ± 57 <sup>b</sup>	1244 ± 9 <sup>ab</sup>
	Shannon	4.67 ± 0.123 <sup>a</sup>	3.57 ± 0.138 <sup>c</sup>	4.11 ± 0.067 <sup>b</sup>	4.37 ± 0.070 <sup>ab</sup>

SC, soybean continuous cropping system; FS, fallow-soybean cropping system; CS, corn-soybean rotation cropping system; and WS, wheat-soybean rotation cropping system. The mean value ± standard deviation (n = 3). For each line, different lowercase letters (a–d) indicate significant differences ( $p < 0.05$ ). Values with the same letter are not significantly different ( $p < 0.05$ ). Chao1 and ACE: abundance of the Chao1 and ACE estimator and operational taxonomic units. Shannon: nonparametric Shannon diversity index.

## Microbial Colony Structure and Diversity Analysis

Bacterial and fungal principal coordinate analysis (PCoA) showed significant differences among the four different cropping systems, but the WS and SC colony structures were similar (Figures 1A,D). The abundance (Chao1/ACE) and diversity (Shannon) of bacteria and fungi in FS were significantly higher ( $p < 0.05$ ). The diversity and abundance of soil bacteria in SC were the lowest among all the cropping systems analyzed, but the diversity of fungi was higher in SC than in the CS and WS (Table 1). The diversity of fungi in CS was the lowest among all the cropping systems analyzed ( $p < 0.05$ ). The archaea in WS exhibited enhanced formation, and the lowest abundance of archaea was found in SC ( $p < 0.05$ ; Figure 1B). The different cropping systems affected the formation of the soil microbial colony structure and diversity. In total, 764 unique bacterial species and 724 unique fungal species were identified in FS, and these values were much higher than those observed in the other treatments. Additionally, 305 bacterial species and 379 fungal species were processed in SC (Figures 1C,F). The FS cropping system was the most conducive to the formation of soil microbial colony diversity among all the cropping systems analyzed, especially in terms of protozoan formation (Figure 1E).

## Microbial Community Abundance Analysis

The bacteria and archaea identified in this study belong to 33 phyla (classified reads = 94%), 61 classes (84%), 73 orders (56%), 164 families (50%), 385 genera (61%), and 626 species (41%). The fungi and protists identified in this study belong to 7 phyla (classified reads = 95%), 31 classes (93%), 88 orders (85%), 191 families (81%), 371 genera (74%), and 453 species (46%; Supplementary Table 2).


The dominant bacteria (i.e., the relative abundance) in the various soil samples were mainly distributed in Proteobacteria, Acidobacteria, Bacteroidetes, Actinobacteria, Planctomycetes, Chloroflexi, Gemmatimonadetes, candidate division WPS-1, Verrucomicrobia, Armatimonadetes, and Nitrospirae. The differences among the microbial communities in the different cropping systems were significant. Proteobacterium was the most abundant in all samples, with a relative abundance of more than 20% of the total number of OTUs (Figure 2A).

The dominant bacteria in the SC soil had significantly more members of the phyla Proteobacteria, Bacteroidetes and Actinobacteria than the other cropping systems, and significantly fewer bacteria from the phyla Acidobacteria, Planctomycetes, Gemmatimonadetes, Chloroflexi, and Nitrospirae were observed.

The dominant bacteria in WS soil included more members from the phyla Planctomycetes and Acidobacteria than in the other cropping systems, and WS had a significantly lower relative abundance of Bacteroidetes than that found in the other cropping systems. Additionally, candidate division WPS-1 was significantly higher and Verrucomicrobia was significantly lower in the WS cropping system than in the other cropping systems analyzed.

Microbial diversity was greater in the FS cropping system, which presented higher relative abundances of Chloroflexi, Verrucomicrobia, Nitrospirae, and Bacteroidetes and a significantly lower relative abundance of Actinobacteria. There was a significant percentage (7.2%) of bacteria in the “No\_Rank” category in FS.

The fungi in all samples were mainly distributed in Ascomycota, Basidiomycota and Zygomycota, encompassing 90% of the total samples. Fungi from Ascomycota were significantly more common in WS than in the other systems. The quantity of Basidiomycota was significantly higher in the



CS cropping system than in the other cropping systems. The relative abundance of Zygomycota in FS was significantly higher (relative abundance of 19.7%) than those in the SC (9.7%) and crop rotation (6.1%) cropping systems (**Figure 2B**).

Proteobacteria was significantly positively correlated with the DTPA Fe and Mn contents in the soil, whereas Acidobacteria was significantly negatively correlated with these factors. Bacteroidetes was significantly correlated with the available N, OM, and Fe contents and significantly negatively correlated with the available P and DTPA Cu contents. Actinobacteria was positively correlated with the available P and DTPA Cu and Mn contents (**Figure 2C**).


Fungal Ascomycota was significantly correlated with the available P and DTPA Cu contents. Zygomycota and Cercozoa (Protista) were significantly negatively correlated with the available P and DTPA Cu contents (**Figure 2E**).

Bacteria in the Vertisol samples were mainly distributed in the *Gp4*, *Gp6*, *Gemmatimonas*, *Gp1*, *Gaiella*, *incertae sedis WPS-1*,

*Flavobacterium*, *Gp3*, *Rhizomicrobium*, *Rhodanobacter*, *Opitutus*, and *Pedobacter* genera, but most bacteria were not identified at the species level (i.e., No\_Rank; **Figure 2D**). The dominant bacteria in the *Gp6* genus in FS was negatively correlated with the DTPA Fe and Mn contents. The dominant bacterial *Gp4* found in the CS was negatively correlated with the available K and DTPA Zn, Fe, and Mn contents. The dominant bacteria in WS were *Gemmatimonas*, *Gaiella*, and *incertae sedis WPS-1*, which were significantly related to the available P and DTPA Cu contents. *Gp1* was negatively related to the available N content.

The dominant bacteria *Rhizomicrobium*, *Rhodanobacter*, and *Pedobacter* in SC were significantly positively correlated with the available K, DTPA Mn and DTPA Fe contents (**Table 2**).

The relative abundance values of fungal genera varied significantly among the samples (**Figure 2F**). The dominant fungus *Mortierella* in the FS was negatively correlated with the DTPA Cu and available P contents, and the dominant fungus *Peziza* was significantly negatively correlated with the available



**FIGURE 2** | and the microbial biomasses among the microbial communities. Each column represents a different soil physico-chemical property, each row represents a different species, and the different colors indicate the sizes of two correlation coefficients. A darker blue indicates a stronger negative correlation, whereas a more intense orange represents a greater positive correlation. \*indicates significance at  $p < 0.05$ , \*\*indicates significance at  $p < 0.01$ . SC, soybean continuous cropping system; FS, fallow-soybean cropping system; CS, corn-soybean rotation cropping system; and WS, wheat-soybean rotation cropping system.

**TABLE 2** | Relationships between dominant bacterial species and soil physico-chemical properties.

Genus	Phylum	Higher	Lower	N	P	K	OM	pH	Cu	Zn	Fe	Mn	B
Gp4	Acidobacteria	CS	SC	-0.66	0.49	-0.94	-0.63	0.08	0.21	-0.86	-0.88	-0.78	0.18
Gp6	Acidobacteria	FS	SC	-0.44	-0.01	-0.62	-0.66	-0.61	-0.21	-0.52	-0.72	-0.92	-0.14
Gp1	Acidobacteria	WS		-0.67	0.37	0.04	-0.31	-0.20	0.46	0.17	-0.42	-0.20	-0.07
Gemmatimonas	Gemmatimonadetes	WS		-0.65	0.73	-0.33	-0.76	0.02	0.65	-0.14	-0.73	-0.35	0.38
WPS-1	candidate_division_WPS-1	WS		-0.81	0.84	-0.38	-0.66	0.20	0.77	-0.16	-0.76	-0.33	0.28
Gaiella	Actinobacteria	WS		-0.68	0.94	-0.36	-0.68	0.29	0.85	-0.29	-0.67	-0.12	0.55
Gp3	Acidobacteria			-0.18	-0.21	0.03	0.16	-0.36	-0.17	-0.04	-0.03	-0.16	-0.22
Rhizomicrobium	Proteobacteria	SC		0.24	0.22	0.71	0.28	0.38	0.47	0.63	0.52	0.87	0.02
Rhodanobacter	Proteobacteria	SC	FS	-0.02	0.19	0.38	0.46	0.46	0.34	0.21	0.45	0.72	0.16
Pedobacter	Bacteroidetes	SC	FS	0.62	-0.17	0.72	0.62	0.33	0.04	0.55	0.80	0.90	-0.10
Opitutus	Verrucomicrobia	FS		0.86	-0.92	0.39	0.67	-0.20	-0.84	0.35	0.75	0.22	-0.30
Flavobacterium	Bacteroidetes	FS	WS	0.59	-0.84	0.25	0.34	-0.35	-0.81	0.35	0.43	-0.14	-0.43

Notes: the most abundant OTUs in each taxonomic level were correlated with the tillage system and soil physico-chemical properties (i.e., pH, OM, available N, P, K, DTPA Fe, Mn, Zn, and Cu) by correlation with a significance level. The red boxes indicate significant positive correlations, and the blue boxes represent significant negative correlations ( $p < 0.01$ ). The red digits represent positive correlations, and the blue digits represent negative correlations ( $p < 0.05$ ). WPS-1: WPS-1\_genera\_incertae\_sedis.

P and DTPA Cu contents. *Chaetomium* was significantly less prominent in FS, and *Chaetomium* was positively correlated with the available P content. The dominant bacterium *Tetracladium* in SC soil was positively correlated with the DTPA Mn content. The dominant bacterium *Gibberella* was positively correlated with the available N content, and the dominant bacterium *Lectera* was positively correlated with the available K and DTPA Zn and Mn contents. In the CS and WS rotation soils, the dominant fungi *Alternaria* and *Hydropisphaera* were positively correlated with the available P and DTPA Cu contents but negatively correlated with the available N and DTPA Fe contents (Table 3).

### Significant Differences in Species

Under the different cropping systems, the genera (Figures 3A,B) that formed were distinct for the different colony structures. The *Geobacter*, *Geothrix*, and *Azonexus* bacteria compositions were significantly different. In FS, significant differences in fungi existed and included *Eocronartium*, *Pluteus*, *Slopeiomyces*, and *Acaulospora*. These microorganisms were significantly correlated with low available P and low DTPA Cu content in the soil.

Microorganisms that produced significant differences in SC were mainly distributed in the bacteria *Sphingobacterium*, *Brevundimonas*, and *Mucilaginibacter* and in *Auxarthron*, the fungi *Entorrhiza*, *unclassified Bionectriaceae*, and *unclassified Corticiaceae*. These microorganisms were significantly associated with a high OM content, high available K content, and high DTPA Fe and Mn contents in SC soil. Microbes and physico-chemical properties have a corresponding relationship (Table 4), and soil physico-chemical property extremes (i.e., extremely high/extremely low) determine the microbial community structure.

Different cropping systems cause significant differences in soil fertility. Thus, the microbial community structure is further influenced. The microbial community structure in FS was significantly affected by the low contents of available P, DTPA Cu and Mn. The soil microbial community structure in the SC cropping system was mainly affected by high available K, DTPA Mn and Fe contents. The microbial community structure in the CS cropping system was mainly affected by low available N and OM contents and high available P and DTPA Cu contents. The soil microbial community structure in the WS cropping system was mainly affected by low available N, OM, and DTPA Fe contents and by high available P and DTPA Cu contents. The crop species and soil characteristics coopted and promoted the growth of microbial communities. The microbial community structure could be altered by regulating the variety of crops and increases and decreases in soil characteristics.

The soil microbial community is well correlated with soil physico-chemical properties (Figures 4A,B; Table 5). The bacterial microbial community was significantly correlated with the available P and DTPA Mn contents in the soil, with a correlation coefficient of 0.93; additionally, the community was correlated (0.91) with the available P, DTPA Cu and Mn contents. The effects of fungi and soil on the available P, K, and DTPA Mn contents were significant, with a correlation coefficient of 0.85 (Table 5).

### Bacterial KEGG Functional Differences

The bacterial microbial communities of the different cropping systems had significant functional differences, while the functional differences in the fungal communities were not obvious (Figures 4C,D). Cyanobacteria were significantly

**TABLE 3** | Relationships between dominant fungal species and soil physico-chemical properties.

Genus	Phylum	Higher	Lower	N	P	K	OM	pH	Cu	Zn	Fe	Mn	B
<i>Mortierella</i>	Zygomycota	FS		0.44	-0.69	0.12	0.33	-0.11	-0.67	0.33	0.28	-0.23	-0.38
<i>Alternaria</i>	Ascomycota	WS CS		-0.73	0.89	-0.51	-0.62	0.20	0.71	-0.53	-0.77	-0.29	0.52
<i>Chaetomium</i>	Ascomycota		FS	-0.34	0.59	0.01	0.11	0.50	0.57	-0.19	-0.05	0.41	0.28
<i>Sistotrema</i>	Basidiomycota	CS		-0.18	0.08	-0.44	0.09	0.29	0.02	-0.40	0.01	-0.01	0.33
<i>Tetracladium</i>	Ascomycota	SC		0.27	0.00	0.52	0.63	0.43	0.16	0.33	0.66	0.85	0.25
<i>Gibberella</i>	Ascomycota	SC CS		0.63	-0.07	0.12	0.19	0.46	-0.09	-0.13	0.32	0.39	-0.02
<i>Peziza</i>	Ascomycota	FS		0.67	-0.93	0.55	0.63	-0.43	-0.79	0.52	0.68	0.16	-0.44
<i>Lectera</i>	Ascomycota	SC		0.33	0.17	0.63	0.16	0.07	0.35	0.61	0.39	0.66	-0.14
<i>Hydropisphaera</i>	Ascomycota	WS CS		-0.78	0.92	-0.37	-0.55	0.25	0.79	-0.39	-0.74	-0.18	0.45

Notes: the most abundant OTUs in each taxonomic level were correlated with the tillage system and soil physico-chemical properties (i.e., pH, OM, available N, P, K, DTPA Fe, Mn, Zn, and Cu) by correlation with a significance level. Higher: indicates that the species is significantly more abundant in that cropping system. Lower: indicates that the species is significantly less abundant in that cropping system ( $p < 0.05$ ). The red boxes represent significant positive correlations, and the blue boxes represent significant negative correlations ( $p < 0.01$ ). The red digits represent positive correlations, and the blue digits represent negative correlations ( $p < 0.05$ ).


more abundant in the SC cropping system than in the other cropping systems (Figure 4E). Additionally, a range of archaeal methanogens, such as *Methanomassiliicoccus* and *Methanosarcina* (both anaerobes of *Euryarchaeota*), were significantly higher in FS than in the other cropping systems. However, these groups were not detected in samples from SC and WS (Figure 4E). *Azotobacter* N-fixing bacteria existed in only FS, and *Nitrospira* and *Anaerolinea* were significantly more abundant in FS than in the other cropping systems. *Azotobacter* bacteria existed in only FS, and *Nitrospira* and *Anaerolinea* were significantly more abundant in FS than in the other cropping systems (Figure 4E).

## DISCUSSION

In this study, the OM and available N contents in soil were highest in SC, which is consistent with previous results for soybean cropping (Lu et al., 2011). In addition, the study showed that SC increased the contents of available K, DTPA Zn, Fe and Mn in soil, which is consistent with previous single-cropping results (Cai et al., 2017). In the soil under the FS cropping system, the available P, DTPA Cu and Mn contents decreased significantly. In contrast, the contents of available P and DTPA Cu increased significantly in the WS soil, but the contents of available N and DTPA Fe decreased significantly. In the soil under the CS cropping system, the contents of available K and DTPA Zn were significantly lower than under other cropping systems. Thus, different cropping systems generated different soil physico-chemical properties, which is consistent with the results of previous studies (Fierer et al., 2009; Xun et al., 2015), demonstrating that the cropping system can regulate soil physico-chemical properties.

The abundance and diversity of soil microbes determine the stability of the soil microbial community and its ability to resist pathogens, which is crucial to the function and sustainable development of the soil ecosystem (Smith et al., 2008; Pang et al., 2017). In this study, the abundance of fungal microorganisms in the SC cropping system increased

significantly, which is consistent with the results of Li et al. (2010). Single cropping may increase the relative abundance of fungi in the soil and increase the occurrence probability of soil-borne diseases. In this study, the FS cropping system was characterized by a more diverse microbial ecosystem, confirming that reducing agricultural cropping can increase soil microbial diversity (Wang et al., 2011). The diversity of soil bacteria in the WC and CS cropping systems was greater than that in the SC cropping system, which is consistent with previous studies (Ai et al., 2015). Proteobacteria and Bacteroidetes can inhibit the occurrence of diseases (Ai et al., 2015), and Acidobacteria has the ability to break down complex soil OM and plant-derived polysaccharides (Ward et al., 2009). The dominant bacteria (i.e., the relative abundance) distributed in various samples of cold-ground Vertisols were previously found to be from Proteobacteria, Acidobacteria, Bacteroidetes, and Actinobacteria (Knelman et al., 2015), and the results of the present study confirmed these conclusions. Proteobacteria was the most abundant in all treatments, accounting for ~20% of the total number of OTUs (Figure 2A), and the Proteobacteria associated with symbiotic rhizobia was significantly more abundant in the SC cropping system. With the accumulation of soil OM and nutrients, the relative abundance of Proteobacteria increased (Freedman and Zak, 2015; Zeng et al., 2017). Acidobacteria was significantly less abundant in SC but significantly more abundant in the WS cropping system. Bacteroidetes was significantly higher in the FS cropping system. Actinobacteria was significantly higher in the SC cropping system but significantly lower in the FS cropping system. Janssen (2006) found that the content of Gemmatimonadetes in soil bacterial communities was 2% (Janssen, 2006), and in the present study, Gemmatimonadetes was significantly more abundant in the WS cropping system and was detected in ~0.91% of reads. We also found that Verrucomicrobia was significantly more abundant in the FS cropping system. Bergmann et al. (2011) studied undisturbed soil and found that 23% of bacterial sequences belonged to Verrucomicrobia microbes (Bergmann et al., 2011). In the present study, the bacteria phylum Chloroflexi was abundant in FS and WS and was nearly twice as abundant as that in



**FIGURE 3** | English letters in the image on the right are represented in each group and play an important role in the outline of the species name level. The names on the outer ring of the letters between groups play an important role in the door at the species name level; thus, each layer of nodes from the inside runs through the phylum, class, order, family, and species. Each species layer represents the process from inside extroversion to the said phylum, class, order, family, genus, and species. The diameter of the node is proportional to the relative abundance. SC, soybean continuous cropping system; FS, fallow-soybean cropping system; CS, corn-soybean rotation cropping system; and WS, wheat-soybean rotation cropping system.

**TABLE 4** | Relationships between species and soil physico-chemical properties.

Community	Taxon	Physico-chemical properties									
		OM	N	P	K	Zn	Fe	Cu	Mn	pH	
FS	Bacteria	<i>Geobacter</i>			-0.70				-0.83		
		<i>Geothrix</i>			-0.80				-0.83		-0.75
		<i>Azonexus</i>			-0.70				-0.80	-0.61	
	Fungi	<i>Eocronartium</i>			-0.63				-0.65		-0.63
		<i>Pluteus</i>									
		<i>Slopeiomyces</i>			-0.73				-0.77		
CS	Bacteria	<i>Segetibacter</i>			0.85				0.80		0.65
		<i>Niastella</i>			0.80				0.71		0.60
		<i>Paralcaligenes</i>			0.65	-0.59	-0.65				
	Fungi	<i>Plenodomus</i>				-0.77	-0.83				
		Unclassified_Xylariaceae									
WC	Bacteria	<i>Bradyrhizobium</i>		-0.62	0.71				0.83		
		<i>Microlunatus</i>	-0.74	-0.81	0.74	-0.61		-0.86	0.61		
		<i>Bacillus</i>	-0.63	-0.69	0.91			-0.61	0.87		
		<i>Angustibacter</i>		-0.67	0.86				0.89		
	Fungi	<i>Terrisporobacter</i>		-0.59							
		Unclassified_Pezizaceae	-0.59	-0.77	0.69				0.74		
SC	Bacteria	<i>Septoglomerus</i>		-0.74	0.64				0.72		
		<i>Sphingobacterium</i>	0.75	0.64		0.82	0.63	0.90		0.92	
		<i>Brevundimonas</i>				0.66		0.71		0.80	
	Fungi	<i>Mucilaginibacter</i>						0.59		0.82	
		<i>Auxarthron</i>	0.68	0.62		0.83	0.67	0.87		0.95	
		<i>Entorrhiza</i>	0.69			0.64		0.69		0.74	
	<i>Unclassified_Bionectriaceae</i>	0.75			0.77	0.61	0.82		0.89		
	<i>Unclassified_Corticaceae</i>	0.58	0.70		0.78	0.64	0.82		0.89		


Notes: the red background represents the highest content of soil physico-chemical properties, and the orange color represents a high content physico-chemical properties. Yellow represents a medium content of physico-chemical properties, and blue represents the lowest content of physico-chemical properties. The value represents the correlation. "+" indicates a positive correlation, and "-" indicates a negative correlation. The higher the value is, the stronger the correlation.

the SC cropping system. These microbes can perform anaerobic photosynthesis and withdraw electrons from hydrogen sulfide (Eisen et al., 2002). Nitrospirae was significantly more abundant in the FS cropping system and significantly lower in SC than in the other cropping systems. Cyanobacteria were more abundant in the SC cropping system; members of this phylum can fix N gas into ammonia (NH<sub>4</sub><sup>+</sup>), nitrite (NO<sub>2</sub><sup>-</sup>) or nitrate (NO<sub>3</sub><sup>-</sup>) as well as significantly reduce carbon aerobically (Flores, 2008).

Thaumarchaeota and a *Candidatus Nitrososphaera* comprise the soil NH<sub>4</sub><sup>+</sup>-oxidizing archaea genera; NH<sub>4</sub><sup>+</sup> is oxidized by producing amnoxase (AMO; Zhahnina et al., 2012). Thaumarchaeota was significantly more abundant in the WS cropping system. Euryarchaeota was significantly abundant in FS, and this group includes extreme halophiles, sulfate-reducing

microbes and anaerobic methanogens as well as many other species (Lei et al., 2017). The genera *Methanomassiliicoccus* and *Methanosarcina* of Euryarchaeota were detected in only FS, but other Euryarchaeota were almost five times more abundant in the SC cropping system. Collectively, these data support the notion that the abundance of anaerobic microorganisms increased in the FS because of less aeration due to low soil disturbance. In addition, the OM in the FS soil maintains a certain water storage capacity, providing anaerobic habitats for anaerobic organisms (Minamisawa et al., 2004).

The fungi in all samples were mainly distributed in Ascomycota, Basidiomycota and Zygomycota, which accounted for 90% of the total sampled fungi. This result confirms previous findings (Xu et al., 2012). Fungi from Ascomycota were



**TABLE 5** | Relationships between microorganisms and soil physico-chemical properties.

Physico-chemical properties combination	Size correlation	Physico-chemical properties combination	Size correlation
	Bacteria		Fungi
P	0.71	P	0.78
P Mn	<b>0.93</b>	P Mn	0.83
P Cu Mn	<b>0.91</b>	P K Cu	<b>0.86</b>
P Cu Fe Mn	0.89	P K Cu Mn	0.85
P K Cu Fe Mn	0.85	P K pH Cu Mn	0.82
P K pH Cu Fe Mn	0.84	N P K pH Cu Mn	0.80
N P K pH Cu Fe Mn	0.82	N P K pH Cu Zn Mn	0.80
N P K OM pH Cu Fe Mn	0.81	N P K pH Cu Zn Fe Mn	0.77
N P K OM pH Cu Zn Fe Mn	0.78	N P K OM pH Cu Zn Fe Mn	0.74
N P K OM pH Cu Zn Fe Mn B	0.74	N P K OM pH Cu Zn Fe Mn B	0.70

Each line represents the level of correlation between each physico-chemical property (i.e., the combination of soil physico-chemical properties) and the community. The greater the correlation absolute value is, the stronger the correlation and the greater the influence on the community. The digits represent *r* values ( $p < 0.05$ ). The red character stands for outstanding numeral.

significantly more common in the WS cropping system. The relative abundance of Basidiomycota did not significantly differ among the samples. The distribution of Zygomycota fungi was significantly greater in the FS soil, with a relative abundance of 19.7%, which was significantly higher than that under the SC (9.7%) and crop rotation (6.1%) cropping systems.

Soil fertility affects soil microbial community structure (Liu et al., 2014; Zhang et al., 2016). The increased OM and N in soil could promote bacterial diversity (Shen et al., 2014; Li et al., 2016; Zeng et al., 2017). An interesting phenomenon was observed in this experiment: in the SC and FS cropping systems, the contents of OM and N were very high, but the abundance and diversity of bacteria differed significantly, indicating that the abundance and diversity of soil bacteria may be related to other factors in addition to OM and N. The diversity of fungi under SC was significantly higher than that under rotation, and Hooper et al. proposed that this might be related to the long-term accumulation of soil nutrients (Hooper et al., 2005).

In the FS cropping system, Bacteroidetes and Verrucomicrobia were significantly positively correlated with the OM content, and Bacteroidetes, Verrucomicrobia, and Latescibacteria were significantly negatively correlated with the available P and DTPA Cu contents. Nitrospirae was significantly negatively correlated with the available P, K, and DTPA Mn contents, and Actinobacteria was significantly positively correlated with the available P, DTPA Cu, and Mn contents. In the SC cropping system, Proteobacteria was significantly positively correlated with the available Fe and DTPA Mn contents. In the WS cropping system, Planctomycetes was negatively correlated with the DTPA Zn content. Zygomycota and Cercozoa of Protista were significantly more abundant in the FS cropping system and were significantly negatively correlated with the P and Cu contents. Ascomycota was significantly more abundant in the WS cropping system and was positively correlated with the DTPA Fe and Cu contents.

These results are inconsistent with those of Cai et al. (2017) and Pang et al. (2017), which may be due to the different types

and amounts of soil samples studied. However, there was a significant correlation between the microbial community and the soil physico-chemical properties.

In this study, microbial community structure was influenced by high or low soil fertility in the soils of different cropping systems. In fact, 33 of the 40 microbial phyla were significantly positively or negatively correlated with one or several soil fertility parameters ( $p < 0.05$ ). The bacterial community structure was significantly correlated with the physico-chemical properties of soil available P, DTPA Cu, and Mn. The fungal community was significantly correlated with available P, K, and DTPA Cu.

These changes in the microbial communities are related to the nutrients and plant biomass available in the soil (Bokhorst et al., 2017). Many studies have concluded that soil and physico-chemical properties shape community structure (Xun et al., 2015; Pang et al., 2017), and as confirmed experimentally, the cropping system, soil fertility, and soil microbial community structure are interrelated and can be adjusted concomitantly (Ofek et al., 2014). In the present study, the cropping system altered the soil properties and the community structure, and we found that there was no perfect cropping system. The FS, WS, and CS systems were capable of balancing soil fertility and microbial stability. Therefore, institutional diversity is most conducive to soil productivity (Kuramae et al., 2012; Yuan et al., 2015). We found that the available P, K, DTPA Cu, and Mn contents were closely related to the microorganism communities.

## CONCLUSION

The cropping system affects the physical and chemical properties of soil and the microbiome structure. (1) Soybean cropping favored the accumulation of OM and the available N, K DTPA Fe, Mn, Zn, and Cu contents in soil but not fixed available P. The WS and CS cropping systems were conducive to the fixed available P, but they consumed OM, DTPA K, and Zn. (2) The FS cropping system could increase the abundance and

diversity of bacteria. The SC cropping system could decrease the abundance and diversity of bacteria but increase the abundance and diversity of fungi. The WS and CS cropping systems could reduce fungal abundance and diversity. (3) There was a significant correlation between microbial community structure and the physico-chemical properties of soil, and the most significant correlation was between the bacterial community and available P, DTPA, and Mn (0.93). The fungal community was most correlated with available P, DTPA K, and Cu, with a correlation coefficient of 0.8. In summary, increasing the diversity of the cropping system was more favorable to maintaining soil fertility. The cropping system changes soil properties and microbial diversity, and the microbial community in turn manipulates nutrient cycling processes and alters soil fertility, plant productivity and environmental sustainability. The results of this study will provide a foundation for further studies on the regulation of soil fertility and microorganism community structure and provide guidance for selecting the best cropping model to protect soil ecology.

## AUTHOR CONTRIBUTIONS

XS and BT planned and arranged the entire study. The main experiments were performed by XS. JG and JL assisted

in the soil testing. XS and BT wrote the manuscript together. GC participated in further discussion. All authors discussed and approved the final version of the manuscript.

## FUNDING

This study was financially supported by the National Key Research and Development Plan of China (2016YFD0200500; 2016YFD070030103).

## ACKNOWLEDGMENTS

We are grateful to the Agriculture College of Northeast Agricultural University, Heilongjiang Academy of Agricultural Sciences, and the numerous staff members who helped protect the test plots and collect the soil samples.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fenvs.2018.00132/full#supplementary-material>

## REFERENCES

- Ai, C., Liang, G., Sun, J., Wang, X., He, P., Zhou, W., et al. (2015). Reduced dependence of rhizosphere microbiome on plant-derived carbon in 32-year long-term inorganic and organic fertilized soils. *Soil Biol. Biochem.* 80, 70–78. doi: 10.1016/j.soilbio.2014.09.028
- Benitez, M. S., Osborne, S. L., and Lehman, R. M. (2017). Previous crop and rotation history effects on maize seedling health and associated rhizosphere microbiome. *Sci. Rep.* 7:15709. doi: 10.1038/s41598-017-15955-9
- Bergmann, G. T., Bates, S. T., Eilers, K. G., Lauber, C. L., Caporaso, J. G., Walters, W. A., et al. (2011). The under-recognized dominance of *Verrucomicrobia* in soil bacterial communities. *Soil Biol. Biochem.* 43, 1450–1455. doi: 10.1016/j.soilbio.2011.03.012
- Bokhorst, S., Kardol, P., Bellingham, P. J., Kooyman, R. M., Richardson, S. J., Schmidt, S., et al. (2017). Responses of communities of soil organisms and plants to soil aging at two contrasting long-term chronosequences. *Soil Biol. Biochem.* 106, 69–79. doi: 10.1016/j.soilbio.2016.12.014
- Bremner, J. M., and Keeney, D. R. (1966). Determination and isotope-ratio analysis of different forms of nitrogen in soils: 3. exchangeable ammonium, nitrate, and nitrite by extraction-distillation methods 1. *Soil Sci. Soc. Am. J.* 30, 577–582. doi: 10.2136/sssaj1966.03615995003000050015x
- Cai, F., Chen, W., Wei, Z., Pang, G., Li, R., Ran, W., et al. (2015). Colonization of *Trichoderma harzianum* strain SQR-T037 on tomato roots and its relationship to plant growth, nutrient availability and soil microflora. *Plant Soil* 388, 337–350. doi: 10.1007/s11104-014-2326-z
- Cai, F., Pang, G., Miao, Y., Li, R., Li, R., Shen, Q., et al. (2017). The nutrient preference of plants influences their rhizosphere microbiome. *Appl. Soil Ecol.* 110, 146–150. doi: 10.1016/j.apsoil.2016.11.006
- Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., and Bushman, F. D., Costello, E. K., et al. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods* 7, 335–336. doi: 10.1038/nmeth.f.303
- Edgar, R. C. (2013). UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat. Methods* 10, 996–998. doi: 10.1038/nmeth.2604
- Eisen, J. A., Nelson, K. E., Paulsen, I. T., Heidelberg, J. F., Wu, M., Dodson, R. J., et al. (2002). The complete genome sequence of *Chlorobium tepidum* TLS, a photosynthetic, anaerobic, green-sulfur bacterium. *Proc. Natl. Acad. Sci. U.S.A.* 99, 9509–9514. doi: 10.1073/pnas.132181499
- Fierer, N., Lauber, C. L., Ramirez, K. S., Zaneveld, J., Bradford, M. A., and Knight, R. (2012). Comparative metagenomic, phylogenetic and physiological analyses of soil microbial communities across nitrogen gradients. *ISME J.* 6, 1007–1017. doi: 10.1038/ismej.2011.159
- Fierer, N., Strickland, M. S., Liptzin, D., Bradford, M. A., and Cleveland, C. C. (2009). Global patterns in belowground communities. *Ecol. Lett.* 12, 1238–1249. doi: 10.1111/j.1461-0248.2009.01360.x
- Flores, F. G. (2008). *The Cyanobacteria: Molecular Biology, Genomics, and Evolution*. Norwich: Horizon Scientific Press.
- Föhse, D., Claassen, N., and Jungk, A. (1991). Phosphorus efficiency of plants. *Plant Soil* 132, 261–272. doi: 10.1007/BF00010407
- Freedman, Z., and Zak, D. R. (2015). Soil bacterial communities are shaped by temporal and environmental filtering: evidence from a long-term chronosequence. *Environ. Microbiol.* 17, 3208–3218. doi: 10.1111/1462-2920.12762
- Hobbs, P. R., Sayre, K., and Gupta, R. (2008). The role of conservation agriculture in sustainable agriculture. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 363, 543–555. doi: 10.1098/rstb.2007.2169
- Hooper, D. U., Chapin, F. S., Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S., et al. (2005). Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecol. Monogr.* 75, 3–35. doi: 10.1890/04-0922
- Janssen, P. H. (2006). Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Appl. Environ. Microbiol.* 72, 1719–1728. doi: 10.1128/AEM.72.3.1719-1728.2006
- Knelman, J. E., Graham, E. B., Trahan, N. A., Schmidt, S. K., and Nemergut, D. R. (2015). Fire severity shapes plant colonization effects on bacterial community structure, microbial biomass, and soil enzyme activity in secondary succession of a burned forest. *Soil Biol. Biochem.* 90, 161–168. doi: 10.1016/j.soilbio.2015.08.004
- Kuramae, E. E., Yergeau, E., Wong, L. C., Pijl, A. S., Veen, J. A., and Kowalchuk, G. A. (2012). Soil characteristics more strongly influence soil bacterial communities than land-use type. *FEMS Microbiol. Ecol.* 79, 12–24. doi: 10.1111/j.1574-6941.2011.01192.x

- Lei, Y., Xiao, Y., Li, L., Jiang, C., Zu, C., Li, T., et al. (2017). Impact of tillage practices on soil bacterial diversity and composition under the tobacco-rice rotation in China. *J. Microbiol.* 55, 349–356. doi: 10.1007/s12275-017-6242-9
- Li, C., Li, X., Kong, W., Wu, Y., and Wang, J. (2010). Effect of monoculture soybean on soil microbial community in the Northeast China. *Plant Soil* 330, 423–433. doi: 10.1007/s11104-009-0216-6
- Li, R., Shen, Z., Sun, L., Zhang, R., Fu, L., Deng, X., et al. (2016). Novel soil fumigation method for suppressing cucumber *Fusarium* wilt disease associated with soil microflora alterations. *Appl. Soil Ecol.* 101, 28–36. doi: 10.1016/j.apsoil.2016.01.004
- Lindsay, W. L., and Norvell, W. L. (1978). Development of DTPA soil test for zinc, iron, manganese and copper. *Soil Sci. Soc. Am. J.* 42, 421–428. doi: 10.2136/sssaj1978.03615995004200030009x
- Liu, J., Sui, Y., Yu, Z., Shi, Y., Chu, H., Jin, J., et al. (2014). High throughput sequencing analysis of biogeographical distribution of bacterial communities in the black soils of Northeast China. *Soil Biol. Biochem.* 70, 113–122. doi: 10.1016/j.soilbio.2013.12.014
- Lu, M., Yang, Y., Luo, Y., Fang, C., Zhou, X., Chen, J., et al. (2011). Responses of ecosystem nitrogen cycle to nitrogen addition: a meta-analysis. *New Phytol.* 189, 1040–1050. doi: 10.1111/j.1469-8137.2010.03563.x
- Luo, S., Yu, L., Liu, Y., Zhang, Y., Yang, W., Li, Z., et al. (2016). Effects of reduced nitrogen input on productivity and N<sub>2</sub>O emissions in a sugarcane/soybean intercropping system. *Eur. J. Agron.* 81, 78–85. doi: 10.1016/j.eja.2016.09.002
- Malekani, K., and Cresser, M. S. (1998). Comparison of three methods for determining boron in soils, plants, and water samples. *Commun. Soil Sci. Plant Anal.* 29, 285–304. doi: 10.1080/00103629809369946
- Minamisawa, K., Nishioka, K., Miyaki, T., Ye, B., Miyamoto, T., You, M., et al. (2004). Anaerobic nitrogen-fixing consortia consisting of clostridia isolated from gramineous plants. *Appl. Environ. Microbiol.* 70, 3096–3102. doi: 10.1128/AEM.70.5.3096-3102.2004
- Mo, A. S., Qiu, Z. Q., He, Q., Wu, H. Y., and Zhou, X. B. (2016). Effect of continuous monocropping of tomato on soil microorganism and microbial biomass carbon. *Commun. Soil Sci. Plant Anal.* 47, 1069–1077. doi: 10.1080/00103624.2016.1165832
- Ofek, M., Voronov-Goldman, M., Hadar, Y., and Minz, D. (2014). Host signature effect on plant root-associated microbiomes revealed through analyses of resident vs. active communities. *Environ. Microbiol.* 16, 2157–2167. doi: 10.1111/1462-2920.12228
- Pang, G., Cai, F., Li, R., Zhao, Z., Li, R., Gu, X., et al. (2017). Trichoderma-enriched organic fertilizer can mitigate microbiome degeneration of monocropped soil to maintain better plant growth. *Plant Soil* 416, 181–192. doi: 10.1007/s11104-017-3178-0
- Petersen, G. W., and Corey, R. B. (1966). A modified Chang and Jackson procedure for routine fractionation of inorganic soil phosphates 1. *Soil Sci. Soc. Am. J.* 30, 563–565. doi: 10.2136/sssaj1966.03615995003000050012x
- R Development Core Team (2008). *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Shen, Z., Wang, D., Ruan, Y., Xue, C., Zhang, J., Li, R., et al. (2014). Deep 16S rRNA pyrosequencing reveals a bacterial community associated with banana *Fusarium* wilt disease suppression induced by bio-organic fertilizer application. *PLoS ONE* 9:e98420. doi: 10.1371/journal.pone.0098420
- Smith, R. G., Gross, K. L., and Robertson, G. P. (2008). Effects of crop diversity on agroecosystem function: crop yield response. *Ecosystems* 11, 355–366. doi: 10.1007/s10021-008-9124-5
- Soil Survey Staff (2014). *Key to Soil Taxonomy*. 12th edn. Washington, DC: United States Department of Agriculture (USDA), and Natural Resource Conservation Service (NRCS).
- Tian, W., Wang, L., Li, Y., Zhuang, K., Li, G., Zhang, J., et al. (2015). Responses of microbial activity, abundance, and community in wheat soil after three years of heavy fertilization with manure-based compost and inorganic nitrogen. *Agric. Ecosyst. Environ.* 213, 219–227. doi: 10.1016/j.agee.2015.08.009
- Walkley, A., and Black, I. A. (1934). An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Sci.* 37, 29–38. doi: 10.1097/00010694-193401000-00003
- Wang, B., Li, R., Ruan, Y., Ou, Y., Zhao, Y., and Shen, Q. (2015). Pineapple–banana rotation reduced the amount of *Fusarium oxysporum* more than maize–banana rotation mainly through modulating fungal communities. *Soil Biol. Biochem.* 86, 77–86. doi: 10.1016/j.soilbio.2015.02.021
- Wang, Y., Tu, C., Cheng, L., Li, C., Gentry, L. F., Hoyt, G. D., et al. (2011). Long-term impact of farming practices on soil organic carbon and nitrogen pools and microbial biomass and activity. *Soil Tillage Res.* 117, 8–16. doi: 10.1016/j.still.2011.08.002
- Ward, N. L., Challacombe, J. F., Janssen, P. H., Henrissat, B., Coutinho, P. M., Wu, M., et al. (2009). Three genomes from the phylum Acidobacteria provide insight into the lifestyles of these microorganisms in soils. *Appl. Environ. Microbiol.* 75, 2046–2056. doi: 10.1128/AEM.02294-08
- Xing, H. Q., Xiao, Z. W., Yan, J. Z., Ma, J. C., and Meng, Y. (2011). Effects of continuous cropping of maize on soil microbes and main soil nutrients. *Pratacult. Sci.* 28, 1777–1780. doi: 10.1081/CSS-200056917
- Xiong, W., Zhao, Q., Xue, C., Xun, W., Zhao, J., Wu, H., et al. (2016). Comparison of fungal community in black pepper-vanilla and vanilla monoculture systems associated with vanilla *Fusarium* wilt disease. *Front. Microbiol.* 7:117. doi: 10.3389/fmicb.2016.00117
- Xu, L., Ravnskov, S., Larsen, J., and Nicolaisen, M. (2012). Linking fungal communities in roots, rhizosphere, and soil to the health status of *Pisum sativum*. *FEMS Microbiol. Ecol.* 82, 736–745. doi: 10.1111/j.1574-6941.2012.01445.x
- Xun, W., Huang, T., Zhao, J., Ran, W., Wang, B., Shen, Q., et al. (2015). Environmental conditions rather than microbial inoculum composition determine the bacterial composition, microbial biomass and enzymatic activity of reconstructed soil microbial communities. *Soil Biol. Biochem.* 90, 10–18. doi: 10.1016/j.soilbio.2015.07.018
- Yuan, J., Chaparro, J. M., Manter, D. K., Zhang, R., Vivanco, J. M., and Shen, Q. (2015). Roots from distinct plant developmental stages are capable of rapidly selecting their own microbiome without the influence of environmental and soil edaphic factors. *Soil Biol. Biochem.* 89, 206–209. doi: 10.1016/j.soilbio.2015.07.009
- Zeng, Q., An, S., and Liu, Y. (2017). Soil bacterial community response to vegetation succession after fencing in the grassland of China. *Sci. Total Environ.* 609, 2–10. doi: 10.1016/j.scitotenv.2017.07.102
- Zhalnina, K., de Quadros, P. D., Camargo, F. A., and Triplett, E. W. (2012). Drivers of archaeal ammonia-oxidizing communities in soil. *Front. Microbiol.* 3:210. doi: 10.3389/fmicb.2012.00210
- Zhang, C., Liu, G., Xue, S., and Wang, G. (2016). Soil bacterial community dynamics reflect changes in plant community and soil properties during the secondary succession of abandoned farmland in the Loess Plateau. *Soil Biol. Biochem.* 97, 40–49. doi: 10.1016/j.soilbio.2016.02.013
- Zhu, C., Ling, N., Guo, J., Wang, M., Guo, S., and Shen, Q. (2016). Impacts of fertilization regimes on arbuscular mycorrhizal fungal (AMF) community composition were correlated with organic matter composition in maize rhizosphere soil. *Front. Microbiol.* 7:1840. doi: 10.3389/fmicb.2016.01840

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Song, Tao, Guo, Li and Chen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.