



## OPEN ACCESS

## EDITED BY

Rubeena Zakar,  
University of the Punjab, Pakistan

## REVIEWED BY

Iñaki Lete,  
Basque Biobank, Spain  
Kornelia Zareba,  
United Arab Emirates University, United Arab  
Emirates

## \*CORRESPONDENCE

Yi Dai  
jacquedai@vip.sina.com  
Xin Sun  
sunxin@wchscu.com

## SPECIALTY SECTION

This article was submitted to Maternal Health, a  
section of the journal Frontiers in Global  
Women's Health

RECEIVED 08 August 2022

ACCEPTED 26 September 2022

PUBLISHED 02 November 2022

## CITATION

Wang J, Deng K, Li L, Dai Y and Sun X (2022)  
Levonorgestrel-releasing intrauterine system vs.  
systemic medication or blank control for  
women with dysmenorrhea: Systematic review  
and meta-analysis of randomized controlled  
trials.  
Front. Glob. Womens Health 3:1013921.  
doi: 10.3389/fgwh.2022.1013921

## COPYRIGHT

© 2022 Wang, Deng, Li, Dai, Sun. This is an  
open-access article distributed under the terms  
of the [Creative Commons Attribution License  
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in  
other forums is permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication in this  
journal is cited, in accordance with accepted  
academic practice. No use, distribution or  
reproduction is permitted which does not  
comply with these terms.

# Levonorgestrel-releasing intrauterine system vs. systemic medication or blank control for women with dysmenorrhea: Systematic review and meta- analysis of randomized controlled trials

Jing Wang<sup>1</sup>, Ke Deng<sup>1</sup>, Ling Li<sup>1</sup>, Yi Dai<sup>2\*</sup> and Xin Sun<sup>1\*</sup>

<sup>1</sup>Chinese Evidence-based Medicine Center, West China Hospital, Sichuan University, Chengdu, China,

<sup>2</sup>Department of Obstetrics and Gynecology, Peking Union College Hospital, Chinese Academy of  
Medical Sciences & Peking Union Medical College, National Clinical Research Center for Obstetrics &  
Gynecologic Diseases, Beijing, China

**Aims:** To compare efficacy and safety of the levonorgestrel-releasing  
intrauterine system (LNG-IUS) with systemic medication or blank control in  
the treatment of dysmenorrhea.

**Methods:** PubMed, EMBASE, the China National Knowledge Infrastructure  
(CNKI) and Wanfang Data were searched to collect randomized controlled  
trials (RCTs) comparing LNG-IUS with systemic medication or blank control  
among women diagnosed with primary dysmenorrhea or secondary  
dysmenorrhea (adenomyosis or endometriosis) from inception to 2020.04.  
Der Simonian-Laird random-effect model was used to pool data.

**Results:** Seventy-one RCTs (6551 patients) were included. Overall bias risk was  
medium. Sixty-two articles enrolled patients with adenomyosis; LNG-IUS  
significantly reduced the visual analogue scale (VAS) score compared with the  
systemic medication group among adenomyosis women at 3 months  
(standardized mean difference (SMD) = -0.81, 95% confidence interval (CI)  
-1.22 to -0.40); 6 months (SMD = -1.25, 95%CI: -1.58 to -0.92); 9 months  
(SMD = -1.23, 95%CI: -1.63 to -0.83); 12 months (SMD = -1.66, 95%CI: -2.14  
to -1.18). No difference was found in the incidence of irregular vaginal  
bleeding (16 RCTs; RR = 0.91, 95%CI: 0.62-1.33,  $P = 0.63$ ,  $I^2 = 4\%$ ) and other  
adverse outcomes. Sensitivity analysis regarding randomization methods was  
robust. Nine RCTs enrolled endometriosis women. Pooling results showed no  
significant difference between LNG-IUS and systemic medication treatment in  
terms of VAS at 6 months (SMD = -0.27, 95% CI: -0.97-0.43). Moreover, LNG-  
IUS was associated with higher risk of irregular vaginal bleeding (26.8% vs. 0).

**Conclusions:** LNG-IUS was associated with a reduced severity of dysmenorrhea  
compared with systemic medication; it was also beneficial for better control of  
menstrual blood loss and fewer adverse outcomes. Owing to small sample  
sizes, further well-designed RCTs are warranted to confirm these findings and  
long-term effects of LNG-IUS in the treatment of dysmenorrhea.

**Systematic Review Registration:** <https://www.crd.york.ac.uk/prospero/>, identifier:  
CRD42021228343.

## KEYWORDS

dysmenorrhea, efficacy, levonorgestrel-releasing intrauterine system, meta-analysis, safety

## Introduction

Dysmenorrhea is the most common gynecologic disorder, and it affects about 60%–90% women of reproductive age (1, 2). It is characterized by crampy lower abdominal pain, nausea, vomiting, and headaches, and it has a significant impact on ability to study, ability to work, and daily life (3). Oral contraceptive pills (OCPs) are used as first-line hormonal therapy for dysmenorrhea. They reduce uterine production of prostaglandin, which causes experienced pain (3). However, owing to the strict dosing cycle required when taking the pills, there is a risk of forgetting to take them.

The levonorgestrel-releasing intrauterine system (LNG-IUS) is an IUD. And LNG-IUS has three models, respectively are Mirena, Kyleena, Jaydess. But only Mirena went into public in China. Mirena releases 20 µg of levonorgestrel per day into the uterine cavity for a period of 5 years. Kyleena contains 19.5 mg of levonorgestrel (LNG) released *in vivo* at a rate of approximately 17.5 mcg/day after 24 days. This rate decreases progressively to 9.8 mcg/day after 1 year and to 7.4 mcg/day after 5 years. Levonorgestrel belongs to the class of medications called progestins, which is a hormone produced by the ovaries. This is a soft, flexible T-shaped contraceptive (birth control) device that is placed inside the uterus (intrauterine device or IUD). The medication is continuously released over a period of 3 years to prevent pregnancy. LNG-IUS was first designed as a contraceptive in 1970; it reduces the experience of pain and avoids the need to regularly take long-term medication. In recent years, LNG-IUS has also been officially approved for the treatment of heavy menstrual bleeding (4) and also been recommended for the treatment of dysmenorrhea when the patients are not currently planning pregnancy (2, 5, 6). But in China, it has not been recommended to women with dysmenorrhea yet. Although several small-sample randomized controlled trials (RCTs) indicate that LNG-IUS is likely beneficial for reducing the pain in women with endometriosis or adenomyosis (7–10), the results were not consistent. Furthermore, there is no existing systematic review about the efficacy and safety of LNG-IUS comparing systemic medication in the treatment of dysmenorrhea. Therefore, we conducted a systematic review and meta-analysis of RCTs to determine the efficacy and safety of LNG-IUS compared to other systemic medication or blank control in women with dysmenorrhea. Additionally, we identify the possible explanations for the heterogeneity through subgroup analysis.

## Methods

### Design

We performed a systematic review with meta-analysis using the protocol registered with PROSPERO (CRD42020203343).

### Eligibility criteria

We included RCTs comparing LNG-IUS with systemic medication or blank control among women diagnosed with primary dysmenorrhea or secondary dysmenorrhea (adenomyosis or endometriosis). We excluded studies that did not report on our interested outcomes and those that were not published in English or Chinese. The primary outcome was defined as the degree of dysmenorrhea (measured using visual analogue scale [VAS] scores) at 3, 6, 9, and 12 months, and blood loss (measured using a pictorial blood loss assessment chart [PBAC] scores, or menstrual blood volume) at 3, 6, 9, and 12 months. The secondary outcome was defined as quality of life (measured using standard scales like short form-questionnaire-36 [SF-36]) and adverse events (irregular vaginal bleeding, amenorrhea, headache, nausea, pelvic pain, acne, ovarian cyst, or weight gain).

### Literature search

We searched PubMed, EMBASE, the China National Knowledge Infrastructure (CNKI) and Wanfang Data, from inception to April 2020, with language limitation to English and Chinese. We used MeSH/Emtree terms and free-text to generate the search strategy, which comprised the terms “levonorgestrel releasing intrauterine system”, “LNG-IUS”, “dysmenorrhea”, “painful menses”, “menstrual pain”, “chronic pelvic pain”, and “dyspareunia”. The details of our search strategy are listed in [Supplementary S1](#).

### Study process

Two paired reviewers who were well-trained in research methodology, independently screened titles, abstracts, and full texts for eligibility; assessed risk of bias; and collected data from eligible studies, using standardized, pilot tested forms with detailed instructions. Reviewers resolved disagreement

through discussion or through adjudication by a third reviewer (LL) if agreement could not be reached.

## Risk of bias assessment

We used the Cochrane Handbook 5.1.0 bias assessment tool (11, 12) to assess the risk of bias of included studies. The items include sequence generation, allocation concealment, blinding of participants, blinding of intervention providers, blinding of outcome assessment, incomplete outcome data, selective outcome reporting, and other source of bias. We assigned items that were addressed “definitely or probably yes” to “high risk of bias”; and items that were addressed “definitely or probably no” to “low risk of bias”; items that did not report on our interested information as “unclear”.

## Data extraction

The following information from all eligible studies were collected:

- Study characteristics: first author, country, publication year, study design, inclusion and exclusion criteria, number of study sites, sample sizes, length of follow-up.
- Patient characteristics: age, parity
- Intervention: details of LNG-IUS and systemic medication of the control group
- Outcomes: degree of dysmenorrhea (VAS score), blood loss volume (PBAC score), quality of life, side effects (number of events and patients included for analyses in each group).

## Data analysis

For dichotomous outcomes, we calculated risk ratio (RR) and their 95% confidence intervals (CIs). For continuous outcomes, we calculated mean difference (MD) or standardized mean difference (SMD) and their 95% CIs. Statistical heterogeneity was examined by  $I^2$  and  $\chi^2$  test. If  $I^2 > 50\%$  or  $\chi^2$  test indicated  $P < 0.1$ , we defined it as statistical significance and we used the DerSimonian-Laird random-effect model to pool data. Subgroup analysis was undertaken to explore the potential heterogeneity based on systemic medication types (mifepristone, gestrinone, methyltestosterone, triptorelin, Desogestrel-Ethinyl Estradiol, drospirosterone-ethinylestradiol, and other types), status of surgery and patient type (primary dysmenorrhea patients vs. secondary dysmenorrhea patients). Sensitivity analysis was also conducted by excluding studies with improper randomization sequence generation or unspecified randomization methods and excluding studies with sample size less than 50. Data analyses were undertaken by Review


Manager 5.3.5. Publication bias was examined by Egger’s test and funnel plots.

## Results

Among 2641 identified publications, reviewers selected 71 RCTs (13–76, 78–84), enrolling 6551 patients, to be included in this meta-analysis (Figure 1).

## Study characteristics

Of 71 included studies, eight (73–76, 78, 82–84) were published in English and 63 (13–72, 79–81, 86) were published in Chinese. The included studies were conducted in China mainland ( $n = 64$ ), Brazil ( $n = 3$ ), Egypt ( $n = 1$ ), Thailand ( $n = 1$ ), Turkey ( $n = 1$ ), Taiwan ( $n = 1$ ). Seventy studies were single center clinical trials, and one study (74) was a multicenter clinical trial. All the LNG-IUS used in article are Mirena. The details of the characteristics of the study are summarized in **Supplementary Table S1**.



## Risk of bias assessment

The results of risk of bias assessment are presented in **Supplementary Table S2**. Twenty-nine studies described adequate randomization methods while the remaining 42 studies did not report randomization methods. Eight studies reported using sealed and opaque envelopes for allocation concealment. One study reported methods for blinding participants and another trial reported methods for blinding outcome assessment. Six studies reported loss of follow-up. The overall risk of bias was medium. Sensitivity analysis considering whether randomization methods were adequate did not show a significant difference (See the online **Supplementary Table S2**).

## Efficacy and safety results in women with adenomyosis

Of the 71 RCTs included, 62 (13–23, 25–36, 38–49, 51–56, 58–72, 76, 78–81, 86) enrolled women patients with adenomyosis, among which 42 (13, 16, 17, 19–23, 25, 27, 31, 32, 34–36, 38–42, 44, 45, 48, 49, 51–53, 55, 56, 58, 60–62, 64, 66, 68–71, 76, 81, 86) studies reported VAS score. Pooling data from these 42 RCTs showed that LNG-IUS significantly reduced the VAS score in the comparison of a medication group among women with adenomyosis at 3 months (17 RCTs; SMD = -0.81, 95% CI = -1.22–(-0.4),  $P < 0.0001$ ;  $I^2 = 93\%$ ); 6 months (36 RCTs; SMD = -1.25, 95% CI: -1.58–(-0.92),  $P < 0.0001$ ;  $I^2 = 94\%$ ); 9 months (2 RCTs; SMD = -1.23, 95% CI: -1.63–(-0.83),  $P < 0.0001$ ,  $I = 0$ ); and 12 months (18 RCTs; SMD = -1.66, 95% CI: -2.14–(-1.18),  $P < 0.0001$ ,  $I^2 = 94\%$ ; **Figure 2**).

Pooling across 16 (16, 17, 27, 32, 36, 49, 53, 55, 56, 60, 62, 65, 70, 71, 78) trials reporting PBAC score showed that the reduction of PBAC score was greater in LNG-IUS groups at 3 months (6 RCTs; MD = -7.33, 95% CI: -11.39–(-3.27),  $P = 0.0004$ ,  $I^2 = 36\%$ ) and 6 months (15 RCTs; MD = -12.41, 95% CI: -16.18–(-8.65),  $P < 0.0001$ ,  $I^2 = 96\%$ ). However, no difference was identified between these groups at 9 months (2 RCTs; MD = -2.78, 95% CI: -6.89–1.34,  $P = 0.19$ ,  $I^2 = 0\%$ ) and 12 months (3 RCTs; MD = -12.13, 95% CI: -25.84–1.58,  $P = 0.08$ ,  $I^2 = 94\%$ ; **Supplementary Figure S1**).

Nineteen RCTs (21, 33–36, 38, 41, 42, 44, 45, 48, 52, 54, 63, 64, 66, 69, 86) that reported menstrual blood loss were included for analysis. The pooling results showed that the reduction in menstrual blood loss was greater in the LNG-IUS group at 3 months (2 RCTs; MD = -29.52, 95% CI: -74.35–15.31),  $P < 0.0001$ ,  $I^2 = 98\%$ ), 6 months (15 RCTs; MD = -18.97, 95% CI: -28.27–(-9.67),  $P < 0.0001$ ,  $I^2 = 96\%$ ), and 12 months (7 RCTs; MD = -40.27, 95% CI: -53.64–(-26.90),  $P < 0.0001$ ,  $I^2 = 96\%$ ; **Supplementary Figure S2**).

As for quality-of-life (QoL) assessment, Feng (18) reported quality of life using the Nottingham Health Profile (NHP, wherein high scores indicated good outcomes) including pain, vigor, sleep, emotion, activity ability, and social loneliness. The results showed no difference in the changes in NHP scores at 6 months from baseline between the LNG-IUS and gestrinone group ( $P > 0.05$ ). Li (28) reported quality of life using the Short-Form 36 Health Survey Questionnaire (SF-36; wherein high scores indicated good outcomes). The results showed that LNG-IUS improved SF-36 score compared to triptorelin in both physical and psychological areas after 6 months follow-up ( $P < 0.001$ ).

There was no significant difference in the incidence of irregular vaginal bleeding (16 RCTs; RR = 0.91, 95% CI: 0.62–1.33,  $P = 0.63$ ,  $I^2 = 4\%$ ), headache (3 RCTs; RR = 0.94, 95% CI: 0.28–3.19,  $P = 0.92$ ,  $I^2 = 7\%$ ), acne (11 RCTs; RR = 0.48, 95% CI: 0.22–1.07,  $P = 0.07$ ,  $I^2 = 21\%$ ), or amenorrhea (7 RCTs; RR = 1.06, 95% CI: 0.34–3.3,  $P = 0.92$ ,  $I^2 = 39\%$ ) between two groups. The systemic medication group was associated with a higher risk of nausea (6 RCTs; RR = 0.23, 95% CI: 0.08–0.64,  $P = 0.005$ ,  $I^2 = 0\%$ ) and menorrhagia (1 RCT; RR = 0.55, 95% CI: 0.30–0.98,  $P = 0.04$ ; **Figure 3**).

Subgroup analysis by type of systemic medications showed a significant difference in VAS score ( $P < 0.0001$ ), while no significant difference was found in subgroup analysis according to whether the patients received surgery or not ( $P = 0.91$ ; **Supplementary Table S4**). Triptorelin could better reduce VAS score compared with LNG-IUS (SMD = 2.17, 95% CI: 1.37–2.96,  $P < 0.0001$ ) after 6 months of treatment.

Sensitivity analysis showed that results were robust after excluding articles with unspecific randomization ( $P < 0.0001$  vs.  $P < 0.001$ ; **Supplementary Table S3**) and excluding studies with sample size is less than 50 ( $P < 0.0001$  vs.  $P < 0.0001$ ; **Supplementary Table S3**). No publication bias was detected using funnel plot and Egger's test with regard to VAS score at 3, 6, and 12 months, PBAC score at 6 months, menstrual blood loss at 6 months, acne, and irregular vaginal bleeding ( $P > 0.05$ ; **Supplementary Table S5**).

## Efficacy and safety results in women with endometriosis

Nine out of all 71 RCTs (37, 50, 57, 73, 74, 82–84, 87) included had enrolled women with (surgically diagnosed) endometriosis, among which six (37, 57, 74, 75, 83, 84) studies reported VAS score. Pooling results showed no significant difference between LNG-IUS and systemic medication treatment in terms of VAS at 6 months (5 RCTs; SMD = -0.27, 95% CI: -0.97–0.43,  $P = 0.45$ ,  $I^2 = 90\%$ ) and 12 months (2 RCTs; SMD = -0.94, 95% CI: -2.16–0.27,  $P = 0.13$ ,  $I^2 = 89\%$ ; **Figure 4**).




FIGURE 2 VAS score among adenomyosis patients receiving treatment with LNG-IUS vs. systemic medication in randomized controlled studies.




FIGURE 3

Adverse events among adenomyosis patients receiving treatments of LNG-IUS vs. systemic medication in randomized controlled studies.




FIGURE 4

VAS score among endometriosis patients receiving treatments of LNG-IUS vs. systemic medication in randomized controlled studies.

As for QoL assessment, Petta (74) used the Psychological General Well-being Index (PGWBI) for the assessment, and no significant difference was identified in the changes in the PGWBI scores from baseline between the LNG-IUS and Triptorelin groups ( $P=0.78$ ). Nelsilene (83) reported on QoL using Endometriosis Health Profile-30 (EHP-30), and their results showed that EHP-30 was significantly higher in a LNG-IUS group in emotional situation ( $P=0.04$ ) and self-image ( $P=0.04$ ) when compared to an etonogestrel implant group. Prasong (75) used the Short-Form 36 Health Survey Questionnaire (SF-36) to assess patients' QoL. The results showed that the SF-36 at 12 months was higher in the LNG-IUS group than in the expectant group in physical domains ( $P<0.05$ ) but there was no difference in psychological domains ( $P=0.229$ ).

Three studies (37, 50, 75) reporting irregular vaginal bleeding were included. LNG-IUS was associated with a higher risk of irregular vaginal bleeding (26.8%), and no similar events were observed in the control group. The results demonstrated that no significant differences were found for acne (2 RCTs; RR = 1.07, 95% CI: 0.69–1.67,  $P=0.07$ ,  $I^2=0\%$ ), nausea (2 RCTs; RR = 0.98, 95% CI: 0.55–1.75,  $P=0.94$ ,  $I^2=0\%$ ), headache (2 RCTs; RR = 0.84, 95% CI: 0.46–1.52,  $P=0.56$ ,  $I^2=55\%$ ), weight gain (3 RCTs; RR = 1.14, 95% CI: 0.76–1.71,  $P=0.53$ ,  $I^2=0\%$ ), or amenorrhea (4 RCTs; RR = 1.41, 95% CI: 0.06–31.51,  $P=0.83$ ,  $I^2=70\%$ ) between the two groups (Figure 5).

Owing to the small sample size of included studies, we did not perform subgroup analysis by systemic medication type. No significance was found in subgroup analysis according to surgical status, which examined whether surgery previously

undergone by patients influenced the efficacy of LNG-IUS. (Supplementary Table S4).

Sensitivity analysis showed that results were robust after excluding articles with unspecific randomization ( $P=0.45$  vs.  $P=0.38$ ; Supplementary Table S3). We did not examine publication bias among women with endometriosis because the number of included articles was less than 10.

## Discussion

### Findings and interpretations

Our study found that LNG-IUS may be more effective in alleviating adenomyosis-associated dysmenorrhea compared to systemic medication at 3, 6, 9, and 12 months. Also, LNG-IUS may be superior to systemic medication in reducing uterine blood loss at 3, 6, and 12 months among women with adenomyosis. However, among women with endometriosis, we observed no significant differences in the VAS score of LNG-IUS vs. systemic medication at 6 and 12 months. Most of the studies included did not report the method of randomization which indicates the low quality of these researches but the sensitivity analysis did not modify the results of our review when we excluded the studies without specific randomization method.

Dysmenorrhea includes both primary and secondary dysmenorrhea. The underlying pathology of primary dysmenorrhea is the increased local secretion of vasopressin and prostaglandins (77). Secondary dysmenorrhea is painful menses caused by pelvic pathology, most commonly




FIGURE 5 Adverse events among endometriosis patients receiving LNG-IUS vs. systemic medication.

endometriosis, followed by adenomyosis, infection, and myomas (2). Although the inclusion criteria for our study contained both primary and secondary dysmenorrhea, after screening abstracts and full-texts, we found no RCTs evaluating the effectiveness of LNG-IUS in primary dysmenorrhea. The probable reason may

be that LNG-IUS's usage is relatively low in primary dysmenorrhea as discussed previously (87).

According to the results, LNG-IUS might be more effective for pain relief in women with adenomyosis, which is in accordance with findings of several existing studies (9, 88).



The exact mechanism of pain control in adenomyosis is still unclear, but it may be associated with the high concentration of LNG on the endometrium (9), which induces glandular atrophy and stromal decidualization, inhibiting prostaglandin synthesis (8). Prostaglandin is a substance that causes pain and uterine contraction while high levels of LNG inhibit prostaglandin synthesis.

Various potential mechanisms have been proposed for this device in women with endometriosis. Whatever the exact mechanism is, the local effect of the progestogen on the endometrium resulting in hypomenorrhoea or amenorrhoea significantly improves the pain of dysmenorrhoea and menorrhagia. A common adverse event of LNG-IUS is irregular vaginal bleeding (75). LNG-IUS groups were associated with higher risk of irregular vaginal bleeding (26.8% vs. 0) in women with endometriosis. However, there were no differences in irregular vaginal bleeding between the systemic medication and LNG-IUS groups in adenomyosis. Furthermore, there were no differences of incidence of headache, acne, or amenorrhea between the systemic medication and LNG-IUS groups. The systemic medication group was associated with a higher risk of nausea (RR:0.23, 95% CI: 0.08–0.64,  $P=0.95$ ) and menorrhagia (RR:0.55, 95% CI: 0.30–0.98,  $P=0.04$ ) in adenomyosis. We could not draw any conclusions according to available data because few studies (18, 28, 74, 75, 83) have reported on quality of life and the scales they adopted differed. More consistent studies are warranted to verify the impact of LNG-IUS on quality of life. Therefore, we suggest LNG-IUS should be offered to women with adenomyosis because it has the ability to alleviate dysmenorrhea in short time (less than 1 year). For women with endometriosis, then medication therapy was more recommended taking into account the higher risk of irregular vaginal bleeding.

## Strengths and limitations

Our study has several strengths. Firstly, to the best of our knowledge, this is the first meta-analysis to systematically compare the efficiency and safety through multiple outcomes of LNG-IUS and systemic medication among dysmenorrhea patients. Secondly, we have included women with both endometriosis and adenomyosis dysmenorrhea and conducted separate analyses. Thirdly, we included multiple control groups, comprehensively comparing the different systemic medication types. Finally, we conducted several pre-specified subgroup analyses to explore the potential source of heterogeneity.

The review also had some limitations. Firstly, most of our studies enrolled Chinese patients, which may limit the generalization of results to other ethnic groups. Majority of the studies included were published in Chinese, which limits their availability for non-Chinese speakers. Secondly, most included

studies had a relatively high risk of bias, which may limit the reliability of meta-analysis results. Thirdly, high heterogeneity exists in our review. Although we have tried some subgroup analysis in terms of systemic medication type and status of surgery, we still failed to find the source of heterogeneity. Insufficient information concerning characteristics of included studies made us unable to conduct further subgroup analyses (e. g. BMI) to explore the source of heterogeneity, which also brought limitations to the reliability of our results. Fourth, we did not include “endometriosis” or “adenomyosis” as search words because we think dysmenorrhea was probably mentioned as their symptoms, which may have greatly limited finding valid studies for the research focus.

Above all, there is still a need for more rigorous, well-designed and high-quality RCTs to investigate the long-term effect of LNG-IUS to confirm the current findings and to examine other vital outcomes such as number of dropouts and withdrawal of treatment.

## Conclusions

In conclusion, our systematic review demonstrated that the use of LNG-IUS was associated with a reduced severity of dysmenorrhea compared with systemic medication. Moreover, LNG-IUS was beneficial for improved control of menstrual blood loss and fewer adverse outcomes. At the same time, further well-designed RCTs are warranted to confirm the current findings and long-term effect of LNG-IUS in the treatment of dysmenorrhea.

## Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## Author contributions

YD and XS conceived the topic, JW and KD screened the articles, JW and LL extracted the data and JW, KD and LL wrote the article. All authors contributed to the article and approved the submitted version. YD and XS are correspondence authors.

## Funding

This article was supported by Sichuan Youth Science and Technology Innovation Research Team (Grant No. 2020JDTD0015); 1-3-5 Project for Disciplines of Excellence, of the West China Hospital, Sichuan University (Grant No.

ZYYC08003); and the Non-profit Central Research Institute Fund of Chinese Academy of Medical Sciences (2021-PT320-001).

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of

their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgwh.2022.1013921/full#supplementary-material>.

## References

- Ju H, Jones M, Mishra G. The prevalence and risk factors of dysmenorrhea. *Epidemiol Rev.* (2014) 36:104–13. doi: 10.1093/epirev/mxt009
- Care COaH. Dysmenorrhea and endometriosis in the adolescent. *Am J Obstet Gynecol.* (2018) 132:e249–58. doi: 10.1097/AOG.0000000000002978
- French L. Dysmenorrhea. *Am Fam Physician.* (2005) 71:285–91.
- Heikinheimo O, Gemzell-Danielsson K. Emerging indications for the levonorgestrel-releasing intrauterine system (LNG-IUS). *Acta Obstet Gynecol Scand.* (2012) 91:3–9. doi: 10.1111/j.1600-0412.2011.01303.x
- National Guideline Alliance (UK). Endometriosis: diagnosis and management. London: National Institute for Health and Care Excellence (NICE) (2017). PMID: 29787038
- Burnett M, Lemyre M. No. 345-primary dysmenorrhea consensus guideline. *J Obstet Gynaecol Can.* (2017) 39:585–95. doi: 10.1016/j.jogc.2016.12.023
- Yoost J, Lajoie AS, Hertweck P, Loveless M. Use of the levonorgestrel intrauterine system in adolescents with endometriosis. *J Pediatr Adolesc Gynecol.* (2013) 26:120–4. doi: 10.1016/j.jpag.2012.11.002
- Bragheto AM, Caserta N, Bahamondes L, Petta CA. Effectiveness of the levonorgestrel-releasing intrauterine system in the treatment of adenomyosis diagnosed and monitored by magnetic resonance imaging. *Contraception.* (2007) 76:195–9. doi: 10.1016/j.contraception.2007.05.091
- Sheng J, Zhang WY, Zhang JP, Lu D. The LNG-IUS study on adenomyosis: a 3-year follow-up study on the efficacy and side effects of the use of levonorgestrel intrauterine system for the treatment of dysmenorrhea associated with adenomyosis. *Contraception.* (2009) 79:189–93. doi: 10.1016/j.contraception.2008.11.004
- Lethaby A, Hussain M, Rishworth JR, Rees MC. Progesterone or progestogen-releasing intrauterine systems for heavy menstrual bleeding. *Cochrane Database Syst Rev.* (2005) 19(4):CD002126. doi: 10.1002/14651858.cd002126pub4
- Higgins JP, Altman DG, Sterne Jonathan AC, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ.* (2011) 343:d5928. doi: 10.1136/bmj.d5928
- Wang Y, Li L, Deng K, Liu J, Liu Y, Zou K, et al. Comparison of the combination of recombinant follicle-stimulating hormone and recombinant luteinizing hormone protocol versus human menopausal gonadotropin protocol in controlled ovarian stimulation: a systematic review and meta-analysis. *J Evid Based Med.* (2020) 13:215–26. doi: 10.1111/jebm.12390
- Zeng CQZGW. Clinical observation of levonorgestrel intrauterine device and pregnenolone in the treatment of uterine adenomyosis. *Shenzhen J Integr Chin and Western Med.* (2016) 26:107–9. doi: 10.16458/j.cnki.1007
- Chen C. Evaluation of the therapeutic effect of manuelle on dysmenorrhea and increased menstrual flow in patients with adenomyosis. *China Health Nutr.* (2018) 31–2.
- Chen YD. Clinical study of levonorgestrel intrauterine delayed-release system in the treatment of dysmenorrhea and increased menstrual flow in patients with adenomyosis. *Pract Med in China.* (2018) 13:142–3. doi: 10.14163/j.cnki.11-5547/r.2018.18.083
- Cheng QH. Clinical efficacy analysis of leuprolide acetate combined with mannitol in the treatment of adenomyosis uteri. *Diet and Health.* (2018) 5:75.
- Dou LL. Effectiveness of intrauterine placement of the intrauterine device (IUD) for the treatment of adenomyosis. *J Henan Higher Inst of Med Sci.* (2019) 31:602–5.
- Feng LL. Clinical analysis of the treatment of adenomyosis by manuelle. *Pract Integr Chin and Western Med Clin.* (2018) 18:146–8.
- Fu F, Chen YY. A comparative study of the efficacy of mannitol and GnRH-a for pain and recurrence of endometriosis. *Med Infor.* (2014) 27:83–4.
- Huang LD, Wu Y, Chen CY, Huang WR, Yang X, Wei LY. Comparison of the efficacy of placing the mannorrhea ring and oral contraceptive pills in the treatment of adenomyosis. *Chin Clin Med.* (2018) 11:632–5.
- Jiang YP. Comparative study of the efficacy of levonorgestrel intrauterine extended-release system and low-dose mifepristone in the treatment of adenomyosis. *China Fam Med.* (2017) 20:111–2.
- Jin YQ. Clinical observation of levonorgestrel intrauterine device system for the treatment of adenomyosis. *Northern Pharm.* (2017) 14:8–9.
- Kang XW, Zhan HY. Clinical observation of levonorgestrel intrauterine extended-release system and pregnenolone in the treatment of adenomyosis. *J Gynaecol Endocrinol.* (2018) 5:126–7.
- Kong CY. Analysis of 126 cases of adenomyosis with dysmenorrhea and increased menstrual flow treated with manuelle. *China Matern and Child Health.* (2015) 30:4067–8.
- Li AG, Wu Q, Huo PW, Gao LL. Effectiveness of the combination of mannorrhea and mafulon in the pharmacological treatment of adenomyosis. *Northern Pharm.* (2018) 15:54.
- Li HH. The therapeutic effect of manuelle on dysmenorrhea and increased menstrual flow in patients with adenomyosis. *Mother and Baby World.* (2018) 2:82–4.
- Li QY, Liu XR, Feng YZ. 49 Cases of leuprolide combined with mannitol in the treatment of uterine adenomyosis. *China Pharmaceutical.* (2015) 24:85–6.
- Li SJ. Clinical value of follow-up treatment of adenomyosis after laparoscopic lesion resection with leuprolide combined with mannitol. *Massage and Rehabil Med.* (2019) 10:44–6.
- Li XY. Effectiveness of manuelle combined with laparoscopic surgery in the treatment of adenomyosis. *Henan Med Res.* (2018) 27:1292–3.
- Li XH. Laparoscopic wedge hysterectomy with postoperative vaginal placement of levonorgestrel intrauterine delayed release system in the treatment of adenomyosis. *Clinical Rational Drug use.* (2018) 11:140–1.
- Li XQ, Tao LN, Ren HY. Clinical efficacy and safety analysis of intrauterine delayed-release system of pregnenolone and levonorgestrel in the treatment of

- uterine adenomyosis. *Med Clin Res.* (2017) 34:46–8. doi: 10.3969/j.issn.1671-7171.2017.01.015
32. Li Y, Ding J, Zhao CM. A comparative clinical study on the treatment of adenomyosis of the uterus with mifepistone in combination with LNG-IUS. *Western Med.* (2018) 30:716–9.
33. Li YF. Clinical observation on the treatment of adenomyosis with intrauterine placement of the mannedel birth control ring. *Health for All.* (2016) 10:173.
34. Li Y Q. Clinical efficacy of levonorgestrel intrauterine extended-release system in the treatment of adenomyosis. *Primary Care Forum.* (2015) 19:32–3.
35. Liu HL, Cai CY, He YL, Et A. Effectiveness of levonorgestrel intrauterine extended-release system in the treatment of adenomyosis. *Modern Drug Appli in China.* (2017) 11:18–9.
36. Liu JN. Clinical effect of levonorgestrel intrauterine device system in the treatment of 86 cases of adenomyosis with dysmenorrhea and increased menstrual flow. *China Health Standards Management.* (2016) 7:101–2. doi: 10.3969/j.issn.1674-9316.2016.01.074
37. Liu X, Zhang LY, Li Y, Et YA. Effectiveness of levonorgestrel intrauterine extended release system in postoperative patients with endometriosis. *Chin Contemp Med.* (2014) 21:28–35.
38. Ma XH. Evaluation of the efficacy of the intrauterine device, mandedol, in the treatment of uterine adenomyosis. *J Henan Sur.* (2016) 22:47–9.
39. Ma YJ, Ma XL. Efficacy of dissemination and analgesia capsules combined with manuelle in the treatment of uterine adenomyosis. *Med Res and Edu.* (2014) 31:16–9.
40. Meng YY, Xiao YB. Clinical study of levonorgestrel intrauterine extended release system for the treatment of adenomyosis. *J Clin Med Lit.* (2018) 5:22.
41. Re D. Clinical efficacy analysis of the combination of levonorgestrel intrauterine device system and mifepristone in the treatment of dysmenorrhea and increased menstrual flow in adenomyosis. *Elect J Pract Gynecol Endocrinol.* (2019) 6:118.
42. Ren FL. Clinical efficacy of manuelle in treating 114 cases of adenomyosis with dysmenorrhea and increased menstrual flow. *World Med News Digest.* (2017) 17:114. doi: 10.3969/j.issn.1671-3141.2017.16.100
43. Shang YHXZB. Clinical study of 100 cases of adenomyosis treated with manuelle. *Chin Pract Med.* (2014) 41:69–71.
44. Shi LH. A comparative study of the efficacy of mannacel ring and mifepristone in the treatment of adenomyosis. *J Contemp Clin Med.* (2016) 29:2533–4.
45. Sun GZ. Clinical analysis of 41 cases of adenomyosis treated with levonorgestrel intrauterine release system. *Chin Pract Med.* (2017) 12:140–1. doi: 10.14163/j.cnki.11-5547/r.2017.26.080
46. Sun LH, Wang XH. The efficacy of leuprolide combined with mannitol in treating 33 cases of adenomyosis. *Health Care Guide.* (2017) 79.
47. Sun W. High-intensity focused ultrasound combined with levonorgestrel intrauterine device system for the treatment of adenomyosis. *China Prescription Drugs.* (2017) 17:87–8.
48. Tao JQ. Effectiveness of levonorgestrel intrauterine device system in the treatment of adenomyosis disease with dysmenorrhea and increased menstrual flow. *Med Food Therapy and Health.* (2019) 4:33–4.
49. Tao YS, Wu H. Clinical observation on the treatment of adenomyosis of the uterus by the combination of the mannorrhea ring and endometrial ablation. *Jilin Med.* (2016) 37:642–3.
50. Tian J. Analysis of the long-term efficacy of levonorgestrel intrauterine extended-release system in the treatment of endometriosis. *Modern Diagn and Treatment.* (2017) 5:852–4.
51. Tu J, Gao Y, Huang ML, Xiao YB. Study on the effect of levonorgestrel intrauterine extended release system on menstruation and ovarian function in patients with adenomyosis. *China Family Med.* (2016) 19:427–30.
52. Tu L, Cao HB, Wu HH. Comparison of the effects of LNG-IUS and GnRHa regimens in the treatment of adenomyosis. *Shandong Med.* (2012) 52:7–9.
53. Wang CB. Exploration of leuprolide in combination with mannitol in adenomyosis with severe menstrual flow and dysmenorrhea. *Systemic Med.* (2017) 2:95–7. doi: 10.19368/j.cnki.2096-1782.2017.24.095
54. Wang HQ, Hu XJ, Liu F. Comparison of the efficacy of ursimine and mannitol in endometriosis. *China Maternal and Child Health Study.* (2018) 29:652–5.
55. Wang H. Analysis of the therapeutic effect of manuelle on dysmenorrhea and increased menstrual flow in patients with adenomyosis. *Laboratory Med and Clin.* (2014) 11:1339–40.
56. Wang J, Xv FL, Yang J. Analysis of the efficacy of levonorgestrel intrauterine extended-release system in patients with adenomyosis. *Primary Care Forum.* (2019) 23:3172–3. doi: 10.19435/j.1672-1721.2019.22.036
57. Wang LN. Analysis of the efficacy and safety of levonorgestrel intrauterine delayed-release system after endometriosis surgery. *Modern Drug Appli in China.* (2016) 10:5–6.
58. Wang QZ, Wu XF, He L. Analysis of the therapeutic effect of levonorgestrel intrauterine extended-release system on adenomyosis. *Hebei Med.* (2015) 21:805–7.
59. Wang YL. Analysis of the efficacy of mannorrhea in combination with mafron in the pharmacological treatment of adenomyosis. *China Health & Nut.* (2018) 202.
60. Wang ZP. Efficacy of laparoscopic lesion resection combined with levonorgestrel intrauterine delayed release system in adenomyosis. *Henan Med Res.* (2018) 27:1476–7.
61. Wu JJ, Feng GN. Clinical efficacy and safety of levonorgestrel intrauterine delayed-release system in the treatment of uterine adenomyosis. *J Mudanjiang Med College.* (2017) 38:47–51.
62. Wu JC. Clinical efficacy of levonorgestrel intrauterine extended release system in patients with adenomyosis. *Modern Diagno and Treatment.* (2018) 4:636–8.
63. Yang LF. Analysis of the effect of levonorgestrel intrauterine extended-release system in the treatment of uterine adenomyosis. *J Pract Gynecolo Endocrinol.* (2016) 3:122–4.
64. Yao Q. Evaluation of the therapeutic effect of manuelle on dysmenorrhea and increased menstrual flow in patients with adenomyosis. *Chin and Foreign Women's Health Study.* (2018) 18:53–4.
65. Sun YHS. Analysis of clinical application of mandedrol in the treatment of adenomyosis. *Med Theory and Pract.* (2017) 30:562–3.
66. Yu R. Clinical effect analysis of levonorgestrel intrauterine extended-release system in the treatment of adenomyosis. *China Med Guide.* (2016) 14(154).
67. Zhang FJ. Effectiveness of levonorgestrel intrauterine device system in the treatment of adenomyosis disease with dysmenorrhea and increased menstrual flow. *Maternal and Child Health.* (2018) 5:66–7.
68. Zhang Q. Clinical study on the treatment of uterine adenomyoma by intrauterine placement and myomectomy with mandedel. *Modern Diagnosis and Treatment.* (2016) 23:4500–1.
69. Zhang XF. Comparison of the efficacy of the levonorgestrel intrauterine extended-release system with pregnenolone in the treatment of adenomyosis. *Chin J Pract Rural Doctor.* (2017) 24:62–5. doi: 10.3969/j.issn.1672-7185.2017.12.026
70. Zhang XG. Effect of laparoscopic lesion excision combined with levonorgestrel intrauterine delayed-release system on dysmenorrhea degree score and disease recurrence rate in patients with adenomyosis. *Med Theory and Pract.* (2019) 32:1040–2.
71. Zhao XJ, Wei K, Wang L, Zhang L. Efficacy of combined ultrasound ablation of LNG-IUS and GnRHa in the treatment of dysmenorrhea in adenomyosis. *Med Clin Res.* (2019) 36:309–11. doi: 10.3969/j.issn.1671-7171.2019.02.032
72. ZY J. Exploring the clinical value of using the levonorgestrel intrauterine delayed-release system in patients with dysmenorrhea due to adenomyosis. *World Med News Digest.* (2015) 15:94–5.
73. Chen YJ, Hsu TF, Huang BS, Tsai HW, Chang YH, Wang PH. Postoperative maintenance levonorgestrel-releasing intrauterine system and endometrioma recurrence: a randomized controlled study. *Am J Obstet Gynecol.* (2017) 216:582 e581–e589. doi: 10.1016/j.ajog.2017.02.008
74. Petta CA, Ferriani RA, Abrao MS, Hassan D, Silva JCRE, Podgaec S, et al. Randomized clinical trial of a levonorgestrel-releasing intrauterine system and a depot GnRH analogue for the treatment of chronic pelvic pain in women with endometriosis. *Hum Reprod.* (2005) 20:1993–8. doi: 10.1093/humrep/deh869
75. Tanmahasamut MR P, Angsuwathana S, Techtraisak K, Indhavivadhana S, Leerasing P. Postoperative levonorgestrel-releasing intrauterine system for pelvic endometriosis-related pain. *American College of Obstet and Gynecol.* (2012) 119:519–26. doi: 10.1097/AOG.0b013e31824264c3
76. Shaaban OM, Ali MK, Sabra AM, Abd El Al DE. Levonorgestrel-releasing intrauterine system versus a low-dose combined oral contraceptive for treatment of adenomyotic uteri: a randomized clinical trial. *Contraception.* (2015) 92:301–7. doi: 10.1016/j.contraception.2015.05.015
77. Wildemeersch D, Jandi S, Pett A, Hasskamp T. Management of primary dysmenorrhea in young women with frameless LNG-IUS. *Open Access Journal of Contraception.* (2014) 5:23–8. doi: 10.2147/OAJC.S52926

78. Wu Q, Lian Y, Chen L, Yu Y, Lin T. Alleviation of symptoms and improvement of endometrial receptivity following laparoscopic adenomyoma excision and secondary therapy with the levonorgestrel-releasing intrauterine system. *Reprod Sci.* (2020) 27:1259–65. doi: 10.1007/s43032-019-00130-4
79. Zhou J. Effectiveness of levonorgestrel intrauterine device system in the treatment of adenomyosis disease with dysmenorrhea and increased menstrual flow and analysis of the effect on VAS scores. *J Pract Gynecol Endocrinol.* (2018) 5:118–9.
80. Wang X. *A controlled clinical study of HIFU combined with LNG-IUS in the treatment of adenomyosis [D]*. Chongqing: Chongqing Medical University (2015). 10–6.
81. Zhu B. Clinical observation of hysteroscopic endometrial resection combined with levonorgestrel intrauterine delayed-release system in the treatment of adenomyosis. *China National Health Med.* (2019) 31:59–60.
82. Bayoglu Tekin Y, Dilbaz B, Altinbas SK, Dilbaz S. Postoperative medical treatment of chronic pelvic pain related to severe endometriosis: levonorgestrel-releasing intrauterine system versus gonadotropin-releasing hormone analogue. *Fertil Steril.* (2011) 95:492–6. doi: 10.1016/j.fertnstert.2010.08.042
83. Carvalho N, Margatho D, Cursino K, Benetti-Pinto CL, Bahamondes L. Control of endometriosis-associated pain with etonogestrel-releasing contraceptive implant and 52-mg levonorgestrel-releasing intrauterine system: randomized clinical trial. *Fertil Steril.* (2018) 110:1129–36. doi: 10.1016/j.fertnstert.2018.07.003
84. Gomes MK, Ferriani RA, Rosa E Silva JC, Japur De Sa Rosa E Silva AC, Vieira CS, Candido Dos Reis FJ. The levonorgestrel-releasing intrauterine system and endometriosis staging. *Fertil Steril.* (2007) 87:1231–4. doi: 10.1016/j.fertnstert.2006.11.044
85. The levonorgestrel-releasing intrauterine system (LNG-IUS) in contraception and reproductive health. *J Fam Plann Reprod Health Care.* (2004) 30:99–109. doi: 10.1783/147118904322995474
86. Kong CY. Analysis of 126 cases of adenomyosis with dysmenorrhea and increased menstrual flow treated with LNG-IUS. *China Mat and Child Health.* (2015) 30:4067–8.
87. Tanmahasamut P, Rattanachaiyanont M, Angsuwathana S, Techatrasak K, Indhavivadhana S, Leerasiri P. Postoperative levonorgestrel-releasing intrauterine system for pelvic endometriosis-related pain: a randomized controlled trial. *Obstet Gynecol.* (2012) 119:519–26. doi: 10.1097/AOG.0b013e31824264c3
88. Wildemeersch D. A frameless LNG-IUD may be preferred over framed LNG-IUD for the treatment of primary dysmenorrhea. *Hum Reprod Update.* (2016) 22:404–7. doi: 10.1093/humupd/dmv061
89. Sun C, Ren XY, Gao Y, Liang ZG, Mou M, Gu HF, et al. Clinical efficacy and safety of Major uterine wall resection and reconstruction of the uterus combined with LNG-IUS for the treatment of severe adenomyosis. *Geburtshilfe Frauenheilkd.* (2020) 80:300–6. doi: 10.1055/a-0995-2200