



## OPEN ACCESS

## EDITED BY

Anastasios Darras,  
University of Peloponnese, Greece

## REVIEWED BY

Pavlos Tsouvaltzis,  
Aristotle University of Thessaloniki, Greece  
Michele Valquíria Reis,  
Universidade Federal de Lavras, Brazil  
Stefanos Hatzilazarou,  
Aristotle University of Thessaloniki, Greece

## \*CORRESPONDENCE

Antonio Ferrante

✉ antonio.ferrante@unimi.it

RECEIVED 10 March 2023

ACCEPTED 05 June 2023

PUBLISHED 19 June 2023

## CITATION

Cavallaro V, Bulgari R, Florio FE,  
Restuccia P, Vinci G, Guffanti D, Vignati S  
and Ferrante A (2023) Postharvest  
strategies for preventing flower wilting and  
leaf yellowing in cut *Ranunculus* flowers.  
*Front. Hortic.* 2:1183754.  
doi: 10.3389/fhort.2023.1183754

## COPYRIGHT

© 2023 Cavallaro, Bulgari, Florio, Restuccia,  
Vinci, Guffanti, Vignati and Ferrante. This is  
an open-access article distributed under the  
terms of the [Creative Commons Attribution  
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or  
reproduction in other forums is permitted,  
provided the original author(s) and the  
copyright owner(s) are credited and that  
the original publication in this journal is  
cited, in accordance with accepted  
academic practice. No use, distribution or  
reproduction is permitted which does not  
comply with these terms.

# Postharvest strategies for preventing flower wilting and leaf yellowing in cut *Ranunculus* flowers

Viviana Cavallaro<sup>1</sup>, Roberta Bulgari<sup>2</sup>, Francesco Elia Florio<sup>1</sup>,  
Pasquale Restuccia<sup>3</sup>, Gianluca Vinci<sup>3</sup>, Davide Guffanti<sup>1</sup>,  
Sara Vignati<sup>1</sup> and Antonio Ferrante<sup>1\*</sup>

<sup>1</sup>Department of Agricultural and Environmental Sciences, Università degli Studi di Milano, Milano, Italy,

<sup>2</sup>Department of Agricultural, Forest and Food Sciences, DISAFA, Vegetable Crops and Medicinal and Aromatic Plants, VEGMAP, University of Torino, Grugliasco, Italy, <sup>3</sup>Florcoop Sanremo Società Cooperativa Agricola (S.C.A.), Taggia, Italy

**Introduction:** Appropriate postharvest treatment, as well as adequate conditions of storage, can be adopted to elongate the lifespan of cut flowers. Thidiazuron (TDZ), a substituted phenylurea, and 1-methylcyclopropane (1-MCP), a non-toxic inhibitor of ethylene perception, are nowadays substances commonly used to prevent early damage caused by senescence and to delay chlorophyll degradation. *Ranunculus asiaticus* L. is cultivated for cut flower production and is highly sensitive to ethylene and leaf yellowing. In this study, the effect of different pulse-treatment in prolonging cut ranunculus vase life was analyzed.

**Methods:** TDZ 10  $\mu$ M, 1-MCP 500 ppb, and a combination of both were applied for 24 hours after harvest. The effect of the treatments was evaluated by performing non-destructive (% loss of fresh weight, chlorophyll a fluorescence, *in vivo* chlorophyll content, and Nitrogen Flavonol Index – NFI) and destructive (chlorophyll, carotenoids, anthocyanins concentration, and phenolic index) analyses at 0, 1, 12, and 14 days from treatments.

**Results and Discussion:** Flower wilting was delayed by 4 days in 1-MCP + TDZ 10  $\mu$ M treatments, which also reduced weight loss and chlorophyll degradation compared to controls. The effectiveness of these compounds in preventing senescence has been confirmed by the decreased biosynthesis of phenolic compounds.

## KEYWORDS

senescence, ethylene, *Ranunculus asiaticus* L., 1-methylcyclopropane, thidiazuron, vase life

## 1 Introduction

The global floricultural industry is nowadays a multi-billion-dollar enterprise in constant growth and with international dimensions (Naing et al., 2022). Consumers are becoming more interested in flower quality and longevity (Darras, 2021); however, during long trade transportation, qualitative characteristics can be compromised by eventual ethylene exposure, causing early petal and bud abscission. Therefore, appropriate control of storage conditions (temperature, humidity, and atmospheric conditions) (Fernandes et al., 2020) and postharvest treatments, which include different chemical or organic substances, can be adopted (van Doorn et al., 2011; da Costa et al., 2015; da Costa et al., 2021; Sun et al., 2022) to meet consumers' preference for prolonged flower longevity and freshness (Naing et al., 2022). The inhibition of ethylene perception in flowers can represent a powerful instrument to prolong their longevity. Silver thiosulfate (STS) was one of the first chemical compounds utilized for this purpose. However, the release of silver ions also represents a serious concern to the environment and public health because of their toxicity (van Doorn et al., 2011; Naing et al., 2022). 1-methylcyclopropene (1-MCP – chemical formula  $C_4H_6$ ) was discovered in 1995 and it is a non-toxic gaseous compound which inhibits ethylene action by binding itself to hormone receptors in an irreversible manner: the ability of 1-MCP to bind receptors is ten times greater compared to ethylene (Blankenship and Dole, 2003; Ferrante and Francini, 2006; Naing et al., 2022). It has been reported that the vase life of various cut flowers such as carnation (*Dianthus carioophyllus* L.) (Ichimura and Shimizu, 2002), stock (*Matthiola incana*) (Çelikel and Reid, 2002), *Consolida ajacis* (Santos et al., 2005), chrysanthemum (*Chrisantemum morifolium* Ram.) (Hassan and Gerzson, 2002), snapdragon (*Antirrhinum majus* L.) (Heffron and Korban, 2022), and *Delphinium* hybrid (cv. Bellamosum) (Ichimura and Shimizu, 2002) can be extended by exposure to 1-MCP (Rani and Singh, 2014). Moreover, treatments with this compound elongate the vase life of sweet pea (Ichimura and Shimizu, 2002) and rose (Chamoni et al., 2005) and prevent bud and floral abscission of *Phalaenopsis* cultivars (Sun et al., 2009). Vase life can be also defined by leaf senescence. It has been demonstrated that treatments with cytokinins and gibberellins inhibit chlorophyll degradation and leaf senescence (Janowska and Andrzejak, 2022). Thidiazuron (TDZ, N-phenyl-N'-1,2,3-thiadiazol-5-ylurea) is a herbicide and defoliant compound of substituted phenylurea with high cytokinin-like activity (Macnish et al., 2010; Ferrante et al., 2012). Different studies on ruscus, *Alstroemeria*, and other cut flowers sensitive to leaf yellowing demonstrated the effectiveness of TDZ treatment in preventing leaf senescence (Ferrante et al., 2002b; Ferrante et al., 2003; Mutui et al., 2005; Bulgari et al., 2015).

*Ranunculus asiaticus* L. is a geophyte with a vegetative phase during cool winters and a quiescence phase in hot summer periods. It is the only *Ranunculus* species cultivated for ornamental scope, especially in the Mediterranean area, where Italy is the largest

producer (300-350 ha, with almost half of them, 132 million stems, produced in Sanremo, Italy). *Ranunculus* represents 0.4% of the total sales of cut flowers and data from Royal FloraHolland, collected during the period January 2015 - May 2017, shows that *Ranunculus* stems come primarily from the Netherlands, Israel, and Italy (Beruto et al., 2018). Flowers belonging to the family of *Ranunculaceae* appear to be highly ethylene-sensitive (Evans et al., 2002; Mensuali Sodi et al., 2002; Scariot et al., 2009) and, considering the importance of ranunculus for the Italian flower market, it appears of extreme importance to find new treatments which can elongate the vase life of cut stems, retarding the appearance of senescence symptoms. In this work, cut ranunculus flowers were treated with TDZ, 1-MCP, and with a combination of both to evaluate their effectiveness in prolongating the flowers' vase life.

## 2 Material and methods

### 2.1 Plant material

Cut ranunculus flowers (*Ranunculus asiaticus* L., “Venere rosa”) were provided by Floorcoop Sanremo (Taggia, IM, Italy) and then they were wet transported to the University of Milan and treated in the 24 hours (h) after harvest as described below.

### 2.2 Treatments and evaluation of stem quality

Forty stems, chosen for uniform growth and stage of development—the “open flower” stage according to Shahri and Tahir (2011)—were divided into four groups and pulse-treated for 24 h in a controlled environment (temperature 20° C, relative humidity 60-70%, and light intensity 10-15  $\mu\text{mol m}^{-2} \text{s}^{-1}$  PPFD for 12 h per day using white fluorescent tube lamps) as follow: distilled water (control), TDZ 10  $\mu\text{M}$ , 1-MCP 500 ppb  $\text{m}^{-3}$ , and 1-MCP + TDZ 10  $\mu\text{M}$ . TDZ was dissolved in 0.1 mM KOH and then pH adjusted with 36% HCl. Flowers of each condition were separated into three different glass bottles containing 750 mL of solution (distilled water or TDZ 10  $\mu\text{M}$ ). 1-MCP application was performed by fumigation in an airtight container, dissolving 1.6 g of powder in 250 mL of distilled water. Concentrations were chosen based on previous experiments performed by Ferrante et al. (2012). At the end of this period, each stem was transferred to a 50 mL tube, rinsed daily with distilled water during the entire experiment, and flowers were kept in the same controlled environment explained previously.

The quality of each flower was assessed by visually determining the presence or absence of senescence symptoms every day. The time duration between the start of the experiment and the day of the appearance of leaf yellowing, flower senescence, and the first petal loss was recorded for every replicate of each treatment (n = 10).

## 2.3 Non-destructive analyses

### 2.3.1 Loss of fresh weight

The fresh weight (FW) of the stems ( $n = 10$ ) for each condition was measured after 1, 7, 12, and 14 days from the treatments, and the percentage of FW loss was estimated as follows:

$$\frac{FWt1 - FWtx}{FWt1} \cdot 100$$

where “t1” refers to the FW measured after 24 h from treatment, and “tx” to all the other time points.

### 2.3.2 *In vivo* chlorophyll content, Nitrogen Flavonol Index, and chlorophyll *a* fluorescence

*In vivo* content of chlorophyll and Nitrogen-Flavonol Index (NFI)—a good indicator of the nutritional status of plants—were measured using a multi-pigment meter MPM-100 (ADC BioScientific Ltd.), randomly choosing ten leaves for each condition ( $n = 10$ ). Chlorophyll *a* fluorescence was measured using a hand-portable fluorimeter (Handy-PEA, Hansatech Instruments), randomly choosing three leaves for each condition ( $n = 3$ ). Before measurements, the leaves were dark-adapted with leaf clips (diameter 4 mm) for 30–40 min, then they were exposed to a saturating light ( $3000 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) provided by an array of three high-intensity light-emitting diodes for 1 s. Information about the structural and functional status of the photosynthetic apparatus was provided by the parameters measured: the maximum quantum efficiency of photosystem II ( $F_v/F_m$ ) and the area (value proportional to the number of electrons transferred by the reaction centers to the quinones -  $Q_A$  - during photosynthesis).

Both analyses were performed at 0, 1, 7, 12, and 14 days after treatments.

## 2.4 Destructive analyses

In order to perform the destructive analyses, three leaves were randomly selected for each treatment from among the ten replicates in all the time points considered to obtain plant materials. The selected stems were weighed at the beginning and after leaf detachment to obtain accurate FW and loss of FW measurements.

### 2.4.1 Chlorophyll and carotenoids concentration

Chlorophylls and carotenoids were extracted from leaf tissues using 5 mL of 99.9% (v/v) methanol. Three leaf disc samples (5 mm diameter, 30 mg FW), for each treatment and time were kept in a dark room for 24 h at 4°C. After that, absorbance readings were measured with a spectrophotometer (Thermo Italy) at 665.2 and 652.4 nm for chlorophylls, and 470 nm for total carotenoids, and pigments levels were calculated by Lichtenthaler's formula. The results were expressed as  $\mu\text{g}$  of

pigments  $\text{g}^{-1}$  FW (Lichtenthaler, 1987). Samplings were conducted at 0, 1, 7, 12, and 14 days after treatment.

### 2.4.2 Phenolic index and anthocyanins concentration

Phenolic index and total anthocyanin were determined from leaf disc samples (5 mm diameter, 30 mg FW). Three leaf samples for each treatment and time were transferred to a tube containing 3 mL of methanol acidified with hydrochloric acid (1% v/v) and were kept in a dark room for 24 h at 4°C. Absorbance readings were measured with a spectrophotometer at 320 nm for total phenols (Ke and Saltveit, 1989), and at 535 nm for anthocyanin determination (Klein and Hagen, 1961). Phenolic index was expressed as  $\text{ABS}_{320} \text{ nm}^{-1} \text{ g}^{-1} \text{ FW}$ . Anthocyanins concentration was expressed in mg cyanidin-3-glucoside equivalents  $100 \text{ g}^{-1} \text{ FW}$  using a molar extinction coefficient ( $\epsilon$ ) of  $29,600 \text{ L M}^{-1} \text{ cm}^{-1}$ . Samplings were conducted at 0, 1, 7, 12, and 14 days after treatment.

## 2.5 Statistical analysis


Data are reported as the means of replicates. Statistics were performed using GraphPad Prism version 8 for Windows (GraphPad Software, La Jolla California USA, [www.graphpad.com](http://www.graphpad.com)). A two-way ANOVA was performed, followed by the Tukey *post-hoc* test ( $p < 0.05$ ), considering the variables of time, treatment, and their interaction. Additional information is reported in the figure legends.

## 3 Results

### 3.1 Loss of FW and quality of the stems

All the treatments showed an increase in FW losses as vase life went on. Only the flowers treated with TDZ  $10 \mu\text{M}$  had no significant differences at 12 and 14 days after treatment. Among the different treatments, it was possible to observe how the treated cut flowers showed, in general, lower FW losses compared with the control group after 7 and 12 days (Figure 1). In particular, flowers treated with TDZ  $10 \mu\text{M}$  and 1-MCP + TDZ  $10 \mu\text{M}$  showed the best results. The interaction between time and treatment was not significant.

Cut flower quality was assessed, evaluating the time duration between the beginning of vase life and the appearance of the first symptoms of senescence, represented by leaf yellowing, flower wilting, and petal fall (Figures 2A, B). Considering leaf yellowing, no significant differences were observed between the control and 1-MCP treatment groups, in which leaf yellowing occurred after 5–6 days. However, both the treatments containing TDZ delayed this phenomenon by 3 days (Figure 2B). All treatments delayed flower senescence by 3 days (Figure 2A), while loss of petals was delayed by 3 days in treatment with only TDZ and 4 days in the other two treatments, compared to the control group (Table 1).



**FIGURE 1** Percentage of fresh weight losses in the different treatments at 7, 12, and 14 days. The interaction of “time x treatment” was not significant. Various lowercase letters indicate significant differences among treatments at the same time after two-way ANOVA ( $p < 0.05$ ) ( $n = 10$ ); various uppercase letters indicate significant differences between the time points for each treatment after two-way ANOVA ( $p < 0.05$ ) ( $n = 10$ ).



**FIGURE 2** Quality of the stems (A) and leaf yellowing (B) in the different treatments after 7 days from the application of TDZ 10 μM, 1-MCP, and 1-MCP + TDZ 10 μM.

**TABLE 1** Days (expressed as mean value) before the appearance of the main symptoms considered in the evaluation of stem quality in the different treatments (leaf yellowing, flower wilting, and loss of petal).

	Leaf Yellowing			
	Control	TDZ 10μM	1-MCP	TDZ 10μM + 1-MCP
	5.4 <sup>b</sup>	8.4 <sup>a</sup>	5.9 <sup>b</sup>	8.4 <sup>a</sup>
Time (days)	Flower wilting			
	Control	TDZ 10μM	1-MCP	TDZ 10μM + 1-MCP
	5.3 <sup>c</sup>	7.8 <sup>b</sup>	8.2 <sup>ab</sup>	8.6 <sup>a</sup>
Time (days)	Loss of petal			
	Control	TDZ 10μM	1-MCP	TDZ 10μM + 1-MCP
	6.8 <sup>c</sup>	10.2 <sup>b</sup>	11 <sup>ab</sup>	11.6 <sup>a</sup>

Various letters indicate significant differences among treatments after one-way ANOVA ( $p < 0.05$ ) ( $n = 10$ ).

### 3.2 *In vivo* chlorophyll content, NFI and chlorophyll a fluorescence

In the cut flowers sector, non-destructive measurements allow the early detection of stress or senescence. In the present

experiment, non-destructive analyses always showed a significant interaction of “time x treatment” ( $p < 0.05$ ), with the exception for NFI values.

*In vivo* chlorophyll concentrations were higher in TDZ and 1-MCP + TDZ 10 μM at day 7 compared with the controls. After this

time point, the values tended to decrease in all the treatments, less evidently in TDZ 10  $\mu$ M stems. At 14 days, 1-MCP showed even lower levels of chlorophyll, compared to controls (Figure 3A). In treatment with TDZ 10  $\mu$ M, higher values of NFI at all time points confirmed a better status of the leaves compared to the control and 1-MCP groups (Figure 3B). While NFI values decreased over time in 1-MCP + TDZ 10  $\mu$ M treatments, at 12 and 14 days the leaves showed intermediate values of between the TDZ 10  $\mu$ M and control groups.

Chlorophyll a fluorescence was measured at different time points to check the health status of the cut flowers. Among the different parameters, Fv/Fm and area were considered for evaluating the effects of the treatments. The maximum quantum efficiency of photosystem II (Fv/Fm), which has an optimal value of around 0.83 in unstressed plants (Maxwell and Johnson, 2000), was not different from controls at 0 and 1 days. Cut flowers treated with TDZ or 1-MCP + TDZ 10  $\mu$ M showed a retention of the PSII functionality until day 12 (Figure 4A). In 1-MCP, the lowest results were recorded at 12 and 14 days after treatment. Area (Figure 4B) tends to decrease as environmental stresses or plant senescence occur (Kalaji and Guo, 2008). As the process of senescence went on, cut flowers of both

treatments which included TDZ 10  $\mu$ M showed higher values compared to the control and 1-MCP groups.

### 3.3 Destructive analyses on leaves of cut flowers

Destructive analyses showed a significant interaction of “time x treatment” ( $p < 0.05$ ).

The total chlorophyll concentration in the flowers at harvest was 1.5-1.7  $\mu$ g/mg FW. Control flowers showed lower concentrations of total chlorophyll starting from 7 days after treatment. According to the different leaf yellowing observed, 1-MCP demonstrated a similar behavior to the controls, while 1-MCP + TDZ 10  $\mu$ M and, particularly, TDZ 10  $\mu$ M showed higher concentrations of chlorophyll with values of 0.9  $\mu$ g/mg FW at the end of the experiment (Figure 5A). A similar trend was observed for leaf carotenoids concentration (Figure 5B). After 12 and 14 days, the carotenoids increased in the TDZ and 1-MCP + TDZ treatments, with values of 0.20-0.25  $\mu$ g/mg FW and 0.14-0.16  $\mu$ g/mg FW, respectively, that were higher than the controls.




FIGURE 3 In vivo chlorophyll content (A) and NFI (B) in the different treatments after 0, 1, 7, 12, and 14 days from treatment. The interaction of “time x treatment” was significant only for chlorophyll content (A); various letters indicate significant differences after two-way ANOVA ( $p < 0.05$ ) ( $n = 10$ ). In the NFI graph (B), various lowercase letters indicate significant differences among treatments at the same time after two-way ANOVA ( $p < 0.05$ ) ( $n = 10$ ); various uppercase letters indicate significant differences between the time points for each treatment after two-way ANOVA ( $p < 0.05$ ) ( $n = 10$ ).





FIGURE 4 Fv/Fm ratio (A) and Area under the chlorophyll a fluorescence induction curve (B) parameters in different treatments after 0, 1, 7, 12, and 14 days from the treatment. The red line (A) indicates the optimal threshold of 0.83. The interaction of “time x treatment” was always significant. Various letters indicate significant differences after two-way ANOVA ( $p < 0.05$ ) ( $n = 3$ ).



Anthocyanins concentration did not show significant differences among the treatments except after 12 days. At this time point, the TDZ showed a higher value of anthocyanins (Figure 5D). The concentration of phenolic compounds increased during the experimental period in control flowers (Figure 5C). Treatment with TDZ did not show an increase in phenolic compounds. In the other treatments the phenolic index increased in time, but less evidently compared to control up to 12 days.

## 4 Discussion

The quality of cut flowers mainly depends on their visual appearance and the longevity of petals and leaves. The cut flower industry is persistently looking for new strategies to enhance postharvest handling practices. Delaying the process of flower and leaf senescence appears to be of pivotal importance, because the vase life is a critical factor in determining commercial and consumer appreciation (Horibe, 2020). The optimization of storage conditions, along with postharvest treatments, are crucial for extending the storage and the vase life of cut flowers. Among the possible strategies, the application of different chemical and organic compounds can help in enhancing the vase life of different species (da Costa et al., 2021).

1-MCP is a non-toxic ethylene action inhibitor that protects cut flowers from ethylene produced by flowers or present in the environment as a pollutant. It has been largely used to delay senescence processes in fruit, leafy vegetables, and cut flowers,

and more than 40 countries have currently approved its use (Grozeff et al., 2010; Dias et al., 2021).

TDZ is a substitute of phenyl urea with high cytokinin-like activity, and its action can be observed in both light and dark conditions. The biosynthesis of cytokinins occurs in roots; therefore, the harvest and detachment of cut flowers from their roots cause the loss of the supply of these plant hormones (Kakimoto, 2003; Sakakibara, 2021; Janowska and Andrzejak, 2022); the reduction of endogenous cytokinins associated with dark conditions postharvest accelerate leaf senescence induction (Eason, 2006; Hönig et al., 2018; Guo et al., 2021). Moreover, TDZ application has shown capability in increasing the level of carotenoids, which can be considered the molecules used for counteracting the senescence process of cut flowers, protecting the flower from excessive damage caused by reactive oxygen species (Ferrante et al., 2002a; Ferrante et al., 2003; Sankhala et al., 2003; Sankhala et al., 2005; Macnish et al., 2010; Hönig et al., 2018). The effectiveness of these compounds has been largely studied, yet little is known about the possible positive effect of the combination of TDZ and 1-MCP.

In this work, pulse applications of TDZ alone or in combination with 1-MCP were studied to preserve the quality and prolong the vase life of cut ranunculus flowers. Our results suggested that all tested treatments strongly inhibited petal abscission and flower senescence, delaying flower wilting by 3 days in TDZ 10  $\mu$ M and 4 days in both 1-MCP and 1-MCP + TDZ 10  $\mu$ M treatments, compared to controls. The effectiveness of these compounds compared to controls in preventing senescence has been confirmed by the decreased biosynthesis of phenolic compounds

in treated flowers. These secondary metabolites are involved in a wide range of biological functions. They are highly accumulated during vase life to counteract the stress generated by flower or leaf senescence (Cavaiuolo et al., 2013). On the other hand, the 1-MCP treatment did not have an effect in preventing leaf yellowing, which occurred after 5 days in both treatments and in controls. The TDZ 10  $\mu\text{M}$  treatment seems to preserve the health status of flowers, as evidenced by the analysis of the Fv/Fm index and by a greater content of chlorophylls, as both *in vivo* and destructive analyses showed. These results suggest a better performance of the photosynthetic apparatus in the TDZ 10  $\mu\text{M}$  thesis, followed by 1-MCP + TDZ 10  $\mu\text{M}$ . A high concentration of carotenoids, with higher levels in TDZ 10  $\mu\text{M}$  treatment followed by 1-MCP + TDZ 10  $\mu\text{M}$  treatment, reveals the important contribution of these pigments in preserving the quality of flowers, acting as photoprotectors and antioxidants able to reduce the oxidation of other molecules (Maoka, 2020; Pérez-Gálvez et al., 2020). The cut flowers treated with TDZ 10  $\mu\text{M}$  and 1-MCP + TDZ 10  $\mu\text{M}$  showed the best results also in terms of less weight loss over the entire period of observation. TDZ is effective at very low concentrations; therefore, it can be a valid postharvest treatment for preserving flowers sensitive to leaf yellowing.

## 5 Conclusion

In contrast to a previous study on stock flowers (Ferrante et al., 2012), our findings showed how the combination of 1-MCP + TDZ 10  $\mu\text{M}$  could represent an efficient postharvest treatment to extend the cut ranunculus vase life, delaying both flower and leaf senescence. In further works, it could be interesting to perform deeper analyses on the senescence process, taking into account the color of flower, the level of total sugars, the ethylene biosynthesis, the leaf gas exchanges, and the activities of senescence associated enzymes. Expanding our knowledge on the interaction between 1-MCP and TDZ can provide further insights into the mechanisms that contribute to the combined beneficial effect observed. These valuable insights can then be utilized as practical tools for implementing postharvest guidelines, enabling us to effectively maintain and enhance the quality of cut flowers throughout the entire supply chain. In fact, delaying the appearance of senescence phenomena, even for few days, could be a significant result for the cut flowers market.

## References

- Beruto, M., Rabaglio, M., Viglione, S., Van Labeke, M.-C., and Dhooge, E. (2018). "Ranunculus," in *Ornamental crops*. Ed. J. V. Huylbroeck (Belle, Belgium: Springer), 649–672.
- Blankenship, S. M., and Dole, J. M. (2003). 1-methylcyclopropene: a review. *Postharvest Biol. Technol.* 28, 1–25. doi: 10.1016/S0925-5214(02)00246-6
- Bulgari, R., Negri, M., and Ferrante, A. (2015). Evaluation of postharvest storage and treatments in cut ruscus foliage. *Adv. Hortic. Sci.* 29, 103–108. doi: 10.13128/ahs-22687
- Cavaiuolo, M., Cocetta, G., and Ferrante, A. (2013). The antioxidants changes in ornamental flowers during development and senescence. *Antioxidants* 2, 132–155. doi: 10.3390/antiox2030132
- Çelikel, F. G., and Reid, M. S. (2002). Postharvest handling of stock (*Matthiola incana*). *HortScience* 37, 144–147. doi: 10.21273/HORTSCI.37.1.144
- Chamoni, E., Kalighi, A., Joyce, D. C., Irving, D. E., Zamani, Z. A., Mostofi, Y., et al. (2005). Ethylene and anti-ethylene treatment effects on cut "First-red" rose. *J. Appl. Horticulture* 7, 3–7. doi: 10.37855/jah.2005.v07i01.01

## Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## Author contributions

VC: Analytical determinations and writing. RB: Training methods. FF: experimental set up. DG: Experimental set up and analysis. SV: non-destructive analysis. AF Supervision, Editing manuscript. All authors contributed to the article and approved the submitted version.

## Funding

Research activities were funded by Psr 2014-2020 Liguria Region action 16.1 "Ottimizzazione della programmazione della fioritura e della conservazione postraccolta di alcune specie da fiore reciso di interesse per la Riviera Ligure di Ponente" – Ottiprogram.

## Conflict of interest

The authors AF, RB, FF declared that they were an editorial board member of Frontiers, at the time of submission. This had no impact on the peer review process and the final decision.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- da Costa, L. C., de Araujo, F. F., Ribeiro, W. S., de Sousa Santos, M. N., and Finger, F. L. (2021). Postharvest physiology of cut flowers. *Ornamental Horticulture* 27, 374–385. doi: 10.1590/2447-536x.v27i3.2372
- da Costa, L. C., Ribeiro, R. R., Carneiro, W. S., Barbosa, J. A., and Finger, F. L. (2015). Postharvest longevity of *Heliconia wagneriana*. *Acta Horticulturae* 1060, 193–199. doi: 10.17660/ActaHortic.2015.1060.28
- Darras, A. (2021). Overview of the dynamic role of specialty cut flowers in the international cut flower market. *Horticulturae* 7, 51–60. doi: 10.3390/horticulturae7030051
- Dias, C., Ribeiro, T., Rodrigues, A. C., Ferrante, A., Vasconcelos, M. W., and Pintado, M. (2021). Improving the ripening process after 1-MCP application: implications and strategies. *Trends Food Sci. Technol.* 113, 382–396. doi: 10.1016/j.tifs.2021.05.012
- Eason, J. R. (2006). Molecular and genetic aspects of flower senescence. *Stewart Postharvest Rev.* 2, 1–7. doi: 10.2212/spr.2006.2.6
- Evans, A. C., Burge, G. K., Littlejohn, R. P., Douglas, M. H., Bicknell, R. A., and Lill, R. E. (2002). Mount cook lily (*Ranunculus lyallii*) - a potential cut flower? *N Z J. Crop Hortic. Sci.* 30, 69–78. doi: 10.1080/01140671.2002.9514200
- Fernandes, L., Casal, S., Pereira, J. A., Saraiva, J. A., and Ramalhosa, E. (2020). An overview on the market of edible flowers. *Food Rev. Int.* 36, 258–275. doi: 10.1080/87559129.2019.1639727
- Ferrante, A., and Francini, A. (2006). "Ethylene and leaf senescence," in *Ethylene action in plants*. Ed. N. A. Khan (Berlin, Germany: Springer), 51–67.
- Ferrante, A., Hunter, D. A., Hackett, W. P., and Reid, M. S. (2002a). Thidiazuron – a potent inhibitor of leaf senescence in alstroemeria. *Postharvest Biol. Technol.* 25, 333–338. doi: 10.1016/S0925-5214(01)00195-8
- Ferrante, A., Serra, G., and Tognoni, F. (2002b). Effects of ethylene and cytokinins on vase life of cut *Eucalyptus parvifolia* cambage branches. *Plant Growth Regul.* 38, 119–125. doi: 10.1023/A:1021212601768
- Ferrante, A., Tognoni, F., Mensuali-Sodi, A., and Serra, G. (2003). Treatment with thidiazuron for preventing leaf yellowing in cut tulips and chrysanthemum. *Acta Horticulturae* 624, 357–363. doi: 10.17660/ActaHortic.2003.624.49
- Ferrante, A., Trivellini, A., and Mensuali Sodi, A. (2012). Interaction of 1-methylcyclopropene and thidiazuron on cut stock flowers vase life. *Open Horticulture J.* 5, 1–5. doi: 10.2174/1874840601205010001
- Grozeff, G. G., Micieli, M. E., Gómez, F., Fernández, L., Guaiamet, J. J., Chaves, A. R., et al. (2010). 1-methylcyclopropene extends postharvest life of spinach leaves. *Postharvest Biol. Technol.* 55, 182–185. doi: 10.1016/j.postharvbio.2009.10.004
- Guo, Y., Ren, G., Zhang, K., Li, Z., Miao, Y., and Guo, H. (2021). Leaf senescence: progression, regulation, and application. *Mol. Horticulture* 1, 5. doi: 10.1186/s43897-021-00006-9
- Hassan, F. A. S., and Gerzson, L. (2002). Effect of 1-MCP (1-methylcyclopropene) on the vase life of *Chrysanthemum* and carnation cut flowers. *Int. J. Hortic. Sci.* 8, 29–32. doi: 10.31421/IJHS/8/3-4/357
- Heffron, L. M., and Korban, S. S. (2022). Evaluation of ethylene mutant snapdragon lines for rooting, gravitropism, 1-MCP, ethylene, and vase-life responses. *Sci. Hortic.* 304, 111274. doi: 10.1016/j.scienta.2022.111274
- Höning, M., Plihalová, L., Husičková, A., Nisler, J., and Doležal, K. (2018). Role of cytokinins in senescence, antioxidant defence and photosynthesis. *Int. J. Mol. Sci.* 19, 4045. doi: 10.3390/ijms19124045
- Horibe, T. (2020). Use of light stimuli as a postharvest technology for cut flowers. *Front. Plant Sci.* 11, 10–13. doi: 10.3389/fpls.2020.573490
- Ichimura, K., and Shimizu, H. (2002). Effect of 1-methylcyclopropene (1-MCP) on the vase life of cut carnation, *Delphinium* and sweet pea flowers. *Bull. Natl. Inst. Flor. Sci.* 2, 1–8.
- Janowska, B., and Andrzejak, R. (2022). The role of cytokinins and gibberellins on post-harvest longevity of florists' greens. *Agriculture* 12, 1–14. doi: 10.3390/agriculture12091375
- Kakimoto, T. (2003). Biosynthesis of cytokinins. *J. Plant Res.* 116, 233–239. doi: 10.1007/s10265-003-0095-5
- Kalaji, H., and Guo, P. (2008). "Chlorophyll fluorescence: a useful tool in barley plant breeding programs," in *Photochemistry research progress*. Eds. A. Sanchez and S. J. Gutierrez (Portland, OR, USA: Nova Science), 447–471.
- Ke, D., and Saltveit, M. E. (1989). Wound-induced ethylene production, phenolic metabolism and susceptibility to russet spotting in iceberg lettuce. *Physiol. Plant* 76, 412–418. doi: 10.1111/j.1399-3054.1989.tb06212.x
- Klein, A. O., and Hagen, C. W. (1961). Anthocyanin production in detached petals of *Impatiens balsamina* L. *Plant Physiol.* 36, 1–9. doi: 10.1104/pp.36.1.1
- Lichtenthaler, H. K. (1987). Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods Enzymol.* 148, 350–382. doi: 10.1016/0076-6879(87)48036-1
- Macnish, A. J., Jiang, C. Z., and Reid, M. S. (2010). Treatment with thidiazuron improves opening and vase life of iris flowers. *Postharvest Biol. Technol.* 56, 77–84. doi: 10.1016/j.postharvbio.2009.11.011
- Maoka, T. (2020). Carotenoids as natural functional pigments. *J. Natural Medicines* 74 (1), 1–16. doi: 10.1007/s11418-019-01364-x
- Maxwell, K., and Johnson, G. N. (2000). Chlorophyll fluorescence—a practical guide. *J. Exp. Bot.* 51, 659–668. doi: 10.1093/jexbot/51.345.659
- Mensuali Sodi, A., Ferrante, A., Serra, G., and Rognoni, F. (2002). Ethylene involvement in *Ranunculus asiaticus* L. longevity. *Proc. Vth Giornate Scientifiche Spoleto* 2, 419–420.
- Mutui, T. M., Mibus, H., and Serek, M. (2005). Effects of thidiazuron, ethylene, abscisic acid and dark storage on leaf yellowing and rooting of *Pelargonium* cuttings. *J. Hortic. Sci. Biotechnol.* 80, 543–550. doi: 10.1080/14620316.2005.11511975
- Naing, A. H., Win, N. M., Kyu, S. Y., Kang, I.-K., and Kim, C. K. (2022). Current progress in application of 1-methylcyclopropene to improve postharvest quality of cut flowers. *Hortic. Plant J.* 8, 1–13. doi: 10.1016/j.hpj.2021.11.014
- Pérez-Gálvez, A., Viera, I., and Roca, M. (2020). Carotenoids and chlorophylls as antioxidants. *Antioxidants* 9 (6), 505. doi: 10.3390/antiox9060505
- Rani, P., and Singh, N. (2014). Senescence and postharvest studies of cut flowers: a critical review. *Pertanika J. Trop. Agric. Sci.* 37, 159–201.
- Sakakibara, H. (2021). Cytokinin biosynthesis and transport for systemic nitrogen signaling. *Plant J.* 105, 421–430. doi: 10.1111/tpj.15011
- Sankhala, N., Mackay, W. A., and Davis, T. D. (2003). Reduction of flower abscission and leaf senescence in cut phlox inflorescence by thidiazuron. *Acta Horticulturae* 628, 837–841. doi: 10.17660/ActaHortic.2003.628.106
- Sankhala, N., Mackay, W. A., and Davis, T. D. (2005). Effect of thidiazuron on senescence of flowers in cut inflorescences of *Lupinus densiflorus* benth. *Acta Horticulturae* 669, 239–243. doi: 10.17660/ActaHortic.2005.669.31
- Santos, V. R., Finger, F. L., Barbosa, J. G., and Barros, R. S. (2005). Influence of ethylene and 1-MCP on senescence and longevity of *Consolida ajacis* inflorescences. *Bragantia* 64, 33–38. doi: 10.1590/S0006-87052005000100004
- Scariot, V., Larcher, F., Caser, M., Costa, E., Beruto, M., and Devecchi, M. (2009). Flower longevity in ten cultivars of cut *Ranunculus asiaticus* L. as affected by ethylene and ethylene inhibitors. *Eur. J. Hortic. Sci.* 74, 137–142.
- Shahri, W., and Tahir, I. (2011). Flower development and senescence in *Ranunculus asiaticus* L. *J. Fruit Ornamental Plant Res.* 19, 123–131.
- Sun, Y., Christensen, B., Liu, F., Wang, H., and Müller, R. (2009). Effects of ethylene and 1-MCP (1-methylcyclopropene) on bud and flower drop in mini-*Phalaenopsis* cultivars. *Plant Growth Regul.* 59, 83–91. doi: 10.1007/s10725-009-9391-y
- Sun, J., Guo, H., and Tao, J. (2022). Effects of harvest stage, storage, and preservation technology on postharvest ornamental value of cut peony (*Paeonia lactiflora*) flowers. *Agronomy* 12 (2), 230. doi: 10.3390/agronomy12020230
- van Doorn, W. G., Perik, R. R. J., Abadie, P., and Harkema, H. (2011). A treatment to improve the vase life of cut tulips: effects on tepal senescence, tepal abscission, leaf yellowing and stem elongation. *Postharvest Biol. Technol.* 61, 56–63. doi: 10.1016/j.postharvbio.2011.02.003