

Corrigendum: A comparison of two sleep spindle detection methods based on all night averages: individually adjusted vs. fixed frequencies

Péter P. Ujma¹, Ferenc Gombos², Lisa Genzel³, Boris N. Konrad⁴, Péter Simor^{5,6}, Axel Steiger⁴, Martin Dresler^{4,7*} and Róbert Bódizs^{1,2}

¹ Institute of Behavioral Science, Semmelweis University, Budapest, Hungary, ² Department of General Psychology, Pázmány Péter Catholic University, Budapest, Hungary, ³ Centre for Cognitive and Neural Systems, University of Edinburgh, Edinburgh, UK, ⁴ Max Planck Institute of Psychiatry, Munich, Germany, ⁵ Department of Cognitive Sciences, Budapest University of Technology and Economics, Budapest, Hungary, ⁶ Nyírő Gyula Hospital, National Institute of Psychiatry and Addictions, Budapest, Hungary, ⁷ Donders Institute, Radboud University Medical Centre, Nijmegen, Netherlands

Keywords: sleep spindles, EEG, individual adjustment method, IAM, algorithm

A Corrigendum on

A comparison of two sleep spindle detection methods based on all night averages: individually adjusted vs. fixed frequencies

by Ujma, P. P., Gombos, F., Genzel, L., Konrad, B. N., Simor, P., Steiger, A., et al. (2015). Front. Hum. Neurosci. 9:52. doi: 10.3389/fnhum.2015.00052

The description of the Individual Adjustment Method (IAM) algorithm for sleep spindle analyses (Ujma et al., 2015) contained an error, which we hereby rectify. On page 5, line 7–8, instead of $f(x) = e^{(-(x - xm)/(w/2))}$, the equation should read as follows:

 $f(x)=e^{\wedge}-(((x-xm)/(w/2))^{\wedge}2)$

References

Ujma, P. P., Gombos, F., Genzel, L., Konrad, B. N., Simor, P., Steiger, A., et al. (2015). A comparison of two sleep spindle detection methods based on all night averages: individually adjusted vs. fixed frequencies. *Front. Hum. Neurosci.* 9:52. doi: 10.3389/fnhum.2015.00052

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Ujma, Gombos, Genzel, Konrad, Simor, Steiger, Dresler and Bódizs. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

OPEN ACCESS

Edited and reviewed by: Simon C. Warby, Stanford University, USA

*Correspondence: Martin Dresler, dresler@mpipsykl.mpg.de

Received: 01 June 2015 **Accepted:** 06 July 2015 **Published:** 21 July 2015

Citation:

Ujma PP, Gombos F, Genzel L, Konrad BN, Simor P, Steiger A, Dresler M and Bódizs R (2015) Corrigendum: A comparison of two sleep spindle detection methods based on all night averages: individually adjusted vs. fixed frequencies. Front. Hum. Neurosci. 9:415. doi: 10.3389/fnhum.2015.00415