



Quorum Sensing by Monocyte-Derived Populations

Jérémy Postat^{1,2*} and Philippe Bousso^{1*}

¹ Dynamics of Immune Responses Unit, Institut Pasteur, INSERM U1223, Paris, France, ² Sorbonne Paris Cité, Cellule Pasteur, University Paris Diderot, Paris, France

OPEN ACCESS

Edited by:

Pierre Gueronprez,
Centre National de la Recherche
Scientifique (CNRS), France

Reviewed by:

Elodie Segura,
Institut Curie, France
Emmanuel L. Gautier,
Institut National de la Santé et de la
Recherche Médicale
(INSERM), France

*Correspondence:

Jérémy Postat
jeremy.postat@mail.mcgill.ca
Philippe Bousso
philippe.bousso@pasteur.fr

† Present address:

Jérémy Postat,
Department of Physiology, McGill
University, Montreal, QC, Canada

Specialty section:

This article was submitted to
Antigen Presenting Cell Biology,
a section of the journal
Frontiers in Immunology

Received: 17 May 2019

Accepted: 27 August 2019

Published: 11 September 2019

Citation:

Postat J and Bousso P (2019)
Quorum Sensing by
Monocyte-Derived Populations.
Front. Immunol. 10:2140.
doi: 10.3389/fimmu.2019.02140

Quorum sensing is a type of cellular communication that was first described in bacteria, consisting of gene expression regulation in response to changes in cell-population density. Bacteria synthesize and secrete diffusible molecules called autoinducers, which concentration varies accordingly with cell density and can be detected by the producing cells themselves. Once autoinducer concentration reaches a critical threshold, all bacteria within the autoinducer-rich environment react by modifying their genetic expression and adopt a coordinated behavior (e.g., biofilm formation, virulence factor expression, or swarming motility). Recent advances highlight the possibility that such type of communication is not restricted to bacteria, but can exist among other cell types, including immune cells and more specifically monocyte-derived cells (1). For such cells, quorum sensing mechanisms may not only regulate their population size and synchronize their behavior at homeostasis but also alter their activity and function in unexpected ways during immune reactions. Although the nature of immune autoinducers and cellular mechanisms remains to be fully characterized, quorum sensing mechanisms in the immune system challenge our traditional conception of immune cell interactions and likely represent an important mode of communication at homeostasis or during an immune response. In this mini-review, we briefly present the prototypic features of quorum sensing in bacteria and discuss the existing evidence for quorum sensing within the immune system. Mainly, we review quorum sensing mechanisms among monocyte-derived cells, such as the regulation of inflammation by the density of monocyte-derived cells that produce nitric oxide and discuss the relevance of such models in the context of immune-related pathologies.

Keywords: monocyte, monocyte-derived cell, quorum sensing (QS), macrophage, nitric oxide, metabolism

QUORUM SENSING: FROM THE BACTERIAL WORLD AND BEYOND

Living cell-based systems, such as multicellular organisms or bacterial biofilms, are very complex collections of biological components (cells) continually interacting with each other. They are organized as coordinated functional communities, which integrity relies on an efficient organization, and therefore, communication. Cells communicate with each other by various mechanisms either dependent on cell contact (e.g., juxtacrine signaling, membrane nanotubes) or dependent on diffusible material (e.g., diffusible signaling molecules, exosomes). Different modes of contact-independent signaling, including autocrine, paracrine, and endocrine signaling, were extensively characterized in the past using mammalian systems (2, 3). Autocrine signaling occurs

when the same cell simultaneously produces and responds to a signaling molecule that stays confined to the cell vicinity, with the help of high-affinity receptors. By contrast, paracrine signaling happens when the responding cell is located at a relatively short distance from the cell producing the signal, and often of a different cell type. Lastly, endocrine signaling is defined when the signaling and the target cells are located at distant sites, the signaling molecule traveling by the bloodstream.

Over 50 years ago, the most well-known bacterial mode of communication named quorum sensing was discovered in the luminous marine bacterial species *Vibrio fischeri* (4). Quorum sensing is a process of chemical communication that bacteria use to orchestrate group behaviors [(5–8); **Figure 1**]. In classic quorum sensing, each bacterium of the same type produces a membrane-diffusive signaling molecule called autoinducer, that can modify bacterial gene expression. The production of autoinducer by individual cells is too low to trigger any significant biological effect. When a threshold of cell density is reached (e.g., by continuous bacterial growth), the autoinducer reaches a sufficient concentration in the local environment to alter gene expression in all cells present in the area. The coordinated alteration of gene expression results in the emergence of group behaviors such as biofilm formation (9), virulence factor expression (10), or swarming motility (11). Therefore, quorum sensing mechanisms in bacteria initiate specific responses only when a sufficient cell density is reached, and rely on a diffusive molecule that acts as a surrogate for cell density (autoinducer). Quorum sensing mechanisms provide the possibility for spatiotemporal regulation of collections of cells, and for the emergence of specific behavior that would be unproductive when undertaken by a single isolated bacterium.

It is only recently that several immune regulatory processes similar to quorum sensing mechanisms have been revealed in the mammalian immune system. Notably, such processes were shown to be triggered locally only when a sufficient number of cells were reached and lead to population-level effects. Also, these mechanisms involve diffusive signaling molecules such as cytokines and chemokines, that resemble autoinducers in that they might be secreted in a reasonably low amount by single cells. Additionally, quorum sensing regulation may occur by mechanisms not only modifying gene expression, but also altering cellular metabolism.

In this mini-review, we will discuss the existence and biological relevance of quorum sensing mechanisms by immune cells at homeostasis and during inflammation, with a particular focus on monocyte-derived cells. We propose that quorum sensing mechanisms are integral to the immune system and that malfunction of such regulatory pathways may lead to uncontrolled monocyte-derived cell accumulation and activation, leading to excessive immune responses and the development of immunopathologies. We will not address the impact of bacterial quorum sensing molecules on monocyte activity, but rather extend the theory originally described in bacteria to the monocyte-derived cells.

MECHANISM OF QUORUM SENSING IN MONOCYTE-DERIVED CELLS

Monocyte-Derived Cell Homeostasis

In mammalian species, the concept of homeostasis establishes that most tissues and organs are made of different cell types whose numbers remain constant at steady-state. Accordingly, monocyte-derived cells such as monocyte-derived macrophages can maintain their cellular density in different organs under physiological conditions (12, 13). Such maintenance is achieved in the different tissues by compensating cell death either by a continuous input of circulating monocytes or by a constant self-renewal of tissue-resident macrophages as for Langerhans cells in the murine epidermis (12, 14–16). However, most of the mechanisms governing monocyte-derived cell homeostasis are still poorly understood. Recently, Antonioli et al. suggested that macrophage homeostasis is controlled under physiological conditions by a quorum sensing mechanism (1, 17). They propose a central role for Colony-Stimulating Factor 1 (CSF1) as the autoinducer for macrophages, as it controls their survival and proliferation at steady-state (18–20). Macrophage density would be controlled by two factors: CSF1 production by stromal cells and its consumption by the entire population. Accordingly, a two-cell circuit-based model between Platelet-Derived Growth Factor (PDGF)-producing macrophages and CSF1-producing fibroblasts was shown to have sufficient stability and robustness to perturbation to allow macrophage/stromal cell homeostasis (21). Further studies will provide new insight regarding the relevance of this model *in vivo* and its potential role not only at steady-state but also during ongoing immune responses. While this particular mechanism somewhat differs from classic microbial quorum sensing in that the autoinducer is produced by stromal cells (which are not the responding cells), it still aims at controlling the macrophage pool size and may be best defined as an indirect quorum sensing mechanism. Finally, the idea that cell density could also affect macrophage behavior at homeostasis has been less investigated. In one study using a model of cultured human monocytes, it was established that the secretion of chondroitin sulfate proteoglycan (structural component of human tissues) by these cells was dependent on their cell density (22).

Quorum Sensing During Immune Reactions

Other mechanisms controlling the size and activity of monocyte-derived cell populations were recently described during immune reactions. For instance, human macrophages infected by the intracellular pathogen *Mycobacterium tuberculosis* restrict bacterial growth more efficiently when cultivated at high density (23). The authors proposed that such finding is compatible with the possibility that high-density cultures release factors that can affect cell behavior [in that case, bacterial behavior]. More recently, TNF- α and IL-10 have been identified as the primary soluble mediators positively and negatively regulating macrophage function, capable of mediating cytokine production in groups vs. in single cells (1). Additionally, it was proposed that the regeneration of hair follicles in response to patterned hair plucking is regulated by a quorum sensing mechanism





FIGURE 1 | Quorum sensing in bacteria. To communicate and synchronize their behavior, bacteria (green rectangles) use quorum sensing. Each bacterium produces a low quantity of a membrane-diffusive molecule called autoinducer (blue triangles), which biological activity is absent at low concentration. Bacterial growth over time increases cell density together with the concentration of the autoinducer in the extracellular environment. Once a sufficient number of bacteria have accumulated, hence reaching a sufficient density, the autoinducer concentration is high enough to initiate biological alterations. In bacteria, the autoinducer often triggers a switch in genetic expression, after binding to transcriptional regulators. Such a genetic switch leads to the emergence of group behaviors such as biofilm formation, increased pathogenicity, or swarming motility.

involving macrophages (24). The authors uncovered a two-step mechanism where CCL2 released from damaged hairs leads to the recruitment of TNF- α -producing macrophages which accumulate and signal to both plucked and unplucked follicles to stimulate their regeneration.

More recently, a quorum sensing mechanism more similar to what described in bacteria was comprehensively described *in vivo* during the immune response against *Leishmania major* [(25); **Figure 2**]. Local infection with this intracellular parasite in murine skin triggers a massive recruitment of immune cells at the site of infection, including circulating monocytes. Recruited monocytes differentiate into mononuclear phagocytes that not only represent the main population of infected cells but also actively participate in controlling the local inflammatory reaction. Such an immune response can be detrimental to the host by inducing irreversible tissue damage if not adequately regulated and terminated on time (26, 27). Controlling mononuclear phagocyte recruitment, activity, and clearance is therefore essential to resolve inflammation concomitantly to parasite elimination. During cutaneous leishmaniasis, recruited mononuclear phagocytes secrete nitric oxide (NO) that acts to adjust and limit the overall inflammation intensity. NO suppresses mononuclear phagocyte accumulation, as well as cytokine and chemokine production by blocking cellular respiration and decreasing the ATP:ADP ratio. Importantly, such mechanism only exists when a sufficient number of mononuclear

phagocytes have accumulated at the site of infection (more than 5,000 cells per mm³) to produce a high quantity of NO in a collective manner (**Figure 2**). NO acts thereafter by diffusing and targeting all mononuclear phagocytes irrespectively of their intrinsic iNOS expression. In this mechanism, the mitochondria, and most probably the cytochrome c oxidase, represents the target of NO (28, 29). Therefore, mononuclear phagocytes not only produce NO but are also regulated in number and activity by this diffusive molecule, establishing a quorum sensing mechanism for the control of the inflammatory reaction, with NO acting as the autoinducer. Furthermore, it is very interesting to note that such mechanism relies on the modification of cellular metabolism. That shows, in contrast to the current paradigm in bacteria, that quorum sensing mechanisms do not necessarily rely on genetic alteration. Targeting cellular metabolism compared to genetic expression could have several advantages, including the possibility to alter cell activity within a very short period of time and in a rapidly reversible manner (25). In addition to dampening oxidative phosphorylation, NO was also shown to accelerate myeloid cell death (30, 31), a phenomenon that could participate in inflammation resolution (32). We propose that this NO-based quorum sensing mechanism is integral to the biology of inflammatory mononuclear phagocytes and could certainly operate in other models of infections or during cancer development. We additionally propose that malfunction of quorum sensing mechanisms may lead to uncontrolled



mononuclear phagocyte accumulation and activation, leading to non-resolving inflammation and therefore immunopathology development (26).

Furthermore, other quorum sensing mechanisms are about to be elucidated during immune reactions. Indeed, using a droplet-based microfluidic approach, Wimmers et al. showed that during human pDCs activation, the fraction of IFN- α -expressing cells at early time points of activation is dependent on their initial cell density. Using microtiter plates, they measured that more than 10^3 pDCs per well should be stimulated together to have a significant percentage of IFN- α -producing cells following activation (33). That suggests that the magnitude of dendritic cell activity significantly increases above a threshold number of cells and that these cells are probably endowed with quorum sensing capability. Similarly, it was recently showed using mathematical modeling that macrophage activation is most probably bimodal, with a proportion of highly activated cells increasing with cell density. Because only a fraction of the population becomes activated, the authors preferably describe such phenomenon as quorum licensing rather than quorum sensing (34). Such studies and early work on human macrophages establish the potential relevance of quorum sensing mechanisms for the control of immune reaction in humans.

Benefits Over and in Conjunction With Autocrine and Paracrine Signaling

Immune cell communication by quorum sensing mechanisms provides many benefits over autocrine or paracrine signaling. Quorum sensing mechanisms can significantly differ from paracrine signaling, a distinction that is not systematically present in the literature. To distinguish the two communication modes, further studies will have to investigate whether the diffusive molecule has any biological effect when the producing cell is present at low density. Indeed, the absence of biological effect of the signaling molecule (autoinducer) when its producer is at too low density is a cornerstone of quorum sensing. In other words, if a diffusive molecule biologically acts when the producer cell is at a negligible density, the communication occurs through paracrine signaling.

A first advantage of quorum sensing is the regulation of the cell number *per se* in tissues during inflammation (35). For instance, a high T cell density is needed for their terminal differentiation into cytotoxic T cells (36). By contrast, an excessive number of cells involved in an immune response can trigger immunopathology (25). Quorum sensing mechanisms may therefore locally adjust the cell density to promote or stop the immune reaction. Next, compared to autocrine signaling,

the use of a diffusive mediator (the autoinducer) that acts simultaneously on numerous cells offers a way to reduce the level of heterogeneity between cells, and therefore coordinate their behavior in their complex microenvironment. For instance, DC populations can synchronize their behavior at late time points of activation provided that a small fraction of them secrete type I interferon rapidly after stimulation (37). Also, compared to paracrine signaling, quorum sensing mechanisms have the potential to temporally and locally adjust cell density and inflammation intensity without the need for external cues or regulatory cells. Paracrine signaling would not allow such self-adjustment to exist because it implies at least two cell types: the producing and the target cells. In a way, quorum sensing mechanisms can be considered as a form of paracrine signaling that depends on the cell population density but in which cells produce both a signaling molecule and its receptor, as in autocrine signaling (38).

Nonetheless, these different types of communication do not seem mutually exclusive. For instance, during the immune reaction against the parasite *L. major* in murine skin, a two-wave immune cell regulation occurs (39). First, a wave of IFN- γ secreted by activated CD4⁺ T cells spreads away from the site of antigen presentation and induces iNOS expression in numerous infected and bystander monocyte-derived cells, by a mechanism resembling paracrine signaling (40). Next, the subsequent collective production of NO allows for both parasite control (41) and regulation of the inflammation intensity at the tissue level by a metabolism-based quorum sensing mechanism (25). Thus, both paracrine signaling and quorum sensing can act in concert to spread a signal originating from a few discrete sites of cell activation to the level of the entire organ, while keeping the inflammatory reaction locally under tight control. Another example of coupling also exists for type I IFN during pDCs activation. In such an event, IFN- α stimulates its own production and alter cellular metabolism via an autocrine amplification loop (33, 42) but also regulates the fraction of IFN- α -producing cell most probably by a quorum sensing mechanism. Indeed, the fraction of activated pDCs is dependent on their cell density and relies on the diffusion of IFN- α that binds most probably IFNAR1 to mediate its biological effects (33, 43). Hence, combining quorum sensing with either autocrine or paracrine signaling appears to be essential to fine-tune immune cell activity.

QUORUM SENSING BY IMMUNE CELLS: PERSPECTIVES

Quorum Sensing by Other Immune Cell Types

While we focused on monocyte-derived cells, quorum sensing mechanisms have been described as well in other immune cells such as T and B cells.

In T cells, interleukin-2 (IL-2) was shown to be a significant autoinducer involved in a quorum sensing regulatory loop stabilizing T cell population density and phenotype (44–48). With the help of mathematical models and *in vitro* experiments, it was shown that a sufficient density of T_{EFF} cells is critical to reach

a minimum threshold of IL-2 above which the phosphorylation of the signal transducer and activator of transcription (STAT) 5 is sustained to allow T cell proliferation (45). It was demonstrated using the same strategy that an excess of IL-2 above a maximal threshold leads to cell death instead of proliferation, participating in the regulation of T cell density to reach homeostasis (46). As well, it was recently evidenced that T cell density can directly modulate T cell differentiation toward T_{EFF} or T_{CM} by a quorum sensing mechanism relying on IL-2 and IL-6 as autoinducers (47). Additionally, during an immune challenge, T cells were shown to establish a negative feedback loop by capturing their cognate pMHC complexes from antigen-presenting cells and presenting them to antigen-experienced CD4⁺ T cells, thereby inhibiting their recruitment into the ongoing response (49). Finally, it has been proposed that T cell activation require a quorum of lymphocytes to happens (50).

Additionally, B cells were also shown to be endowed with a quorum sensing regulatory mechanism (51). After challenge, activated B cells secrete soluble immunoglobulin G (IgG) which concentration rises in the serum and is detected by the inhibitory receptor Fc γ RIIB, present on all B cells. At a sufficient concentration of IgG, the binding on this receptor is enough to trigger intracellular inhibitory pathways and prevent further IgM-secreting B cell activation. As a result, the number of IgM-secreting B cells is kept under control (51). So far, only IgG molecules were shown to act as autoinducer for B cells.

Outstanding Questions and Perspectives

We envision that quorum sensing is integral to the regulation of immune responses. In this respect, future studies are critically needed to extend our understanding of this mode of communication. Several outstanding questions remain open: How autoinducer concentrations evolve in tissues during immune reactions? How can the physical tissue architecture and organ structure impact autoinducer diffusion, distribution, and access to the cells? How accurately can a cell population sense its own density based on an autoinducer concentration? Are there other unpredicted advantages of quorum sensing in immune cells? Do regulatory mechanisms for quorum sensing, such as quorum sensing quenching (52), can be triggered by invading pathogens or by the immune cells themselves? Can we target quorum sensing mechanisms in the context of immunotherapies (53). Further studies in both bacterial and mammalian systems will help answer these questions (54–57).

Given the complexity of the immune system and the vast number of quorum sensing mechanisms described in bacteria, we anticipate systems immunology to be essential in unraveling new quorum sensing mechanisms in immune cells (58). Mathematical modeling and computational simulations should help identify new quorum sensing mechanisms in immune cells, as it has been done for T cells (46, 59, 60) and more recently in macrophages (34). Furthermore, the development of new biological tools to measure autoinducer concentration and diffusion in complex tissue microenvironment are needed. For instance, reliable tools to map IL-2 or nitric oxide gradient *in vivo* would be of great help to better decipher quorum sensing mechanisms and characterize the effect of tissue architecture of its efficacy.

CONCLUDING REMARKS

While quorum sensing is the norm in the bacteria world, it is only recently that similar mechanisms were shown to exist in the immune system, including in monocyte-derived cells. Quorum sensing mechanisms provide a way to regulate immune cell activity concurrently in a spatial and temporal manner complementary to what autocrine or paracrine signaling can achieve. They also favor the emergence of group behaviors and synchronized responses, two features decreasing the sensitivity of the system to external perturbations and therefore increasing its robustness. Alteration of immune quorum sensing mechanisms by impaired access or dysregulated response to the autoinducer may certainly trigger the emergence of immunopathology, as demonstrated in the context of the infection by the parasite

L. major. Future studies are needed to extend the concept to other cell types and models to provide a better understanding of how this unique mode of communication integrates within the complexity of immune reactions.

AUTHOR CONTRIBUTIONS

All authors listed drafted, wrote, and edited the manuscript until the final version was approved and contributed to figures.

FUNDING

The work was supported by Institut Pasteur, INSERM and an Advanced grant (ENLIGHTEN) from the European Research Council.

REFERENCES

- Antonioli L, Blandizzi C, Pacher P, Williams M, Haskó G. Rethinking communication in the immune system: the quorum sensing concept. *Trends Immunol.* (2019) 40:88–97. doi: 10.1016/j.it.2018.12.002
- Reece JB, Urry LA, Cain ML, Wasserman SA, Minorsky PV, Jackson RB. *Campbell Biology*. London: Pearson (2014).
- Murphy K, Weaver C. *Janeway's Immunobiology*. Oxfordshire: Garland Science/Taylor & Francis Group, LLC (2016).
- Neelson KH, Hastings JW. Bacterial bioluminescence: its control and ecological significance. *Microbiol Rev.* (1979) 43:496–518.
- Eickhoff MJ, Bassler BL. SnapShot: bacterial quorum sensing. *Cell.* (2018) 174:1328.e1. doi: 10.1016/j.cell.2018.08.003
- Miller MB, Bassler BL. Quorum sensing in bacteria. *Annu Rev Microbiol.* (2001) 55:165–99. doi: 10.1146/annurev.micro.55.1.165
- Ng WL, Bassler BL. Bacterial quorum-sensing network architectures. *Annu Rev Genet.* (2009) 43:197–222. doi: 10.1146/annurev-genet-102108-134304
- Waters CM, Bassler BL. Quorum sensing: cell-to-cell communication in bacteria. *Annu Rev Cell Dev Biol.* (2005) 21:319–46. doi: 10.1146/annurev.cellbio.21.012704.131001
- Solano C, Echeverez M, Lasa I. Biofilm dispersion and quorum sensing. *Curr Opin Microbiol.* (2014) 18:96–104. doi: 10.1016/j.mib.2014.02.008
- Rutherford ST, Bassler BL. Bacterial quorum sensing: its role in virulence and possibilities for its control. *Cold Spring Harb Perspect Med.* (2012) 2:a012427. doi: 10.1101/cshperspect.a012427
- Daniels R, Vanderleyden J, Michiels J. Quorum sensing and swarming migration in bacteria. *FEMS Microbiol Rev.* (2004) 28:261–89. doi: 10.1016/j.femsre.2003.09.004
- Ginhoux F, Williams M. Tissue-resident macrophage ontogeny and homeostasis. *Immunity.* (2016) 44:439–49. doi: 10.1016/j.immuni.2016.02.024
- Raff MC. Size control: the regulation of cell numbers in animal development. *Cell.* (1996) 86:173–5. doi: 10.1016/S0092-8674(00)80087-2
- Scott CL, Henri S, Williams M. Mononuclear phagocytes of the intestine, the skin, and the lung. *Immunol Rev.* (2014) 262:9–24. doi: 10.1111/imr.12220
- Tamoutounour S, Williams M, Montanana Sanchis F, Liu H, Terhorst D, Malosse C, et al. Origins and functional specialization of macrophages and of conventional and monocyte-derived dendritic cells in mouse skin. *Immunity.* (2013) 39:925–38. doi: 10.1016/j.immuni.2013.10.004
- Yona S, Kim KW, Wolf Y, Mildner A, Varol D, Breker M, et al. Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. *Immunity.* (2013) 38:79–91. doi: 10.1016/j.immuni.2012.12.001
- Antonioli L, Blandizzi C, Pacher P, Williams M, Haskó G. Quorum sensing in the immune system. *Nat Rev Immunol.* (2018) 18:537. doi: 10.1038/s41577-018-0040-4
- Hamilton JA, Cook AD, Tak PP. Anti-colony-stimulating factor therapies for inflammatory and autoimmune diseases. *Nat Rev Drug Discov.* (2017) 16:53–70. doi: 10.1038/nrd.2016.231
- Jenkins SJ, Hume DA. Homeostasis in the mononuclear phagocyte system. *Trends Immunol.* (2014) 35:358–67. doi: 10.1016/j.it.2014.06.006
- Dai XM, Ryan GR, Hapel AJ, Dominguez MG, Russell RG, Kapp S, et al. Targeted disruption of the mouse colony-stimulating factor 1 receptor gene results in osteopetrosis, mononuclear phagocyte deficiency, increased primitive progenitor cell frequencies, and reproductive defects. *Blood.* (2002) 99:111–20. doi: 10.1182/blood.V99.1.111
- Zhou X, Franklin RA, Adler M, Jacox JB, Bailis W, Shyer JA, et al. Circuit design features of a stable two-cell system. *Cell.* (2018) 172:744–57.e17. doi: 10.1016/j.cell.2018.01.015
- Uhlin-Hansen L, Kolset SO. Cell density-dependent expression of chondroitin sulfate proteoglycan in cultured human monocytes. *J Biol Chem.* (1988) 263:2526–31.
- Boechat N, Bouchonnet F, Bonay M, Grodet A, Pelicic V, Gicquel B, et al. Culture at high density improves the ability of human macrophages to control mycobacterial growth. *J Immunol.* (2001) 166:6203–11. doi: 10.4049/jimmunol.166.10.6203
- Chen CC, Wang L, Plikus MV, Jiang TX, Murray PJ, Ramos R, et al. Organ-level quorum sensing directs regeneration in hair stem cell populations. *Cell.* (2015) 161:277–90. doi: 10.1016/j.cell.2015.02.016
- Postat J, Olekhovitch R, Lemaître F, Bousso P. A metabolism-based quorum sensing mechanism contributes to termination of inflammatory responses. *Immunity.* (2018) 49:654–65.e5. doi: 10.1016/j.immuni.2018.07.014
- Nathan C, Ding A. Nonresolving inflammation. *Cell.* (2010) 140:871–82. doi: 10.1016/j.cell.2010.02.029
- Ortega-Gómez A, Perretti M, Soehnlein O. Resolution of inflammation: an integrated view. *EMBO Mol Med.* (2013) 5:661–74. doi: 10.1002/emmm.201202382
- Brown GC. Regulation of mitochondrial respiration by nitric oxide inhibition of cytochrome c oxidase. *Biochim Biophys Acta.* (2001) 1504:46–57. doi: 10.1016/S0005-2728(00)00238-3
- Poderoso JJ, Helfenberger K, Poderoso C. The effect of nitric oxide on mitochondrial respiration. *Nitric Oxide.* (2019) 88:61–72. doi: 10.1016/j.niox.2019.04.005
- Everts B, Amiel E, van der Windt GJ, Freitas TC, Chott R, Yarasheski KE, et al. Commitment to glycolysis sustains survival of NO-producing inflammatory dendritic cells. *Blood.* (2012) 120:1422–31. doi: 10.1182/blood-2012-03-419747
- Amiel E, Everts B, Fritz D, Beauchamp S, Ge B, Pearce EL, et al. Mechanistic target of rapamycin inhibition extends cellular lifespan in dendritic cells by preserving mitochondrial function. *J Immunol.* (2014) 193:2821–30. doi: 10.4049/jimmunol.1302498
- Gautier EL, Ivanov S, Lesnik P, Randolph GJ. Local apoptosis mediates clearance of macrophages from resolving inflammation in mice. *Blood.* (2013) 122:2714–22. doi: 10.1182/blood-2013-01-478206
- Wimmers F, Subedi N, van Buuringen N, Heister D, Vivié J, Beeren-Reinieren I, et al. Single-cell analysis reveals that stochasticity and paracrine signaling

- control interferon-alpha production by plasmacytoid dendritic cells. *Nat Commun.* (2018) 9:3317. doi: 10.1038/s41467-018-05784-3
34. Muldoon JJ, Chuang Y, Bagheri N, Leonard JN. Macrophages employ quorum licensing to regulate collective activation. *bioRxiv.* (2019) 610592. doi: 10.1101/610592
 35. Perié L, Aru J, Kourilsky P, Slotine JJ. Does a quorum sensing mechanism direct the behavior of immune cells? *C R Biol.* (2013) 336:13–6. doi: 10.1016/j.crvi.2013.01.006
 36. Thaventhiran JE, Hoffmann A, Magiera L, de la Roche M, Lingel H, Brunner-Weinzierl M, et al. Activation of the Hippo pathway by CTLA-4 regulates the expression of Blimp-1 in the CD8+ T cell. *Proc Natl Acad Sci USA.* (2012) 109:E2223–9. doi: 10.1073/pnas.1209115109
 37. Shalek AK, Satija R, Shuga J, Trombetta JJ, Gennert D, Lu D, et al. Single-cell RNA-seq reveals dynamic paracrine control of cellular variation. *Nature.* (2014) 510:363–9. doi: 10.1038/nature13437
 38. Doganer BA, Yan LKQ, Youk H. Autocrine signaling and quorum sensing: extreme ends of a common spectrum. *Trends Cell Biol.* (2016) 26:262–71. doi: 10.1016/j.tcb.2015.11.002
 39. Olekhnovitch R, Bousoo P. Induction, propagation, and activity of host nitric oxide: lessons from *Leishmania* infection. *Trends Parasitol.* (2015) 31:653–64. doi: 10.1016/j.pt.2015.08.001
 40. Müller AJ, Filipe-Santos O, Eberl G, Aebischer T, Späth GF, Bousoo P. CD4+ T cells rely on a cytokine gradient to control intracellular pathogens beyond sites of antigen presentation. *Immunity.* (2012) 37:147–57. doi: 10.1016/j.immuni.2012.05.015
 41. Olekhnovitch R, Ryffel B, Müller AJ, Bousoo P. Collective nitric oxide production provides tissue-wide immunity during *Leishmania* infection. *J Clin Invest.* (2014) 124:1711–22. doi: 10.1172/JCI172058
 42. Wu D, Sanin DE, Everts B, Chen Q, Qiu J, Buck MD, et al. Type I interferons induce changes in core metabolism that are critical for immune function. *Immunity.* (2016) 44:1325–36. doi: 10.1016/j.immuni.2016.06.006
 43. Pehler J, Thomas C, Garcia KC, Schreiber G. Structural and dynamic determinants of type I interferon receptor assembly and their functional interpretation. *Immunol Rev.* (2012) 250:317–34. doi: 10.1111/imr.12001
 44. Almeida AR, Amado IF, Reynolds J, Berges J, Lythe G, Molina-Paris C, et al. Quorum-sensing in CD4+ T cell homeostasis: a hypothesis and a model. *Front Immunol.* (2012) 3:125. doi: 10.3389/fimmu.2012.00125
 45. Feinerman O, Jentsch G, Tkach KE, Coward JW, Hathorn MM, Sneddon MW, et al. Single-cell quantification of IL-2 response by effector and regulatory T cells reveals critical plasticity in immune response. *Mol Syst Biol.* (2010) 6:437. doi: 10.1038/msb.2010.90
 46. Hart Y, Reich-Zeliger S, Antebi YE, Zaretsky I, Mayo AE, Alon U, et al. Paradoxical signaling by a secreted molecule leads to homeostasis of cell levels. *Cell.* (2014) 158:1022–32. doi: 10.1016/j.cell.2014.07.033
 47. Polonsky M, Rimer J, Kern-Perets A, Zaretsky I, Miller S, Bornstein C, et al. Induction of CD4 T cell memory by local cellular collectivity. *Science.* (2018) 360:eaaj1853. doi: 10.1126/science.aaj1853
 48. Savir Y, Waysbort N, Antebi YE, Tlusty T, Friedman N. Balancing speed and accuracy of polyclonal T cell activation: a role for extracellular feedback. *BMC Syst Biol.* (2012) 6:111. doi: 10.1186/1752-0509-6-111
 49. Helft J, Jacquet A, Joncker NT, Grandjean I, Dorothée G, Kissenpfennig A, et al. Antigen-specific T-T interactions regulate CD4 T-cell expansion. *Blood.* (2008) 112:1249–58. doi: 10.1182/blood-2007-09-114389
 50. Al-Yassin GA, Bretscher PA. Does T cell activation require a quorum of lymphocytes? *J Immunol.* (2018) 201:2855–61. doi: 10.4049/jimmunol.1800805
 51. Montaudouin C, Anson M, Hao Y, Duncker SV, Fernandez T, Gaudin E, et al. Quorum sensing contributes to activated IgM-secreting B cell homeostasis. *J Immunol.* (2013) 190:106–14. doi: 10.4049/jimmunol.1200907
 52. Pietschke C, Treitz C, Forêt S, Schultze A, Künzel S, Tholey A, et al. Host modification of a bacterial quorum-sensing signal induces a phenotypic switch in bacterial symbionts. *Proc Natl Acad Sci USA.* (2017) 114:E8488–97. doi: 10.1073/pnas.1706879114
 53. Jayatilaka H, Tyle P, Chen JJ, Kwak M, Ju J, Kim HJ, et al. Synergistic IL-6 and IL-8 paracrine signalling pathway infers a strategy to inhibit tumour cell migration. *Nat Commun.* (2017) 8:15584. doi: 10.1038/ncomms15584
 54. Bagnall J, Boddington C, England H, Brignall R, Downton P, Alsoufi Z, et al. Quantitative analysis of competitive cytokine signaling predicts tissue thresholds for the propagation of macrophage activation. *Sci Signal.* (2018) 11:eaaf3998. doi: 10.1126/scisignal.aaf3998
 55. Whiteley M, Diggle SP, Greenberg EP. Progress in and promise of bacterial quorum sensing research. *Nature.* (2017) 551:313–20. doi: 10.1038/nature24624
 56. Oylar-Yaniv A, Oylar-Yaniv J, Whitlock BM, Liu Z, Germain RN, Huse M, et al. A tunable diffusion-consumption mechanism of cytokine propagation enables plasticity in cell-to-cell communication in the immune system. *Immunity.* (2017) 46:609–20. doi: 10.1016/j.immuni.2017.03.011
 57. Assen FP, Sixt M. The dynamic cytokine niche. *Immunity.* (2017) 46:519–20. doi: 10.1016/j.immuni.2017.04.006
 58. Villani AC, Sarkizova S, Hacohen N. Systems immunology: learning the rules of the immune system. *Annu Rev Immunol.* (2018) 36:813–42. doi: 10.1146/annurev-immunol-042617-053035
 59. Youk H, Lim WA. Secreting and sensing the same molecule allows cells to achieve versatile social behaviors. *Science.* (2014) 343:1242782. doi: 10.1126/science.1242782
 60. Youk H, Lim WA. Sending mixed messages for cell population control. *Cell.* (2014) 158:973–5. doi: 10.1016/j.cell.2014.08.013

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Postat and Bousoo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.