



Corrigendum: Monomeric C-Reactive Protein Binds and Neutralizes Receptor Activator of NF-κB Ligand-Induced Osteoclast Differentiation

OPEN ACCESS

Edited and reviewed by:

Mark Slevin,
Manchester Metropolitan University,
United Kingdom

*Correspondence:

Yi Wu
wuy@lzu.edu.cn
Shang-Rong Ji
jsr@lzu.edu.cn

Specialty section:

This article was submitted to
Inflammation,
a section of the journal
Frontiers in Immunology

Received: 22 October 2020

Accepted: 30 October 2020

Published: 24 November 2020

Citation:

Jia Z-K, Li H-Y, Liang Y-L, Potempa LA, Ji S-R and Wu Y (2020) Corrigendum: Monomeric C-Reactive Protein Binds and Neutralizes Receptor Activator of NF-κB Ligand-Induced Osteoclast Differentiation. *Front. Immunol.* 11:619847. doi: 10.3389/fimmu.2020.619847

Zhe-Kun Jia¹, Hai-Yun Li², Yu-Lin Liang¹, Lawrence A. Potempa³, Shang-Rong Ji^{1*} and Yi Wu^{2,4*}

¹ MOE Key Laboratory of Cell Activities and Stress Adaptations, School of Life Sciences, Lanzhou University, Lanzhou, China, ² MOE Key Laboratory of Environment and Genes Related to Diseases, School of Basic Medical Sciences, Xi'an Jiaotong University, Xi'an, China, ³ Roosevelt University College of Pharmacy, Schaumburg, IL, United States, ⁴ The Affiliated Children's Hospital of Xi'an Jiaotong University, Xi'an, China

Keywords: inflammation, rheumatoid arthritis, osteoclast, receptor activator of NF-κB ligand, C-reactive protein

A Corrigendum on

Monomeric C-Reactive Protein Binds and Neutralizes Receptor Activator of NF-κB Ligand-Induced Osteoclast Differentiation

By Jia Z-K, Li H-Y, Liang Y-L, Potempa LA, Ji S-R and Wu Y (2018). *Front. Immunol.* 9:234. doi: 10.3389/fimmu.2018.00234.

In the original article, there was a mistake in **Figure 4B** as published. The lower panel of **Figure 4B** was erroneously assigned. The corrected **Figure 4B** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2020 Jia, Li, Liang, Potempa, Ji and Wu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 4 | mCRP neutralizes the effects of receptor activator of NF- κ B ligand (RANKL). Bone marrow-derived macrophages (BMDMs) were treated with 10 ng/mL of M-CSF and 50 ng/ml RANKL in the presence or absence of 100 μ g/mL of mCRP. After treatment for 2 days, the expression of TRAP and Cathepsin K were determined by q-PCR (**A**). After treatment for 6 days, cells were stained for TRAP (**B**) and counted for the number of osteoclasts (**C**). Their bone resorption activities were evaluated by toluidine blue staining (**D**) and the quantification of eroded surface (**E**). The potent effects of RANKL on BMDMs were absent when treated together with mCRP. Comparable results were also obtained with Raw cells (**B–E**). * $p < 0.05$; *** $p < 0.001$.