



Therapeutic Potential of Anti-Interferon α Vaccination on SjS-Related Features in the MRL/lpr Autoimmune Mouse Model

Martin Killian^{1,2}, Fabien Colaone³, Philippe Haumont³, Carole Nicco⁴, Olivier Cerles⁴, Sandrine Chouzenoux⁴, Pascal Cathébras^{1,2}, Nicolas Rochereau¹, Blandine Chanut¹, Mireille Thomas⁵, Norbert Laroche⁵, Fabien Forest⁶, Géraldine Grouard-Vogel^{3†}, Frédéric Batteux^{4†} and Stéphane Paul^{1*†}

OPEN ACCESS

Edited by:

Bertrand Kaeffer,
Institut National de recherche pour
l'agriculture, l'alimentation et
l'environnement (INRAE), France

Reviewed by:

Gwenny M. Verstappen,
University Medical Center
Groningen, Netherlands
Peter Delputte,
University of Antwerp, Belgium

*Correspondence:

Stéphane Paul
stephane.paul@chu-st-etienne.fr

[†]These authors jointly supervised
this work

Specialty section:

This article was submitted to
Vaccines and Molecular Therapeutics,
a section of the journal
Frontiers in Immunology

Received: 09 February 2021

Accepted: 21 October 2021

Published: 17 November 2021

Citation:

Killian M, Colaone F, Haumont P,
Nicco C, Cerles O, Chouzenoux S,
Cathébras P, Rochereau N, Chanut B,
Thomas M, Laroche N, Forest F,
Grouard-Vogel G, Batteux F and
Paul S (2021) Therapeutic Potential of
Anti-Interferon α Vaccination on
SjS-Related Features in the MRL/lpr
Autoimmune Mouse Model.
Front. Immunol. 12:666134.
doi: 10.3389/fimmu.2021.666134

¹ Centre International de Recherche en Infectiologie (CIRI), Team Groupe Immunité des Muqueuses et Agents Pathogènes (GIMAP), Université de Lyon, Université Jean Monnet, Université Claude Bernard Lyon 1, Inserm, U1111, Centre National de la Recherche Scientifique (CNRS), UMR530, Saint-Etienne, France, ² Internal Medicine Department, Saint-Etienne University Hospital, Saint-Etienne, France, ³ NEOVACS S.A., Paris, France, ⁴ Team Stress Oxydant, Prolifération Cellulaire et Inflammation, Institut National de la Santé Et de la Recherche Médicale (INSERM) U1016 Institut Cochin, Paris, France, ⁵ Institut National de la Santé Et de la Recherche Médicale (INSERM) U1059-Sainbiose, Université de Lyon, Saint Priest en Jarez, France, ⁶ Department of Pathology, Saint-Etienne University Hospital, Saint-Etienne, France

Sjögren's syndrome (SjS) is a frequent systemic autoimmune disease responsible for a major decrease in patients' quality of life, potentially leading to life-threatening conditions while facing an unmet therapeutic need. Hence, we assessed the immunogenicity, efficacy, and tolerance of IFN-Kinoid (IFN-K), an anti-IFN α vaccination strategy, in a well-known mouse model of systemic autoimmunity with SjS-like features: MRL/MpJ-Faslpr/lpr (MRL/lpr) mice. Two cohorts (with ISA51 or SWE01 as adjuvants) of 26 female MRL/lpr were divided in parallel groups, "controls" (not treated, PBS and Keyhole Limpet Hemocyanin [KLH] groups) or "IFN-K" and followed up for 122 days. Eight-week-old mice received intra-muscular injections (days 0, 7, 28, 56 and 84) of PBS, KLH or IFN-K, emulsified in the appropriate adjuvant, and blood samples were serially collected. At sacrifice, surviving mice were euthanized and their organs were harvested for histopathological analysis (focus score in salivary/lacrimal glands) and IFN signature evaluation. SjS-like features were monitored. IFN-K induced a disease-modifying polyclonal anti-IFN α antibody response in all treated mice with high IFN α neutralization capacities, type 1 IFN signature's reduction and disease features' (ocular and oral sicca syndrome, neuropathy, focus score, glandular production of BAFF) improvement, as reflected by the decrease in Murine Sjögren's Syndrome Disease Activity Index (MuSSDAI) modelled on EULAR Sjögren's Syndrome Disease Activity Index (ESSDAI). No adverse effects were observed. We herein report on the strong efficacy of an innovative anti-IFN α vaccination strategy in a mouse model of SjS, paving the way for further clinical development (a phase IIb trial has just been completed in systemic lupus erythematosus with promising results).

Keywords: Sjögren's syndrome, vaccination, interferon α (IFN- α), adjuvant, kinoid

INTRODUCTION

Sjögren's syndrome (SjS) is a frequent and predominantly female (10:1) systemic autoimmune disease, affecting 0.01-0.3% of the adult population worldwide (1). Sicca syndrome (i.e. dry eyes and/or mouth), lymphocytic infiltration of the salivary and lacrimal glands and anti-Ro/Sjögren's Syndrome A (SSA) and anti-La/Sjögren's Syndrome B (SSB) autoantibodies are typical in SjS (2). Apart from these classical features, the clinical spectrum is broad with systemic (i.e. extra-glandular) manifestations in one third of cases (3, 4). SjS's burden in terms of quality of life is well-known but the disease is also responsible for increased mortality with recent estimates of the adjusted standardized mortality ratio of 4.66 (95% CI 3.85 to 5.60) (5). Despite these worrisome elements, therapeutic resources are scarce, artificial tears and secretagogues only offer temporary relief, and no immunomodulatory treatment was superior to placebo in randomized controlled trials to date (6), partly because of the difficulty in the choice of the right outcome (7). However, a growing body of evidence indicates a major role for type 1 interferons (IFNs) in Sjögren's pathogenesis (8), susceptibility (9), and disease activity/complications (10, 11), hence becoming a potential therapeutic target. Among these findings, the identification of a type 1 IFN signature – the overexpression of type 1 IFN-inducible genes – in the blood and salivary glands of more than 50% of SjS patients (9, 10), its correlation with disease activity and B cell activating factor (BAFF) – which is type 1 IFN-inducible – overexpression (10, 12), the description of potentially protective endogenous anti-IFN α autoantibodies in SjS (13), and the promising results of recent studies evaluating anti-B cells therapies such as the anti-BAFF mAbs, belimumab (14), alone or combined with rituximab (15, 16), or ianalumab (17) (whose phase III trial is about to start in 2021), as well as a preclinical trial with filgotinib (18), a JAK1 inhibitor (with effects on IFN pathways, including BAFF production), are noteworthy. We previously showed that an active immunotherapy strategy, IFN-Kinoid (IFN-K), made up of conjugated IFN α and Keyhole Limpet Hemocyanin (KLH), was successful in preventing systemic lupus erythematosus (SLE)-manifestations and improving survival in New Zealand Black and White (NZB/W) mice (19), by inducing neutralizing anti-IFN α antibodies (Abs). Phase I/IIa (20, 21) and IIb (22) trials, in human SLE showed encouraging results, allowing the pursuit of IFN-K's clinical development.

Given the pivotal role of type 1 IFN in SjS and the urgent need for treatments (23), we investigated in a preclinical study using the MRL/MpJ-Fas^{lpr/lpr} (MRL/lpr) model, a well-recognized mouse model of systemic autoimmunity with SjS-like features (24), the potential benefits of the murine surrogate of the human IFN-K vaccine.

MATERIALS AND METHODS

Mouse Model

The MRL/lpr mouse model is a well-recognized congenital model of SjS (25–27), whose autoimmunity was found to be

particularly caused by an acquired mutation in the lymphoproliferation (lpr) gene, leading to a defect in Fas-mediated apoptosis of auto-reactive T cells, prompting the proliferation of double negative (CD4⁻ CD8⁻) T cells [which were previously found to be involved in the pathogenic mechanisms leading to SjS (28, 29)].

Preparation of the Immunogen

IFN-K was prepared by complexing murine IFN α 3 (PBL Assay Science, Piscataway, NJ) with KLH, by aldehyde treatment, as previously reported (19, 30).

Study Approval

All experiments received approval of the INSERM and Université Paris Descartes CEEA34 Ethics committee on Animal Experimentation, under the number 15-012.

Experimental Protocol

Two cohorts of 26 female MRL/MpJ-Fas^{lpr/lpr} (MRL/lpr, bred in our colony) mice were followed simultaneously (**Figure 1A**). One cohort, injected with ISA51 (Seppic, Paris, France), a water-in-oil adjuvant, was randomly divided into 4 parallel groups: NT (n=6), PBS/ISA51 (n=6), KLH/ISA51 (n=7) and IFN-K/ISA51 (n=7). The other cohort, whose adjuvant was SWE01 (University of Lausanne, Switzerland), an MF59-like oil-in-water emulsion, was also randomly divided into 4 parallel groups: NT (n=6), PBS/SWE01 (n=6), KLH/SWE01 (n=7) and IFN-K/SWE01 (n=7). NT, PBS and KLH-receiving mice were considered as Controls of IFN-K-treated mice.

Except for NT mice, 8-weeks-old-mice received 5 i.m. injections (0.1 mL) of PBS, KLH (10 μ g) or IFN-K (10 μ g), emulsified in the appropriate adjuvant at a ratio of 1:1, at day 0 (i.e. 54th day of life), 7, 28, 56 and 84.

Blood samples were collected in all mice by retro-orbital bleeding, at day -11 (pre-immune), 38, 66, 94 and 122 (final bleed). At final bleed, surviving mice were euthanized and their organs were harvested.


The stimulated salivary flow rate measurement, phenol red thread test, Von Frey test, cold plate assay, open field test and forced swim test were all performed 2 weeks before, and 8 and 16 weeks after immunization. The sucrose consumption test was performed 1 week before, and 9 and 17 weeks after immunization.

Anti-IFN α /KLH Titer Assay

Serum anti-IFN α and anti-KLH antibody titers were determined by ELISA, and tested in duplicate, as previously described (19). Briefly, pre-coated ELISA plates with 100 ng of murine IFN α 3 (PBL Assay Science, Piscataway, NJ) per well or 100 ng of KLH per well were incubated with serial dilutions of the collected sera. Specific IgG were detected by using anti-mouse IgG-HRP Conjugate (Zymed, 61-6520). Titers were calculated by interpolating the O.D.max/2 on the x axis and the equation used was $y=ax+b$ from the straight line passing by the two points surrounding the O.D.max/2. Titers were expressed as serum dilutions.

IFN α Neutralizing Capacity Bioassay

The collected sera capacity to neutralize antiviral activity of IFN α was assessed by using the L929 cells with Encephalomyocarditis



virus (EMCV) cytopathic effect inhibition assay, as previously described (19). Briefly, sera were tested in duplicate, diluted in assay medium (RPMI supplemented with 4% FBS 2 mM glutamine, 1 mM Sodium Pyruvate, and 1 mM HEPES) and filtered on spinX 0.22 μ m tubes. Two-fold serum sample dilution or anti-murine IFN α polyclonal antibody (PBL Assay Science, Piscataway, NJ) as a positive control, and murine IFN α 3 (10 U/well) were added to the wells and incubated for 1 h at room temperature. Then, 20,000 L929 cells were added per well and plates were incubated for 18h at 37°C and 5% CO₂ in a humidified incubator. After medium removal and PBS washes, the cell culture was infected with EMCV (diluted to 100 Cell Culture Infectious Dose 50). Plates were incubated for 48h until cell survival determination with a 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium/phenazine methosulfate (MTS/PMS) solution dye (Promega; Madison, Wisconsin, USA) according to the manufacturer's instructions. The 50% neutralization capacity (NC50) value (i.e. serum dilution resulting in 50% neutralization of virus-mediated cell mortality) was assessed.

Anti-IFN α B-Cell Response Assay

Mouse IgG ELISpot^{BASIC} (Mabtech, France) was used for the quantification of anti-IFN α -IgG-producing cells among splenocytes (300 000 cells/well), under different stimulation conditions (no stimulation, Concanavalin A, KLH, IFN α or IFN-K, at a final concentration of 10 μ g/ml), tested in duplicates. The assays were performed according to the manufacturer's protocol and an automated reader (BIOSYS, Germany) was used.

SjS-Related Autoantibodies

Mice sera collected at -11 (1.5 months of age) and 122 days (6 months of age) were used for ELISA detection of autoantibodies and tested in duplicate. Briefly, precoated ELISA plates with murine Ro52 or Ro60 were incubated with 100-fold diluted sera. Specific IgG were detected with HRP-conjugated anti-mouse IgG and binding quantified using 3,3',5,5'-tetramethylbiphenyl-4,4'-diamine (TMB) substrate by absorbance measurement at 490 nm. Threshold for positivity was determined as the mean titer + 5 standard deviations in 6-month-old C57Bl/6 mice (n=5).

Salivary Flow Rate Measurement

Mice were anesthetized (i.p. injection of a 60mg/ml ketamine and 8mg/ml xylazine solution), and their stimulated (s.c. pilocarpine hydrochloride at 0.5 μ g/g of body weight; Santa Cruz Biotechnologies, Dallas, USA) 10-minute-Salivary Flow Rates (SFR) were measured. During anesthesia, saliva was collected continuously for 10 min with a hematocrit capillary tube (Siemens Healthcare Diagnostics, Tarrytown, USA) and transferred to a 1.5 ml tube. After a brief centrifugation, the volume of saliva was determined using a micropipette and standardized against the weight (in grams) of the individual mouse.

Phenol Red Thread Test

Following saliva collection, phenol red threads (Danyang Nuorishi Optical Co., Jiangsu, China) were used to measure tear production in mice, as previously described (31). Briefly, the bent end of a thread was carefully placed at the intercanthus of one eye and held in place with sterile forceps for 20 seconds. On removal of the thread, the length of the red area was measured (in millimeters), using the scale provided, and recorded.

Von Frey Test

Tactile sensitivity was evaluated using the von Frey test (32). The principle of this test is standardized. Once the mouse is calm and motionless, a hind paw is touched with the tip of a flexible fiber of given length and diameter (Bioseb, Vitrolles, France). The force of application increases as long as the investigator is pushing the probe and until the fiber bends. This principle makes it possible for the investigator to apply a reproducible force to the skin surface. The scale of force used ranged from 0.008 to 1 g. The force required to induce a withdrawal paw movement was recorded.

Cold Plate Assay

Cold hypoalgesia was evaluated during 5 min using a cold plate set at $2 \pm 0.2^\circ\text{C}$ (Bioseb, Vitrolles, France) as previously described (33). The total number of brisk lifts of either hind paw was counted, and two observers were tasked with counting to ensure accuracy. Normal locomotion was distinct, involving coordinated movements of all four limbs that were excluded. A maximal cut-off time of 5 min was used to prevent tissue damage.

Open Field Test

To evaluate anxiety-like behavior (34), mice were placed in a field of 40 x 20 cm (surrounded by walls to prevent escape) including a new object in the center, for 10 min. The amount of time the mouse spent in the arena center (15 x 7.5 cm around the new object, i.e. the "exposed field") and in the arena periphery (the rest of the field, i.e. the "protected field") were measured using an automated video-tracking device (Smart, Panlab Harvard Apparatus, Barcelona, Spain). Data recorded during the last 5 min were analyzed.

Forced-Swim Test

To evaluate depressive-like behavior (34), mice were placed in Plexiglas cylinders (20 cm high, 15 cm in diameter) filled with 35°C water to a depth of 15 cm (preventing mice from touching

the bottom of the cylinder). During 6 min, the duration of immobility (i.e. the time during which mice made only the smallest movements necessary to keep their heads above water) was scored using an automated video-tracking device and the last 4-minute-data were analyzed.

Sucrose Consumption Test

Because reduced sensitivity to palatable stimulus is considered to reflect anhedonia in mice (34), their responsiveness to a 4% sucrose solution was evaluated. Three days prior to the beginning of the test, mice were placed in individual cages. Water (1st phase of the test), then 4% sucrose solution (2nd phase) intakes were measured (by reweighting pre-weighted bottles) over two 48h-periods, and average sucrose solution/water intake ratios were analyzed.

Lymph Nodes Measurement

To evaluate their level of lymphoproliferation, mice's main lymph nodes (cervical, axillary, brachial and inguinal areas) were measured before sacrifice. A semi-quantitative gradation was performed (0, not palpable; 1, ≤ 5 mm; 2, > 5 mm).

Dipstick Test for Proteinuria

Mice were observed for proteinuria at weeks 0, 3, 6, 9, 12 and 16 from first immunization, by using urine dipstick tests (Siemens Healthcare Diagnostics, Tarrytown, USA) which allowed a semi-quantitative gradation, according to the manufacturer's instructions (0, negative to traces; 1: 0.3 g/l; 2: 1 g/l; 3: ≥ 3 g/l).

Murine Sjögren's Syndrome Disease Activity Index (MuSSDAI)

Modelled on the EULAR Sjögren's Syndrome Disease Activity Index (35) (ESSDAI), MuSSDAI was designed as a global activity score using all the applicable features in MRL/lpr mice. The weight of each organ-specific domain in MuSSDAI was chosen to be as close as possible to the human index.

Lymphadenopathy (weight = 4) and renal (weight = 5) domains' scores respectively ranged from 0 to 2 and 0 to 3, using the aforementioned semi-quantitative gradation.

For the other selected domains (glandular, peripheral nervous system [PNS] and central nervous system [CNS]), the results of their respective tests were converted into a semi-quantitative gradation system ranging from 0 (no activity) to 3 (high activity). The limits of each activity class were determined, using the tests' results in all 52 premorbid MRL/lpr mice, as follows: no activity (=0) when the result was included in the 95% confidence interval (CI95) of the mean; low activity (=1) when the result was between the upper (length of the fibers [Von Frey] and % of immobility [forced-swim test]) or the lower (SFR, phenol red thread test, cold plate assay, % in the center zone [open field test] and sucrose solution/water intake ratio) limit of the CI95 and the same limit $\pm 15\%$ of the mean; moderate activity (=2) when the result was between the upper/lower limit of the CI95 $\pm 15\%$ of the mean and the same limit $\pm 30\%$ of the mean; high activity (=3) when the result was beyond the upper/lower limit of the CI95 $\pm 30\%$ of the mean.

Glandular domain's weight (=3) was equally divided between salivary and lacrimal functions assessments. Similarly, PNS (=5)

and CNS (=6) domains' weights were respectively and equally divided between mechanical and thermal nociception evaluations, and between depressive-like behavior, anxiety-like behavior and anhedonia evaluations.

MuSSDAI was calculated for each mouse as the sum of every organ-specific domain's score multiplied by its own weight and ranged from 0 to 65.

Histology

Submandibular glands (SMG), lacrimal glands (LG), lungs and kidneys harvested at the time of sacrifice were fixed in 10% neutral buffered formalin and embedded in paraffin wax. Kidneys (4 μm sections) were stained with periodic acid/Schiff light green. SMG, LG and lungs (6 μm sections) were stained with hematoxylin/eosin. For each sample, 2 to 6 sections of both (left and right) organs were blindly examined.

For SMG and LG, the focus score (36), based on enumeration of the number of foci (i.e. infiltrates of at least 50 mononuclear cells/ mm^2), was determined. For lungs, perivascular (medium to large vessels surrounded by inflammatory cells/total number of medium to large sized vessels) and peribronchiolar (medium to large bronchioles surrounded by inflammatory cells/total number of medium to large bronchioles) infiltrates were scored (37).

Immunohistochemistry

Tissue sections of formalin-fixed, paraffin-embedded SMG were immersed in a Proteinase K (Euromedex, France) antigen retrieval solution at 5 $\mu\text{g}/\text{ml}$ for 10 min at 37°C. Endogenous peroxidase activity was quenched with 3% H_2O_2 in TB. Non-specific binding was blocked using 1% Bovine Serum Albumin (BSA) in PBS for 1 hour 40 min at 4°C. Tissues were first incubated with rabbit anti-mouse BAFF (ALX-210-799, Enzo Life Sciences, France) at 10 $\mu\text{g}/\text{ml}$ or rabbit isotype control (1:200) antibodies overnight at 4°C; they were then stained with HRP-conjugated goat anti-rabbit secondary antibody (Abcam, Cambridge, UK) for 30 min at room temperature. Staining was developed using 3,3'-diaminobenzidine (DAB) chromogen. Tissue sections were counterstained with hematoxylin. For quantification of local epithelial production of BAFF in SMG, positive zones (brown spots) were blindly counted out of 3 non-overlapping microscope fields (x40) per sample (from IFN-K and KLH groups; n=5 per group).

Interferon-Inducible Genes Expression Profiling

RNeasy Protect Mini Kit (Qiagen, Maryland, USA) was used for RNA extraction of mandibular lymph nodes, according to the manufacturer's instructions. Quality was assessed using an Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, USA).

The Mouse Type 1 Interferon Response RT² Profiler PCR Array, RT² First Strand Kit and RT² Real-Time SYBR Green PCR Master Mix (Qiagen, Maryland, USA) were used according to the manufacturer's instructions and using the data analysis web portal at <http://www.qiagen.com/geneglobe>. Quantitative real-

time PCR array was performed (using 0.5 μg of extracted RNA/mouse) on the ABI 7500 Real-Time PCR system (Applied Biosystems, Foster City, USA). Samples were assigned to Controls (NT, PBS and KLH mice) and test (IFN-K-treated mice) groups. CT values were normalized based on IFNA4 expression (automatically selected from the full panel because of its stable expression). Fold change/regulation was calculated using $\Delta\Delta\text{CT}$ method (38), with a cut off of 2.

Statistics

Means (\pm standard error of the mean) of MuSSDAI, SFR, phenol red thread, Von Frey, cold plate, sucrose consumption, forced-swim and open field tests, as well as histological scoring were analyzed between control (NT, PBS and KLH groups) and treated (IFN-K) mice by the Mann-Whitney U test. These parameters and cells' response assays results were also analyzed by the Kruskal-Wallis test, mainly to assess the lack of differences between the 3 control groups, followed by Dunn tests to compare individually the treated groups with their respective control groups. Survival and proteinuria data were analyzed using Kaplan-Meier graphs and log-rank tests. For IFN-inducible genes expression, the p values were calculated based on a Student's t-test of the replicate $2^{(-\Delta\text{CT})}$ values for each gene in the Control and IFN-K-treated groups.

The data analyses were performed using GraphPad PRISM 5.0 (Intuitive Software for Science, USA).

RESULTS

Immunogenicity and Neutralization

Immunization with IFN-K adjuvanted with SWE01 or ISA51 (Figures 1B, C) efficiently elicited high titers of anti-IFN α Abs in all treated mice by day+38 post immunization (respectively 1/27172 \pm 1/2856 and 1/16768 \pm 1/3861, p=0.0973) with persistent titers at day+66 (1/25631 \pm 1/2674 and 1/14137 \pm 1/3726, p=0.0728) day+94 (1/25386 \pm 1/3718 and 1/8068 \pm 1/2197, p=0.0047) and day+122 (1/16313 \pm 1/2304 and 1/9654 \pm 1/2963, p=0.1014). KLH/ISA51-treated mice belatedly developed low titers of anti-IFN α Abs starting at day+94 (1/1682 \pm 1/492) and still present at day+122 (1/1643 \pm 1/308), without detectible neutralizing properties.

Neutralizing anti-IFN α Abs were only detected in IFN-K-treated mice, with higher neutralizing capacities in the IFN-K/SWE01 group than in the IFN-K/ISA51 group (Figure 1D), at all time points: day+38 (1/287 \pm 1/61 versus 1/56 \pm 1/36, p=0.0095), day+66 (1/2226 \pm 1/799 versus 1/147 \pm 1/77, p=0.0045), day+94 (1/2168 \pm 1/663 versus 1/151 \pm 1/59, p=0.0079) and day+122 (1/2363 \pm 1/548 versus 1/95 \pm 1/44, p=0.0032).

The quantification of anti-IFN α -IgG-secreting splenocytes (see Supplementary Figure S1 for the detailed analysis) was superior in IFN-K/SWE01-treated mice (Figure 1E) than in their controls (183 spots \pm 11 versus 31 spots \pm 4, p=0.0079), regardless of the restimulation conditions. The difference was not significant between IFN-K/ISA51-treated mice (Figure 1F) and their controls (32 spots \pm 3 versus 26 spots \pm 1, p=0.1508).

Only mice that received KLH (i.e. KLH and IFN-K groups) developed anti-KLH Abs (see **Supplementary Figure S2**)

Type 1 IFN Signature

Type 1 IFN-inducible genes expression (**Figure 2** and see **Supplementary Figure S3** for a more detailed analysis) was reduced in IFN-K-treated mice's mandibular lymph nodes when compared to their controls, with a superior effect with SWE01. Seventy-three transcripts were downregulated (with \log_2 [fold change] ranging from -6.9 to -2.11) in the IFN-K/SWE01 group, and 10 (with \log_2 [fold change] ranging from -3.02 to -2.01) in the IFN-K/ISA51 group. Among the more downregulated transcripts in the IFN-K/SWE01 group were classical IFN-inducible genes: *Mx1* (-6.9), *Cxcl-10* (-6.62), *Isg15* (-6.27), *Ccl2* (-6.02), *Stat1* (-5.57), *Oas1a* (-5.54), or *Tlr7* (-5.4). Among these transcripts, *Cxcl-10*, *Isg15*, *Ccl2*, and *Oas1a* were downregulated in both IFN-K-treated groups.

Global Disease Activity

The Murine Sjögren's Syndrome Disease Activity Index (MuSSDAI), a systemic disease activity score we modelled on the EULAR Sjögren's Syndrome Disease Activity Index (ESSDAI) (**Supplementary Table S1**), was statistically improved in IFN-K/SWE01 (24.8 ± 2.3 versus 37.8 ± 3.5 , $p=0.0059$) and IFN-K/ISA51-treated (22.1 ± 2.6 versus 36.2 ± 2.2 , $p=0.0012$) mice when compared to their respective controls (**Figures 3A, B**).

Glandular Features


IFN-K improved saliva and tear productions in MRL/lpr mice, when compared to control groups. In SWE01-adjuvanted groups, the SFR (**Figure 3G**) were 3.4 ± 0.8 ml/g versus $2.0 \pm$


0.7 ml/g ($p=0.0026$) and 4.0 ± 1.2 ml/g versus 2.3 ± 0.5 ml/g ($p=0.0044$), while the phenol red thread tests (**Figure 3H**) were 3.3 ± 0.5 mm versus 1.3 ± 0.2 mm ($p=0.0017$) and 2.6 ± 0.2 mm versus 0.8 ± 0.1 mm ($p=0.0002$), at 8 and 16 weeks post-immunization, respectively. In ISA51-adjuvanted groups, the SFR (**Figure 3K**) were 4.0 ± 0.4 ml/g versus 2.1 ± 0.2 ml/g ($p=0.0012$) and 3.1 ± 0.3 ml/g versus 1.3 ± 0.2 ml/g ($p=0.0004$), and the phenol red thread tests (**Figure 3L**) were 2.9 ± 0.8 mm versus 1.3 ± 0.2 mm ($p=0.0027$) and 2.3 ± 0.2 mm versus 0.9 ± 0.1 mm ($p=0.0001$) at the same time points.

At the histological level, the focus scores were improved in SMG (0.22 focus/mm $^2 \pm 0.07$ versus 0.85 focus/mm $^2 \pm 0.12$, $p=0.0010$) and LG (0.44 focus/mm $^2 \pm 0.17$ versus 2.90 focus/mm $^2 \pm 0.23$, $p=0.0006$) of IFN-K/SWE01-treated mice (**Figures 3C, D**), when compared to SWE01-adjuvanted controls. The same beneficial effect was observed in IFN-K/ISA51-treated mice's SMG (0.28 focus/mm $^2 \pm 0.10$ versus 0.75 focus/mm $^2 \pm 0.07$, $p=0.0068$), but in their LG (2.08 focus/mm $^2 \pm 0.35$ versus 3.15 focus/mm $^2 \pm 0.38$, $p=0.0602$) the downward trends did not reach statistical significance (**Figures 3E, F**). Representative microphotographs are shown in **Figures 4** for SMG, and **Figure 5** for LG. Accordingly, immunocytochemistry experiments showed a significant reduction in immunostaining for BAFF in SMG from IFN-K-treated mice when compared to SMG from KLH-treated mice (**Figure 6**) in SWE01 (18.5 spots ± 1.6 versus 27.3 spots ± 1.3 , $p=0.0003$) as was as in ISA51 (13.7 spots ± 1.8 versus 21.9 spots ± 2.1 , $p=0.0099$) cohorts (**Figure 4**).

Extra-Glandular Features

Mechanically-induced nociception was not altered in IFN-K-treated mice when compared to mice from the control groups





whose hypoesthesia significantly worsened with aging. In SWE01-adjuvanted groups (Figure 3I), the threshold of withdrawal of the touched paw was lower in IFN-K-treated mice than in their controls, at 8 weeks (0.03 ± 0.01 g versus 0.15 ± 0.03 g, $p=0.0007$) and 16 weeks (0.19 ± 0.06 g versus 0.89 ± 0.12 g, $p=0.0005$) post-immunization. In ISA51-adjuvanted groups (Figure 3M), the threshold of withdrawal of the touched paw was also lower in IFN-K-treated mice than in their controls, at 8 weeks (0.04 ± 0.01 g versus 0.22 ± 0.04 g, $p=0.0004$) and 16 weeks (0.11 ± 0.05 g versus 0.55 ± 0.06 g, $p=0.0006$) post-immunization.

As for thermally-induced nociception (Figures 3J, N), earlier effects were observed in IFN-K/SWE01-treated mice (with a higher mean number of brisk lifts of either hind paw), starting at 8 weeks (8.71 ± 0.68 paw lifts versus 3.74 ± 0.52 paw lifts, $p=0.0005$) post-immunization and maintained at 16 weeks (9.14 ± 1.14 paw lifts versus 4.43 ± 0.48 paw lifts, $p=0.0016$), when beneficial effects were also significant in IFN-K/ISA51-treated mice (6.00 ± 0.98 paw lifts versus 2.32 ± 0.30 paw lifts, $p=0.0021$). No difference was observed between IFN-K/ISA51-

treated mice and their controls at 8 weeks post-immunization (5.43 ± 0.95 paw lifts versus 4.32 ± 0.47 paw lifts, $p=0.41$).

Neuropsychiatric tests and lung infiltrated were not affected by IFN-K (Figure 7).

Proteinuria-free survival (Supplementary Figure S4) was not statistically different between groups.

Anti-Ro60 and anti-Ro52 autoantibodies (Supplementary Figure S5) were respectively positive in 44.4% and 68.9% of the 1.5-month-old mice, and 100% of the 6-month-old mice, but they were not statistically different between groups.

DISCUSSION

IFN-K induced high levels of anti-IFN α Abs, regardless of its adjuvantation protocol, with higher trends in IFN-K/SWE01-treated mice. However, the difference was only significant at one time point. In both IFN-K-vaccinated groups, both glandular and PNS manifestations of the disease were significantly improved. These improvements were concordant at the




FIGURE 4 | Salivary histopathological features. Representative low (left; x10) and high (right; x40) magnification images of hematoxylin and eosin-stained sections of the submandibular glands harvested at the end of follow-up in SWE01/KLH (n=5), SWE01/IFN-K (n=6), ISA51/KLH (n=7) and ISA51/IFN-K (n=7) living mice, are shown.

histological level, as shown by the reduced inflammatory cell infiltrates in the glands of IFN-K-treated mice. In IFN-K/ISA51-treated mice, a similar beneficial effect was observed, but did not reach statistical significance in lacrimal glands, which is although not incompatible with a normalized tear production (39).

Regarding the anti-IFN α neutralizing capacity, there was a clear superiority in IFN-K/SWE01-treated mice, with concordant results in the large downregulation of type 1 IFN signature and the quantification of anti-IFN α -IgG-secreting splenocytes. As a MF59-like adjuvant, SWE01's superiority may be due to its known positive effect on T follicular helper cells' (TFH) response and thus the magnitude of the germinal center B cell response (40). MF59 has previously been shown to be able to induce a greater adjuvant efficacy (with different vaccine antigens: influenza, tetanus toxoid, hepatitis B, Group B and C Meningococcal bacteria) than any other adjuvant (41), including ISA51 (42).

The fact that KLH/ISA51-treated mice belatedly developed low titers of non-neutralizing-anti-IFN α Abs starting at day+94, and that all control mice had low counts of anti-IFN α -IgG-secreting splenocytes, indicate that, IFN-K induces an anti-IFN α Abs production by increasing or promoting the maturation of pre-existing autoreactive B cells as previously described in SLE patients (20, 43).

As for the other extra-glandular manifestations, the neuropsychiatric manifestations of the disease were not affected by IFN-K, but the behavioral tests evaluating these features are known to be complex with many confounding factors (44). Furthermore, proteinuria-free survival was also not different between groups. However, our trial was not specifically designed to evaluate this

aspect of the disease (because glomerulonephritis in the MRL/lpr model is more a lupus-like feature than a SjS-like one) and the occurrence of these events was lower than expected (45). Finally, about the fact that pulmonary disease was unaffected by IFN-K, we think that a non-IFN-dependent pathologic process might be mediating lung damage, as it has previously been improved in MRL/lpr mice by inhibiting Tumor Necrosis Factor- α (37). This element will have to be kept in mind during IFN-K development.

The main limitation that one could see in this study is the use of the MRL/lpr mouse model, because it is usually referred to as a secondary mouse model of SjS. That is because it presents with glomerulonephritis (which was absent in our mice) and CNS manifestations, which are rare but not absent in SjS (4). However, many authors (46–48) now criticize the primary/secondary SjS dichotomy which misrepresents the potential manifestations and risks of the syndrome. As Professor Moutsopoulos (among others), who coined the concept of secondary SjS more than 35 years ago, we think that the term “secondary”, in human and murine SjS, should be replaced by a more descriptive term (*i.e.* “SjS associated with”) or by the term “polyautoimmunity” (47, 48). Moreover, the MRL/lpr mouse model is included in all review papers about murine models of SjS, including the ones focusing on primary SjS and it has all key features of SjS: female sex predilection, decreased salivary flow rate (49) and tear production, lymphocytic infiltrates in the salivary and lacrimal glands, and anti-Ro60, anti-Ro52 and anti-La autoantibodies. On this latter point, we detected a much higher rate of anti-Ro60 and/or anti-Ro52-positive mice




FIGURE 5 | Lacrimal histopathological features. Representative low (left; x10) and high (right; x40) magnification images of hematoxylin and eosin-stained sections of the lacrimal glands harvested at the end of follow-up in SWE01/KLH (n=5), SWE01/IFN-K (n=6), ISA51/KLH (n=7) and ISA51/IFN-K (n=7) living mice, are shown.





FIGURE 6 | In-situ reduction of B cell activating factor (BAFF) production. Local expression of BAFF was analyzed by immunocytochemistry in formalin-fixed, paraffin-embedded submandibular glands (SMG) harvested at the end of follow-up in SWE01/KLH, SWE01/IFN-K, ISA51/KLH and ISA51/IFN-K mice (n=5 per group). BAFF-positive zones (brown spots) in SMG were counted out of 3 non-overlapping microscope fields (x40) per sample and compared between controls and treated mice in (A) SWE01 cohort and (B) ISA51 cohort. Representative images of BAFF-immunostained sections and (C) negative control (rabbit IgG) are shown (original magnification x40). Error bars indicate mean \pm S.E.M. **p < 0.01 and ***p < 0.001.



than what had been historically described in the MRL/lpr mouse model (6% and 30% respectively) (34, 35), using the same stringent threshold. The two main elements to explain these differences are probably the type of antigens which were used [full-length murine proteins versus partial human proteins for the publication dating from 1994 (50)] and the age of the mice which were tested (6-month-old mice in our study versus 4-5-month-old mice previously). In addition to the rare aforementioned CNS and renal manifestations these mice also develop classical systemic SjS manifestations: peripheral neuropathy (rare in SLE), arthritis, pneumonitis, cryoglobulinemia and signs of lymphoproliferation (lymphadenopathies, splenomegaly) (34).

As we noted in a recent correspondence (51), no perfect model for SjS exists, but we think that the MRL/lpr mouse model should be considered as a model of choice for preclinical trials evaluating new immunological treatments in SjS, because of its unique systemic manifestations. Another limitation could be the use of mandibular lymph nodes to evaluate type 1 IFN signature, however, even if we had to use most of SMG tissues to perform the more rigorous and accurate evaluation of the focus score, we were able to demonstrate that local epithelial production of BAFF – which is a pivotal type 1 IFN-inducible cytokine – was reduced in SMG from IFN-K-treated mice compared to their controls'. In line with our findings, other teams have previously

reported that type 1 IFNs and plasmacytoid dendritic cells were involved in the pathogenesis of the MRL/lpr mouse model (52). It is also important to note that the draining lymph nodes were previously described as important elements in SjS (53, 54).

Anti-Ro60 or anti-Ro52 titers did not differ between groups, which is consistent with the absence of fluctuation in anti-Ro titers with disease activity in human SjS (55).

As reported in previous preclinical studies (19) no signs of autoreactive anti-IFN α T cell response were detected using IFN- γ ELISpot^{BASIC} (data not shown).

We herein describe for the first time, a proof-of-concept for the efficacy of IFN-K in a mouse model of systemic SjS. Its beneficial effects on sicca syndrome and peripheral neuropathy pave the way for its clinical development in SjS, which is still facing an unmet therapeutic need. Due to its substantial advantages when compared to monoclonal Abs (absence of anti-drug Abs formation, neutralization of all IFN α subtypes (56), better predictable patient compliance because of a low number of administrations (21), lower production and health care costs...) and its good safety profile (20, 21), IFN-K is the good candidate for Sjögren's syndrome's treatment, but of course, further studies are required to confirm this proof-of-concept.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**. Further inquiries can be directed to the corresponding author.

ETHICS STATEMENT

The animal study was reviewed and approved by PLEXAN.

REFERENCES

- Qin B, Wang J, Yang Z, Yang M, Ma N, Huang F, et al. Epidemiology of Primary Sjögren's Syndrome: A Systematic Review and Meta-Analysis. *Ann Rheumatol Dis* (2015) 74:1983–9. doi: 10.1136/annrheumdis-2014-205375
- Kyriakidis NC, Kapsogeorgou EK, Tzioufas AG. A Comprehensive Review of Autoantibodies in Primary Sjögren's Syndrome: Clinical Phenotypes and Regulatory Mechanisms. *J Autoimmun* (2014) 51:67–74. doi: 10.1016/j.jaut.2013.11.001
- Ramos-Casals M, Brito-Zerón P, Sisó-Almirall A, Bosch X. Primary Sjogren Syndrome. *BMJ* (2012) 344:e3821. doi: 10.1136/bmj.e3821
- Ramos-Casals M, Brito-Zerón P, Solans R, Camps M-T, Casanovas A, Sopena B, et al. Systemic Involvement in Primary Sjögren's Syndrome Evaluated by the EULAR-SS Disease Activity Index: Analysis of 921 Spanish Patients (GEAS-SS Registry). *Rheumatology* (2014) 53:321–31. doi: 10.1093/rheumatology/ket349
- Brito-Zerón P, Kostov B, Solans R, Fraile G, Suárez-Cuervo C, Casanovas A, et al. Systemic Activity and Mortality in Primary Sjögren Syndrome: Predicting Survival Using the EULAR-SS Disease Activity Index (ESSDAI) in 1045 Patients. *Ann Rheumatol Dis* (2014) 1–8. doi: 10.1136/annrheumdis-2014-206418
- Saroux A, Pers J-O, Devauchelle-Pensec V. Treatment of Primary Sjögren Syndrome. *Nat Rev Rheumatol* (2016) 12:456–71. doi: 10.1038/nrrheum.2016.100
- Seror R, Theander E, Bootsma H, Bowman SJ, Tzioufas A, Gottenberg J-E, et al. Outcome Measures for Primary Sjögren's Syndrome: A Comprehensive Review. *J Autoimmun* (2014) 51:51–6. doi: 10.1016/j.jaut.2013.12.010
- Maria NI, Steenwijk EC, Ijpma AS, Helden-Meeuwse CG, Vogelsang P, Beumer W, et al. Contrasting Expression Pattern of RNA-Sensing Receptors TLR7, RIG-I and MDA5 in Interferon-Positive and Interferon-Negative Patients With Primary Sjögren's Syndrome. *Ann Rheumatol Dis* (2017) 76:721–30. doi: 10.1136/annrheumdis-2016-209589
- Yao Y, Liu Z, Jallal B, Shen N, Rönnblom L. Type I Interferons in Sjögren's Syndrome. *Autoimmun Rev* (2013) 12:558–66. doi: 10.1016/j.autrev.2012.10.006
- Brkic Z, Maria NI, van Helden-Meeuwse CG, van de Merwe JP, van Daele PL, Dalm VA, et al. Prevalence of Interferon Type I Signature in CD14 Monocytes of Patients With Sjogren's Syndrome and Association With Disease Activity and BAFF Gene Expression. *Ann Rheumatol Dis* (2013) 72:728–35. doi: 10.1136/annrheumdis-2012-201381
- Nezos A, Papageorgiou A, Tassidou A, Kapsogeorgou EK, Voulgarelis M, Koutsilieris M, et al. Type I and II Interferon Signatures in Sjogren's Syndrome Pathogenesis: Contributions in Distinct Clinical Phenotypes and Sjogren's Related Lymphomagenesis. *J Autoimmun* (2015) 63:47–58. doi: 10.1016/j.jaut.2015.07.002
- Nezos A, Papageorgiou A, Fragoulis G, Ioakeimidis D, Koutsilieris M, Tzioufas AG, et al. B-Cell Activating Factor Genetic Variants in Lymphomagenesis Associated With Primary Sjogren's Syndrome. *J Autoimmun* (2014) 51:89–98. doi: 10.1016/j.jaut.2013.04.005
- Burbelo PD, Browne S, Holland SM, Iadarola MJ, Alevizos I. Clinical Features of Sjögren's Syndrome Patients With Autoantibodies Against Interferons. *Clin Transl Med* (2019) 8:1. doi: 10.1186/s40169-018-0218-1

AUTHOR CONTRIBUTIONS

MK, GG-V, FB, and SP designed the study. MK, FC, PH, CN, OC, SC, NR, BC, MT, NL and FF conducted experiments. MK acquired data. MK, FC, OC, SC, BC, MT, FF, GG-V, FB, and SP analyzed data. NL, FF, GG-V, FB, and SP provided reagents. MK wrote the manuscript. FC, OC, PC, GG-V, FB, and SP reviewed the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by the Association Française du Gougerot Sjögren (AFGS) and NEOVACS S.A. The funder was not involved in the study design, collection, analysis, interpretation of data, the writing of this article or the decision to submit it for publication.

ACKNOWLEDGMENTS

The authors thank Ms. Béatrice Drouet for her help in writing and reviewing the manuscript, and Pr. Driss Zoukhri for his technical advice. Seppic supplied ISA51 adjuvant and the University of Lausanne supplied the SWE01 adjuvant.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2021.666134/full#supplementary-material>

14. Mariette X, Seror R, Quartuccio L, Baron G, Salvin S, Fabris M, et al. Efficacy and Safety of Belimumab in Primary Sjögren's Syndrome: Results of the BELISS Open-Label Phase II Study. *Ann Rheumatol Dis* (2015) 74:526–31. doi: 10.1136/annrheumdis-2013-203991
15. Cornec D, Costa S, Devauchelle-Pensec V, Jousse-Joulin S, Marcorelles P, Berthelot J-M, et al. Blood and Salivary-Gland BAFF-Driven B-Cell Hyperactivity is Associated to Rituximab Inefficacy in Primary Sjögren's Syndrome. *J Autoimmun* (2016) 67:102–10. doi: 10.1016/j.jaut.2015.11.002
16. Verstappen GM, van Nimwegen JF, Vissink A, Kroese FGM, Bootsma H. The Value of Rituximab Treatment in Primary Sjögren's Syndrome. *Clin Immunol Orlando Fla* (2017) 182:62–71. doi: 10.1016/j.clim.2017.05.002
17. Dörner T, Posch MG, Li Y, Petricoul O, Cabanski M, Milojevic JM, et al. Treatment of Primary Sjögren's Syndrome With Ianalumab (VAY736) Targeting B Cells by BAFF Receptor Blockade Coupled With Enhanced, Antibody-Dependent Cellular Cytotoxicity. *Ann Rheumatol Dis* (2019) 78:641–7. doi: 10.1136/annrheumdis-2018-214720
18. Lee J, Lee J, Kwok S-K, Baek S, Jang SG, Hong S-M, et al. JAK-1 Inhibition Suppresses Interferon-Induced BAFF Production in Human Salivary Gland: Potential Therapeutic Strategy for Primary Sjögren's Syndrome. *Arthritis Rheumatol Hoboken NJ* (2018) 70:2057–66. doi: 10.1002/art.40589
19. Zagury D, Le Buanec H, Mathian A, Larcier P, Burnett R, Amoura Z, et al. IFN α Kinoid Vaccine-Induced Neutralizing Antibodies Prevent Clinical Manifestations in a Lupus Flare Murine Model. *Proc Natl Acad Sci U S A* (2009) 106:5294–9. doi: 10.1073/pnas.0900615106
20. Lauwerys BR, Hachulla E, Spertini F, Lazaro E, Jorgensen C, Mariette X, et al. Down-Regulation of Interferon Signature in Systemic Lupus Erythematosus Patients by Active Immunization With Interferon α -Kinoid. *Arthritis Rheumatol* (2013) 65:447–56. doi: 10.1002/art.37785
21. Ducreux J, Houssiau FA, Vandepapelière P, Jorgensen C, Lazaro E, Spertini F, et al. Interferon α Kinoid Induces Neutralizing Anti-Interferon α Antibodies That Decrease the Expression of Interferon-Induced and B Cell Activation Associated Transcripts: Analysis of Extended Follow-Up Data From the Interferon α Kinoid Phase I/II Study. *Rheumatol Oxf Engl* (2016) 55:1901–5. doi: 10.1093/rheumatology/kew262
22. Houssiau FA, Thanou A, Mazur M, Ramitterre E, Gomez Mora DA, Misterska-Skora M, et al. IFN- α Kinoid in Systemic Lupus Erythematosus: Results From a Phase IIb, Randomised, Placebo-Controlled Study. *Ann Rheumatol Dis* (2020) 79:347–55. doi: 10.1136/annrheumdis-2019-216379
23. Cornec D, Jamin C, Pers J-O. Sjögren's Syndrome: Where do We Stand, and Where Shall We Go? *J Autoimmun* (2014) 51:109–14. doi: 10.1016/j.jaut.2014.02.006
24. Lavoie TN, Lee BH, Nguyen CQ. Current Concepts: Mouse Models of Sjögren's Syndrome. *J Biomed Biotechnol* (2011) 2011:549107. doi: 10.1155/2011/549107
25. Park Y-S, Gauna AE, Cha S. Mouse Models of Primary Sjögren's Syndrome. *Curr Pharm Des* (2015) 21:2350–64. doi: 10.2174/1381612821666150316120024
26. Delaleu N, Nguyen CQ, Peck AB, Jonsson R. Sjögren's Syndrome: Studying the Disease in Mice. *Arthritis Res Ther* (2011) 13:217. doi: 10.1186/ar3313
27. Hayashi Y, Haneji N, Hamano H. Cytokine Gene Expression and Autoantibody Production in Sjögren's Syndrome of MRL/lpr Mice. *Autoimmunity* (1996) 23:269–77. doi: 10.3109/08916939608995349
28. Alunno A, Carubbi F, Bistoni O, Caterbi S, Bartoloni E, Bigerna B, et al. CD4 (-)CD8(-) T-Cells in Primary Sjögren's Syndrome: Association With the Extent of Glandular Involvement. *J Autoimmun* (2014) 51:38–43. doi: 10.1016/j.jaut.2014.01.030
29. Brandt D, Hedrich CM. Tcr $\alpha\beta$ +CD3+CD4-CD8- (Double Negative) T Cells in Autoimmunity. *Autoimmun Rev* (2018) 17:422–30. doi: 10.1016/j.autrev.2018.02.001
30. Le Buanec H, Delavallée L, Bessis N, Paturance S, Bizzini B, Gallo R, et al. TNF α Kinoid Vaccination-Induced Neutralizing Antibodies to TNF α Protect Mice From Autologous TNF α -Driven Chronic and Acute Inflammation. *Proc Natl Acad Sci U S A* (2006) 103:19442–7. doi: 10.1073/pnas.0604827103
31. Doyle ME, Boggs L, Attia R, Cooper LR, Saban DR, Nguyen CQ, et al. Autoimmune Dacryoadenitis of NOD/LtJ Mice and its Subsequent Effects on Tear Protein Composition. *Am J Pathol* (2007) 171:1224–36. doi: 10.2353/ajpath.2007.070388
32. Bradman MJG, Ferrini F, Salio C, Merighi A. Practical Mechanical Threshold Estimation in Rodents Using Von Frey Hairs/Semmes-Weinstein Monofilaments: Towards a Rational Method. *J Neurosci Methods* (2015) 255:92–103. doi: 10.1016/j.jneumeth.2015.08.010
33. Ta LE, Low PA, Windebank AJ. Mice With Cisplatin and Oxaliplatin-Induced Painful Neuropathy Develop Distinct Early Responses to Thermal Stimuli. *Mol Pain* (2009) 5:9. doi: 10.1186/1744-8069-5-9
34. Jeltsch-David H, Muller S. Neuropsychiatric Systemic Lupus Erythematosus and Cognitive Dysfunction: The MRL-Lpr Mouse Strain as a Model. *Autoimmun Rev* (2014) 13:963–73. doi: 10.1016/j.autrev.2014.08.015
35. Seror R, Ravaud P, Bowman SJ, Baron G, Tzioufas A, Theander E, et al. EULAR Sjogren's Syndrome Disease Activity Index: Development of a Consensus Systemic Disease Activity Index for Primary Sjogren's Syndrome. *Ann Rheumatol Dis* (2010) 69:1103–9. doi: 10.1136/ard.2009.110619
36. Jonsson R, Tarkowski A, Bäckman K, Holmdahl R, Klareskog L. Sialadenitis in the MRL-L Mouse: Morphological and Immunohistochemical Characterization of Resident and Infiltrating Cells. *Immunology* (1987) 60:611–6.
37. Kim N, Ussin L, Cheng X, Murali R, Sullivan KE. TNF α Inhibition in MRL/lpr Mice Ameliorates Pulmonary But Not Renal Disease. *J Autoimmun* (2002) 19:215–22. doi: 10.1006/jaut.2002.0617
38. Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2⁻(Delta Delta C(T)) Method. *Methods San Diego Calif* (2001) 25:402–8. doi: 10.1006/meth.2001.1262
39. Kosenda K, Ichii O, Otsuka S, Hashimoto Y, Kon Y. BXSJ/MpJ-Yaa Mice Develop Autoimmune Dacryoadenitis With the Appearance of Inflammatory Cell Marker Messenger RNAs in the Lacrimal Fluid. *Clin Experiment Ophthalmol* (2013) 41:788–97. doi: 10.1111/ceo.12083
40. Mastelic Gavillet B, Eberhardt CS, Auderset F, Castellino F, Seubert A, Tregoning JS, et al. MF59 Mediates Its B Cell Adjuvanticity by Promoting T Follicular Helper Cells and Thus Germinal Center Responses in Adult and Early Life. *J Immunol Baltim Md 1950* (2015) 194:4836–45. doi: 10.4049/jimmunol.1402071
41. Ko E-J, Kang S-M. Immunology and Efficacy of MF59-Adjuvanted Vaccines. *Hum Vaccines Immunother* (2018) 14:3041–5. doi: 10.1080/21645515.2018.1495301
42. Zhang N, Channappanavar R, Ma C, Wang L, Tang J, Garron T, et al. Identification of an Ideal Adjuvant for Receptor-Binding Domain-Based Subunit Vaccines Against Middle East Respiratory Syndrome Coronavirus. *Cell Mol Immunol* (2015) 13:180–90. doi: 10.1038/cmi.2015.03
43. Gupta S, Tatouli IP, Rosen LB, Hasni S, Alevizos I, Manna ZG, et al. Distinct Functions of Autoantibodies Against Interferon in Systemic Lupus Erythematosus: A Comprehensive Analysis of Anticytokine Autoantibodies in Common Rheumatic Diseases. *Arthritis Rheumatol Hoboken NJ* (2016) 68:1677–87. doi: 10.1002/art.39607
44. Hänell A, Marklund N. Structured Evaluation of Rodent Behavioral Tests Used in Drug Discovery Research. *Front Behav Neurosci* (2014) 8. doi: 10.3389/fnbeh.2014.00252
45. Andrews BS, Eisenberg RA, Theofilopoulos AN, Izui S, Wilson CB, McConahey PJ, et al. Spontaneous Murine Lupus-Like Syndromes. Clinical and Immunopathological Manifestations in Several Strains. *J Exp Med* (1978) 148:1198–215. doi: 10.1084/jem.148.5.1198
46. Mavragani CP, Moutsopoulos HM. Primary Versus Secondary Sjögren Syndrome: Is It Time To Reconsider These Terms? *J Rheumatol* (2019) 46:665–6. doi: 10.3899/jrheum.180392
47. Amador-Patarroyo MJ, Arbelazay JG, Mantilla RD, Rodriguez-Rodriguez A, Cárdenas-Roldán J, Pineda-Tamayo R, et al. Sjögren's Syndrome at the Crossroad of Polyautoimmunity. *J Autoimmun* (2012) 39:199–205. doi: 10.1016/j.jaut.2012.05.008
48. Rojas-Villarraga A, Amaya-Amaya J, Rodriguez-Rodriguez A, Mantilla RD, Anaya J-M. Introducing Polyautoimmunity: Secondary Autoimmune Diseases No Longer Exist. *Autoimmune Dis* (2012) 2012:254319. doi: 10.1155/2012/254319
49. Sato M, Kuroda S, Mansjur KQ, Khaliunaa G, Nagata K, Horiuchi S, et al. Low-Intensity Pulsed Ultrasound Rescues Insufficient Salivary Secretion in Autoimmune Sialadenitis. *Arthritis Res Ther* (2015) 17:278. doi: 10.1186/s13075-015-0798-8

50. Wahren M, Skarstein K, Blange I, Pettersson I, Jonsson R. MRL/lpr Mice Produce Anti-Ro 52,000 MW Antibodies: Detection, Analysis of Specificity and Site of Production. *Immunology* (1994) 83:9–15.
51. Killian M, Batteux F, Paul S. The MRL/lpr Mouse Model: An Important Animal Model for Systemic Sjögren Syndrome and Polyautoimmunity. *J Rheumatol* (2019) 47(1):157. doi: 10.3899/jrheum.190820
52. Zhou Z, Ma J, Xiao C, Han X, Qiu R, Wang Y, et al. Phenotypic and Functional Alterations of pDCs in Lupus-Prone Mice. *Sci Rep* (2016) 6:20373. doi: 10.1038/srep20373
53. Saito M, Otsuka K, Ushio A, Yamada A, Arakaki R, Kudo Y, et al. Unique Phenotypes and Functions of Follicular Helper T Cells and Regulatory T Cells in Sjögren's Syndrome. *Curr Rheumatol Rev* (2018) 14:239–45. doi: 10.2174/1573397113666170125122858
54. Puñet-Ortiz J, Sáez Moya M, Cuenca M, Caleiras E, Lazaro A, Engel P. Ly9 (CD229) Antibody Targeting Depletes Marginal Zone and Germinal Center B Cells in Lymphoid Tissues and Reduces Salivary Gland Inflammation in a Mouse Model of Sjögren's Syndrome. *Front Immunol* (2018) 9. doi: 10.3389/fimmu.2018.02661
55. Praprotnik S, Bozic B, Kveder T, Rozman B. Fluctuation of Anti-Ro/SS-A Antibody Levels in Patients With Systemic Lupus Erythematosus and Sjögren's Syndrome: A Prospective Study. *Clin Exp Rheumatol* (1999) 17:63–8.
56. Mathian A, Amoura Z, Adam E, Colaone F, Hoekman MFM, Dhellin O, et al. Active Immunisation of Human Interferon α Transgenic Mice With a Human Interferon α Kinoid Induces Antibodies That Neutralise Interferon α in Sera

From Patients With Systemic Lupus Erythematosus. *Ann Rheumatol Dis* (2011) 70:1138–43. doi: 10.1136/ard.2010.141101

Conflict of Interest: Authors FC, PH and GG-V were employed by company NEOVACS S.A.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Killian, Colaone, Haumont, Nicco, Cerles, Chouzenoux, Cathébras, Rochereau, Chanut, Thomas, Laroche, Forest, Grouard-Vogel, Batteux and Paul. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.