



Corrigendum: High-Dose Intravenous Immunoglobulin in Severe Coronavirus Disease 2019: A Multicenter Retrospective Study in China

Wei Cao^{1†}, Xiaosheng Liu^{1,2,3†}, Ke Hong⁴, Zhiyong Ma⁵, Yuelun Zhang⁶, Ling Lin¹, Yang Han¹, Yong Xiong⁵, Zhengyin Liu¹, Lianguo Ruan⁴ and Taisheng Li^{1,2,3*}

¹ Department of Infectious Diseases, Peking Union Medical College Hospital, Peking Union Medical College and Chinese Academy of Medical Sciences, Beijing, China, ² Department of Basic Medical Sciences, School of Medicine, Tsinghua University, Beijing, China, ³ Tsinghua-Peking Center for Life Sciences, Beijing, China, ⁴ Department of Infectious Diseases, JinYin-tan Hospital, Wuhan, China, ⁵ Department of Infectious Diseases, Zhongnan Hospital of Wuhan University, Wuhan, China, ⁶ Medial Research Center, Peking Union Medical College Hospital, Peking Union Medical College and Chinese Academy of Medical Sciences, Beijing, China

OPEN ACCESS

Edited and reviewed by
Christoph T. Berger,
University of Basel, Switzerland

***Correspondence:**
Taisheng Li
lits@263.net

[†]These authors have contributed equally to this work

Specialty section:
This article was submitted to
Vaccines and Molecular Therapeutics,
a section of the journal
Frontiers in Immunology

Received: 23 February 2021
Accepted: 04 March 2021
Published: 22 March 2021

Citation:

Cao W, Liu X, Hong K, Ma Z, Zhang Y, Lin L, Han Y, Xiong Y, Liu Z, Ruan L and Li T (2021) Corrigendum: High-Dose Intravenous Immunoglobulin in Severe Coronavirus Disease 2019: A Multicenter Retrospective Study in China. *Front. Immunol.* 12:671443. doi: 10.3389/fimmu.2021.671443

Keywords: COVID-19, high-dose intravenous immunoglobulin, immunomodulation, 28-day mortality, inflammatory markers

A Corrigendum on

High-Dose Intravenous Immunoglobulin in Severe Coronavirus Disease 2019: A Multicenter Retrospective Study in China

By Cao W, Liu XS, Hong K, Ma ZY, Zhang YL, Lin L, et al. (2021). *Front. Immunol.* doi: 10.3389/fimmu.2021.627844

ERROR IN TABLE

In the original article, there was a mistake in **Table 1** as published. The author identified the calculating errors in the percentage of comorbidity, symptoms, and treatment. The corrected **Table 1** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2021 Cao, Liu, Hong, Ma, Zhang, Lin, Han, Xiong, Liu, Ruan and Li. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

TABLE 1 | Clinical and relevant baseline characteristics of patients.

Characteristic	Total (n = 115)	IVIg group (n = 26)	Control group (n = 89)	SD (%)
Demographics				
Age, median (IQR)	59 (47–69)	58 (42–65)	59 (48–70)	16
Male sex, No. (%)	76 (66)	19 (73)	57 (64)	18
Any comorbidity, No. (%)				
Hypertension	52 (45)	10 (38)	42 (47)	9
Diabetes	24 (21)	2 (8)	22 (25)	39
Chronic cardiac disease	18 (16)	5 (19)	13 (15)	11
Chronic respiratory disease	11 (10)	2 (8)	9 (10)	4
Any symptoms, No. (%)				
Fever	111 (97)	22 (85)	89 (100)	31
Cough	97 (84)	22 (85)	75 (84)	8
Dyspnea	96 (83)	22 (85)	74 (83)	10
Fatigue or myalgia	52 (45)	9 (35)	43 (48)	19
Diarrhea	30 (26)	4 (15)	26 (29)	24
Vital signs				
Systolic BP, mm Hg	132 ± 19	131 ± 14	133 ± 20	5
HR./min	88 (80–102)	85 (81–100)	89 (80–105)	9
RR, breaths/min	22 (20–25)	22 (20–23)	22 (20–25)	2
SPO ₂ , %	90 (87–94)	91 (89–94)	89 (86–92)	6
Complete blood count				
WBCs, × 10 ⁹ /L	6.59 (4.38–9.64)	6.25 (4.15–10.28)	6.60 (4.42–9.56)	4
LYM, × 10 ⁹ /L	0.76 (0.57–1.02)	0.90 (0.62–1.05)	0.73 (0.56–0.91)	19
PLT, × 10 ⁹ /L	191 (148–269)	172 (144–279)	199 (150–270)	18
Inflammatory biomarkers				
IL-6, pg/mL	13.0 (8.0–31.5)	15.5 (10.5–34.0)	11.0 (7.2–32.2)	15
IL-8, pg/mL	15.2 (6.0–27.3)	16.2 (5.8–23.8)	15.2 (5.9–28.4)	3
IL-10, pg/mL	5.2 (5.0–10.3)	5.0 (5.0–8.5)	5.5 (5.0–11.2)	2
hsCRP, mg/mL	48 (17–94)	28 (7–91)	45 (17–101)	14
Ferritin, ng/mL	807.0 (473.4–1383.3)	774.1 (444.2–1525.4)	838.8 (501.5–1351.1)	18
ESR, mm/h	50 (25–65)	35 (19–67)	50 (27–65)	14
Immunoglobulins				
IgA, g/L	2.2 (1.8–3.0)	2.7 (1.9–4.6)	2.2 (1.6–3.0)	2
IgG, g/L	11.1 (9.0–13.1)	11.8 (8.5–15.7)	11.1 (9.3–13.1)	2
IgM g/L	1.0 (0.7–1.2)	1.0 (0.9–1.1)	0.9 (0.7–1.3)	3
Seven-category scales at admission				
3, No. (%)	20 (17)	7 (27)	13 (15)	24
4, No. (%)	79 (69)	17 (65)	62 (70)	7
5, No. (%)	11 (10)	1 (4)	10 (11)	25
6, No. (%)	5 (4)	1 (4)	4 (4)	3
Murray lung injury scores at admission				
3.93 ± 0.22	3.90 ± 0.28	3.94 ± 0.20	11	
Time to admission after onset, median (IQR)				
Earlier (< 7 days), No. (%)	10 (7–12)	10 (7–12)	10 (8–12)	5
Later (between 7–14 days), No. (%)	29 (25)	7 (27)	22 (25)	12
86 (75)	19 (73)	67 (75)	12	
Treatment during hospitalization, No. (%)				
Antiviral treatment	100 (87)	22 (85)	78 (88)	7
Arbidol	75 (65)	15 (58)	60 (67)	16
IFN-α	34 (30)	14 (54)	20 (22)	53
LPV/r	31 (27)	9 (35)	22 (25)	17
RBV	12 (10)	3 (12)	9 (10)	4
OSV	33 (29)	8 (31)	25 (28)	5
Antibiotic treatment	100 (87)	25 (96)	75 (84)	37
Moxifloxacin	73 (63)	15 (58)	58 (65)	12
Cefoperazone and tazobactam	47 (41)	13 (50)	34 (38)	19
Antifungal treatment	11 (10)	5 (19)	6 (7)	29
TCM	58 (50)	15 (58)	43 (48)	15
LMWH	19 (16)	4 (15)	15 (17)	3
Glucocorticoids	71 (62)	18 (69)	53 (60)	17
IVIg therapy				
Initiation time, days, median (IQR)	—	2 (1–4)	—	—
Course, days, median (IQR)	—	5 (5–9)	—	—
Amount, g, median (IQR)	—	122.5 (95.0–213.8)	—	—

SD, standard deviation; Tmax, maximal temperature; BP, blood pressure; HR, heart rate; RR, respiratory rate; SPO₂, pulse oximeter O₂ saturation; WBC, white cell counts; LYM, lymphocyte counts; PLT, platelet counts; IL, interleukin; CRP, C-reactive protein; ESR, erythrocyte sedimentation rate; IFN-α, Interferon-alfa; LPV/r, lopinavir/ritonavir; RBV, ribavirin; OSV, oseltamivir; TCM, traditional Chinese medicine; LMWH, low molecular weight heparin.