



# Oral Administration of *Lactococcus lactis* Producing Interferon Type II, Enhances the Immune Response Against Bacterial Pathogens in Rainbow Trout

Alvaro Santibañez<sup>1,2</sup>, Diego Paine<sup>1,2</sup>, Mick Parra<sup>1,2</sup>, Carlos Muñoz<sup>1,2</sup>, Natalia Valdes<sup>1,2</sup>, Claudia Zapata<sup>1,2</sup>, Rodrigo Vargas<sup>1</sup>, Alex Gonzalez<sup>3</sup> and Mario Tello<sup>1,2,4\*</sup>

<sup>1</sup> Departamento de Biología, Laboratorio de Metagenómica Bacteriana, Facultad de Química y Biología, Universidad de Santiago de Chile, Santiago, Chile, <sup>2</sup> Consorcio Tecnológico de Sanidad Acuicola, Ictio Biotechnologies S.A., Santiago, Chile, <sup>3</sup> Laboratorio de Microbiología Ambiental y Extremófilos, Departamento de Ciencias Biológicas, Universidad de los Lagos, Osorno, Chile, <sup>4</sup> IctioBiotic SpA, Santiago, Chile

## OPEN ACCESS

### Edited by:

Peter Bossier,  
Ghent University, Belgium

### Reviewed by:

Kerry Lynne Bartie,  
University of Stirling, United Kingdom  
Patricia Díaz-Rosales,  
Centro de Investigación en Sanidad  
Animal (CISA), Spain

### \*Correspondence:

Mario Tello  
mario.tello@usach.cl

### Specialty section:

This article was submitted to  
Comparative Immunology,  
a section of the journal  
Frontiers in Immunology

**Received:** 17 April 2021

**Accepted:** 08 June 2021

**Published:** 25 June 2021

### Citation:

Santibañez A, Paine D, Parra M,  
Muñoz C, Valdes N, Zapata C,  
Vargas R, Gonzalez A and Tello M  
(2021) Oral Administration of  
*Lactococcus lactis* Producing  
Interferon Type II, Enhances the  
Immune Response Against Bacterial  
Pathogens in Rainbow Trout.  
*Front. Immunol.* 12:696803.  
doi: 10.3389/fimmu.2021.696803

Lactic acid bacteria are a powerful vehicle for releasing of cytokines and immunostimulant peptides at the gastrointestinal level after oral administration. However, its therapeutic application against pathogens that affect rainbow trout and Atlantic salmon has been little explored. Type II interferon in Atlantic salmon activates the antiviral response, protecting against viral infection, but its role against bacterial infection has not been tested *in vivo*. In this work, through the design of a recombinant lactic acid bacterium capable of producing Interferon gamma from Atlantic salmon, we explore its role against bacterial infection and the ability to stimulate systemic immune response after oral administration of the recombinant probiotic. Recombinant interferon was active *in vitro*, mainly stimulating IL-6 expression in SHK-1 cells. *In vivo*, oral administration of the recombinant probiotic produced an increase in IL-6, IFN $\gamma$  and IL-12 in the spleen and kidney, in addition to stimulating the activity of lysozyme in serum. The challenge trials indicated that the administration of the IFN $\gamma$ -producing probiotic doubled the survival in fish infected with *F. psychrophilum*. In conclusion, our results showed that the oral administration of lactic acid bacteria producing IFN $\gamma$  managed to stimulate the immune response at a systemic level, conferring protection against pathogens, showing a biotechnological potential for its application in aquaculture.

**Keywords:** *Lactococcus lactis*, interferon type II, *Flavobacterium psychrophilum*, salmonids, oral administration

## INTRODUCTION

Interferons are a group of cytokines originally discovered by its antiviral properties (1). In mammals, these cytokines can be classified in three families according their homology, receptors, structure and function (2). Interferon types I and III are multi-gene families (17 type I and four type III) strongly induced by viruses, playing a major role controlling viral infection (3–5). Interferon

type I also show a strong pro-apoptotic and anti-proliferative activity, being currently applied in anticancer and antiviral therapies (6).

In contrast, interferon type II or gamma Interferon (IFN $\gamma$ ) is a single gene encoded immunoregulatory cytokine, produced mainly by activated Natural Killer (NK) and T lymphocytes (T cells), although also can be secreted by B lymphocytes (B cells) and professional antigen-presenting cells (APCs) (7–9). Besides its antiviral properties, IFN $\gamma$  also allows the activation of macrophages, improves the antigen presentation and induction of Major Histocompatibility Complex (MHC)-peptides complexes, regulates the T cell polarization, regulates the CD8 T cell homeostasis, stimulates apoptosis and improves the antimicrobial mechanism against intracellular bacteria or parasites (8, 10–12). IFN $\gamma$  is clinically used against infection in patients with Chronic Granulomatous Disease, although it is also effective against *Mycobacterium tuberculosis*, *Salmonella* and fungal infection (13–16).

Teleost and mammalian immune systems show several parallels in innate and acquired immunity. Bioactivity assays using recombinant IFN $\gamma$  from *Salmo salar* or *Oncorhynchus mykiss* have shown this cytokine is able to induce the expression of MHC-I, MHC-II and proteins related to the processing of the antigens and increase the respiratory burst of phagocytic cells (17–20). *In vitro*, IFN $\gamma$  from *S. salar* also has antiviral properties against Salmonid Alphavirus (SAV) and Infectious Pancreatic Necrosis Virus (IPNV) (21). Moreover, in *O. mykiss* cells, IFN $\gamma$  is able to reduce *in vitro* the load of *Piscirickettsia salmonis*, suggesting that in salmonids, as in mammals, IFN $\gamma$  also has antibacterial properties (22).

IFN $\gamma$  has the potential to be used as a therapeutic or prophylactic treatment against bacterial and viral pathogens from salmonid aquaculture. However, the application of these cytokines has been limited due to the lack of a feasible vehicle of delivery compatible with the fish physiology.

In the last decades, Lactic Acid Bacteria (LAB) have risen as a feasible vehicle for the *in situ* delivery of cytokines and bioactive peptides inside of the gastrointestinal tract of mammals (23, 24). Alpha Interferon (IFN $\alpha$ ) (25–27), beta Interferon (IFN $\beta$ ) (28), IFN $\gamma$  (29, 30), and IL-10 (31) have been successfully expressed in LAB under its biologically functional form, producing local and systemic effects after administration to animals.

In *S. salar* and *O. mykiss* the use of LAB for *in situ* delivery of bioactive proteins has been poorly explored, being used only for delivery of antigenic peptides (32, 33). In this work we use *Lactococcus lactis* for the expression of a functional IFN $\gamma$  from *S. salar*. The oral administration of this modified LAB reduces the susceptibility of treated fish against *F. psychrophilum* infection, an important bacterium that affects the salmonid industry.

## MATERIALS AND METHODS

### Construction of MT009 Strain

The gene encoding for the recombinant IFN $\gamma$  from *S. salar* (rIFN $\gamma$ ) was designed *in silico* using the protein sequence of type II

interferon from *S. salar* (FJ263446.1), the Usp45 Signal peptide (SP), the tag GGGHHHHHH, and the promoter P1 of *L. lactis*. The Usp45 SP, type II interferon and tag sequence, were arranged in frame downstream of the P1 promoter. The sequence was synthesized in GenScript and codogenically optimized according to the codon usage of *L. lactis*. This sequence was cloned in the pNZ8149 plasmid (MoBiTec GmbH, Germany) between the sites NcoI and XbaI. The ligation product was electroporated into *L. lactis* NZ3900. The identity of the new construct, prIFN $\gamma$ , was corroborated by sequencing. The *L. lactis* strain NZ3900 containing the plasmid prIFN $\gamma$ , was named MT009.

The preparation of electrocompetent *L. lactis* NZ3900 cells was performed based on the protocol suggested by Mobitec GmbH (34). A colony of *L. lactis* NZ3900 was inoculated in 5 ml of SG-GM17 medium (M17 medium containing 0.5 M sucrose, 2.5% glycine, and 0.5% glucose) and cultured overnight at 30°C without agitation. The culture was inoculated in 40 ml of SG-GM17 medium and grown for 16 h at 30°C without shaking. The next day, this culture was inoculated in 400 ml of SG-GM17 medium and grown to an OD600 between 0.2 and 0.3. Subsequently, the culture was centrifuged at 6,000 $\times$ g for 20 min at 4°C and the collected pellet was washed three times in cold wash buffer (0.5 M sucrose, 10% glycerol, 4°C). In each step, the pellet was collected by centrifugation at 6,000 $\times$ g for 20 min at 4°C and was resuspended by vortex in the corresponding buffer. After the final wash, the pellet was resuspended in 3 ml of wash buffer, aliquoted into 200  $\mu$ l fractions, and then stored at –80°C.

### MT009 Culture Conditions for Hybridization and Biological Activity Assays

From an isolated colony of MT009 a pre-culture was prepared in 0.5% M17-Lactose medium, after incubating overnight, this culture was used to inoculate (2%) 40 ml of M17-Lactose 0.5% medium. After reaching an optical density at 600 nm between 0.4 and 0.6, the culture was induced with nisin 10 ng/ml for 2 h. Bacteria were separated from the supernatant by centrifugation at 6,000 $\times$ g for 20 min at 4°C. The obtained pellet was used in western blot and biological functionality tests, while the supernatant was used in dot blot assays. The MT009 cultures were carried out at 30°C without shaking.

### Preparation of Cytoplasmic Extracts

From the MT009 culture, the bacterial pellet was resuspended in 1 ml of 1 $\times$  PBS supplemented with 1 mM protease inhibitor Phenylmethylsulfonyl fluoride (PMSF). For their rupture, the resuspended cells were kept on ice and sonicated (ultrasonic processor, Sonic Vibracell) for 2 min divided into EIGHT pulses (130 W, 20 kHz, 100%, 2 mm Cv188 stem) of 15 s with intervals of 1 min. After treatment, the cellular debris was separated by centrifugation (13,000 $\times$ g, for 10 min at 4°C). The supernatant containing the cytoplasmic proteins was removed and stored at –20°C. Total protein concentration was determined by the Bradford method.

## SDS-PAGE and Western Blot

For the detection of rIFN $\gamma$  in cytoplasmic extracts, proteins were separated according to their mass by SDS-PAGE (Stacking gel: 8% Acrylamide/Bisacrylamide 29: 1, pH 6.8; Resolving gel: 10% Acrylamide/Bisacrylamide 29: 1, pH 8.8). Ten  $\mu$ g of total protein extracts from MT009 and control strains MT005 (*L. lactis* NZ3900 with pNZ8149) were loaded, together with the BenchMark™ His-tagged Protein Standard. The samples were subjected to electrophoresis for 90 min at 100 V. After electrophoresis the proteins were electrotransferred (300 mA for 2 h at 16°C) to a nitrocellulose membrane. Subsequently, the membrane was blocked for 1 h in a 2% BSA solution and washed three times with 1 $\times$  PBS-Tween 20 (0.5%). The membrane was incubated with Rabbit polyclonal anti-His antibody (Abcam) (dilution 1/5,000) for 1 h at 37°C and then washed three times with PBS-Tween 20 (0.5%). Subsequently, the membrane was incubated with the polyclonal anti-rabbit IgG antibody conjugated to Horseradish Peroxidase (HRP) (1/5,000 dilution) for 1 h at 37°C. Then the membrane was washed three times with PBS-Tween 20 (0.5%). The membrane was incubated with 10 ml chemiluminescent developer solution (Pierce™ ECL Western Blotting Substrate) and exposed to photographic film. For the detection of rIFN $\gamma$  in the extracellular protein concentrate, a dot blot was performed. The nitrocellulose membrane was loaded with 10  $\mu$ l of extracts to give 1  $\mu$ g of total protein per sample. Once air-dried, the membrane was hybridized and developed following the same protocol described for the western blot.

## Culture Cells

Cultures of the Atlantic salmon head kidney cell line (SHK-1) (Sigma-Aldrich) were prepared up to 80% confluence in L15 medium supplemented with 4 mM glutamine and 5% fetal bovine serum. These cultures were incubated for 8 h at 16°C with 0 to 200 ng/ml of *L. lactis* cytoplasmic extracts containing rIFN $\gamma$  (MT009). The total protein concentration was adjusted to the same value with cytoplasmic protein extract of the control strain MT005. After the treatment with cytoplasmic extracts, SHK-1 cell cultures were processed to extract total RNA.

## RT-qPCR

Total RNA from cell cultures was extracted using total RNA kit (Omega Bio-tek), while the total RNA from spleen and kidney was extracted with TRIZOL using approximately 30 mg of each organ. In both cases total RNA integrity was evaluated using agarose gels (1%, TAE 1X) and quantified by absorbance at 260 nm, using the Tecan Infinite 200 PRO equipment or a Synergy™ 2.0 multi-well reader (Biotek). The RNA (2  $\mu$ g) was treated with 1 U of RQ1 RNase-Free DNase (Promega, USA) for 30 min at 37°C in a 10  $\mu$ l volume. The RNase-Free DNase was inactivated adding 1  $\mu$ l of 25 mM EDTA and incubating at 65°C during 10 min. The RT reaction was performed with 1  $\mu$ g of total RNA previously treated with RNase-Free DNase, 200 Units of M-MLV reverse transcriptase (Promega, USA) and 100 pmol of oligo-dT 18-mer, in a 25  $\mu$ l volume. The RT reaction was performed at 42°C for 1 h and stopped by denaturation at 70°C for 15 min. The qPCR reaction was set up using the SYBR

Fast Universal qPCR kit (Kapa Biosystem USA), in a 20  $\mu$ l volume, using 2  $\mu$ l of the RT reaction, and 10 pmol of each primer. To detect and quantify expression of Stat-1, gamma IP10, IFN $\gamma$ , IL-1 $\beta$ , TNF $\alpha$ , TGF $\beta$ , IL-12, IL-6, Mx and eF1 $\alpha$ , the thermal program consisted of 40 cycles of 15 s at 95°C, 15 s at the annealing temperature, and 30 s at 72°C. The qPCR reactions were performed in duplicate on a Stratagene Mx3000P qPCR cycler. The expression of the genes was normalized with respect to the control condition and the expression of the eF1 $\alpha$  gene using the  $\Delta\Delta$ Ct method described by Pfaffl (35). The primers used together the annealing temperature are listed in **Table 1**.

## Oral Administration of MT009 and Challenge Trials

To analyze the expression of genes that respond to IFN $\gamma$  during oral administration of MT009, *O. mykiss* of approximately 25 g were placed in two experimental groups of 12 fish into 27 l tanks. After 7 d of acclimatation, one group was fed with food supplemented with MT009 and the other group continued with a normal diet. Samples of three fish were taken from each group at the beginning of the experiment (T0) and at 2, 4 and 6 d during the treatment with MT009. Each sampled fish was euthanized (see section *Fish Maintenance and Euthanize Protocols*) and processed to extract blood, spleen and kidney.

To analyze the effect of MT009 on the expression of genes that respond to IFN $\gamma$  after oral administration, we followed a similar design as above, with fish placed into three experimental groups of 12 fish into 27 l tanks. After one week of acclimatation, one group was fed 7 d with feed supplemented with MT009, the second group was fed with feed supplemented with the control strain (MT005) and the third group continued with a normal diet. After administration of MT005 and MT009 fish were fed with normal feed. Samples of three fish were taken from each group at 1, 3, 5 and 7 d after the end of the treatment with the test strain. The fish were euthanized and processed to extract blood, spleen and kidney tissues.

**TABLE 1** | Primers used in qPCR experiments.

Gene	Sequence	Ta	Ref
EF1a	F: 5'-GGGTGAGITTTGAGGCTGGTA-3' R: 5'-TTCTGGATCTCCTCAAACCG-3'	60°C	(36)
IL-12	F: 5'-TGACGCTTTTTCTCACCGTTGT-3' R: 5'-ACGCTTTGCAGCATGAGCTTGA-3'	60°C	(37)
IFN- $\gamma$	F: 5'-CCGTACACCGATTGAGGACT-3' R: 5'-GCGGCATTACTCCATCCTAA-3'	60°C	(36)
TGF- $\beta$	F: 5'-AGCTCTCGGAAGAACGACA-3' R: 5'-AGTAGCCAGTGGGTTTCATGG-3'	60°C	(36)
IL-1 $\beta$	F: 5'-CCCCATTGAGACTAAAGCCA-3' R: 5'-GCAACCTCCTCTAGGTGCAG-3'	60°C	(36)
IL-6	F: 5'- CCTTGCGGAACCAACAGTTTG-3' R: 5'- CCTCAGCAACCTTCATCTGGTC- 3'	60°C	(38)
STAT-1	F: 5'- GACCAGCGAACC CAAGAACCTGAA-3' R: 5'-CACAAAGCCAGGATGCAACCAT-3'	60°C	(38)
Gamma IP-10	F: 5'-GTGTCTGAATCCAGAGGCTCCA-3' R: 5'-TCTCATGGTGCTCTCTGTTC-3'	60°C	This work
rpoc	F: 5'-AGG GAG ACT GCC GGT GAT A-3' R: 5'-ACTACGAGGCGCTTTCTCA-3'	55°C	(39)

For the challenge assay with *Flavobacterium psychrophilum* strain 10094 (ETECMA) (40), six tanks of 27 l with 12 specimens of *O. mykiss* of approximately 25 g were arranged in three experimental groups as for the feed trials above. The 6th day post treatment, the groups fed with MT005 (one tank) and MT009 (two tanks), and the untreated fish (two tanks) were challenged by intraperitoneal injection with  $10^8$  CFU of *F. psychrophilum* suspended in 100  $\mu$ l of sterile PBS. As control, one tank of untreated fish was injected with 100  $\mu$ l of sterile PBS.

The RPS of the MT009 treatment was calculated according to the following formula:

$$\text{RPS} = \left( 1 - \frac{\%M^{\text{MT009}}}{\%M^{\text{NF}}} \right) \times 100$$

Where  $\%M^{\text{MT009}}$  is the percentage of mortality of the group fed with MT009 and  $\%M^{\text{NF}}$  is the percentage of mortality of the group fed with normal feed.

The *F. psychrophilum* load in surviving fish was determined by qPCR using absolute quantification of *rpoC* gene with a standard curve. Approximately 30 mg of spleen were processed to extract total DNA using Wizard<sup>®</sup> Genomic DNA Purification Kit (Promega). The DNA was quantified with an Infinite<sup>®</sup> 200 PRO NanoQuant (TECAN) and adjusted to 50 ng/ $\mu$ l. The qPCR was set up using the SensiFAST<sup>™</sup> SYBR<sup>®</sup> No-ROX Kit (BioLine), in a 10  $\mu$ l volume, using 50 ng of DNA from spleen, 2.5 pmol of *rpoC*-Fp forward primer, and 2.5 pmol of *rpoC*-Fp reverse primer (Table 1) (39). The thermal program consisted of one initial denaturation of 3 min at 95°C followed by 35 cycles of 30 s at 95°C, 15 s at 55°C, and 20 s at 72°C. The standard curve was set up using a plasmid that contain cloned the *rpoC* gene of *F. psychrophilum* (the plasmid was donated by Ictio Biotechnologies S.A). The plasmid was diluted in a range between  $10^1$  and  $10^9$  copies/ $\mu$ l.

## Lysozyme Activity

The lysozyme activity in serum was evaluated using the *Micrococcus lysodeikticus* assay. Samples of blood were coagulated at room temperature and centrifugated at 6,000 $\times$ g for 15 min at 4°C. The serum was separated in a clean tube and stored at -80°C. Lysozyme activity was measured according to the protocol described by Shugar (41) adapted in a microtiter plate. Briefly, 20  $\mu$ l of serum was mixed with 180  $\mu$ l of 0.015% [w/v] *M. lysodeikticus* cell suspension in 50 mM Potassium Phosphate Buffer, pH 6.24. The changes in the  $A_{450}$  of samples (S) and blank (B) were measured for 5 min. The slopes ( $\Delta A_{450}/\text{min}$ ) of sample and blank were used to calculate the lysozyme activity according to the following formula:

$$\text{Lysozyme activity} \left[ \frac{\text{U}}{\text{ml}} \right] = \frac{\left( \frac{\Delta A_{450}^{\text{S}}}{\text{min}} - \frac{\Delta A_{450}^{\text{B}}}{\text{min}} \right) \times \text{Df}}{2 \times 10^{-5} [\text{ml}]}$$

## Preparation of the Pathogen Inoculum for Challenge Assays

For the challenge assay with *F. psychrophilum* strain 10094, the bacteria were grown in 10 ml of Tryptone Yeast Extract Salts

(TYES) medium pH 7.2 at 15°C with constant agitation (200 rpm) for 72 h. Five milliliters of this pre-inoculum were used to inoculate 95 ml of sterile TYES broth pH 7.2, then the bacteria were cultured for 4 d at 15°C with constant agitation (200 rpm). At the end of the culture final density was adjusted with TYES to  $10^9$  CFU/ml and 100  $\mu$ l of this culture was used to infect the fish intraperitoneally. *F. psychrophilum* strain 10094 was isolated on November 6th, 2015 from a specimen of *O. mykiss* in Chiloé, Chile.

## Fish Maintenance and Euthanize Protocols

Fish were acclimated for one week before treatment at 12°C in freshwater aquariums with a biomass not higher than 14 g/l, with continuous aeration, and fed with commercial pellets (EWOS MICRO<sup>™</sup> 2 mm) at 1% of body weight. The fish were maintained in freshwater with a pH between 6.6 and 7, the salinity was adjusted to 6 PSU with NaCl to prevent fungal infection, and total ammonia was maintained in a range below 0.02 mg/l. Seventy percent of the water in all the aquaria were changed every day after feeding. Water parameters were monitored daily prior to and after changing the water. Feeding, changing the water, and measuring water parameters were all done manually. The *L. lactis* strains (MT009 or MT005) were administered to the fish together with feed. The bacterial pellet of the cultures was washed with 1 ml of M17 medium, collected by centrifugation at 6,000 $\times$ g for 10 min at 4°C and resuspended in M17 medium in a volume equivalent to 1/10 of the volume of the original culture. The bacteria suspension was mixed with edible oil in a 2:1 ratio and emulsified by vortexing. The emulsion obtained was mixed with the feed, homogenized by shaking in a plastic container and incubated for 1 h at 30°C to allow absorption and drying on feed pellets.

To avoid unnecessary suffering of fish during the challenge and sampling, fish were anesthetized with benzocaine 40 mg/l no longer than 2 min before the intraperitoneal injection, while fish sampled were euthanized with an overdose of anesthesia with benzocaine 40 mg/l for 5–10 min. The fish were maintained in accordance with the ethical standards of the Institutional Ethics Committee of the Universidad de Santiago de Chile and the relevant legislation in force.

## RESULTS

### Expression of rIFN $\gamma$ in *L. lactis* NZ3900


The interferon type II encoded in the *S. salar* genome is a 21.2 kDa protein in its immature form. The modification introduced to allow its expression and secretion in *L. lactis*, included the replacement of the eukaryotic signal peptide (SP) by the prokaryotic SP of Usp45 protein, the introduction of a GGGHHHHHH tag in the C-terminal, the codon optimization to the codon usage of *L. lactis* and the inclusion of a P1 promoter to allow a basal expression. This *in silico* design produces a theoretical protein (rIFN $\gamma$ ) of 22.2 kDa. This sequence was synthesized and cloned into pNZ8149 using *L. lactis* NZ3900 as host. The plasmid containing the rIFN $\gamma$  gene in pNZ8149 was

named rIFN $\gamma$  and the strain of the *L. lactis* containing this plasmid was named MT009.

The expression of rIFN $\gamma$  was evaluated in the cytoplasmic and in extracellular extracts of the MT009 strain. The expression was evaluated in the presence of different concentration of nisin (1, 5, and 10 ng/ml), which is the inducer of the pNIS promoter present in the plasmid pNZ8149. We also analyzed the expression of rIFN $\gamma$  in cultures of MT009 without nisin. As a negative control, we used the cytoplasmic or extracellular extract prepared from cultures of the MT005 strain (*L. lactis* NZ3900). We identified a protein of approximately 23–24 kDa close to the theoretical 22.2 kDa. This protein was present in the cytoplasmic extract from MT009 with and without presence of the inducer, suggesting that promoter P1 acts constitutively and is able to quench the pNIS promoter (**Figure 1**). When the supernatant was analyzed, no band was identified, so the supernatant was concentrated and, after this, a thin band at 21 kDa was identified (data not shown), suggesting poor extracellular secretion.

### Bioactivity of rIFN $\gamma$

In order to test if the rIFN $\gamma$  present in the cytoplasmic extracts was functional, we designed a test using SHK-1 cell cultures. These cells were exposed to several concentrations (0 to 200 ng/ml)




of cytoplasmic protein extracted from MT009 cultures. To control the effect on the cells of different concentrations of bacterial proteins from cytoplasm of *L. lactis* NZ3900, the total concentration of cytoplasmic bacterial proteins was adjusted to 200 ng/ml using cytoplasmic protein extracts from MT005 cultures grown under the same condition. We evaluated the effect over the expression of genes under the activation pathway of IFN $\gamma$ , STAT1, gamma IP10, IL-6 and IFN $\gamma$ . We also evaluated the effect on the expression of general anti-inflammatory and inflammatory cytokines TGF- $\beta$  and IL-1 $\beta$ , respectively. We observed a dose dependent tendency in the expression of all the genes that were downstream in the pathway of genes activated by IFN $\gamma$ , particularly in the expression of IFN $\gamma$  and IL-6. We also observed that the cytoplasmic extracts of MT009 showed no effects on the expression of anti-inflammatory cytokine TGF- $\beta$ , and in the increase of the expression of IL-1 $\beta$  suggesting that cells respond to other bacterial components present in the MT009 protein extract (**Figure 2**).

### *In vivo* Effects During the Oral Administration of MT009 on the Expression of IFN $\gamma$ Related Genes


Our initial results showed that rIFN $\gamma$  was biologically active *in vitro* stimulating the genes related with the IFN $\gamma$  response. To determine if the MT009 strain stimulate the interferon  $\gamma$  pathway when administered orally to rainbow trout, we designed an experiment where fish were fed with  $10^7$  CFU of MT009 per day for one week. Tissues from fish were taken every 2 d during the feeding. The expression of IFN $\gamma$ , STAT-1, gamma IP-10, and IL-6 was evaluated in the spleen and kidney. We observed an increase in the expression of IFN $\gamma$  in kidney between days 2 and 4 after MT009 feeding began, returning to the initial values on day 6. In the spleen, the increase in the expression of IFN $\gamma$  began later, on day 4 and remained high until day 6. A similar pattern was observed with gamma IP10 which increased its expression in the kidney between days 2 and 4, and returned to values close to those observed in the control fish on day 6. In the spleen, we observed an increase in the expression of gamma IP10 only on day 2. STAT-1 also increased its expression mainly in kidney after day 4. IL-6 only increased its expression in kidney, such as is observed in SHK-1 cells (**Figure 3**). MT009 also increased the expression of IL-12, a cytokine secreted by macrophages in response to IFN $\gamma$ , in the spleen and in a lower extent in kidney (**Supplementary Figure 2B**). Interestingly, the expression on IL-1 $\beta$  showed a slight increase in the spleen throughout the time analyzed, but in the kidney showed a strong increase after day 2 (**Supplementary Figure 2A**). Our results show that the kidney and spleen responded differentially to MT009, with the kidney an organ with a more robust and complex response than the spleen.

### *In Vivo* Effects Post Oral Administration of MT009 on the Expression of IFN $\gamma$ Related Genes

The previous results show that MT009 was able to induce the expression of genes related to the IFN $\gamma$  pathway. These effects could be due to the bacterial host or the associated rIFN $\gamma$ . In



**FIGURE 2** | Effect of MT009 protein extract on the expression of IFN $\gamma$  related genes in SHK-1 cells. The figure shows the relative expression of genes related with the downstream IFN $\gamma$ -response pathway Stat-1 (A), gamma IP10 (B), IFN $\gamma$  (C) and IL-6 (D) after exposure to the cytoplasmic extract of MT009 (*L. lactis* NZ3900 prIFN $\gamma$ ). The figure also shows the effects of these extracts on the expression of (E) inflammatory (IL-1 $\beta$ ) and (F) anti-inflammatory (TGF- $\beta$ ) cytokines. The expression was normalized with respect to the expression of eF1 $\alpha$  (NRE). The concentration of bacterial cytoplasmic proteins extracts (CPE) was adjusted with extracts of MT005 (*L. lactis* pNZ8149) to 200 ng/ml. The NRE of each gene was compared against the NRE of the control (cells without treatment). The assays were performed using biological and technical replicates. The significance was analyzed using the Mann-Whitney test (\* $p < 0.05$ ).



**FIGURE 3** | Expression of IFN $\gamma$  related genes during feeding with MT009. The figure shows the relative expression of STAT1 (A), gamma IP10 (B), IFN $\gamma$  (C) and IL-6 (D). The expression was evaluated in the spleen and kidney at the beginning of the treatment (T0) and every two days during treatment (T2, T4, T6). The expression was normalized with respect to the expression of eF1 $\alpha$  gene and by the expression of the genes in the control condition at the same time point. The expression values were compared with the expression at T0. The significance was analyzed using the Mann-Whitney test (\* $p < 0.05$ ).


order to determine whether MT009 influenced the immune system due to prIFN $\gamma$  and if these effects are present post feed with MT009, we evaluated the expression of IFN $\gamma$ , STAT-1, IL-6 and gamma IP10 in the spleen and kidney of fish fed with feed

supplemented with MT009, MT005 (*L. lactis* NZ3900 with pNZ8149) and without any bacterial supplementation. Samples were taken every 2 d for a week, starting the first day after feeding with MT005 or MT009. Our results showed that *L. lactis* was also

able to induce the expression of IFN $\gamma$  after feeding, but in the fish fed with *L. lactis* expressing rIFN $\gamma$ , the induction of IFN $\gamma$  remained high on day 7 post feeding compared to MT005 (~4 fold in spleen and ~1.2 fold in kidney). A similar pattern was observed with the expression of gamma IP10. A more prominent effect was observed in the expression of STAT-1 and IL-6 where maximum expression was observed on days 3 and 5 post treatment, respectively. This effect was observed in spleen and kidney and in both the expression decreased drastically on day 7 post treatment (Figure 4). Interestingly, both MT005 and MT009 were able to induce the expression of TGF- $\beta$ , suggesting that the *L. lactis* possesses anti-inflammatory properties, besides the increase in the expression of IL-6 (Supplementary Figure 3).

### Effect of MT009 on the Lysozyme Activity in Serum After Treatment

The previous finding indicated that IL-6 was strongly induced after treatment with MT009 and this effect was not present in the parental strain MT005, suggesting that is an effect related to the production of IFN $\gamma$  by MT009. Taken into account that IFN $\gamma$  and IL-6 induce the expression of lysozyme in mammals, we evaluated lysozyme activity in serum after treatment with MT009 and MT005. Our results showed a strong induction of the lysozyme activity in serum after treatment with MT009, but not with MT005, indicating that the induction of lysozyme activity is related to the presence of the prIFN $\gamma$  in the strain MT009 (Figure 5). This induction was not observed in serum during the initial administration of MT009 (Supplementary Figure 4), suggesting that induction of lysozyme activity is consequence of a complex cascade and is probably achieved by




**FIGURE 5** | Lysozyme activity in serum post treatment. The figure shows the lysozyme activity in serum of fish treated during seven days with MT009 (solid blue line) and MT005 (dashed blue line). Samples were taken at 1, 3, 5 and 7-days post treatment (PT). The assay was performed using three fish per day and the lysozyme activity was measure in triplicated. The significance was analyzed using the Mann–Whitney test (\* $p < 0.05$ , \*\* $p < 0.01$ ).

the constant accumulation in serum due to the stimulation of the immune system.

### Protective Activity of MT009 Against Bacterial Infection

In order to test if the immunostimulation produced by MT009 is able to confer protection against bacterial pathogens, we challenged *O. mykiss* with *F. psychrophilum*. After



**FIGURE 4** | Expression of IFN $\gamma$  related genes after oral administration of MT009 and MT005. The figure shows the relative expression of STAT1 (A), gamma IP10 (B), IFN $\gamma$  (C), and IL-6 (D) in fish treated with MT009 (solid blue line) and fish treated with the *L. lactis* strain containing the plasmid pNZ8149 (MT005) (dashed blue line). Gene expression was evaluated in the spleen and kidney at the beginning of the treatment (T0) and every two days after treatment (T1, T3, T5, T7). The expression was normalized with respect to the expression of eF1 $\alpha$  gene and by the expression of the genes in the control condition at the same time point. The expression values were compared with the expression at T0. The significance was analyzed using the Mann–Whitney test (\* $p < 0.05$ ).

intraperitoneal injection of *F. psychrophilum* strain 10094, the mortality started at day 6 in all groups challenged with the pathogen and remained similar up to day 10 when survival in the treated group with MT009 stabilized reaching an average of 73.3% at day 17 post-challenge, while the fish treated with MT005 and without treatment reached a 26.6 and 36.6% of survival, respectively. Kaplan–Meier analyses (42) of the survival of fish treated with MT009 showed a statistical difference with the survival curve of fish treated with MT005 ( $p = 0.0052$ , Log-rank (Mantel–Cox) Test;  $p = 0.0169$ , Gehan–Breslow–Wilcoxon Test) and without treatment ( $p = 0.0077$ , Log-rank (Mantel–Cox) Test;  $p = 0.0180$ , Gehan–Breslow–Wilcoxon Test). We were unable to identify statistically significant differences in the survival curve of fish treated with MT005 and without treatment. Thus, the Relative Percentage Survival (RPS) of fish treated with MT009 corresponded to 57.8% with respect to fish without treatment (Figure 6). The quantification by qPCR of the *F. psychrophilum* (39) in the spleen of survivors and in moribund fish indicated that the former had a bacterial load at least 10 times lower than the moribund fish. The surviving fish without treatment showed a bacterial load that was at least three orders lower than the bacterial load of moribund fish, while surviving fish treated with MT009 showed bacterial loads at least two orders lower than moribund fish (Supplementary Figure 5).

## Comparison With Interferon Type II From Salmonids

The results above implies that rIFN $\gamma$  from *S. salar* can also has biological effect on *O. mykiss*, a salmonid closely related to *S. salar*. Previous studies using IFN $\gamma$  from *O. mykiss* have indicated that this interferon has a biological effect on *S. salar* kidney cell lines SHK-1, ASK, and TO (17, 21). These findings allow us to hypothesize that the MT009 strain may also have biological effects


on other salmonids depending on the degree of relatedness among the type II interferon present in the species. To evaluate this possible range of action, a BLASTP search was performed using the IFN $\gamma$  sequence of *S. salar* (NP\_001117030.1) as query, and a subsequent phylogenetic reconstruction of these results was applied. BLAST analysis indicated that the closest type II interferons were found in *Salmo trutta* with 98.3% (XP\_029599439.1) and 97.8% (XP\_029599441.1) identity, followed by interferon present in *Salvelinus namaycush* with 94.3% (XP\_038851623.1) and *O. mykiss* with 93.3% (CAR95729.1) identity. These results suggest that MT009 could also stimulate the response to type II interferon in *S. trutta* and *S. namaycush* (Figure 7).

## DISCUSSION


The current work explores the hypothesis that probiotic microorganisms modified to express type II interferon are able to modify the immune status of the fish and confer protection against pathogens when orally administrated to fish. This protection would result from stimulation of the innate immunity, improving the serological or cellular components. These kind of probiotics, that stimulate the mucosa and beneficially regulate the immune system improving performance against microbial pathogens are termed immunobiotics (47). Some immunobiotics mainly belonging to *Lactobacillus* and *Bifidobacterium* genus, stimulate the expression of interferons (Types I and II) (48, 49) triggering a cascade that confers protection against viral pathogens, such as influenza (50), rotavirus (51) and respiratory syncytial virus (52), and also intracellular parasites like *Toxoplasma* (53). A common characteristic of these immunobiotics is their capacity to stimulate the cellular immune response mediated by Th1 lymphocytes (48), NK (54, 55) and macrophages (52, 53) which is initiated and amplified by the production of type II interferon.

The type II interferon is a pleotropic cytokine produced mainly by Th1 lymphocytes and NK cells in response to the secretion of IL-12, IL-18, and IL-1 $\beta$  produced by the APC (56). Type II interferon stimulates its own expression on Th1 cells polarizing the Th1/Th2 balance toward a Th1 response (57). Type II interferon also produces a polarization of macrophages M1/M2 balance, favoring the M1 cells phenotype (58). The polarization toward a Th1 and M1 response produces metabolic changes in these cells favoring a glycolytic metabolism, with increased expression of enzymes related to the production of Reactive Oxygen Species (ROS) and Reactive Nitrogen Species (RNS), and the expression of autophagy mediators, especially in M1 macrophages (59–61). These metabolic changes enhance the microbicidal capacity of the cells, improving cell-autonomous immunity against intracellular microorganisms (62). The M1 phenotype of macrophages also secretes IL-6 and TNF- $\alpha$  (63, 64), which in turn induces the expression of serum lysozyme, improving the antimicrobial activity in serum against extracellular pathogens (65, 66).

In teleost, although type II interferon has been much less studied, increasing evidence shows that type II interferon plays a similar role as in mammals sharing activation pathways and functions that are



**FIGURE 6** | Challenge assay with *Flavobacterium psychrophilum* strain 10094. The figure shows the Kaplan–Meier survival plot of *Oncorhynchus mykiss* specimens challenged intraperitoneally with  $10^8$  CFU of *F. psychrophilum* strain 10094. A total of three diets were analyzed, (1) fish fed with *L. lactis* NZ3900 producer of rIFN $\gamma$  (MT009), (2) fish fed with *L. lactis* NZ3900 containing the plasmid pNZ3900 (MT005), and (3) fish fed with normal feed (NF, and Control). All conditions were challenged with *F. psychrophilum* 10094, with exception of the control. The challenge of fish treated with MT009 and NF was performed in duplicate.



conserved and complexed with paralogous genes of non-redundant function (IFN $\eta$ ) and diverse receptors (67).

Our work used a Gram-positive bacterial model to express IFN $\gamma$ , an eukaryotic protein, which is predicted to be glycosylated, as its mammalian counterpart (20). In mammals and teleost specimens like rainbow trout, recombinant IFN $\gamma$  has been produced in *Escherichia coli* (20). This recombinant IFN $\gamma$  has shown to be functional, indicating that glycosylation is not necessary for the activity, although glycosylation increases human interferon half-life protecting against proteolytic degradation (68).

In this work, we detected the rIFN $\gamma$  mainly in the cytoplasmic fraction of MT009 cultures, despite our original design incorporating an Usp45 signal peptide to allow secretion of the rIFN $\gamma$ . Low levels of secretion have been also detected in the recombinant production of human (25) and Atlantic salmon type I Interferons (unpublished data) using the Usp45 signal peptide. In human type I IFN this problem was resolved by introducing a spacer sequence between the Usp45 signal peptide and the rest of the interferon protein (25). It remains to determine whether this spacer sequence improves the secretion of rIFN $\gamma$  by MT009 or not.

Others work have used recombinant overexpression of type II interferon in *E. coli* with further purification to study its function in fish (20). However, *L. lactis*, as an alternative recombinant protein system, may not offer as efficient a protein producer as *E. coli*, and purification of recombinant protein from *L. lactis* cultures may be limited (in yields). To overcome this, in this study we decided to evaluate the ability of different amounts of cytoplasmic extracts of the MT009 strain to stimulate the

expression of genes that respond to type II interferon, but taking care to maintain the total amount of bacterial protein constant to evaluate if the effect was caused by the presence of rIFN $\gamma$  or by some other bacterial component present in the extract.

This approach showed that the genes previously described in rainbow trout and mammals that respond to type II interferon (Stat1, gamma IP10, IFN $\gamma$ , and IL-6) (20, 63, 67) are also induced in SHK-1, dose-dependent of the MT009 extract, unlike the gene expression pattern of IL-1 $\beta$  that appeared to respond to the presence of a common component of *L. lactis*.

An interesting point is that *L. lactis* as a Gram-positive bacterium lacks LPS, an inducer of IFN $\gamma$  expression, that supports that the activation of these genes is due to the presence of rIFN $\gamma$  in the cytoplasmic extracts of MT009. Although it is not possible to rule out that the overall response is the product of the simultaneous interaction of rIFN $\gamma$  and some other component of the bacterium. In mammals, type II interferon has been reported to improve the response of keratinocytes and macrophages after stimulation of Toll-Like Receptors (TLR) with microorganisms or agonist, respectively (69, 70).

Once the effect of Type II interferon was tested *in vitro*, we evaluated whether the administration of strain MT009 could induce the expression of genes markers from the response to interferon in the central immune organs of rainbow trout, the spleen and kidney. Although the first encounter of the MT009 strain with the fish immune system is through the digestive mucosa, we hypothesized that the immune cells (lymphocytes and macrophages) present in the GALT respond to type II

interferon and stimulate the central immune response, similar to how oral vaccines stimulate the immune response at the local and systemic level (71).

However, the mechanism of how GALT stimulation produces changes at the level of the central immune organs remains poorly understood in teleost. In the case of MT009, this could involve a) stimulation by rIFN $\gamma$  of the immune cells of the GALT, which in turn responds by secreting cytokines that stimulate the spleen and kidney, b) diffusion of rIFN $\gamma$  from the intestine to the central immune organs, or both. An interesting point is that during the administration of MT009, we identified an early response in the kidney of the IFN $\gamma$  response genes, STAT-1, gamma IP10, and IFN $\gamma$  which was out of phase with the later response observed in spleen. This difference could be the consequence of the presence of a population of immune cells with different activation thresholds by IFN $\gamma$  or by another cytokine secreted in response to IFN $\gamma$ . The transient increase in the expression of IFN $\gamma$  in the spleen and kidney during feeding suggests an initial polarization of lymphocytes to a Th1 phenotype.

After feeding with MT009, changes were also observed in the expression of the genes in response to IFN $\gamma$ , in particular, an increase in the expression of STAT-1 at 3 days post treatment with MT009. This change in expression could be a consequence of additive effect of rIFN $\gamma$  and endogenous interferon produced during the administration of MT009. An experiment carried out in our laboratory indicated that *L. lactis* remains in the gut up to 4 days after oral administration has ended (data not shown), a situation that supports the proposal of a combined effect of rIFN $\gamma$  and endogenous IFN $\gamma$ .

An interesting point to note is that we also observed a slight increase in IFN $\gamma$  expression in fish fed with *L. lactis* NZ3900 (MT005), but that it was not associated with a significant increase in other IFN $\gamma$  response markers. This could be a result of the stimulating effect of *L. lactis*, as occurs in the case of some immunobiotics or lactic acid bacteria in finfish (47, 72). However, in fish fed with MT009, the effect on IFN $\gamma$  expression was sustained until day 7, which suggests that the effect of rIFN $\gamma$  administration initiates a cascade that is maintained even after feeding has ended. This is also reflected in the increased expression of IL-6, a cytokine that is secreted by the M1 phenotype of macrophages in mammals and teleosts (73) and also in the increased activity of serum lysozyme, which is induced specifically by strain MT009 and not by MT005.

Taking into consideration that M1 macrophages are producers of IL-6 (73) and lysozyme (65, 66), and that IL-6 also promotes differentiation of monocyte to macrophages in mammals (74), the results suggest that the administration of MT009 promotes macrophages with the phenotype M1 increasing the microbicidal activity at the cellular and humoral level.

This interpretation is consistent with the increase in resistance to extracellular infection such as that produced by *F. psychrophilum*, and potentially could aid the treatment or prevention of infection by intracellular pathogens such as *P. salmonis*. Taking into consideration that the challenge test was carried out by intraperitoneal injection, bypassing the mucosal barriers, the stimulation at the mucosal level by the MT009 strain

would improve the ability of the fish to resolve the infection, increasing the fish survival.

The biological functionality of rIFN $\gamma$  from *S. salar* was tested in *S. salar* cells and in *O. mykiss*, based that the IFN $\gamma$  of both species presents 93% identity and 96% similarity. It therefore plausible that the MT009 strain could present activity in other salmonids whose type II interferon are closer to that of *S. salar*, such as the IFN $\gamma$  of *S. trutta*, and *S. namaycush*. It however, remains to be determined whether MT009 actually has a stimulating effect on salmonids other than *O. mykiss* and *S. salar*.

## CONCLUSIONS

Our results indicate that in *O. mykiss*, the oral administration of rIFN $\gamma$ -producing *L. lactis* (MT009) stimulates the expression of genes that participate in the IFN $\gamma$ -mediated response at the systemic level, producing a state of protection against infection of extracellular bacterial pathogens. This protection could be mediated by polarization towards the Th1 and M1 phenotypes. The process by which stimulation at the mucosal level with MT009 is connected to a change at the systemic level that provides protection remains to be elucidated. Further characterization of the IFN $\gamma$  response genes and immune cell populations in GALT could help clarify this mechanism.

## DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## ETHICS STATEMENT

Ethical review and approval were not required for the animal study because the applied projects financed by CORFO do not require the approval of an ethics committee for their execution.

## AUTHOR CONTRIBUTIONS

Conceptualization, MT. Methodology, AS, DP, MP, CM, NV, CZ and RV. Investigation, AS, DP, MP, CM, NV, CZ and RV. Resources, MT. Writing—original draft preparation, MT. Writing, review and editing, MT, RV, MP and AG. All authors contributed to the article and approved the submitted version.

## FUNDING

This research was funded by CORFO-INNOVA, grant number 13CTI21257. The APC was funded by USACH. AS was supported by National Doctorate scholarship no. 21180465 (ANID; Government of Chile).

## ACKNOWLEDGMENTS

We would like to thank the administrative team of Consorcio Tecnológico de Sanidad Acuicola S.A for its support in the execution of this research.

## REFERENCES

- Isaacs A, Lindenmann J. Virus Interference. I. The Interferon. *Proc R Soc Lond B Biol Sci* (1957) 147:258–67. doi: 10.1098/rspb.1957.0048
- Pestka S, Krause CD, Walter MR. Interferons, Interferon-Like Cytokines, and Their Receptors. *Immunol Rev* (2004) 202:8–32. doi: 10.1111/j.0105-2896.2004.00204.x
- Lazear HM, Schoggins JW, Diamond MS. Shared and Distinct Functions of Type I and Type III Interferons. *Immunity* (2019) 50:907–23. doi: 10.1016/j.immuni.2019.03.025
- McNab F, Mayer-Barber K, Sher A, Wack A, O'Garra A. Type I Interferons in Infectious Disease. *Nat Rev Immunol* (2015) 15:87–103. doi: 10.1038/nri3787
- Teijaro JR. Type I Interferons in Viral Control and Immune Regulation. *Curr Opin Virol* (2016) 16:31–40. doi: 10.1016/j.coviro.2016.01.001
- Antonelli G, Scagnolari C, Moschella F, Proietti E. Twenty-Five Years of Type I Interferon-Based Treatment: A Critical Analysis of its Therapeutic Use. *Cytokine Growth Factor Rev* (2015) 26:121–31. doi: 10.1016/j.cytogfr.2014.12.006
- Zimonjic DB, Rezanka LJ, Evans CH, Polymeropoulos MH, Trent JM, Popescu NC. Mapping of the Immune Interferon Gamma Gene (IFNG) to Chromosome Band 12q14 by Fluorescence in Situ Hybridization. *Cytogenet Cell Genet* (1995) 71:247–8. doi: 10.1159/000134119
- Kearney S, Delgado C, Lenz LL. Differential Effects of Type I and II Interferons on Myeloid Cells and Resistance to Intracellular Bacterial Infections. *Immunol Res* (2013) 55:187–200. doi: 10.1007/s12026-012-8362-y
- Frucht DM, Fukao T, Bogdan C, Schindler H, O'Shea JJ, Koyasu S. IFN-Gamma Production by Antigen-Presenting Cells: Mechanisms Emerge. *Trends Immunol* (2001) 22:556–60. doi: 10.1016/s1471-4906(01)02005-1
- Tewari K, Nakayama Y, Suresh M. Role of Direct Effects of IFN-Gamma on T Cells in the Regulation of CD8 T Cell Homeostasis. *J Immunol* (2007) 179:2115–25. doi: 10.4049/jimmunol.179.4.2115
- Boehm U, Klamp T, Groot M, Howard JC. Cellular Responses to Interferon- $\gamma$ . *Annu Rev Immunol* (1997) 15:749–95. doi: 10.1146/annurev.immunol.15.1.749
- Xu H, Wipasa J, Yan H, Zeng M, Makobongo MO, Finkelman FD, et al. The Mechanism and Significance of Deletion of Parasite-Specific CD4(+) T Cells in Malaria Infection. *J Exp Med* (2002) 195:881–92. doi: 10.1084/jem.20011174
- Ingram JP, Brodsky IE, Balachandran S. Interferon- $\gamma$  in Salmonella Pathogenesis: New Tricks for an Old Dog. *Cytokine* (2017) 98:27–32. doi: 10.1016/j.cyto.2016.10.009
- Miller CHT, Maher SG, Young HA. Clinical Use of Interferon- $\gamma$ . *Ann N Y Acad Sci* (2009) 1182:69–79. doi: 10.1111/j.1749-6632.2009.05069.x
- Armstrong-James D, Teo IA, Shrivastava S, Petrou MA, Taube D, Dorling A, et al. Exogenous Interferon-Gamma Immunotherapy for Invasive Fungal Infections in Kidney Transplant Patients. *Am J Transplant* (2010) 10:1796–803. doi: 10.1111/j.1600-6143.2010.03094.x
- Razaghi A, Owens L, Heimann K. Review of the Recombinant Human Interferon Gamma as an Immunotherapeutic: Impacts of Production Platforms and Glycosylation. *J Biotechnol* (2016) 240:48–60. doi: 10.1016/j.jbiotec.2016.10.022
- Martin SAM, Taggart JB, Seear P, Bron JE, Talbot R, Teale AJ, et al. Interferon Type I and Type II Responses in an Atlantic Salmon (*Salmo Salar*) SHK-1 Cell Line by the Salmon TRAITs/SGP Microarray. *Physiol Genomics* (2007) 32:33–44. doi: 10.1152/physiolgenomics.00064.2007
- Martin SAM, Zou J, Houlihan DF, Secombes CJ. Directional Responses Following Recombinant Cytokine Stimulation of Rainbow Trout (*Oncorhynchus Mykiss*) RTS-11 Macrophage Cells as Revealed by Transcriptome Profiling. *BMC Genomics* (2007) 8:150. doi: 10.1186/1471-2164-8-150
- Martin SAM, Mohanty BP, Cash P, Houlihan DF, Secombes CJ. Proteome Analysis of the Atlantic Salmon (*Salmo Salar*) Cell Line SHK-1 Following Recombinant IFN-Gamma Stimulation. *Proteomics* (2007) 7:2275–86. doi: 10.1002/pmic.200700020
- Zou J, Carrington A, Collet B, Dijkstra JM, Yoshiura Y, Bols N, et al. Identification and Bioactivities of IFN- $\gamma$  in Rainbow Trout *Oncorhynchus Mykiss*: The First Th1-Type Cytokine Characterized Functionally in Fish. *J Immunol* (2005) 175:2484–94. doi: 10.4049/jimmunol.175.4.2484
- Sun B, Skjaeveland I, Svingerud T, Zou J, Jørgensen J, Robertsen B. Antiviral Activity of Salmonid Gamma Interferon Against Infectious Pancreatic Necrosis Virus and Salmonid Alphavirus and Its Dependency on Type I Interferon. *J Virol* (2011) 85:9188–98. doi: 10.1128/JVI.00319-11
- Álvarez C, Marshall S, Mercado L. IFN-Gamma-Mediated Control of Piscirickettsia Salmonis Survival in Trout Macrophages. *Fish Shellfish Immunol* (2016) 53:112. doi: 10.1016/j.fsi.2016.04.084
- Steidler L, Rottiers P, Coulie B. Actobiotics™ as a Novel Method for Cytokine Delivery. *Ann N Y Acad Sci* (2009) 1182:135–45. doi: 10.1111/j.1749-6632.2009.05067.x
- Bahey-El-Din M, Gahan CGM, Griffin BT. Lactococcus Lactis as a Cell Factory for Delivery of Therapeutic Proteins. *Curr Gene Ther* (2010) 10:34–45. doi: 10.2174/156652310790945557
- Zhang Q, Zhong J, Liang X, Liu W, Huan L. Improvement of Human Interferon Alpha Secretion by Lactococcus Lactis. *Biotechnol Lett* (2010) 32:1271–7. doi: 10.1007/s10529-010-0285-x
- Bayat O, Baradaran A, Ariff A, Mohamad R, Rahim RA. Intracellular Production of IFN-alpha 2b in Lactococcus Lactis. *Biotechnol Lett* (2014) 36:581–5. doi: 10.1007/s10529-013-1390-4
- Ma S, Li K, Li X-S, Guo X-Q, Fu P-F, Yang M-F, et al. Expression of Bioactive Porcine Interferon-Alpha in Lactobacillus Casei. *World J Microbiol Biotechnol* (2014) 30:2379–86. doi: 10.1007/s11274-014-1663-7
- Zhuang Z, Wu Z-G, Chen M, Wang PG. Secretion of Human Interferon-Beta 1b by Recombinant Lactococcus Lactis. *Biotechnol Lett* (2008) 30:1819–23. doi: 10.1007/s10529-008-9761-y
- Rupa P, Monedero V, Wilkie BN. Expression of Bioactive Porcine Interferon-Gamma by Recombinant Lactococcus Lactis. *Vet Microbiol* (2008) 129:197–202. doi: 10.1016/j.vetmic.2007.11.010
- Bernúdez-Humarán LG, Cortes-Perez NG, L'Haridon R, Langella P. Production of Biological Active Murine IFN-gamma by Recombinant Lactococcus Lactis. *FEMS Microbiol Lett* (2008) 280:144–9. doi: 10.1111/j.1574-6968.2007.01038.x
- Steidler L, Hans W, Schotte L, Neiryck S, Obermeier F, Falk W, et al. Treatment of Murine Colitis by Lactococcus Lactis Secreting Interleukin-10. *Science* (2000) 289:1352–5. doi: 10.1126/science.289.5483.1352
- Min L, Li-Li Z, Jun-Wei G, Xin-Yuan Q, Yi-Jing L, Di-Qiu L. Immunogenicity of Lactobacillus-expressing VP2 and VP3 of the Infectious Pancreatic Necrosis Virus (IPNV) in Rainbow Trout. *Fish Shellfish Immunol* (2012) 32:196–203. doi: 10.1016/j.fsi.2011.11.015
- Li-Li Z, Min L, Jun-Wei G, Xin-Yuan Q, Yi-Jing L, Di-Qiu L. Expression of Infectious Pancreatic Necrosis Virus (IPNV) VP2–VP3 Fusion Protein in Lactobacillus Casei and Immunogenicity in Rainbow Trouts. *Vaccine* (2012) 30:1823–9. doi: 10.1016/j.vaccine.2011.12.132
- Mobitec GmbH. *Nice® Expression System for Lactococcus Lactis The Effective & Easy-to-Operate NIsin Controlled Gene Expression System* (2015). Available at: www.mobitec.com (Accessed March 20, 2021).
- Pfaffl MW. A New Mathematical Model for Relative Quantification in Real-Time RT-PCR. *Nucleic Acids Res* (2001) 29:e45. doi: 10.1093/nar/29.9.e45
- Reyes-Cerpa S, Reyes-López F, Toro-Ascuy D, Montero R, Maisey K, Acuña-Castillo C, et al. Induction of Anti-Inflammatory Cytokine Expression by IPNV in Persistent Infection. *Fish Shellfish Immunol* (2014) 41:172–82. doi: 10.1016/j.fsi.2014.08.029
- Parra M, Espinoza D, Valdes N, Vargas R, Gonzalez A, Modak B, et al. Microbiota Modulates the Immunomodulatory Effects of Filifolinone on Atlantic Salmon. *Microorganisms* (2020) 8:1320. doi: 10.3390/microorganisms8091320
- Maisey K, Montero R, Corripio-Miyar Y, Toro-Ascuy D, Valenzuela B, Reyes-Cerpa S, et al. Isolation and Characterization of Salmonid

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2021.696803/full#supplementary-material>

- CD4+ T Cells. *J Immunol* (2016) 196:4150–63. doi: 10.4049/jimmunol.1500439
39. Strepaparava N, Wahli T, Segner H, Petrini O. Detection and Quantification of Flavobacterium Psychrophilum in Water and Fish Tissue Samples by Quantitative Real Time PCR. *BMC Microbiol* (2014) 14:105. doi: 10.1186/1471-2180-14-105
  40. Valdés N, Gonzalez A, Garcia V, Tello M. Analysis of the Microbiome of Rainbow Trout (*Oncorhynchus Mykiss*) Exposed to the Pathogen Flavobacterium Psychrophilum 10094. *Microbiol Resour Announc* (2020) 9:e01562–19. doi: 10.1128/mra.01562-19
  41. Shugar D. The Measurement of Lysozyme Activity and the Ultra-Violet Inactivation of Lysozyme. *Biochim Biophys Acta* (1952) 8:302–9. doi: 10.1016/0006-3002(52)90045-0
  42. Kaplan EL, Meier P. Nonparametric Estimation From Incomplete Observations. *J Am Stat Assoc* (1958) 53:457. doi: 10.2307/2281868
  43. Jones DT, Taylor WR, Thornton JM. The Rapid Generation of Mutation Data Matrices From Protein Sequences. *Bioinformatics* (1992) 8:275–82. doi: 10.1093/bioinformatics/8.3.275
  44. Felsenstein J. Confidence Limits on Phylogenies: An Approach Using the Bootstrap. *Evolution (N Y)* (1985) 39:783. doi: 10.2307/2408678
  45. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary Genetics Analysis Across Computing Platforms. *Mol Biol Evol* (2018) 35:1547–9. doi: 10.1093/molbev/msy096
  46. Stecher G, Tamura K, Kumar S. Molecular Evolutionary Genetics Analysis (MEGA) for Macos. *Mol Biol Evol* (2020) 37:1237–9. doi: 10.1093/molbev/msz312
  47. Villena J, Kitazawa H. Editorial: Immunobiotics-interactions of Beneficial Microbes With the Immune System. *Front Immunol* (2017) 8:1580. doi: 10.3389/fimmu.2017.01580
  48. Villena J, Chiba E, Vizoso-Pinto MG, Tomosada Y, Takahashi T, Ishizuka T, et al. Immunobiotic Lactobacillus Rhamnosus Strains Differentially Modulate Antiviral Immune Response in Porcine Intestinal Epithelial and Antigen Presenting Cells. *BMC Microbiol* (2014) 14:126. doi: 10.1186/1471-2180-14-126
  49. Abt MC, Osborne LC, Monticelli LA, Doering TA, Alenghat T, Sonnenberg GF, et al. Commensal Bacteria Calibrate the Activation Threshold of Innate Antiviral Immunity. *Immunity* (2012) 37:158–70. doi: 10.1016/j.immuni.2012.04.011
  50. Zelaya H, Alvarez S, Kitazawa H, Villena J. Respiratory Antiviral Immunity and Immunobiotics: Beneficial Effects on Inflammation-Coagulation Interaction During Influenza Virus Infection. *Front Immunol* (2016) 7:633. doi: 10.3389/fimmu.2016.00633
  51. Villena J, Vizoso-Pinto MG, Kitazawa H. Intestinal Innate Antiviral Immunity and Immunobiotics: Beneficial Effects Against Rotavirus Infection. *Front Immunol* (2016) 7:563. doi: 10.3389/fimmu.2016.00563
  52. Garcia-Castillo V, Tomokiyo M, Raya Tonetti F, Islam MA, Takahashi H, Kitazawa H, et al. Alveolar Macrophages are Key Players in the Modulation of the Respiratory Antiviral Immunity Induced by Orally Administered Lactobacillus Rhamnosus CRL1505. *Front Immunol* (2020) 11:568636. doi: 10.3389/fimmu.2020.568636
  53. Salas-Lais AG, Robles-Parteras A, Balderas-López JA, Bautista-de Lucio VM. Immunobiotic and Probiotic Potential Effect of Lactobacillus Casei in a Systemic Toxoplasmosis Murine Model. *Microorganisms* (2020) 8:113. doi: 10.3390/microorganisms8010113
  54. de Vrese M, Winkler P, Rautenberg P, Harder T, Noah C, Laue C, et al. Effect of Lactobacillus Gasseri PA 16/8, Bifidobacterium Longum SP 07/3, B. Bifidum MF 20/5 on Common Cold Episodes: A Double Blind, Randomized, Controlled Trial. *Clin Nutr* (2005) 24:481–91. doi: 10.1016/j.clnu.2005.02.006
  55. Namba K, Hatano M, Yaeshima T, Takase M, Suzuki K. Effects of Bifidobacterium Longum BB536 Administration on Influenza Infection, Influenza Vaccine Antibody Titer, and Cell-Mediated Immunity in the Elderly. *Biosci Biotechnol Biochem* (2010) 74:939–45. doi: 10.1271/bbb.90749
  56. Schoenborn JR, Wilson CB. Regulation of Interferon- $\gamma$  During Innate and Adaptive Immune Responses. *Adv Immunol* (2007) 96:41–101. doi: 10.1016/S0065-2776(07)96002-2
  57. Schulz EG, Mariani L, Radbruch A, Höfer T. Sequential Polarization and Imprinting of Type 1 T Helper Lymphocytes by Interferon- $\gamma$  and Interleukin-12. *Immunity* (2009) 30:673–83. doi: 10.1016/j.immuni.2009.03.013
  58. Martinez FO, Gordon S. The M1 and M2 Paradigm of Macrophage Activation: Time for Reassessment. *Front Immunol* (2014) 5:13. doi: 10.3389/fimmu.2014.00013
  59. Su X, Yu Y, Zhong Y, Giannopoulou EG, Hu X, Liu H, et al. Interferon- $\gamma$  Regulates Cellular Metabolism and mRNA Translation to Potentiate Macrophage Activation. *Nat Immunol* (2015) 16:838–49. doi: 10.1038/ni.3205
  60. Palmer CS, Ostrowski M, Balderson B, Christian N, Crowe SM. Glucose Metabolism Regulates T Cell Activation, Differentiation, and Functions. *Front Immunol* (2015) 6:1. doi: 10.3389/fimmu.2015.00001
  61. Almeida L, Lochner M, Berod L, Sparwasser T. Metabolic Pathways in T Cell Activation and Lineage Differentiation. *Semin Immunol* (2016) 28:514–24. doi: 10.1016/j.smim.2016.10.009
  62. MacMicking JD. Interferon-Inducible Effector Mechanisms in Cell-Autonomous Immunity. *Nat Rev Immunol* (2012) 12:367–82. doi: 10.1038/nri3210
  63. Delneste Y, Charbonnier P, Herbault N, Magistrelli G, Caron G, Bonnefoy JY, et al. Interferon- $\gamma$  Switches Monocyte Differentiation From Dendritic Cells to Macrophages. *Blood* (2003) 101:143–50. doi: 10.1182/blood-2002-04-1164
  64. Beyer M, Mallmann MR, Xue J, Staratschek-Jox A, Vorholt D, Krebs W, et al. High-Resolution Transcriptome of Human Macrophages. *PLoS One* (2012) 7:e45466. doi: 10.1371/journal.pone.0045466
  65. Krishnaraju K, Hoffman B, Liebermann DA. The Zinc Finger Transcription Factor Egr-1 Activates Macrophage Differentiation in M1 Myeloblastic Leukemia Cells. *Blood* (1998) 92:1957–66. doi: 10.1182/blood.v92.6.1957
  66. Lewis CE, McCarthy SP, Lorenzen J, McGee JO. Differential Effects of LPS, IFN-Gamma and TNF Alpha on the Secretion of Lysozyme by Individual Human Mononuclear Phagocytes: Relationship to Cell Maturity. *Immunology* (1990) 69:402–8.
  67. Pereira P, Figueras A, Novoa B. Insights Into Teleost Interferon-Gamma Biology: An Update. *Fish Shellfish Immunol* (2019) 90:150–64. doi: 10.1016/j.fsi.2019.04.002
  68. Sareneva T, Pirhonen J, Cantell K, Julkunen I. N-Glycosylation of Human Interferon- $\gamma$  Glycans at Asn-25 are Critical for Protease Resistance. *Biochem J* (1995) 308:9–14. doi: 10.1042/bj3080009
  69. Kajita A, Morizane S, Takiguchi T, Yamamoto T, Yamada M, Iwatsuki K. Interferon-Gamma Enhances Tlr3 Expression and Anti-Viral Activity in Keratinocytes. *J Invest Dermatol* (2015) 135:2005–11. doi: 10.1038/jid.2015.125
  70. Müller E, Christopoulos PF, Halder S, Lunde A, Beraki K, Speth M, et al. Toll-Like Receptor Ligands and Interferon- $\gamma$  Synergize for Induction of Antitumor M1 Macrophages. *Front Immunol* (2017) 8:1383. doi: 10.3389/fimmu.2017.01383
  71. Muñoz-Atienza E, Diaz-Rosales P, Tafalla C. Systemic and Mucosal B and T Cell Responses Upon Mucosal Vaccination of Teleost Fish. *Front Immunol* (2021) 11:622377. doi: 10.3389/fimmu.2020.622377
  72. Ringø E, Hoseinifar SH, Ghosh K, Van Doan H, Beck BR, Song SK. Lactic Acid Bacteria in finfish-An Update. *Front Microbiol* (2018) 9:1818. doi: 10.3389/fmicb.2018.01818
  73. Wentzel AS, Petit J, van Veen WG, Fink IR, Scheer MH, Piazzon MC, et al. Transcriptome Sequencing Supports a Conservation of Macrophage Polarization in Fish. *Sci Rep* (2020) 10:13470. doi: 10.1038/s41598-020-70248-y
  74. Chomarat P, Banchereau J, Davoust J, Palucka AK. IL-6 Switches the Differentiation of Monocytes From Dendritic Cells to Macrophages. *Nat Immunol* (2000) 1:510–4. doi: 10.1038/82763
- Conflict of Interest:** Author MT was employed by company Ictio Biotechnologies SA. The authors declare that the research carried out as well as the results present in this work have been in common agreement with Ictio Biotechnologies SA and the licensee Ictiobiotic SpA. Part of these results have been included in the patent application number 2897-2017 requested in the Chilean industrial property registry (INAPI).
- The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Copyright © 2021 Santibañez, Paine, Parra, Muñoz, Valdes, Zapata, Vargas, Gonzalez and Tello. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.