



Neurological Infection, Kynurenine Pathway, and Parasitic Infection by *Neospora caninum*

Ana Elisa Del'Arco¹, Deivison Silva Argolo², Gilles Guillemin^{3†},
Maria de Fátima Dias Costa^{2†}, Silvia Lima Costa^{2*†} and Alexandre Moraes Pinheiro^{1*†}

OPEN ACCESS

Edited by:

Guochang Hu,
University of Illinois at Chicago,
United States

Reviewed by:

J. Alex Grizzell,
University of Colorado Boulder,
United States
Auriel Willette,
National Institute on Aging (NIH),
United States

*Correspondence:

Silvia Lima Costa
costasl@ufba.br
Alexandre Moraes Pinheiro
amp@ufrb.edu.br

[†]These authors have contributed
equally to this work and share
last authorship

Specialty section:

This article was submitted to
Inflammation,
a section of the journal
Frontiers in Immunology

Received: 24 May 2021

Accepted: 31 December 2021

Published: 26 January 2022

Citation:

Del'Arco AE, Argolo DS, Guillemin G,
Costa MFD, Costa SL and Pinheiro AM
(2022) Neurological Infection,
Kynurenine Pathway, and Parasitic
Infection by *Neospora caninum*.
Front. Immunol. 12:714248.
doi: 10.3389/fimmu.2021.714248

¹ Laboratory of Biochemistry and Veterinary Immunology, Center of Agrarian, Environmental and Biological Sciences, Federal University of Recôncavo of Bahia (UFRB), Cruz das Almas, Brazil, ² Laboratory of Neurochemistry and Cellular Biology, Department of Biochemistry and Biophysics, Institute of Health Sciences, Federal University of Bahia (UFBA), Bahia, Brazil, ³ Neuroinflammation Group, Macquarie Medicine School, Macquarie University, Sydney, NSW, Australia

Neuroinflammation is one of the most frequently studied topics of neurosciences as it is a common feature in almost all neurological disorders. Although the primary function of neuroinflammation is to protect the nervous system from an insult, the complex and sequential response of activated glial cells can lead to neurological damage. Depending on the type of insults and the time post-insult, the inflammatory response can be neuroprotective, neurotoxic, or, depending on the glial cell types, both. There are multiple pathways activated and many bioactive intermediates are released during neuroinflammation. One of the most common one is the kynurenine pathway, catabolizing tryptophan, which is involved in immune regulation, neuroprotection, and neurotoxicity. Different models have been used to study the kynurenine pathway metabolites to understand their involvements in the development and maintenance of the inflammatory processes triggered by infections. Among them, the parasitic infection *Neospora caninum* could be used as a relevant model to study the role of the kynurenine pathway in the neuroinflammatory response and the subset of cells involved.

Keywords: kynurenic acid, glia, neuroinflammation, *Neospora caninum*, quinolinic acid

INTRODUCTION

As the world population is aging, the rates of diagnosable neurological disorders have increased accordingly, indicating an overall adverse impact on health and quality of life (1). Aiming to understand the etiopathogeneses of an array of neurological disorders, many studies seek to elucidate the potential roles of neuroinflammation (2). Although inflammatory processes may not trigger such disorders alone, the immune system nonetheless can greatly influence symptom severity and progression. Scientists are actively looking for therapeutic targets that may efficiently control the exacerbated immune responses associated with neuroinflammation in such conditions (3, 4).

Neurons and glial cells in the central nervous system (CNS) form complex and coordinated networks, of which a key function is to maintain homeostasis (5). Among glia, both astrocytes and microglia constantly assess the CNS environment for potential insult (6). Astrocytes particularly play key roles in maintaining the integrity of the blood–brain barrier (BBB), regulating CNS metabolism, and releasing antioxidants and trophic factors, as well as participating in the process of synaptic communication (7). On the other hand, microglia are considered the resident immune cells of the CNS, contributing to the pro- and anti-inflammatory immune response as they constantly scrutinize the brain parenchyma to eliminate metabolic waste, abnormal cells and proteins, infectious agents, and damaged tissue (8). In this way, astrocytes and microglia become activated and undergo morphological as well as functional transformations in response to different microenvironmental signals. With the aim to maintain homeostasis, the crosstalk among astrocytes and microglia supports neuronal function and plasticity (9).

Infections, traumatic or ischemic injuries, and accumulation of toxic metabolites often induce dysregulation of brain homeostatic processes. At early stages and/or lower levels of activation, astrocytes and microglia can be neuroprotective (polarized as A2 and M2, respectively) as they initiate coordinated responses to restore homeostasis and limit neurotoxicity by rapidly triggering acute inflammation. This might contribute to tissue repair and neurogenesis, as well as clearance of cellular debris, infectious agents, and abnormal proteins (10, 11). However, at chronic and/or high levels of activation (polarized as A1 and M1), these glial cells can become neurotoxic and contribute to neurodegenerative processes (12). While much has been described, researchers are still trying to fully understand the molecular and cellular triggers for this functional switch (2).

One of the most studied pathways in recent years is the kynurenine pathway (KP), which produces a variety of neuroactive metabolites (13, 14). During neuroinflammation, the KP catabolizes approximately 95% of tryptophan to profoundly decrease serotonin and melatonin production and, instead, generate a host of neurotoxic, neuroprotective, and immune-modulating molecules that play key roles in various brain diseases [reviewed by (15–18)].

Many studies underscore the necessity to better characterize the complex yet coordinated glial response, as well as their relevant communications with neurons in neuroinflammatory-implicated dysfunction (3, 9, 19). In that regard, experimental models able to mimic these interactions may grant opportunities to shed new light on the involvement of tryptophan metabolism in neuroinflammation. For example, exposing external factors such as infectious agents to *in vivo* animal models, *ex vivo* brain tissue slices, and *in vitro* freshly dissociated brain cells can potentially be used to obtain much-needed answers (20). However, effective models should permit recognition by and activation of astrocytes and microglia as well as parasite persistence mechanisms allowing the survival and proliferation of the infectious agents. This immune-escape mechanism might

bring information about alternative routes to understand cellular function and linkage of different biochemical pathways, such as KP, and activation and release of neuroprotective factors.

As such, the infection by the parasite *Neospora caninum* appears to be a promising model to study relevant neuroinfectious processes and may contribute to improve the understanding of crosstalk mechanisms between neurons, astrocytes, and microglia. *N. caninum* is an obligate intracellular protozoan, belonging to the phylum *Apicomplexa*, which forms cysts in the CNS and has been shown to lead to abortions in cattle as well as neurological symptoms in dogs (21, 22). Thus, the aim of this review is to highlight the current knowledge about the complex interactions between neuroinflammation and the KP, and to discuss the relevance of the *N. caninum* infection model.

NEUROINFLAMMATION AND THE KYNURENINE PATHWAY

The KP has been widely studied in the CNS over the last three decades, especially with regard to its interactions with the immune system [reviewed by (23–26)]. That said, the KP is highly dynamic. For instance, many CNS cell types display different KP profiles that depend on the disease and region affected (27–29). Microglial activation rapidly occurs during neuroinflammation and is characterized by structural changes from a putatively resting, surveilling, ramified cell toward an activated, spheroidal one (M1 type) producing proinflammatory mediators such as cytokines (30). These M1 microglia also activate astrocytes, which become reactive themselves (A1 type). Together, these intercellular signals stimulate the release of many proinflammatory mediators, such as cytokines (IL-1 β , IL-6, IL-12, IL-23, and TNF- α), chemokine (CCL5 and CCL2), adenosine triphosphate (ATP), reactive oxygen species (ROS), and growth factors (2, 9, 31).

Activation of the KP is associated with induction of the regulatory enzyme, indoleamine 2,3, dioxygenase 1 (IDO-1), and many proinflammatory mediators can stimulate IDO-1 activity. This includes synergistic actions between, for example, TNF- α , IL-1 β , and IL-6 (32–34). Other studies have shown that the induction of IDO-1 may also occur in monocyte/macrophage-like cells, even in the absence of IFN- γ (34–36).

After IDO-1, another key step in the KP is the activity of kynurenine mono oxygenase (KMO), an enzyme highly expressed in microglia, which converts kynurenine (Kyn) into 3-hydroxyanthranilic acid (37). The former regulates apoptosis/necrosis pathways in macrophages and has immunoregulatory and T-cell survival properties. Activation of KMO further leads to the formation of quinolinic acid (QA). The best-known action of QA is as an agonist of NMDA receptors in the nervous system (14) and a potent neuro- and gliotoxin (38).

Interestingly, astrocytes do not express KMO while microglia express all enzymatic components of the KP (29, 39, 40). Thus, microglial activation by inflammatory mediators has a fundamental role in increasing the production of QA (41). In homeostatic situations, the production of kynurenic acid (KA) by

astrocytes antagonizes, to a certain extent, the excitotoxic effects of the QA produced by the microglia, through its antagonism of NMDA receptors (28). In inflammatory conditions, astrocytes produce large amounts of Kyn that can be taken up and used by microglia as additional substrates to produce QA (29, 42–44). The neurotoxicity of QA is observed through at least five different mechanisms including excitotoxicity by NMDA receptor activation, ROS formation, and cytoskeletal destabilization (45, 46). The imbalance in the production of QA and KA, with accumulation of QA, increases the neurotoxic effects by blocking the glutamate uptake by astrocytes (38). Consequently, this process cyclically stimulates ROS production, disturbs the BBB, and increases phosphorylation of structural proteins such as Tau, neurofilament (NF), and glial fibrillary protein (GFAP), which in turn leads to cellular cytoskeletal destabilization (39, 47, 48).

Together with QA, other catabolites of the KP have synergistic neurotoxic effects. O'Farrell et al. (49) observed a reduced neurite outgrowth and complexity after treatment of neuron cultures with conditioned media derived from BV-2 microglia stimulated with IFN- γ . They also observed an increased concentration of tryptophan, Kyn, and 3-hydroxykynurenine (3-HK) in the conditioned media. When the authors used KP inhibitors, the neuronawfi 2l atrophy was fully prevented.

MODELS OF EXOGENOUSLY ACTIVATED NEUROINFLAMMATION AND KP

Many studies have tried to clarify the role of astrocytes and microglia in neuroinflammation and neuroprotection, and each of the mechanisms involved in neuroinflammation are yet to be fully characterized. As above, one common approach to studying neuroinflammation in cell culture and rodent models is exposure to the Gram-negative bacterial lipopolysaccharide (LPS). *In vitro*, LPS induces IFN- γ production and consequently results in the activation of IDO-1 and thus triggers the KP (28, 29, 50, 51). Systemic LPS administration does the same, inducing IDO-1 activity alongside production of brain TNF- α and IL-6 (52).

Despite the effectiveness of stimulating neuroinflammatory processes though, systemic challenge with LPS has failed to fully clarify the mechanisms of KP (53). The response elicited by LPS administration can activate pathogen-associated molecular pattern receptors (PAMPs) and consequently stimulate signaling pathways leading to the production of inflammatory cytokines. Nonetheless, the neuroinflammatory processes triggered in the CNS by infection with bacteria, viruses, and parasites appear to be far more dynamic than the more uniform responses observed following LPS alone.

On the other hand, there are several models of neuroinfection using microorganisms such as HIV or *Toxoplasma gondii*. The model of infection with *N. caninum* is interesting because it is not infectious to humans, which makes it a safer agent to use in medical research. Furthermore, this parasite is easy to cultivate

(54, 55). Importantly, *N. caninum* has the capacity to activate neuroinflammatory processes and grant a viable alternative path to study brain cell interactions and KP activity.

This parasite belongs to the phylum *Apicomplexa*, which are unicellular and spore-forming parasites. Parasites from this group activate the immune response with an associated INF- γ production leading to IDO-1 activation and associated depletion of tryptophan in the host cells. Infection by these groups of parasites also induces an increase in TNF- α and IL-1 β production (56, 57). Infection by *Apicomplexa* parasites also triggers an increased production of Kyn, 3-HK, and QA (17, 58).

N. caninum infection leads to nervous symptoms in cattle and canids related to infection sites in the CNS [reviewed by (59)]. During its infection, the initial recognition by the immune system involves the toll-like receptors, cytosolic sensors such as nucleotide ligand oligomerization domain-like receptors, and NLR family sensors containing pyrin (60–65). Some studies have shown that the activation of these receptors can lead to an increase of Kyn production *via* the NF- κ B signaling pathway (66, 67). During *N. caninum* infection, the lymphocyte T helper 1 (Th1) response is effective to limit the multiplication of the parasite and consequently induces the formation of parasitic cysts in the host. The involvement of lymphocytes T CD4+ and CD8+ is crucial for the development of the anti-parasitic response in mice and is strongly influenced by the systemic increase of IFN- γ (68, 69). Mice treated with recombinant IL-12, which directly is mediated by IFN- γ activity endogenously, had decreased markers of encephalitis as well as brain parasite load 3 weeks later (70). The effectiveness of IFN- γ in protecting against *N. caninum* infection *in vivo* is further supported in a study of mouse strains. Long et al. (71) demonstrated that BALB/c and C57Bl/6 mice were both highly susceptible to the development of *N. caninum*-induced encephalitis, whereas B10.D2 mice were highly resistant. Importantly, splenocytes from B10.D2-infected mice also displayed high antigen-stimulated INF- γ to IL-4 ratios while these ratios were much lower in the other two strains, which indicates that peripheral immune responses favoring INF- γ production might contribute to *N. caninum* protection in select rodent strains *in vivo* (71).

To demonstrate the steps of *N. caninum* infection within the CNS, Yamane et al. (72) confirmed that the parasite proliferation in cultured primary bovine brain cells was controlled by IFN- γ as well as TNF- α . Similarly, we have found in mixed cultures of astrocytes and microglia that *N. caninum* induces the production of TNF- α , IL-10, IL-6, and nitric oxide (NO) (55, 73–75). Interestingly, when glia-neuron co-cultures were infected by tachyzoites for 72 h, we observed a *N. caninum*-induced retraction of neurites but no hypothesized neuronal loss (76). However, application of IFN- γ to the medium restored neurite outgrowth of infected cells (76), which is consistent with the IFN- γ -mediated neuroprotection described in the *in vivo* models above. Taken together, we proposed that *N. caninum* infection triggers a local inflammatory response by way of TNF- α and NO production, but without IFN- γ application, the presence of IL-10 and IL-6 may trigger a switch to a Th2 profile, which could help preserve the environment.

Neuro-glia co-cultures infected with *N. caninum* also induced astroglioses, which were characterized by an increased GFAP expression and also induced the mRNA expression for IL-10 (77). At the same time, the infection induced the mRNA expression for brain-derived neurotrophic factor (BDNF) and neuronal growth factor (NGF), which facilitate actions such as synapses plasticity and formation (78). Additionally, treatment of the glia-neuron co-culture with the medium of mixed cultures of astrocytes and microglia infected by this parasite also induced neurite outgrowth (77).

In bovine endothelial cells infected with *N. caninum*, IDO-1 activation was observed in the presence of IFN- γ (79). In a glia-neuron co-culture model, IDO-1 activation was associated with the control of the parasitic proliferation, since inhibiting IDO-1 with 1-MT increased tachyzoite proliferation (80). Also, in the absence of IFN- γ , IDO-1 was activated by the infection, inducing a 50% increase in Kyn compared to uninfected co-cultures. Recently, Argolo et al. (81) demonstrated that, despite observation of neurite outgrowth and release of neurotrophic factors, the infection increased levels of QA and of CCL5 and CCL2 mRNA expression.

These chemokines recruit cells from immune system and activate microglia to control the parasite infection. Aside from its neurotoxic role as an NMDA agonist, QA also contributes to the production of NAD⁺, which together modulate the production of inflammatory cytokines IL-1 β , TNF- α , and IL-6 and facilitate the change toward a pro-inflammatory profile and a resolutive response by the release of IL-10 (29, 82, 83). However, dysregulation of QA production has been seen during infectious models and it is still unclear whether NAD production is altered in these processes. It is possible that a portion of KP metabolites, such as QA, are directed toward NAD⁺ production in response to infection, but the mechanism(s) mediating this process remain unclear (84).

It is possible that some effects of *N. caninum* infection, such as Th2 cytokine production and release of neurotrophic factors, evidence an atypical immune response associated with parasite persistence. However, these findings should not be confounded with universal neuroprotection as infection progression triggers astrocyte death and neurological impairment. Other studies demonstrated that the parasite could change the immune response to favor its persistence, such as increased population of T CD8⁺ regulatory cells (68), inhibition of IL-12p40 production (85), and the inhibition of Th1 response by STAT3 phosphorylation in the invasion process (86, 87). Taken together, these varied immune responses to *N. caninum* infection underscore the viability of this model in aiding discovery and further characterization of the dynamic and context-dependent function of neuroinflammatory processes related to the KP and, ultimately, brain function in normal and dysfunctional conditions.

CONCLUSION

The KP, the major route of tryptophan catabolism, produces NAD⁺ and several intermediates, which have neuroactive properties. In recent decades, studies of the KP have not only brought new understanding about interactions of these

intermediates and the function of CNS, but also highlighted the influence of the unbalanced production of these bioactive catabolites and their potential impact on various neurological disorders. At the same time, many scientists have used *in vitro* experimental models to study cellular mechanisms and pathways involved in both physiological and pathological conditions. The complex and multi-factorial crosstalk between glial cells and neurons have been studied to better understand the processes of brain homeostasis and neuroinflammation, which is a common feature in most brain diseases and disorders.

This paper describes a potentially relevant and novel approach utilizing neuroinfection *via* the parasite *N. caninum* to study KP activation as well as astrocyte/glia crosstalk, particularly given the atypical immune response following infection. Although *N. caninum* triggers an acute inflammatory response marked also by astrogliosis, proinflammatory activation of microglia, and QA production, it also triggers concomitant neurite outgrowth and neurotrophic factor release in culture. This is likely due to the production of neurotrophins and immuno-modulating cytokine IL-10. This highlights the importance of assessing the KP profile and its relationship with other inflammatory molecules in neurological disorders associated with infection by non-LPS factors, such as viruses and parasites, with the aim to understand the consequences of lesser characterized biochemical interactions.

The infectious process most commonly begins in the periphery, resulting in the dysregulation of KP metabolism and alteration of the immune system before propagating in a secondary stage to affect the CNS. The multifactorial and complex interactions between periphery and CNS, the KP, and the alteration of BBB integrity should all be taken into consideration when using models of neuroinfection. The systematic and comprehensive characterization of this response presents another step toward a better understanding of cellular and molecular communication mechanisms between all the protagonists and the inflammatory response triggered by the parasitic infection. With this concluded, some questions remain unanswered: (1) how the mechanism of neural protection occurs, even with the increase in QA, and whether this relationship may bring new insights to understand the CNS response to external insult; (2) how CNS homeostasis is disrupted after a systemic challenge with *N. caninum*, where the KP intermediates are produced, and by which brain cells; (3) what other metabolic pathways would be associated with KP to justify a neuroprotective response; (4) do the atypical immune responses described in cell culture studies translate fully to *in vivo* models or provide additional novelty for understanding relevant immune activity; and (5) can the *N. caninum* infection model revolutionize our understanding of the cellular crosstalk in the CNS, highlight new processes worthy of investigation, and ultimately facilitate the development of more effective therapeutic interventions for immune-related dysfunction of the brain.

AUTHOR CONTRIBUTIONS

AE, MF, and AP performed the literature search and drafted the manuscript. DA wrote a section of the manuscript. MF, GG, SC,

and AP contributed to the exchange of knowledge, and wrote, edited, and critically revised the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by the National Council for Scientific and Technological Development (CNPq, Process 431927/2016-2; INCT) and the Foundation for Research Support of the State of

Bahia (FAPESB, Processes APP 0058/2015). GG is funded by the NHMRC.

ACKNOWLEDGMENTS

We thank the Postgraduate Program in Immunology of the Federal University of Bahia and FAPESB for fellowship to DSA (Process N° BOL0474/2018).

REFERENCES

- Feigin VL, Abajobir AA, Abate KH, Abd-Allah F, Abdulle AM, Abera SF, et al. Global, Regional, and National Burden of Neurological Disorders During 1990–2015: A Systematic Analysis for the Global Burden of Disease Study 2015. *Lancet Neurol* (2017) 16(11):877–97. doi: 10.1016/S1474-4422(17)30299-5
- Jha MK, Lee WH, Suk K. Functional Polarization of Neuroglia: Implications in Neuroinflammation and Neurological Disorders. *Biochem Pharmacol* (2016) 103:1–16. doi: 10.1016/j.bcp.2015.11.003
- Mishra A, Bandopadhyay R, Singh PK, Mishra PS, Sharma N, Khurana N. Neuroinflammation in Neurological Disorders: Pharmacotherapeutic Targets From Bench to Bedside. *Metab Brain Dis* (2021) 36(7):1–36. doi: 10.1007/s11011-021-00806-4
- Gillhus NE, Deuschl G. Neuroinflammation—A Common Thread in Neurological Disorders. *Nat Rev Neurol* (2019) 15(8):429–30. doi: 10.1038/s41582-019-0227-8
- Vainchtein ID, Molofsky AV. Astrocytes and Microglia: In Sickness and in Health. *Trends Neurosci* (2020) 43(3):144–54. doi: 10.1016/j.tins.2020.01.003
- Michinaga S, Koyama Y. Dual Roles of Astrocyte-Derived Factors in Regulation of Blood-Brain Barrier Function After Brain Damage. *Int J Mol Sci* (2019) 20(3):571. doi: 10.3390/ijms20030571
- Araque A, Parpura V, Sanzgiri RP, Haydon PG. Tripartite Synapses: Glia, the Unacknowledged Partner. *Trends Neurosci* (1999) 22(5):208–15. doi: 10.1016/S0166-2236(98)01349-6
- Nimmerjahn A, Kirchhoff F, Helmchen F. Resting Microglial Cells are Highly Dynamic Surveillants of Brain Parenchyma *In Vivo*. *Science* (2005) 308(5726):1314–8. doi: 10.1126/science.1110647
- Jha MK, Jo M, Kim JH, Suk K. Microglia-Astrocyte Crosstalk: An Intimate Molecular Conversation. *Neuroscientist* (2019) 25(3):227–40. doi: 10.1177/1073858418783959
- Shahab T, Khanabdalil R, Moghadamtousi SZ, Kadir HA, Mohan G. Neuroinflammation Pathways: A General Review. *Int J Neurosci* (2017) 127(7):624–33. doi: 10.1080/00207454.2016.1212854
- Russo MV, McGavern DB. Inflammatory Neuroprotection Following Traumatic Brain Injury. *Science* (2016) 353(6301):783–5. doi: 10.1126/science.aaf6260
- Block ML, Zecca L, Hong JS. Microglia-Mediated Neurotoxicity: Uncovering the Molecular Mechanisms. *Nat Rev Neurosci* (2007) 8(1):57–69. doi: 10.1038/nrn2038
- Jones SP, Guillemin GJ, Brew BJ. The Kynurenine Pathway in Stem Cell Biology. *Int J Tryptophan Res* (2013) 6:IJTR-S12626. doi: 10.4137/IJTR.S12626
- Stone TW. Neuropharmacology of Quinolinic and Kynurenic Acids. *Pharmacol Rev* (1993) 45(3):309–79.
- Huang YS, Ogbechi J, Clanchy FI, Williams RO, Stone TW. IDO and Kynurenine Metabolites in Peripheral and CNS Disorders. *Front Immunol* (2020) 11:388. doi: 10.3389/fimmu.2020.00388
- Cervenka I, Agudelo LZ, Ruas JL. Kynurenines: Tryptophan's Metabolites in Exercise, Inflammation, and Mental Health. *Science* (2017) 357(6349):eaaf9794. doi: 10.1126/science.aaf9794
- Hunt NH, Too LK, Khaw LT, Guo J, Hee L, Mitchell AJ, et al. The Kynurenine Pathway and Parasitic Infections That Affect CNS Function. *Neuropharmacology* (2017) 112:389–98. doi: 10.1016/j.neuropharm.2016.02.029
- O'Farrell K, Harkin A. Stress-Related Regulation of the Kynurenine Pathway: Relevance to Neuropsychiatric and Degenerative Disorders. *Neuropharmacology* (2017) 112:307–23. doi: 10.1016/j.neuropharm.2015.12.004
- Yang QQ, Zhou JW. Neuroinflammation in the Central Nervous System: Symphony of Glial Cells. *Glia* (2019) 67(6):1017–35. doi: 10.1002/glia.23571
- Schwarcz R, Stone TW. The Kynurenine Pathway and the Brain: Challenges, Controversies and Promises. *Neuropharmacology* (2017) 112:237–47. doi: 10.1016/j.neuropharm.2016.08.003
- Dubey JP. Recent Advances in *Neospora* and Neosporosis. *Vet Parasitol* (1999) 84(3-4):349–67. doi: 10.1016/S0304-4017(99)00044-8
- Dubey JP, Lindsay DS. A Review of *Neospora Caninum* and Neosporosis. *Vet Parasitol* (1996) 67(1-2):1–59. doi: 10.1016/S0304-4017(96)01035-7
- Tóth F, Cseh EK, Vécsei L. Natural Molecules and Neuroprotection: Kynurenic Acid, Pantethine and α -Lipoic Acid. *Int J Mol Sci* (2021) 22(1):403. doi: 10.3390/ijms22010403
- Nagy EE, Frigy A, Szász JA, Horváth E. Neuroinflammation and Microglia/Macrophage Phenotype Modulate the Molecular Background of Post-Stroke Depression: A Literature Review. *Exp Ther Med* (2020) 20(3):2510–23. doi: 10.3892/etm.2020.8933
- Venkatesan D, Iyer M, Narayanasamy A, Siva K, Vellingiri B. Kynurenine Pathway in Parkinson's Disease—An Update. *Neurologicalsci* (2020) 10:100270. doi: 10.1016/j.ensci.2020.100270
- Savitz J. The Kynurenine Pathway: A Finger in Every Pie. *Mol Psychiatry* (2020) 25(1):131–47. doi: 10.1038/s41380-019-0414-4
- Guillemin GJ, Cullen KM, Lim CK, Smythe GA, Garner B, Kapoor V, et al. Characterization of the Kynurenine Pathway in Human Neurons. *J Neurosci* (2007) 27(47):12884–92. doi: 10.1523/JNEUROSCI.4101-07.2007
- Guillemin GJ, Smythe G, Takikawa O, Brew BJ. Expression of Indoleamine 2, 3-Dioxygenase and Production of Quinolinic Acid by Human Microglia, Astrocytes, and Neurons. *Glia* (2005) 49(1):15–23. doi: 10.1002/glia.20090
- Guillemin GJ, Kerr SJ, Smythe GA, Smith DG, Kapoor V, Armati PJ, et al. Kynurenine Pathway Metabolism in Human Astrocytes: A Paradox for Neuronal Protection. *J Neurochem* (2001) 78(4):842–53. doi: 10.1046/j.1471-4159.2001.00498.x
- Franco R, Fernandez-Suarez D. Alternatively Activated Microglia and Macrophages in the Central Nervous System. *Prog Neurobiol* (2015) 131:65–86. doi: 10.1016/j.pneurobio.2015.05.003
- Farina C, Aloisi F, Meinl E. Astrocytes are Active Players in Cerebral Innate Immunity. *Trends Immunol* (2007) 28(3):138–45. doi: 10.1016/j.it.2007.01.005
- Zunszain PA, Anacker C, Cattaneo A, Choudhury S, Musaelyan K, Myint AM, et al. Interleukin-1 β : A New Regulator of the Kynurenine Pathway Affecting Human Hippocampal Neurogenesis. *Neuropsychopharmacology* (2012) 37(4):939–49. doi: 10.1038/npp.2011.277
- O'Connor JC, André C, Wang Y, Lawson MA, Szegedi SS, Lestage J, et al. Interferon- γ and Tumor Necrosis Factor- α Mediate the Upregulation of Indoleamine 2, 3-Dioxygenase and the Induction of Depressive-Like Behavior in Mice in Response to *Bacillus Calmette-Guérin*. *J Neurosci* (2009) 29(13):4200–9. doi: 10.1523/JNEUROSCI.5032-08.2009
- Fujigaki H, Saito K, Fujigaki S, Takemura M, Sudo K, Ishiguro H, et al. The Signal Transducer and Activator of Transcription 1 α and Interferon Regulatory Factor 1 are Not Essential for the Induction of Indoleamine 2, 3-Dioxygenase by Lipopolysaccharide: Involvement of P38 Mitogen-Activated Protein Kinase and Nuclear Factor- κ B Pathways, and Synergistic Effect of Several Proinflammatory Cytokines. *J Biochem* (2006) 139(4):655–62. doi: 10.1093/jb/mvj072

35. Fujigaki S, Saito K, Sekikawa K, Tone S, Takikawa O, Fujii H, et al. Lipopolysaccharide Induction of Indoleamine 2, 3-Dioxygenase Is Mediated Dominantly by an IFN- γ -Independent Mechanism. *Eur J Immunol* (2001) 31(8):2313–8. doi: 10.1002/1521-4141(200108)31:8<2313::AID-IMMU2313>3.0.CO;2-S
36. Wang Y, Lawson MA, Dantzer R, Kelley KW. LPS-Induced Indoleamine 2, 3-Dioxygenase Is Regulated in an Interferon- γ -Independent Manner by a JNK Signaling Pathway in Primary Murine Microglia. *Brain Behav Immun* (2010) 24(2):201–9. doi: 10.1016/j.bbi.2009.06.152
37. Jones SP, Franco NF, Varney B, Sundaram G, Brown DA, de Bie J, et al. Expression of the Kynurenine Pathway in Human Peripheral Blood Mononuclear Cells: Implications for Inflammatory and Neurodegenerative Disease. *PLoS One* (2015) 10(6):e0131389. doi: 10.1371/journal.pone.0131389
38. Guillemin GJ. Quinolinic Acid, The Inescapable Neurotoxin. *FEBS J* (2012) 279(8):1356–65. doi: 10.1111/j.1742-4658.2012.08485.x
39. Garrison AM, Parrott JM, Tuñon A, Delgado J, Redus L, O'Connor JC. Kynurenine Pathway Metabolic Balance Influences Microglia Activity: Targeting Kynurenine Monooxygenase to Dampen Neuroinflammation. *Psychoneuroendocrinology* (2018) 94:1–10. doi: 10.1016/j.psyneuen.2018.04.019
40. Chiarugi A, Calvani M, Meli E, Traggiai E, Moroni F. Synthesis and Release of Neurotoxic Kynurenine Metabolites by Human Monocyte-Derived Macrophages. *J Neuroimmunol* (2001) 120(1-2):190–8. doi: 10.1016/S0165-5728(01)00418-0
41. Cogo A, Mangin G, Maier B, Callebort J, Mazighi M, Chabriat H, et al. Increased Serum QUIN/KYNA Is a Reliable Biomarker of Post-Stroke Cognitive Decline. *Mol Neurodegener* (2021) 16(1):1–20. doi: 10.1186/s13024-020-00421-4
42. Garcez ML, Tan VX, Heng B, Guillemin GJ. Sodium Butyrate and Indole-3-Propionic Acid Prevent the Increase of Cytokines and Kynurenine Levels in LPS-Induced Human Primary Astrocytes. *Int J Tryptophan Res* (2020) 13:1178646920978404. doi: 10.1177/1178646920978404
43. Herédi J, Cseh EK, Berkó AM, Veres G, Zádori D, Toldi J, et al. Investigating KYN Production and Kynurenergic Manipulation on Acute Mouse Brain Slice Preparations. *Brain Res Bull* (2019) 146:185–91. doi: 10.1016/j.brainresbull.2018.12.014
44. Notarangelo FM, Beggato S, Schwarz R. Assessment of Prenatal Kynurenine Metabolism Using Tissue Slices: Focus on the Neosynthesis of Kynurenine Acid in Mice. *Dev Neurosci* (2019) 41(1-2):102–11. doi: 10.1159/000499736
45. Muller FL, Song W, Jang YC, Liu Y, Sabia M, Richardson A, et al. Denervation-Induced Skeletal Muscle Atrophy Is Associated With Increased Mitochondrial ROS Production. *Am J Physiol Regul Integr Comp Physiol* (2007) 293:R1159–68. doi: 10.1152/ajpregu.00767.2006
46. Pierozan P, Zamoner A, Krombauer-Soska A, Silvestrin R, Loureiro S, Heimfarth L, et al. Acute Intrastriatal Administration of Quinolinic Acid Provokes Hyperphosphorylation of Cytoskeletal Intermediate Filament Proteins in Astrocytes and Neurons of Rats. *Exp Neurol* (2010) 224:188–96. doi: 10.1016/j.expneurol.2010.03.009
47. Kaindl AM, Degos V, Peineau S, Gouadon E, Chhor V, Liron G, et al. Activation of Microglial N-Methyl-D-Aspartate Receptors Triggers Inflammation and Neuronal Cell Death in the Developing and Mature Brain. *Ann Neurol* (2012) 72(4):536–49. doi: 10.1002/ana.23626
48. Guillemin GJ, Croitoru-Lamoury J, Dormont D, Armati PJ, Brew BJ. Quinolinic Acid Upregulates Chemokine Production and Chemokine Receptor Expression in Astrocytes. *Glia* (2003) 41(4):371–81. doi: 10.1002/glia.10175
49. O'Farrell K, Fagan E, Connor TJ, Harkin A. Inhibition of the Kynurenine Pathway Protects Against Reactive Microglial-Associated Reductions in the Complexity of Primary Cortical Neurons. *Eur J Pharmacol* (2017) 810:163–73. doi: 10.1016/j.ejphar.2017.07.008
50. Lestage J, Verrier D, Palin K, Dantzer R. The Enzyme Indoleamine 2, 3-Dioxygenase Is Induced in the Mouse Brain in Response to Peripheral Administration of Lipopolysaccharide and Superantigen. *Brain Behav Immun* (2002) 16(5):596–601. doi: 10.1016/S0889-1591(02)00014-4
51. Cesura AM, Alberati-Giani D, Köhler C, Ricciardi-Castagnoli P. Regulation of Enzymes of the Kynurenine Pathway by Interferon-Gamma in Murine Macrophages and Microglia. *Pharmacol Res* (1995) 31:125. doi: 10.1016/1043-6618(95)86760-0
52. O'Connor JC, Lawson MA, Andre C, Moreau M, Lestage J, Castanon N, et al. Lipopolysaccharide-Induced Depressive-Like Behavior Is Mediated by Indoleamine 2, 3-Dioxygenase Activation in Mice. *Mol Psychiatry* (2009) 14(5):511–22. doi: 10.1038/sj.mp.4002148
53. Batista CRA, Gomes GF, Candelario-Jalil E, Fiebich BL, de Oliveira ACP. Lipopolysaccharide-Induced Neuroinflammation as a Bridge to Understand Neurodegeneration. *Int J Mol Sci* (2019) 20(9):2293. doi: 10.3390/ijms20092293
54. Débare H, Moiré N, Ducournau C, Schmidt J, Laakmann JD, Schwarz RT, et al. *Neospora Caninum* Glycosylphosphatidylinositols Used as Adjuvants Modulate Cellular Immune Responses Induced In Vitro by a Nanoparticle-Based Vaccine. *Cytokine* (2021) 144:155575. doi: 10.1016/j.cyto.2021.155575
55. Pinheiro AM, Costa SL, Freire SM, Almeida MAOD, Tardy M, El Bachá R, et al. Astroglial Cells in Primary Culture: A Valid Model to Study *Neospora Caninum* Infection in the CNS. *Vet Immunol Immunopathol* (2006) 113(1-2):243–7. doi: 10.1016/j.vetimm.2006.05.006
56. Ufermann CM, Domröse A, Babel T, Tersteegen A, Cengiz SC, Eller SK, et al. Indoleamine 2, 3-Dioxygenase Activity During Acute Toxoplasmosis and the Suppressed T Cell Proliferation in Mice. *Front Cell Infect Microbiol* (2019) 9:184. doi: 10.3389/fcimb.2019.00184
57. Schmid M, Lehmann MJ, Lucius R, Gupta N. Apicomplexan Parasite, *Eimeria Falciformis*, Co-opts Host Tryptophan Catabolism for Life Cycle Progression in Mouse. *J Biol Chem* (2012) 287(24):20197–207. doi: 10.1074/jbc.M112.351999
58. Notarangelo FM, Wilson EH, Horning KJ, Thomas MAR, Harris TH, Fang Q, et al. Evaluation of Kynurenine Pathway Metabolism in *Toxoplasma Gondii*-Infected Mice: Implications for Schizophrenia. *Schizophr Res* (2014) 152(1):261–7. doi: 10.1016/j.schres.2013.11.011
59. Dubey JP. Review of *Neospora Caninum* and Neosporosis in Animals. *Korean J Parasitol* (2003) 41(1):1–16. doi: 10.3347/kjip.2003.41.1.1
60. Wang X, Gong P, Zhang X, Li S, Lu X, Zhao C, et al. NLRP3 Inflammasome Participates in Host Response to *Neospora Caninum* Infection. *Front Immunol* (2018) 9:1791. doi: 10.3389/fimmu.2018.01791
61. Wang X, Gong P, Zhang N, Li L, Chen S, Jia L, et al. Inflammasome Activation Restrains the Intracellular *Neospora Caninum* Proliferation in Bovine Macrophages. *Vet Parasitol* (2019) 268:16–20. doi: 10.1016/j.vetpar.2019.02.008
62. Jin X, Gong P, Zhang X, Li G, Zhu T, Zhang M, et al. Activation of ERK Signaling via TLR11 Induces IL-12p40 Production in Peritoneal Macrophages Challenged by *Neospora Caninum*. *Front Microbiol* (2017) 8:1393. doi: 10.3389/fmicb.2017.01393
63. Davoli-Ferreira M, Fonseca DM, Mota CM, Dias MS, Lima-Junior DS, da Silva MV, et al. Nucleotide-Binding Oligomerization Domain-Containing Protein 2 Prompts Potent Inflammatory Stimuli During *Neospora Caninum* Infection. *Sci Rep* (2016) 6(1):1–14. doi: 10.1038/srep29289
64. Beiting DP, Peixoto L, Akopyants NS, Beverley SM, Wherry EJ, Christian DA, et al. Differential Induction of TLR3-Dependent Innate Immune Signaling by Closely Related Parasite Species. *PLoS One* (2014) 9(2):e88398. doi: 10.1371/journal.pone.0088398
65. Mineo TW, Oliveira CJ, Gutierrez FR, Silva JS. Recognition by Toll Like Receptor 2 Induces Antigen-Presenting Cell Activation and Th1 Programming During Infection by *Neospora Caninum*. *Immunol Cell Biol* (2010) 88(8):825–33. doi: 10.1038/ichb.2010.52
66. Mithaiwala MN, Santana-Coelho D, Porter GA, O'Connor JC. Neuroinflammation and the Kynurenine Pathway in CNS Disease: Molecular Mechanisms and Therapeutic Implications. *Cells* (2021) 10(6):1548. doi: 10.3390/cells10061548
67. Farzi A, Reichmann F, Meinitzer A, Mayerhofer R, Jain P, Hassan AM, et al. Synergistic Effects of NOD1 or NOD2 and TLR4 Activation on Mouse Sickness Behavior in Relation to Immune and Brain Activity Markers. *Brain Behav Immun* (2015) 44:106–20. doi: 10.1016/j.bbi.2014.08.011
68. Correia A, Ferreira P, Costa AA, Dias J, Melo J, Costa R, et al. Mucosal and Systemic T Cell Response in Mice Intragastrically Infected With *Neospora Caninum* Tachyzoites. *Vet Res* (2013) 44(1):1–13. doi: 10.1186/1297-9716-44-69
69. Tanaka T, Hamada T, Inoue N, Nagasawa H, Fujisaki K, Suzuki N, et al. The Role of CD4+ or CD8+ T Cells in the Protective Immune Response of BALB/c Mice to *Neospora Caninum* Infection. *Vet Parasitol* (2000) 90(3):183–91. doi: 10.1016/S0304-4017(00)00238-7

70. Baszler TV, Long MT, McElwain TF, Mathison BA. Interferon- γ and Interleukin-12 Mediate Protection to Acute *Neospora Caninum* Infection in BALB/c Mice. *Int J Parasitol* (1999) 29(10):1635–46. doi: 10.1016/S0020-7519(99)00141-1
71. Long MT, Baszler TV, Mathison BA. Comparison of Intracerebral Parasite Load, Lesion Development, and Systemic Cytokines in Mouse Strains Infected With *Neospora Caninum*. *J Parasitol* (1998) 84(2):316–20. doi: 10.2307/3284489
72. Yamane I, Kitani H, Kokuho T, Shibahara T, Haritani M, Hamaoka T, et al. The Inhibitory Effect of Interferon Gamma and Tumor Necrosis Factor Alpha on Intracellular Multiplication of *Neospora Caninum* in Primary Bovine Brain Cells. *J Vet Med Sci* (2000) 62(3):347–51. doi: 10.1292/jvms.62.347
73. Pinheiro AM, Costa SL, Freire SM, Meyer R, Almeida MAOD, Tardy M, et al. *Neospora Caninum*: Infection Induced IL-10 Overexpression in Rat Astrocytes *In Vitro*. *Exp Parasitol* (2006) 112(3):193–7. doi: 10.1016/j.exppara.2005.10.008
74. Pinheiro AM, Costa SL, Freire SM, Ribeiro CSO, Tardy M, El-Bachá RDS, et al. *Neospora Caninum*: Early Immune Response of Rat Mixed Glial Cultures After Tachyzoites Infection. *Exp Parasitol* (2010) 124(4):442–7. doi: 10.1016/j.exppara.2009.12.018
75. Jesus EEV, Pinheiro AM, Santos AB, Freire SM, Tardy MB, El-Bachá RDS, et al. Effects of IFN- γ , TNF- α , IL-10 and TGF- β on *Neospora Caninum* Infection in Rat Glial Cells. *Exp Parasitol* (2013) 133(3):269–74. doi: 10.1016/j.exppara.2012.11.016
76. De Jesus EEV, Santos ABD, Ribeiro CSO, Pinheiro AM, Freire SM, El-Bachá RS, et al. Role of IFN- γ and LPS on Neuron/Glial Co-Cultures Infected by *Neospora Caninum*. *Front Cell Neurosci* (2014) 8:340. doi: 10.3389/fncel.2014.00340
77. Grangeiro MS, Santos CCD, Borges JMP, Sousa CDS, Freitas S, Argolo D, et al. Neuroprotection During *Neospora Caninum* Infection Is Related to the Release of Neurotrophic Factors BDNF and NGF. *J Parasitol* (2019) 105(2):313–20. doi: 10.1645/18-81
78. Chao MV. Neurotrophins and Their Receptors: A Convergence Point for Many Signalling Pathways. *Nat Rev Neurosci* (2003) 4(4):299–309. doi: 10.1038/nrn1078
79. Spekker K, Czesla M, Ince V, Heseler K, Schmidt SK, Schares G, et al. Indoleamine 2, 3-Dioxygenase Is Involved in Defense Against *Neospora Caninum* in Human and Bovine Cells. *Infect Immun* (2009) 77(10):4496–501. doi: 10.1128/IAI.00310-09
80. Jesus LB, Santos AB, Jesus EEV, Santos RGD, Grangeiro MS, Bispo-da-Silva A, et al. IDO, COX and iNOS Have an Important Role in the Proliferation of *Neospora Caninum* in Neuron/Glia Co-Cultures. *Vet Parasitol* (2019) 266:96–102. doi: 10.1016/j.vetpar.2019.01.003
81. Argolo D, Borges JM, Freitas L, Grangeiro MS, Pina G, da Silva VD, et al. Activation of the Kynurenine Pathway and Production of Inflammatory Cytokines by Astrocytes and Microglia Infected With *Neospora Caninum*. *Int J Tryptophan Res* (In Press). doi: 10.1177/11786469211069946. ID: TRY-2020-0033.RV2.
82. Tullius SG, Bieffer HR, Li S, Trachtenberg AJ, Edtinger K, Quante M, et al. NAD⁺ Protects Against EAE by Regulating CD4⁺ T-Cell Differentiation. *Nat Commun* (2014) 5:5101. doi: 10.1038/ncomms6101
83. Wang J, Zhao C, Kong P, Sun H, Sun Z, Bian G, et al. Treatment With NAD⁺ Inhibited Experimental Autoimmune Encephalomyelitis by Activating AMPK/SIRT1 Signaling Pathway and Modulating Th1/Th17 Immune Responses in Mice. *Int Immunopharmacol* (2016) 39:287–94. doi: 10.1016/j.intimp.2016.07.036
84. Groth B, Venkatakrisnan P, Lin SJ. NAD⁺ Metabolism, Metabolic Stress, and Infection. *Front Mol Biosci* (2021) 8:686412. doi: 10.3389/fmolb.2021.686412
85. Mota CM, Oliveira A, Davoli-Ferreira M, Silva MV, Santiago FM, Nadipuram SM, et al. *Neospora Caninum* Activates P38 MAPK as an Evasion Mechanism Against Innate Immunity. *Front Microbiol* (2016) 7:1456. doi: 10.3389/fmicb.2016.01456
86. Ma L, Liu G, Liu J, Li M, Zhang H, Tang D, et al. *Neospora Caninum* ROP16 Play an Important Role in the Pathogenicity by Phosphorylating Host Cell STAT3. *Vet Parasitol* (2017) 243:135–47. doi: 10.1016/j.vetpar.2017.04.020
87. Bradley PJ, Sibley LD. Rhoptries: An Arsenal of Secreted Virulence Factors. *Curr Opin Microbiol* (2007) 10(6):582–7. doi: 10.1016/j.mib.2007.09.013

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Del'Arco, Argolo, Guillemín, Costa, Costa and Pinheiro. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.