



Corrigendum: Eosinophils and Neutrophils Eliminate Migrating *Strongyloides ratti* Larvae at the Site of Infection in the Context of Extracellular DNA Trap Formation

Alexandra Ehrens¹, Nikolas Rüdiger², Lennart Heepmann², Lara Linnemann², Wiebke Hartmann², Marc P. Hübner^{1,3*} and Minka Breloer^{2,4*}

¹ Institute for Medical Microbiology, Immunology and Parasitology, University Hospital Bonn, Bonn, Germany, ² Section of Molecular Biology and Immunology, Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany, ³ German Center for Infection Research (DZIF), Partner site Bonn-Cologne, Bonn, Germany, ⁴ Department of Biology, University of Hamburg, Hamburg, Germany

OPEN ACCESS

Edited and reviewed by:

Tamás Laskay,
University of Lübeck, Germany

*Correspondence:

Minka Breloer
Breloer@bnitm.de
Marc P. Hübner
huebner@uni-bonn.de

Specialty section:

This article was submitted to
Parasite Immunology,
a section of the journal
Frontiers in Immunology

Received: 18 February 2022

Accepted: 09 March 2022

Published: 21 March 2022

Citation:

Ehrens A, Rüdiger N, Heepmann L,
Linnemann L, Hartmann W,
Hübner MP and Breloer M (2022)
Corrigendum: Eosinophils and
Neutrophils Eliminate Migrating
Strongyloides ratti Larvae at the
Site of Infection in the Context of
Extracellular DNA Trap Formation.
Front. Immunol. 13:878640.
doi: 10.3389/fimmu.2022.878640

Keywords: nematodes, L3, eosinophils, neutrophils, extracellular DNA traps, ETosis, MPO, *Strongyloides*

A Corrigendum on:

Eosinophils and Neutrophils Eliminate Migrating *Strongyloides ratti* Larvae at the Site of Infection in the Context of Extracellular DNA Trap Formation

By Ehrens A, Rüdiger N, Heepmann L, Linnemann L, Hartmann W, Hübner MP and Breloer M (2021). *Front. Immunol.* 12:715766. doi: 10.3389/fimmu.2021.715766

In the original article, there was a mistake in **Figure 6** as published. **The units of the axis was indicated in µg/ml and should be corrected to ng/ml.** The corrected **Figure 6** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Ehrens, Rüdiger, Heepmann, Linnemann, Hartmann, Hübner and Breloer. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 6 | *S. ratti* L3 induce release of MPO and MMP-9-containing granules. **(A)** MPO and **(B)** MMP-9 concentration measured in the supernatant of neutrophils stimulated with plasma from *S. ratti*-infected or naive animals, *S. ratti* L3 larvae alone or in combination with plasma. Plasma has been left untreated or antibody-depleted (Δ Ab). Shown are box plots with tukey with $n=5$. One out of two independent experiment. One-Way ANOVA with Bonferroni post-hoc test. Only statistically significant comparisons are depicted, all other comparisons were statistically not significant. $p < 0.01$ **, $p < 0.001$ ***.