Check for updates

OPEN ACCESS

EDITED AND REVIEWED BY Yanpei Zhuang, Jimei University, China

*CORRESPONDENCE Jakob Karl Giesler Øjakob.giesler@awi.de

RECEIVED 25 February 2025 ACCEPTED 04 March 2025 PUBLISHED 21 March 2025

CITATION

Giesler JK, Lemley DA, Adams JB and Moorthi SD (2025) Corrigendum: Interactive effects of salinity, temperature and food web configuration on performance and harmfulness of the raphidophyte *Heterosigma akashiwo*. *Front. Mar. Sci.* 12:1583268. doi: 10.3389/fmars.2025.1583268

COPYRIGHT

© 2025 Giesler, Lemley, Adams and Moorthi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Interactive effects of salinity, temperature and food web configuration on performance and harmfulness of the raphidophyte *Heterosigma akashiwo*

Jakob Karl Giesler (1,2*, Daniel Alan Lemley (1,2*, Janine Barbara Adams (1,2*, and Stefanie Devi Moorthi (1,2*)

¹ICBM, Institute for Chemistry and Biology of the Marine Environment, Carl von Ossietzky University of Oldenburg, Oldenburg, Germany, ²AWI, Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany, ³DSI-NRF Research Chair in Shallow Water Ecosystems, Department of Botany, Nelson Mandela University, Gqeberha, South Africa, ⁴Institute for Coastal and Marine Research, Nelson Mandela University, Gqeberha, South Africa

KEYWORDS

Heterosigma akashiwo, harmful algal bloom, estuaries, temperature, salinity, mucus production, selective grazing

A Corrigendum on

Interactive effects of salinity, temperature and food web configuration on performance and harmfulness of the raphidophyte *Heterosigma akashiwo*

By Giesler JK, Lemley DA, Adams JB and Moorthi SD (2023) Front. Mar. Sci. 10:1244639. doi: 10.3389/fmars.2023.1244639

In the published article, there was an error in **Figure 5** as published. The figure showed population grazing rates instead of mortality rates. The corrected **Figure 5** and its caption "Mortality of *B. plicatilis* over time when exposed to *H. akashiwo*, *H. rotundata* or both algae species in combination (indicated by color) across two levels of salinity (15 and 30) represented by facet grid columns" appear below.

In the published article, there was an error. The research permit number was missing in the Acknowledgements.

A correction has been made to the **Acknowledgments** section, Paragraph 1. This sentence previously stated:

"We thank Heike Rickels (ICBM Terramare, Wilhelmshaven, Germany) for analysis of dissolved and particulate nutrient samples as well as technical support in the lab. Furthermore, we also thank Jan-Claas Dajka (HIFMB, Oldenburg, Germany) for advice in data analysis and Cedric Meunier (AWI, Helgoland, Germany) for providing the *A. tonsa* eggs."

The corrected sentence appears below:

"We thank Heike Rickels (ICBM Terramare, Wilhelmshaven, Germany) for analysis of dissolved and particulate nutrient samples as well as technical support in the lab. We also thank Jan-Claas Dajka (HIFMB, Oldenburg, Germany) for advice in data analysis and Cedric Meunier (AWI, Helgoland, Germany) for providing the *A. tonsa* eggs. Furthermore, we thank the South African department of forestry, fisheries & the environment for permitting this research (permit reference number: RES2023/62)."

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.



Mortality of *B. plicatilis* over time when exposed to *H. akashiwo, H. rotundata* or both algae species in combination (indicated by color) across two levels of salinity (15 and 30) represented by facet grid columns.