



Corrigendum: Variations in Vaginal, Penile, and Oral Microbiota After Sexual Intercourse: A Case Report

Miguel Carda-Diéguez¹, Nivia Cárdenas², Marina Aparicio², David Beltrán³, Juan M. Rodríguez² and Alex Mira^{1,4*}

OPEN ACCESS

Edited and reviewed by:

Zhongheng Zhang,
Zhejiang University, China

*Correspondence:

Alex Mira
mira_ale@gva.es

Specialty section:

This article was submitted to
Infectious Diseases - Surveillance,
Prevention and Treatment,
a section of the journal
Frontiers in Medicine

Received: 18 October 2019

Accepted: 29 November 2019

Published: 09 January 2020

Citation:

Carda-Diéguez M, Cárdenas N,
Aparicio M, Beltrán D, Rodríguez JM
and Mira A (2020) Corrigendum:
Variations in Vaginal, Penile, and Oral
Microbiota After Sexual Intercourse: A
Case Report. *Front. Med.* 6:294.
doi: 10.3389/fmed.2019.00294

¹ Department of Health and Genomics, Center for Advanced Research in Public Health, FISABIO, Valencia, Spain,

² Department of Nutrition and Food Science, Complutense University of Madrid, Madrid, Spain, ³ Centro de Diagnóstico Médico, Ayuntamiento de Madrid, Madrid, Spain, ⁴ Network of Epidemiology and Public Health, CIBERESP, Madrid, Spain

Keywords: microbiota, vagina, penile, oral, oral sex, bacterial vaginosis, *Lactobacillus*

A Corrigendum on

Variations in Vaginal, Penile, and Oral Microbiota After Sexual Intercourse: A Case Report
by Carda-Diéguez, M., Cárdenas, N., Aparicio, M., Beltrán, D., Rodríguez, J. M., and Mira, A. (2019).
Front. Med. 6:178. doi: 10.3389/fmed.2019.00178

In the original article, there was a mistake in **Figure 1** as published. The penile swab compositions for H₂O₂ and 15H₂O₂ were mistakenly swapped. Similarly, the vaginal swab compositions for H₂O₂ and 15H₂O₂ were also mistakenly swapped. The corrected **Figure 1** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2020 Carda-Diéguez, Cárdenas, Aparicio, Beltrán, Rodríguez and Mira. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 1 | Microbial composition of penile and vaginal swabs. Most abundant genus were plotted in different colors. NoS, microbiota without sexual intercourse; BeS, immediately before sexual encounter; AfUS, 4 days after unprotected sexual encounter, no oral sex and with ejaculation inside the vagina; AfPS, immediately after vaginal sex with condom and oral sex; H₂O₂ after 1 day of H₂O₂ treatment; 15H₂O₂, after 15 days of intermittent H₂O₂ treatment. Male and female samples are indicated with the corresponding symbol.