



Integrated Care for Older People Screening Tool for Measuring Intrinsic Capacity: Preliminary Findings From ICOPE Pilot in China

Lina Ma^{1*}, Jagadish K. Chhetri², Yaxin Zhang¹, Pan Liu¹, Yumeng Chen¹, Yun Li¹ and Piu Chan^{1,2*}

¹ Department of Geriatrics, National Research Center for Geriatric Medicine, Xuanwu Hospital, Capital Medical University, Beijing, China, ² Department of Neurology and Neurobiology, Xuanwu Hospital, Capital Medical University, Beijing, China

OPEN ACCESS

Edited by:

Leocadio Rodriguez-Mañas,
Hospital de Getafe, Spain

Reviewed by:

John Edward Morley,
Saint Louis University, United States
Hidenori Arai,
National Center for Geriatrics and
Gerontology (NCGG), Japan
Luis Miguel Gutierrez Robledo,
Instituto Nacional de Geriatria, Mexico

*Correspondence:

Piu Chan
pbchan90@gmail.com
Lina Ma
malina0883@126.com

Specialty section:

This article was submitted to
Geriatric Medicine,
a section of the journal
Frontiers in Medicine

Received: 25 June 2020

Accepted: 30 October 2020

Published: 30 November 2020

Citation:

Ma L, Chhetri JK, Zhang Y, Liu P,
Chen Y, Li Y and Chan P (2020)
Integrated Care for Older People
Screening Tool for Measuring Intrinsic
Capacity: Preliminary Findings From
ICOPE Pilot in China.
Front. Med. 7:576079.
doi: 10.3389/fmed.2020.576079

Objectives: The World Health Organization (WHO) proposed the Integrated Care for Older People (ICOPE) screening tool to identify older people with priority conditions associated with declines in intrinsic capacity (IC). We aimed to determine the clinical utility of the WHO ICOPE screening tool in a Chinese population.

Method: A total of 376 adults aged 68.65 ± 11.41 years participated in the study. IC was assessed with the WHO ICOPE screening tool, covering five domains: cognitive, locomotor, sensory, vision, and psychological capacity. We assessed the activities of daily living (ADL); instrumental activities of daily living (IADL); the Fried frailty phenotype; FRAIL scale; Strength, Assistance With Walking, Rising From chair, Climbing Stairs, and Falls (SARC-F) scale; Mini-mental State Examination (MMSE); Geriatric Depression Scale (GDS); social frailty; and quality of life.

Results: There were 260 (69.1%) participants who showed declines in one or more IC dimensions. The percentages of decline in mobility, cognition, vitality, hearing, vision, and psychological capacity were 25.3, 46.8, 16.2, 15.4, 11.7, and 12.0%, respectively. IC decreased with increasing age. After adjusting for age, sex, and multimorbidity, participants with declines in IC were more likely to be older, frail, and disabled. They also had worse physical, mental, and overall health. There was a higher prevalence of declines in IC in participants with frailty. After adjusting for age, IC was positively correlated with walking speed, resilience score, and MMSE score and negatively correlated with frailty, SARC-F score, IADL score, GDS score, and physical and mental fatigue. The IC score was not associated with body composition variables such as fat-free mass, body fat percentage, or visceral fat area. Higher IC was associated with better quality of life. The area under the curve of the receiver operating characteristic (AUC-ROC) for the ICOPE screening tool vs. Fried phenotype, FRAIL, ADL disability, IADL disability, and SARC-F were 0.817, 0.843, 0.954, 0.912, and 0.909, respectively.

Conclusion: Our research affirms that the ICOPE screening tool is useful to identify adults with poor physical and mental function in a Chinese sample. This tool may assist in identifying declines in IC in an integrative care model and help slow down function decline and onset of care dependence.

Keywords: intrinsic capacity, frailty, older adults, screening, integrated care

INTRODUCTION

The increase in the aging population has emerged as a major global phenomenon. Frailty is an important construct that has been used in geriatric medicine over the past 20 years to conceptualize the age-related increase in vulnerability at an individual level (1, 2). Frailty is defined as a geriatric syndrome characterized by the reduced ability of an individual to maintain physiological homeostasis and increased risks of adverse clinical outcomes (3). Although there is a growing interest regarding frailty related to all geriatric syndromes and diseases, integration of this concept in clinical practice has been impeded due to a lack of unified frailty instruments and variations in research outcomes (4–6).

The disease concept is increasingly being replaced by a functional approach to address the healthcare needs of the older people (7). To overcome the weaknesses of the frailty concept and to better disseminate geriatric care to the aging population, the World Health Organization (WHO) recently proposed a novel model for healthy aging oriented around trajectories of functional ability (8). This model holds great potential to make a significant public health impact, particularly by aiding in healthy aging. The World Report on Aging and Health defines healthy aging as the process of developing and maintaining the functional ability that fosters well-being in old age. Functional ability is proposed to be determined by an individual's environmental factors, physical and mental attributes known as intrinsic capacity (IC) and the interaction between them (9). As the composite of all physical and mental capacities, IC could decline with age due to underlying disease and the aging process. While traditional healthcare models focus on the identification and treatment of medical conditions, IC emphasizes on the positive attributes of individuals, i.e., focuses on maintaining an individual's functional capacities throughout life course rather than treating a particular illness.

IC can be evaluated using five domains: cognition, locomotion, vitality, sensory, and psychological capacities (10). The five domains of IC are related to the reduction in physical and mental capacities and independently predict mortality and care dependence in older adults (10). Thus, it could be intuitive to consider that enhancing IC through life could potentially prevent adverse events and geriatric conditions such as frailty in old age. The WHO Guidelines on Integrated Care for Older People (ICOPE) offers evidence-based recommendations for effective interventions to address declines in physical and mental capacities in older people so as to optimize IC (11). However, questions pertaining to translate theory into practice arise because there is less agreement on how the domains of IC are assessed (12). Very few evidence on the clinical utility of the

ICOPE tool is available. International validation studies of the instrument and how to incorporate IC assessment in the new care model are ongoing. It remains unclear how the specific domains, assessed by the new tool, relate to physical and mental functions as well as underlying biological pathways. Thus, in this pilot study, we aimed to determine if the WHO ICOPE screening tool can identify adults with poor physical and mental function in a Chinese population.

METHODS

Study Design and Participants

This is a cross-sectional study. A total of 376 participants aged 50–97 years admitted to the Department of Geriatrics in Xuanwu Hospital Capital Medical University during 2019 were included. The participants consisted of 224 male and 152 female, with an average age of 68.65 ± 11.41 years. Inclusion criteria consisted of relatively healthy participants without acute illness and aged ≥ 50 years who were admitted for a physical examination and completed the WHO ICOPE screening assessment. Participants who suffered from acute heart failure, acute infection, acute cerebrovascular disease, severe cardiac, liver or kidney dysfunction, dementia, Parkinson's disease, and mental illnesses or were unable to provide informed consent because of aphasia, deafness, and blindness were excluded. All the participants underwent physical examination and comprehensive geriatric assessment. The clinical and demographic variables were collected using a face-to-face questionnaire by trained staff. All subjects gave written informed consent in accordance with the Declaration of Helsinki, and the study was approved by the ethical review board of Xuanwu Hospital Capital Medical University.

Measures

IC Construct

IC was assessed by the WHO ICOPE screening tool and included the following five domains (11):

- (1) Cognitive decline was determined if participants provided an incorrect response to either of the two questions on orientation in time and space or if they could not recall the three words they were asked to remember.
- (2) Limited mobility was defined as being unable to complete five chair rises within 14 s.
- (3) Malnutrition was defined as weight loss (more than 3 kg over the previous 3 months) or appetite loss.
- (4) Sensory loss: (a) Visual impairment was defined as any problems experienced with their eyes, difficulties in seeing far, reading, eye diseases, or currently under medical treatment; (b) hearing loss was defined as failing to hear whispers in the whisper test.
- (5) Depressive symptoms were defined as the participants being bothered by feeling down, feeling depressed or hopeless, or having little interest or pleasure in doing things over the preceding 2 weeks. The impairment of each item was scored as 0 points.

The IC score ranged from 0 to 6, with a higher score representative of better IC, i.e., for the six items indicated above

Abbreviations: ADL, activities of daily living; BFP, body fat percentage; BMD, bones mineral density; BMI, body mass index; CCI, Charlson Comorbidity Index; CRP, C-reactive protein; DBP, diastolic blood pressure; EQ-VAS, the EQ visual analog scale; GDS, Geriatric Depression Scale; IADL, instrumental activities of daily living; IC, intrinsic capacity; ICOPE, Integrated Care for Older people; MMSE, Mini-mental State Examination; PEF, peak expiratory flow; RS, resilience scale; SARC-F, Strength, Assistance With Walking, Rising From Chair, Climbing Stairs, and Falls; SBP, systolic blood pressure; SF, social frailty; WHO, World Health Organization.

participants were scored either 0 (representing decline) or 1 (representing no decline).

Frailty Assessment

Frailty was measured with the frailty phenotype assessment that includes five components: slowness, weakness, exhaustion, weight loss, and low activity (13). Participants who exhibited three or more of these components were classified as frail. The cutoff value for weakness was determined by grip strength after adjusting for sex and body mass index (BMI) (14). Walking speed was measured by a 4-m walk test, and the cutoff value for slowness was determined after adjustment for sex and height (15). Exhaustion was indicated by self-rated responses to questions from the Center for Epidemiological Studies Depression Scale: “I felt that everything I did was an effort” or “I could not get going” (16). Weight loss was defined as a self-reported unintentional weight loss of 3 kg in the past 6 months or as having a BMI <18.5 kg/m². Low activity was defined as self-reported exercise for <3 h/week over the past 12 months (17). The FRAIL scale was used to identify frailty, with individuals scoring three and above being identified as frail (18).

Physical Function Assessment

Functional ability was assessed by the Barthel index for activities of daily living (ADL) and instrumental activities of daily living (IADL). An individual’s performance on each item was classified as independent, partially dependent, and completely dependent and scored as 1, 2, and 3, respectively. Chronic diseases were defined as self-reported history of chronic disease diagnosed by a doctor. Charlson Comorbidity Index (CCI) was calculated (19). Polypharmacy was defined as taking more than five drugs. The Strength, Assistance with walking, Rising from chair, Climbing stairs, and Falls (SARC-F) questionnaire was used to assess sarcopenia, with higher scores indicating more severe sarcopenia (20).

Psychological Function Assessment

Depression was assessed with the 30-item Geriatric Depression Scale (GDS) (21). Cognitive function was evaluated using the Mini-mental State Examination (MMSE) (22). The 25-item Resilience Scale (RS) was used to assess psychological resilience (23), with higher scores reflecting greater resilience.

Social Function Assessment

Social frailty (SF) was assessed by the HALFT scale, which was validated in the Beijing Longitudinal Study of Aging and consists of the following five items: unhelpful to others, limited social participation, loneliness, financial difficulty, and not having anyone to talk to. The total score ranges from 0 to 5 points: a score of 0 was considered no SF or robust social function, 1–2 was considered pre-SF, and a score of ≥ 3 indicated SF (24).

Quality of Life

The health-related quality of life questionnaire 5-level EQ-5D (EQ-5D-5L) and the EQ visual analog scale (EQ-VAS) were used to measure generic health status; a higher EQ-VAS score indicates better health status (25). The Pittsburgh Fatigability Scale was used to measure physical and mental fatigue (26).

Organ Function Assessment

Heart rate was measured and recorded once every 20 min for 24 h using the active electrocardiography and disassembled by professionals. Peak expiratory flow (PEF) was determined using a spirometer. Bones mineral density (BMD) was determined using a dual-energy X-ray absorptiometry (DEXA) scan (LUNAR iDXA, USA). A single experienced technologist performed all the DEXA scans. BMD was measured at the lumbar spine vertebra L2–L4, femoral neck, and total hip. Bioelectrical impedance analysis, to estimate body composition, was carried out with the ioi353 Composition Analyzer (JAWON, Korea) with 8 points of tactile electrodes according to the manufacturer’s guidelines; fat mass, fat-free mass, body fat percentage (BFP), and visceral fat area were obtained.

Blood Test

Peripheral blood samples were collected after overnight fasting; serum samples were isolated and were stored at -80°C before use. C-reactive protein (CRP) levels were measured using immunoturbidimetry.

Statistical Analysis

Data were presented as mean \pm standard deviation or mean rank or number and percentage. The difference in the characteristics between the two groups was evaluated using the chi-square test for categorical variables, independent *t*-test for continuous variables, and Kruskal–Wallis comparisons for the abnormally distributed continuous variables. A multivariate logistic regression analysis—adjusted for age, sex, and multimorbidity—was used to identify the factors associated with lower IC. Spearman correlation coefficients were calculated to assess the relationship between IC and the other factors. The area under the curve of the receiver operating characteristic (AUC-ROC) was calculated. All statistical analyses were performed with SPSS (Chicago, IL, USA, version 20.0) or GraphPad Prism 7.0 software (GraphPad Software Inc., CA, USA). *p*-value was considered significant at <0.05 .

RESULTS

Of the 376 participants, the average IC score was 4.73 ± 1.27 , 116 participants (30.9%) were categorized as having no decline in IC. Of the participants, 137 (36.4%), 64 (17.0%), 35 (9.3%), 14 (3.7%), 7 (1.9%), and 3 (0.8%) showed decline in one, two, three, four, five, and six domains, respectively. The percentage of decline in mobility, cognition, vitality, hearing, vision, and psychological domains were 25.3, 46.8, 16.2, 15.4, 11.7, and 12.0%, respectively. Participants’ IC score decreased with increasing age, from 5.32 ± 0.79 at age 50–59 to 4.01 ± 1.56 at age 80 and older (Figure 1A). Slow walking speed and low grip strength were associated with worse IC (Figures 1C,D). There was no difference observed in IC between men and women (Figure 1B). Those with physical frailty and social frailty had decreased IC when compared to the nonfrail participants (Figures 1E,F). After adjusting for age,




FIGURE 1 | Comparison of intrinsic capacity score between different groups. **(A)** Comparison of intrinsic capacity score among the different age groups; **(B)** comparison of intrinsic capacity score between male and female; **(C)** comparison of intrinsic capacity score between weakness (low grips strength) and normal grip strength; **(D)** comparison of intrinsic capacity score between slowness (slow walking speed) and normal walking speed; **(E)** comparison of intrinsic capacity score among robust, prefrailty, and frailty, assessed by Fried phenotype; **(F)** comparison of intrinsic capacity score among normal, presocial frailty, and social frailty, assessed by HALFT scale. IC, intrinsic capacity; SF, social frailty.

these associations remained unchanged, except for grip strength ($p = 0.128$).

As shown in **Table 1**, participants with declines in IC were older, had more chronic diseases, and had worse general health as indicated by the EQ-VAS. Participants with declines in IC had worse physical function as indicated by lower Barthel index,

walk speed, grip strength, and physical fatigue. They had worse mental function indicated by lower MMSE scores, had higher GDS scores, more mental fatigue, and had worse social function as indicated by higher SF score. There was a higher prevalence of declines in IC in participants with polypharmacy and frailty. However, IC was not affected by sex, CCI, and resilience score.

TABLE 1 | Characteristics of the DIC and non-DIC participants.

	Non-DIC (n = 116)	DIC (n = 260)	x ² /t/z	p-value	Adjusted p-value*
Age (years)	62.47 ± 9.40	71.41 ± 11.16	-8.026	<0.001	-
Gender (n, %)					
Male	71 (31.7)	153 (68.3)	0.186	0.667	
Female	45 (29.6)	107 (70.4)			
Number of chronic diseases	4.03 ± 2.91	4.97 ± 2.96	-2.815	0.005	-
CCI	3 (1,4)	3 (2,4)	1.464	0.226	-
EQ-VAS	74.81 ± 12.27	68.63 ± 16.17	6.177	<0.001	0.031
Barthel index	98.30 ± 3.53	91.24 ± 15.39	4.474	<0.001	0.735
IADL score	7.04 ± 0.30	8.34 ± 2.95	-6.992	<0.001	0.039
Walking speed (m/s)	1.00 (0.85, 1.14)	0.92 (0.72, 1.08)	8.298	0.003	0.995
Grip strength (kg)	31.0 (25.2, 40.7)	25.1 (19.9, 35.3)	13.392	<0.001	0.531
MMSE	29.23 ± 1.31	27.22 ± 2.65	9.700	<0.001	<0.001
GDS	2 (1,4)	4 (2,7)	18.898	<0.001	<0.001
Resilience	140.21 ± 15.82	136.83 ± 20.86	1.623	0.106	-
Polypharmacy (n, %)					
Yes	38 (20.2)	150 (79.8)			
No	78 (41.5)	110 (58.5)	19.947	<0.001	0.173
Fried frailty (n, %)					
Robust	59 (43.4)	77 (56.6)			
Prefrail	55 (29.7)	130 (70.3)	36.168	<0.001	0.014
Frail	2 (3.6)	53 (96.4)			
Social frailty score	0.90 ± 0.88	1.26 ± 1.01	-3.198	0.002	0.120
Physical fatigue	13 (8.5, 23)	23 (14,33)	-5.556	<0.001	0.046
Mental fatigue	11 (0, 21)	14 (1,30)	-2.921	0.004	0.824
FRAIL score	0 (0, 1)	1 (0, 1)	24.428	<0.001	0.018
SARC-F score	0 (0, 0)	0 (0, 2)	10.628	0.001	0.353

Data were expressed as mean (standard deviation), median (interquartile range) or n (%).

*Adjusted by age, sex, and multimorbidity.

DIC, decline in intrinsic capacity; CCI, Charlson Comorbidity Index; EQ-VAS, the EQ visual analog scale; IADL, instrumental activities of daily living; MMSE, the Mini-mental State Examination; GDS, the Geriatric Depression Scale; SARC-F, Strength, Assistance With Walking, Rising From Chair, Climbing Stairs, and Falls.

After adjusting for age, sex, and multimorbidity, a decline in IC was associated with declining physical function (Fried frailty, FRAIL score, IADL score, and physical fatigue), mental health (MMSE and GDS), and quality of life (EQ-VAS).

Next, we explored whether declines in IC affects organ function. **Table 2** shows that participants with declines in IC had higher systolic blood pressure (SBP), higher 24 h heart rate, higher CRP, and lower diastolic blood pressure (DBP). However, no difference was observed after adjusting for age, sex, and multimorbidity. There was no difference in EF, PEF, BMD, and body composition.

We further examined the correlation between IC and physical (**Figure 2**) and mental functions (**Figure 3**). IC was significantly positively correlated with the Barthel index, grip strength, walking speed, resilience score, and MMSE score and significantly negatively correlated with age, IADL score, frailty, GDS score, physical fatigue, mental fatigue, and worse quality of life. Further analysis based on chronic diseases showed that the IC score was positively correlated with DBP and negatively correlated with CCI, number of chronic diseases, number of drugs, and SBP (**Figure 4**). However, after adjusting for age, IC

was positively correlated with walking speed ($r = 0.168$, $p = 0.002$), resilience score ($r = 0.316$, $p < 0.001$), and MMSE ($r = 0.358$, $p < 0.001$), while it was negatively correlated with IADL score ($r = -0.446$, $p < 0.001$), Fried frailty score ($r = -0.398$, $p < 0.001$), FRAIL score ($r = -0.365$, $p < 0.001$), SARC-F score ($r = -0.347$, $p < 0.001$), physical fatigue ($r = -0.278$, $p < 0.001$), mental fatigue ($r = -0.195$, $p = 0.001$), and GDS ($r = -0.552$, $p < 0.001$). We further explored the relationship between IC and body composition; however, IC was not associated with fat-free mass, BFP, or visceral fat area.

AUC-ROC was performed to explore the performance of the ICOPE screening tool with respect to frailty, IADL, and ADL. As shown in **Table 3**, using a cutoff of 4, the sensitivity and specificity of the ICOPE screening tool in identifying frailty defined by Fried phenotype were 90.3 and 50.9%, respectively, and the AUC-ROC for the ICOPE screening tool vs. Fried phenotype was 0.817 ($p < 0.001$). The sensitivity, specificity, and AUC-ROC for the ICOPE screening tool vs. ADL disability were 85.4%, 100%, and 0.954 ($p < 0.001$), respectively, while the sensitivity, specificity, and AUC-ROC for the ICOPE screening tool vs. IADL disability were 90.8%, 73.7%, and 0.912 ($p < 0.001$), respectively.


TABLE 2 | Comparison of organ functions between the DIC and non-DIC participants.

	Non-DIC (n = 116)	DIC (n = 260)	x ² /t/z	p-value	Adjusted p-value*
SBP (mmHg)	132.43 ± 15.37	137.44 ± 18.79	-2.719	0.007	0.189
DBP (mmHg)	77.66 ± 11.42	74.86 ± 11.58	2.179	0.030	0.705
BMI (kg/m ²)	25.55 ± 3.99	25.58 ± 3.69	-0.068	0.946	-
EF (%)	66.24 ± 6.22	65.91 ± 6.87	0.420	0.675	-
24 h HR (/min)	70.53 ± 10.63	66.86 ± 9.37	2.770	0.006	0.071
Fat (kg)	20.08 ± 4.94	20.39 ± 6.25	-0.306	0.760	-
Fat-free mass (kg)	46.51 ± 8.50	46.88 ± 10.50	-0.222	0.825	-
BFP (%)	28.81 ± 5.18	28.40 ± 6.18	0.445	0.657	-
Visceral fat area (cm ²)	116.08 ± 39.77	124.18 ± 42.82	-1.132	0.259	-
PEF (L/s)	5.46 (4.34,8.11)	5.05 (3.58,6.55)	0.159	0.690	-
BMD (g/cm ²)	0.86 ± 0.20	0.82 ± 0.17	0.908	0.367	-
CRP (mg/L)	2.30 (1.60,3.40)	2.67 (1.65,4.33)	3.955	0.026	0.239

Data were expressed as mean (standard deviation), median (interquartile range) or n (%).

*Adjusted by age, sex and multimorbidity.


DIC, decline in intrinsic capacity; SBP, systolic blood pressure; DBP, diastolic blood pressure; BMI, body mass index; EF, ejection fraction; HR, heart rate; BFP, body fat percentage; PEF, peak expiratory flow; BMD, bones mineral density; CRP, C-reactive protein.



DISCUSSION

In this pilot study, two-thirds of participants aged 50 years and above experienced one or more declines in IC. IC decreased with increasing age, which is in accordance with the IC model and rationale that an individual's capacities will decline with aging (27). Our results showed that participants with lower IC scores were more likely to be frail, disabled, and have worse quality of

life. Frailty due to multiple causes decreases homeostatic reserves and leads to accentuated vulnerability to negative outcomes in older adults (1). The Fried phenotype of frailty considered frailty as an underlying biological syndrome (13), while the frailty index describes frailty as age-related deficits accumulation (28). Interestingly, impairments in some domains in the IC construct are the same as some components of frailty. However, IC and frailty might represent the opposite sides of the same coin.



As we have emphasized before, the concept of IC was related to an individual's positive functional status, which is crucial to healthy aging.

This study showed that the IC score was positively correlated with physical function as demonstrated by measures such as walking speed and grip strength, which is consistent with findings from previous research (29). However, IC score was not associated with body composition variables such as fat-free mass, BFP, and visceral fat area; this further indicates that IC may have higher association with muscle function but not muscle mass. Individual's physical decline has been shown to predict mortality and has been considered as an "additional vital sign" in older adults (30). Grip strength predicts decline in physical and mental capacities in older adults. In accordance with our findings, a recent study in Colombia showed that participants with greater handgrip strength had better IC and lower risks of adverse events (29). Moreover, gait speed has a linear relationship with the risk of negative outcomes (31).

Our findings showed that the IC score was negatively correlated with the number of chronic diseases and CCI; moreover, participants with polypharmacy had lower IC. However, after adjusting for age, these factors were no longer statistically different. Previous research has shown that chronic diseases are common in older adults and associated with disability and mortality and that aging increased risk of multimorbidity (32). These factors may have a significant impact on the healthcare burden. However, a lack of geriatric professionals may lead to poor management (under/over

treatment) of chronic disease in areas with limited geriatric care resources. Monitoring the trajectories of individuals' physical and mental capacities by collecting preclinical parameters relative to the health profile instead of trying to manage an array of disease diagnoses might overcome such limitations. Data from the English Longitudinal Study on Aging showed that, adjusted for multimorbidity, IC was associated with subsequent functioning (10). The ICOPE approach may enable such a mode of healthcare management when changes in IC suggest the need for clinical intervention.

In our study, IC was positively correlated with MMSE scores and negatively correlated with GDS scores and mental fatigue. Declines in cognitive function are associated with high risks of negative outcomes, and previous studies have shown that cognition can directly affect the other domains of IC. Depressive symptoms have been shown to be strongly associated with functional status (33) and is a risk for subsequent physical decline, even among individuals with no ADL difficulties or mobility disability (34). SF was associated with physical functioning, cognition, and depression and was found to predict mortality in the Beijing Longitudinal Study of Aging (24). This implies that the interactions between mental capacity and social factors may largely contribute to the construct of IC.

Although the present study found that participants with lower IC had higher levels of CRP, the association was lost after adjusting for age. Identifying biomarkers of declines in IC may potentially help develop strategies aimed at preventing the incidence of age-related functional decline and to slow down




FIGURE 4 | Correlation between intrinsic capacity and chronic diseases. Spearman correlation coefficients were calculated between intrinsic capacity score and **(A)** Charlson Comorbidity Index, **(B)** Disease disturb, **(C)** number of chronic diseases, **(D)** number of drugs, **(E)** systolic blood pressure, **(F)** diastolic blood pressure. IC, intrinsic capacity; CCI, Charlson Comorbidity Index; SBP, systolic blood pressure; DBP, diastolic blood pressure.

the progression toward future disability and care dependency. CRP was significantly associated with worse physical function in Colombia, Brazil, and Canada (35), while we previously showed no association between CRP and frailty (36). Data from the Multidomain Alzheimer Preventive Trial showed that baseline low-grade inflammation was related to a 5-year declined IC in older adults (37). In addition, another study found allostatic load to be incrementally associated with IC, even when adjusting for socioeconomic variables and chronic diseases (38). Further studies with large samples on the chronic inflammation markers are needed.

This is the first study in the Chinese population implementing the ICOPE screening tool recommended by the WHO to measure IC. All five domains used in the study were based on the ICOPE guidelines, while in many other studies, the measures included

are neither complete nor unified. Furthermore, an inflammation marker was explored since few have operationalized this physiological condition by measuring CRP as we did. We also used the same simple approaches to determine functional deterioration—for example, using the FRAIL and SARC-F scales as validation instruments—and showed that the WHO ICOPE tool is effective in identifying older adults with frailty and sarcopenia. Future studies to determine the utility of the Rapid Geriatric Assessment tool (39) for determining declined IC are warranted. We also showed for the first time the AUC-ROC for the ICOPE screening tool vs. Fried phenotype, FRAIL, ADL disability, IADL disability, and SARC-F to be equally good. However, certain limitations that affect the interpretation of the results should be noted. First, although the participants recruited were relatively healthy and without acute illness, the inpatient

TABLE 3 | The estimating sensitivity, specificity, and AUC-ROC of the ICOPE screening tool.

	Sensitivity	Specificity	AUC	p-value
Frailty (FP)	0.903	0.509	0.817	<0.001
Frailty (FRAIL)	0.875	0.600	0.843	<0.001
ADL disability	0.854	1.000	0.954	<0.001
IADL disability	0.908	0.737	0.912	<0.001
Sarcopenia (SARC-F)	0.867	0.800	0.909	<0.001

AUC-ROC, under the curve of the receiver operating characteristic; ICOPE, the Integrated Care for Older People; FP, Fried phenotype; ADL, activities of daily living; IADL, instrumental activities of daily living; SARC-F, The Strength, Assistance With Walking, Rising From Chair, Climbing Stairs, and Falls.

population—with an average age of 68.65 ± 11.41 years—may not be representative of the older population in China. Although IC total score was positively correlated with the resilience score, there was no difference in resilience score between individuals with declines in IC and individuals without declines in IC; one potential explanation may be the participants we recruited were relatively healthy. Second, the study is cross-sectional and used a limited sample. Future follow-up studies with a large population should be conducted to validate the WHO ICOPE screening tool as well as explore the longitudinal changes in IC and its relevance to clinical outcomes such as incident disability and other adverse events.

CONCLUSION

Early identification of declines in IC is essential to implement interventions that maintain individuals' physical and mental capacities and function or to slow down or reverse the declines in older adults. Our research revealed the WHO ICOPE screening tool can identify those adults with poor physical and mental function in Chinese adults; this indicates that measurement of IC with commonly used instruments is feasible and provides useful information on an individual's physical and mental functioning. By using an integrated care approach, it may be possible to maintain IC and slow down the declines in mobility, nutrition, sensory functions, cognition, and mood, which, in turn, is

REFERENCES

- Dent E, Martin FC, Bergman H, Woo J, Romero-Ortuno R, Walston JD. Management of frailty: opportunities, challenges, and future directions. *Lancet*. (2019) 394:1376–86. doi: 10.1016/S0140-6736(19)31785-4
- Rodriguez-Mañas L, Fried L. Frailty in the clinical scenario. *Lancet*. (2015) 385:e7–9. doi: 10.1016/S0140-6736(14)61595-6
- Cesari M, Calvani R, Marzetti E. Frailty in older persons. *Clin Geriatr Med*. (2017) 33:293–303. doi: 10.1016/j.cger.2017.02.002
- Hoogendijk EO, Afilalo J, Ensrud KE, Kowal P, Onder G, Fried LP. Frailty: implications for clinical practice and public health. *Lancet*. (2019) 394:1365–75. doi: 10.1016/S0140-6736(19)31786-6
- Walston J, Robinson TN, Ziemann S, McFarland F, Carpenter CR, Althoff KN, et al. Integrating frailty research into the medical specialties—Report from a U13 Conference. *J Am Geriatr Soc*. (2017) 65:2134–9. doi: 10.1111/jgs.14902

important to prevent or delay the onset of care dependence. Further studies on the monitoring of IC trajectories and ICOPE approaches are warranted in multiple socioeconomic settings.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Ethical Review Board of Xuanwu Hospital Capital Medical University. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

LM and PC contributed to the design of the work. LM drafted the manuscript and wrote it together with JC and PC. PL, YZ, YC, and YL recruited the participants. LM, JC, PC, and YL contributed to the analysis and interpretation of data. All authors contributed to writing the paper and revising it critically and gave final approval of this version.

FUNDING

This work was supported by the following grants: National Key R&D Program of China (2018YFC1312001, 2017YFC0840105), Beijing Municipal Hospital Scientific Research Training Project (PX2020036), Beijing Municipal Administration of Hospitals' Mission Plan (SML20150803), and Beijing Municipal Science & Technology Commission (Z171100000117013).

ACKNOWLEDGMENTS

We thank Drs. Shijie Li, Ying Li, Tong Ji, Yu Song, Ou Zhao, Bixi Li, and Chuan Liu for their assistance in data collection, entry, and study coordination and all member of the team. We acknowledge all the people who participated in the study.

- The Lancet. Bringing frailty into all realms of medicine. *Lancet*. (2019) 394:1298. doi: 10.1016/S0140-6736(19)32279-2
- Michel JP, Graf C, Ma L, Ecartot F. The end of the disease concept in geriatric medicine. *Ageing Med Healthc*. (2020) 11:3–9. doi: 10.33879/AMH.2020.032-2002.006
- World Health Organization. *Ageing and Health*. (2018). Available online at: <https://www.who.int/ageing/health-systems/icope/en/> (accessed June 23, 2018).
- Beard JR, Officer A, De Carvalho IA, Sadana R, Pot AM, Michel JP, et al. The World report on ageing and health: a policy framework for healthy ageing. *Lancet*. (2016) 387:2145–54. doi: 10.1016/S0140-6736(15)0516-4
- Beard JR, Jotheeswaran AT, Cesari M, Araujo De Carvalho I. The structure and predictive value of intrinsic capacity in a longitudinal study of ageing. *BMJ Open*. (2019) 9:e026119. doi: 10.1136/bmjopen-2018-026119

11. WHO. WHO | *Integrated Care for Older People (ICOPE): Guidance for Person-Centred Assessment and Pathways in Primary Care*. (2019). Available online at: <https://www.who.int/ageing/health-systems/icope/en/> (accessed June 16, 2020).
12. de Barreto PS, Guyonnet S, Ader I, Andrieu S, Casteilla L, Davezac N, et al. The inspire research initiative: a program for geroscience and healthy aging research going from animal models to humans and the healthcare system. *J Frailty Aging*. (2020) 1–8. doi: 10.14283/jfa.2020.18
13. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol Ser A Biol Sci Med Sci*. (2001) 56:M146–57. doi: 10.1093/gerona/56.3.M.146
14. Wu C, Smit E, Xue Q-L, Odden MC. Prevalence and correlates of frailty among community-dwelling Chinese older adults: the China health and retirement longitudinal study. *J Gerontol A Biol Sci Med Sci*. (2017) 73:102–8. doi: 10.1093/geroni/igx004.895
15. Ma L, Zhang L, Sun F, Li Y, Tang Z. Cognitive function in Prefrail and frail community-dwelling older adults in China. *BMC Geriatr*. (2019) 19:53. doi: 10.1186/s12877-019-1056-8
16. Radloff L. The CES-D scale : a self-report depression scale for research in the general population. *Appl Psychol Meas*. (1977) 1:385–401. doi: 10.1177/014662167700100306
17. Ma L, Tang Z, Chan P, Walston JD. Novel frailty screening questionnaire (FSQ) predicts 8-year mortality in older adults in China. *J Frailty Aging*. (2019) 8:33–8. doi: 10.14283/jfa.2018.38
18. Morley JE, Malmstrom TK, Miller DK. A simple frailty questionnaire (FRAIL) predicts outcomes in middle aged African Americans. *J Nutr Heal Aging*. (2012) 16:601–8. doi: 10.1007/s12603-012-0084-2
19. Chan TC, Luk JKH, Chu LW, Chan FHW. Validation study of Charlson Comorbidity Index in predicting mortality in Chinese older adults. *Geriatr Gerontol Int*. (2014) 14:452–7. doi: 10.1111/ggi.12129
20. Woo J, Leung J, Morley JE. Validating the SARC-F: a suitable community screening tool for sarcopenia? *J Am Med Dir Assoc*. (2014) 15:630–4. doi: 10.1016/j.jamda.2014.04.021
21. Chan ACM. Clinical validation of the Geriatric Depression Scale (GDS): Chinese version. *J Aging Health*. (1996) 8:238–53. doi: 10.1177/089826439600800205
22. Li H, Jia J, Yang Z. Mini-mental state examination in elderly Chinese: a population-based normative study. *J Alzheimers Dis*. (2016) 53:487–96. doi: 10.3233/JAD-160119
23. Wagnild GM, Young HM. Development and psychometric evaluation of the resilience scale. *J Nurs Meas*. (1993) 1:165–78.
24. Ma L, Sun F, Tang Z. Social frailty is associated with physical functioning, cognition, and depression, and predicts mortality. *J Nutr Health Aging*. (2018) 22:989–95. doi: 10.1007/s12603-018-1054-0
25. Yang Z, Busschbach J, Liu G, Luo N. EQ-5D-5L norms for the urban Chinese population in China 11 Medical and Health Sciences 1117 Public Health and Health Services. *Health Qual Life Outcomes*. (2018) 16:210. doi: 10.1186/s12955-018-1036-2
26. Glynn NW, Santanasto AJ, Simonsick EM, Boudreau RM, Beach SR, Schulz R, et al. The Pittsburgh fatigability scale for older adults: Development and validation. *J Am Geriatr Soc*. (2015) 63:130–5. doi: 10.1111/jgs.13191
27. Cesari M, Araujo de Carvalho I, Amuthavalli Thiyagarajan J, Cooper C, Martin FC, Reginster J-Y, et al. Evidence for the domains supporting the construct of intrinsic capacity. *J Gerontol Ser A*. (2018) 73:1653–60. doi: 10.1093/gerona/gly011
28. Mitnitski AB, Mogilner AJ, Rockwood K. Accumulation of deficits as a proxy measure of aging. *Sci World J*. (2001) 1:323–36. doi: 10.1100/tsw.2001.58
29. Ramírez-Vélez R, Correa-Bautista JE, García-Hermoso A, Cano CA, Izquierdo M. Reference values for handgrip strength and their association with intrinsic capacity domains among older adults. *J Cachexia Sarcopenia Muscle*. (2019) 10:278–86. doi: 10.1002/jcsm.12373
30. Studenski S, Perera S, Wallace D, Chandler JM, Duncan PW, Rooney E, et al. Physical performance measures in the clinical setting. *J Am Geriatr Soc*. (2003) 51:314–22. doi: 10.1046/j.1532-5415.2003.51104.x
31. Sourial N, Wolfson C, Bergman H, Zhu B, Karunanathan S, Quail J, et al. A correspondence analysis revealed frailty deficits aggregate and are multidimensional. *J Clin Epidemiol*. (2010) 63:647–54. doi: 10.1016/j.jclinepi.2009.08.007
32. Prince MJ, Wu F, Guo Y, Gutierrez Robledo LM, O'Donnell M, Sullivan R, et al. The burden of disease in older people and implications for health policy and practice. *Lancet*. (2015) 385:549–62. doi: 10.1016/S0140-6736(14)61347-7
33. Vaughan L, Goveas J, Corbin A. Depression and frailty in later life: a systematic review. *Clin Interv Aging*. (2015) 10:1947. doi: 10.2147/CIA.S69632
34. Penninx BWJH, Guralnik JM, Ferrucci L, Simonsick EM, Deeg DJH, Wallace RB. Depressive symptoms and physical decline in community-dwelling older persons. *J Am Med Assoc*. (1998) 279:1720–6. doi: 10.1001/jama.279.21.1720
35. Matheï C, Vaes B, Wallemacq P, Degryse J. Associations between cytomegalovirus infection and functional impairment and frailty in the BELFRAIL Cohort. *J Am Geriatr Soc*. (2011) 59:2201–8. doi: 10.1111/j.1532-5415.2011.03719.x
36. Ma L, Sha G, Zhang Y, Li Y. Elevated serum IL-6 and adiponectin levels are associated with frailty and physical function in Chinese older adults. *Clin Interv Aging*. (2018) 13:2013–20. doi: 10.2147/CIA.S180934
37. Giudici KV, de Souto Barreto P, Guerville F, Beard J, Araujo de Carvalho I, Andrieu S, et al. Associations of C-reactive protein and homocysteine concentrations with the impairment of intrinsic capacity domains over a 5-year follow-up among community-dwelling older adults at risk of cognitive decline. (MAPT Study). *Exp Gerontol*. (2019) 127:110716. doi: 10.1016/j.exger.2019.110716
38. Gutiérrez-Robledo LM, García-Chanes RE, Pérez-Zepeda MU. Allostatic load as a biological substrate to intrinsic capacity: a secondary analysis of CRELES. *J Nutr Heal Aging*. (2019) 23:788–95. doi: 10.1007/s12603-019-1251-5
39. Merchant RA, Hui RJY, Kwek SC, Sundram M, Tay A, Jayasundram J, et al. Rapid geriatric assessment using mobile app in primary care: prevalence of geriatric syndromes and review of its feasibility. *Front Med*. (2020) 7:261. doi: 10.3389/fmed.2020.00261

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Ma, Chhetri, Zhang, Liu, Chen, Li and Chan. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.