



OPEN ACCESS

EDITED BY

Weiren Luo,
The Second Affiliated Hospital of
Southern University of Science and
Technology, China

REVIEWED BY

Ankush Kawali,
Narayana Nethralaya, India
Laura De Luca,
University of Messina, Italy

*CORRESPONDENCE

Giorgio Zauli
giorgio.zauli@unife.it

SPECIALTY SECTION

This article was submitted to
Ophthalmology,
a section of the journal
Frontiers in Medicine

RECEIVED 24 March 2022

ACCEPTED 27 June 2022

PUBLISHED 14 July 2022

CITATION

Zauli G, AlHilali S, Al-Swailem S,
Secchiero P and Voltan R (2022)
Therapeutic potential of the MDM2
inhibitor Nutlin-3 in counteracting
SARS-CoV-2 infection of the eye
through p53 activation.
Front. Med. 9:902713.
doi: 10.3389/fmed.2022.902713

COPYRIGHT

© 2022 Zauli, AlHilali, Al-Swailem,
Secchiero and Voltan. This is an
open-access article distributed under
the terms of the [Creative Commons
Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use,
distribution or reproduction in other
forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which
does not comply with these terms.

Therapeutic potential of the MDM2 inhibitor Nutlin-3 in counteracting SARS-CoV-2 infection of the eye through p53 activation

Giorgio Zauli^{1*}, Sara AlHilali¹, Samar Al-Swailem¹,
Paola Secchiero² and Rebecca Voltan³

¹Research Department, King Khaled Eye Specialistic Hospital, Riyadh, Saudi Arabia, ²Department of Translational Medicine and LTTA Centre, University of Ferrara, Ferrara, Italy, ³Department of Environmental and Prevention Sciences and LTTA Centre, University of Ferrara, Ferrara, Italy

Starting from the beginning of the severe acute respiratory syndrome Coronavirus-2 (SARS-CoV-2) global pandemic, most of the published data has concentrated on the respiratory signs and symptoms of Covid-19 infection, underestimating the presence and importance of ocular manifestations, such as conjunctivitis, usually reported in SARS-CoV-2 infected patients. With the present review we intend to resume the ocular involvement in SARS-CoV-2 infection and the recent discoveries about the different cell types and tissues of the eye that can be directly infected by SARS-CoV-2 and propagate the infection. Moreover, reviewing literature data about p53 expression in normal and diseased eye tissues, we hypothesize that the pleiotropic protein p53 present at high levels in cornea, conjunctiva and tear film might play a protective role against SARS-CoV-2 infection. Since p53 can be easily up-regulated by using small molecule non-genotoxic inhibitors of MDM2, we propose that topical use of Nutlin-3, the prototype member of MDM2 inhibitors, might protect the anterior surface of the eye from SARS-CoV-2 infection, reducing the spreading of the virus.

KEYWORDS

SARS-CoV-2, p53, Nutlin-3, MDM2 inhibitors, eye, cornea, tear film

Introduction

Under physiological conditions, p53 protein is maintained at low level in most normal tissues through a variety of mechanisms, mainly depending by the control mediated by its principal inhibitor murine double minute 2 (MDM2), also known as human double minute 2 (HDM2) in humans. MDM2 acts on p53 through a well-characterized negative feedback loop, and, when the negative regulatory role of MDM2 is lost, as often it happens in response to viral infections (1), the intracellular p53 levels increase, leading to biological effects which depend on the duration and the strength of p53 activation (2). p53 is a pleiotropic molecule deeply analyzed for its several functions

and involvement in different pathways, ranging from cell cycle arrest and apoptosis induction to senescence modulation.

In the context of viral infection, it is emerging that human coronaviruses have developed several specific molecular ways to interfere with p53-mediated activities in infected cells. It has been shown that the viral papain-like proteases, PLPs, can act as MDM2 stabilizer *via* deubiquitination, leading to accelerated p53 proteasomal degradation (3, 4), and to a cascade of events which ensures the rapid growth of virus-infected cells due to the p53 loss. Analogously, the PLPs' mediated effect was demonstrated also on RCHY1, another E3 ubiquitin ligase involved in p53 ubiquitination, resulting in enhanced p53 degradation inside infected cells (5). Moreover, it was observed that knockout of *TP53* gene promoted viral replication, and that, on the opposite, the expression of p53 served as antiviral cellular molecule able to downregulate SARS-CoV replication (5). More recently, another group, screening the impact of SARS-CoV-2 proteins in several signaling pathways involved in viral infection, has demonstrated that the viral protease nsp5 can functionally repress p53 by interfering with its transcriptional activity, and has suggested p53 as an "intrinsic host restriction factor for the virus" (6). Similarly to Ma-Lauer, they have also observed that the overexpression of p53 significantly reduced virus production, sustaining the hypothesis of a fundamental role of p53 in managing the cellular antiviral defenses. These observations highlight the central role of p53 in controlling coronavirus replication inside infected cells. On these bases, we have recently suggested that Nutlin-3, or its oral version Idasanutlin, a potent and selective small-molecule inhibitor of MDM2, that promotes stabilization of p53, might be beneficial to treat the pulmonary infection induced by SARS-CoV-2 (7). Similarly, following the concept that restoring/inducing p53 may provide beneficial therapeutic results for infectious diseases, a group of the Georgetown University has proposed a *TP53* gene therapy approach for patients affected by COVID-19, hypothesizing the repurposing of the SGT-53 agent, an immunoliposome carrying a plasmid vector for *TP53* expression, currently in phase II human trials for pancreatic cancer (8).

In this review, we aimed to focus the attention on the eye and investigate the potential role of the MDM2/p53 pathway, modulated by Nutlin-3, in counteracting/preventing SARS-CoV-2 ocular manifestations and diseases. To support this rationale, we reviewed the recent literature about the involvement of eye's tissues, cells and tears during SARS-CoV-2 infection, and resumed the data about MDM2 and p53 expression in ocular health and disease situations. Moreover, we analyzed the observations about the feasibility of p53 modulation through MDM2 inhibition in preclinical ocular models. Finally, we proposed a topical use of Nutlin-3 as a therapeutic approach to protect the anterior surface of the eye and to contrast SARS-CoV-2 infection and spreading. A

reasoning on importance of cytokines modulation was also included.

The anterior surface of the eye represents a gateway for SARS-CoV-2 infection

The main receptor for SARS-CoV-2, the angiotensin-converting enzyme 2, ACE2, is naturally present in the epithelial lining of the lower respiratory tract and digestive system (9). It serves as the main portal of entry for Covid-19 virus into the human body, leading to a variety of respiratory and gastrointestinal symptoms, but it has been detected in several cell types, with highest levels of expression in type II alveolar pneumocytes (10). Increasing experimental evidence has recently demonstrated that also ocular tissues express ACE2. Indeed, ACE2 receptor expression at the mRNA and protein level has been shown in multiple cell types involved in human vision, including cell types of the external surface of the eye (11–13). However, the proportion of cells in the ocular surface having greater-than-zero expression of ACE2 does not exceed 10% and the mean level of mRNA expression is approximately 0.6% (14). Although the expression level of ACE2 receptor seemed to be relatively low compared to the type II alveolar pneumocytes (15), epidemiological studies have reported ocular surface infection among the clinical manifestations of patients with COVID-19 (16); moreover, conjunctivitis has been reported as the initial presenting symptom in several cases (17). A systematic review and meta-analysis reported that approximately 10% of COVID-19 patients has ocular manifestations with at least one ocular symptom (such as dry eye, conjunctival injection/chemosis, tearing, itching and foreign body sensation), indicating also that attention to these manifestations is important in the detection of COVID-19 infection in the population (18).

Such evidence underscores the importance of the eye as a gateway for SARS-COV-2. Therefore, we will explore experimental and clinical evidence supporting the hypothesis that the ocular surface is a potential route for SARS-CoV-2 transmission.

The tissues of the anterior surface of the eye express ACE2 and TMPRSS2 receptors

ACE2 has been detected in the epithelial cells of both skin and cornea, tissues that share the role of primary barrier against the external environment, having both an external stratified squamous epithelium, and that govern several patho-physiological responses including inflammation (19). In

the anterior surface of the eye, as well as in the cornea, ACE2 has been found in the conjunctiva and limbus (20). Furthermore, it has been shown that cells of the same ocular surface tissues expressing ACE2 also express the co-receptor transmembrane serine protease 2 (TMPRSS2) (21), resulting susceptible to SARS-CoV-2 infection from a molecular point of view. Interestingly, from the clinical point of view, it has been shown that some SARS-CoV-2 patients presented conjunctivitis as first symptom, and sometimes as unique manifestation of the infection. Moreover, SARS-CoV-2 was detectable on the ocular surface (conjunctival swabs and tears) of both symptomatic and asymptomatic patients, and the isolated virus was infectious if used to infect cells *in vitro*, demonstrating that the eye can be infected and at the same time can support virus replication and spreading (21). Consistently with the hypothesis that the anterior surface of the eye might represent a potential route of infection, Casagrande detected SARS-CoV-2 genomic RNA in corneas of deceased patients affected by COVID-19 (22). Furthermore, Singh showed that SARS-CoV-2 can infect human conjunctival tissues, especially of inflamed conjunctiva (23), and observed that diseased conjunctival tissues, such as nevi, cyst, papilloma and polyps, overexpress ACE2 as compared to normal conjunctival tissues (24). Another study used a single-cell sequencing approach to demonstrate that both ACE2 and TMPRSS2 genes were highly co-expressed in the goblet cells of the conjunctiva, along with genes involved in immunity process (25). Once established that the anterior surface of the eye represents a potentially important route of infection, it is noteworthy that the highest levels of SARS-CoV-2 viral entry-related proteins are present in the limbus (26, 27). ACE2 and TMPRSS2 were also described in post-mortem SARS-CoV-2 patient ocular surface tissues, in association with productive viral replication, most notably in limbal regions (26). Furthermore, the same authors described in whole-eye organoid model that conjunctival and limbal cells can be infected by the virus and that they support active viral replication. It is noteworthy that transcriptional analysis of *ex-vivo* infected ocular surface cells revealed robust induction of NF- κ B in infected cells as well as diminished type I interferon signaling (26). These findings are particularly interesting since it has been clearly demonstrated the key role of p53 in activating type I interferon signaling and in down-regulating the NF- κ B pro-inflammatory pathway (28).

In summary, corneal, limbal and conjunctival epithelia express ACE2 and TMPRSS2, can be infected by SARS-CoV-2 and support a productive infection, suggesting that the ocular surface is a potential route for the transmission of the virus, and it represents a possible risk of viral transmission, in particular for healthcare workers working at proximity to the eye and for recipient patients during the corneal transplantation procedures (29).

Retinal cells can be infected by SARS-CoV-2

Although most of the experimental data available on this topic suggest the anterior surface of the eye as a possible site for SARS-CoV-2 entry and infection, other evidence suggests that also the retina can be infected. In this respect, Zhou demonstrated that ACE2 is expressed in several neuro-retinal cells and retinal vessels analyzed in post-mortem retinopathy specimens donated by non-diabetic and diabetic patients (30). In the same specimens, TMPRSS2 co-expression was detected in retinal neuronal cells, vascular and perivascular cells (30). Additional evidence of retinal involvement was provided by the detection of SARS-CoV-2 genomic RNA in retinal biopsies of deceased COVID-19 patients (31). These findings could explain the presence of retinal changes such as cotton wool spots and dot-blot hemorrhages reported in the literature (32). In line with the findings of Zhou, that observed a higher expression of ACE2 receptor in vessels of diabetic patients, Landecho showed that retinal microangiopathy might represent an *in vivo* biomarker of systemic vascular disease during SARS-CoV-2 infection (33). In fact, 6 out of 27 evaluated patients involved in the study showed cotton wool exudates, a marker of vascular disease severity in diabetes, associated with increased risk for acute vascular events. At present, it is not clear if these events can be caused directly by the virus acting in the endothelium of the vasculature or if they can be provoked by a disseminated intravascular hypercoagulable state (33). Moreover, if infection of the retina occurs and if it derives directly from the eye, from the upper respiratory tract or from another site are still open questions. Nonetheless, in the course of SARS-CoV-2 infection, it is very likely that ocular manifestations that affect the retina are modulated by an alteration of the blood-retinal barrier, BRB (34). This could occur at the level of the retinal endothelial cells of the inner barrier, iBRB, or at the level of retinal pigment epithelial, RPE, cells of the outer barrier, oBRB, or at both levels.

Role of the tears/tear film and of the nasolacrimal duct morphology in the transmission of SARS-CoV-2 infection

During the first months of the pandemic, the viral RNA was not clinically evaluated in patients' tears, due to the fast spreading of the emergency, and rare eye symptoms were reported probably because underestimated compared to the more severe systemic symptoms (16, 35). Literature of later periods investigated the eye as possible route of infection, searching the presence of SARS-CoV-2 virus in tears, but results appeared highly variable (Table 1), leading authors to different conclusions about the relevance of SARS-CoV-2 RNA in tears.

TABLE 1 Overview of relevant literature data investigating SARS-CoV-2 in tears/tear film of infected patients.

Aim of the study	Patients (n)	COVID-19 severity	Sampling method	Sampling time*	SARS-CoV-2 RNA in samples	Eye symptoms	Refs
SARS-CoV-2 on ocular surfaces of sub-intensive patients with pneumonia	9	Severe	Conjunctival swab	<48h	72%	Bilateral conjunctivitis (78%)	Troisi et al. (36)
SARS-CoV-2 on ocular surfaces	91	Severe	Conjunctival swab	<48h	57.1%	Hyperemia (5.8%), secretions (5.8%), blepharitis (7.7%), other signs (13.5%)	Azzolini et al. (37)
SARS-CoV-2 on corneal disks	11	Severe	Conjunctival swab (post-mortem)	NA	45%	NA	Casagrande et al. (22)
SARS-CoV-2 in tears of patients with moderate and severe COVID-19	75	Moderate and severe	Conjunctival swab and/or Schirmer's test strip	<48 h	24%	None	Arora et al. (38)
SARS-CoV-2 on ocular surfaces (observational)	243	Mild, moderate and severe	Conjunctival swab	1–17 days	7%	NA	Gijs et al. (39)
Viral RNA in conjunctival secretion	49	Asymptomatic, mild and moderate (treated with antivirals)	Conjunctival swab	2–27 days	8.2%	None	Li et al. (40)
SARS-CoV-2 in tears	30	Mild and severe (treated with antivirals)	Conjunctival swab	1–16 days	3.3%	Conjunctivitis (3.3%)	Xia et al. (41)
Characteristics of ocular findings (retrospective)	38	Mild, moderate and severe	Conjunctival swab	NA	5.3%	Conjunctivitis (31.6%)	Wu et al. (42)
Viral RNA in conjunctival secretion	37	Mild and severe	NA	NA	2.7%	Conjunctival congestion (8.1%), eye inflammation (91.9%)	Liang et al. (43)
Viral shedding in tears	17	Mild	Schirmer's test strip	3–20 days	0%	None	Seah et al. (44)

*Time after positive naso/oropharyngeal swab.

More recently, when data originated from studies where tears analysis was the main specific research target, and methods were standardized, the situation was clearer, indicating that tears/tear film collected by conjunctival swabs from moderate to severe COVID-19 affected patients contain viral RNA (Table 1). In the work of Arora and colleagues, SARS-CoV-2 RNA was detected in tears of 24% of patients with a confirmed COVID-19 infection, despite the absence of any ocular manifestation (38). In the study of Azzolini, viral RNA was detected in a larger part of the investigated cohort (57,1%), reporting low rate of ocular signs and indicating that in some patients the ocular surface can be positive when nasopharyngeal swab is negative (37). Interestingly, a small study reported preliminary data indicating tear film positivity in 72% of analyzed patients (36). This incredible high percentage derives probably by the characteristics of the investigated cohort that included patients with severe Covid-19 treated with full-mask or oxygen helmet. Indeed, the authors hypothesized that the dryness and eye irritation due to wearing the mask could facilitate the ocular surface infection by the high viral load present in the breath of pneumonia affected patients. Together, these evidence supports the hypothesis that viral shedding through the tear film is a potential route of SARS-CoV-2 transmission, even if tears are characterized by a specific content of constitutive antimicrobial molecules, such as lactoferrin and lysozyme. In particular, lactoferrin has been shown to possess antiviral activity against SARS-CoV-2 (45) and to be able to upregulate p53 through the activation of NF- κ B (46), but its capacity of modulating viral spreading at physiological concentration needs to be evaluated.

The nasolacrimal ducts constitute the anatomical connection between the eye and the nose, permitting the direct communication between the ocular mucosal immune system and the associated lymphoid tissue of the nasal cavity, and contributing to the immunological interdependence between the ocular and the respiratory systems. Thus, the nasolacrimal system may function as a route for virus migration, either through drainage of tears through the duct to the respiratory tract or, vice-versa, from the upper respiratory tract through the nasolacrimal duct to the eye. Indeed, it has been hypothesized that after infecting the ocular surface, SARS-CoV-2 virus could enter the nasolacrimal ducts to infect the high respiratory tract (47). Moreover, other investigators indicated that the nasolacrimal system may provide an additional route of entry and infection of other tissues, including the epithelium of the lacrimal canaliculi, the nasolacrimal drainage system, the nasal passage, and the upper respiratory tract (48–50).

MDM2 and p53 expression in healthy ocular tissues

Once established that the receptors for SARS-CoV-2 infection are present in different ocular tissues, it is of interest

to elucidate whether these tissues express p53 and/or its major inhibitor MDM2. Although only few studies addressed the issue of the basal p53 expression in the eye, it is noteworthy that Tendler and Panshin recently demonstrated the presence of significant p53 protein levels both in the corneal epithelium as well as in the corneal tear film (51). Of note, they observed that while the concentration of p53 was low in the cytoplasm of most normal cell types, due to the potential harmful role of elevated p53 concentrations in inducing cell cycle arrest and/or apoptosis, abnormally high p53 content was detected by immunohistochemistry, Western blot analysis and electronic microscope examinations in corneal epithelial cells. These data confirmed previous results that reported strong p53 cytoplasmic expression both in corneal as well as in conjunctival epithelium of mice (52). Differently from corneal epithelium, minute amount of p53 was found in retina, lens and iris. Consistently, MDM2 was identified in the retina, lens and iris while it was absent in the corneal epithelium (51). The abnormally high levels of p53 in the corneal epithelium and the absence of its negative regulator adds to corneal immune privilege (53, 54). In addition, it was demonstrated that a significant number of exosomes and other microvesicles containing p53 were present in the corneal mucin layer of the tear film (51). In a previous study, the same authors showed that, after ultraviolet irradiation, the cytoplasmic p53 protein in corneal epithelium cells became functionally active, following phosphorylation in Ser15, and moved from cytoplasm to nucleus (55).

Summarizing, it is possible to indicate that p53 is present at significant levels in the normal tissues of the anterior surface of the eye, with elevated concentration in the cornea, and it appears to be functional. Moreover, as mentioned above, high levels of p53 are expressed not only in the anterior surface of the eye but are present also in tear film, where p53 might play anti-neoplastic as well as anti-infection activities. Interestingly, a crosstalk has been demonstrated between the p53 pathway and lactoferrin (46), and both proteins are components of the tears. Together, they might represent a natural barrier against the development of tumors in the anterior surface of the eye (56), as well as against infections, and in particular against SARS-CoV-2 (57, 58).

Modulation of p53 activity in pathological ocular tissues following MDM2-inhibition by Nutlin-3

In this section of the review we summarize literature data suggesting that the p53 activator Nutlin-3 can modulate the MDM2/p53 axis in the ocular tissues where co-expression of both ACE2 and TMPRSS2 has been demonstrated. As mentioned earlier, it is particularly noteworthy that p53 is abundantly expressed in normal eye tissues of cornea,

limbus and conjunctiva, it's present in corneal tear film, and it may play a protective role against viral infections. In this respect, Nutlin-3 has been proposed as a potential pharmacological approach against pterygium, an anomalous non-tumoral proliferation of the conjunctiva invading limbus and cornea (59). Experiments based on primary cell lines, obtained after explant of pterygium tissues, have demonstrated that p53, highly expressed but not active and relegated to cytoplasm, translocated to the nucleus after treatment with Nutlin-3 and showed transcriptional activity and apoptotic effect. More recently, the therapeutic use of Nutlin-3 was showed in another disease of the conjunctiva, the conjunctival melanoma that typically express p53 in wild-type status. These authors demonstrated that Nutlin-3 was able to reactivate p53 and reduced viability in several models of conjunctival melanoma, both *in vitro* in cell lines and 3D spheroids, as well as *in vivo* in zebrafish xenografts (60). Considering the retina, an ocular formulation of Nutlin-3 was used in preclinical models of retinoblastoma, a pediatric disease characterized for expressing high levels of MDM2/HDMX (61). In this study, subconjunctival administration of Nutlin-3 improved intraocular penetration, respect to oral and iv administration, and exhibited specific p53-mediated antitumor effects. More recently, the use of Nutlin-3 was proposed to target and eliminate senescent RPE cells involved in the retinal degeneration phenomena occurring during the progressive and multifactorial disease known as age-related macular degeneration (AMD) (62). In this work, senescent RPE cells were shown to express high level of p53 and other expression markers of senescence like p21, p16, IL-1 β , IL-6, and Mmp-3. Nutlin-3 treatment selectively killed more than 50% of senescent RPE cells, with no cytotoxicity on non-senescent cells, also decreasing the levels of released inflammatory cytokines, such as IL-6. This pharmacological approach was efficient both *in vitro* in Dox-induced senescent ARPE-19 cells, as well as *in vivo* in a mouse model of RPE senescence treated intravitreally with Nutlin-3. Interestingly, the authors observed also retinal regeneration and improving of retinal function in a model of aged mice, hypothesizing a possible therapeutic use of MDM2 inhibitors for this disease. Moreover, these data confirm the ability of Nutlin-3 to reduce the secretion of cytokines during senescence-associated secretory phenotype (SASP). Interestingly, in the context of retinal detachment secondary to proliferative vitreoretinopathy, Nutlin-3 was efficient in upregulate p53 expression of hyper proliferating RPE cells, limiting their growth and preventing retinal detachment (63). This observation leads to the important consideration that Nutlin-3 can modulate the proliferation of RPE cells when they are in an altered state, such as senescence or hyperproliferation, restoring a physiological situation.

So far, to our knowledge, beside conjunctiva, Nutlin-3 or other MDM2 inhibitors were not evaluated specifically to treat pathological cornea or limbus, the other ocular tissues expressing high levels of SARS-CoV-2 receptors. Differently,


experimental uses of Nutlin-3 in diseases affecting other ocular tissues have been recently reviewed and further sustain the feasibility of its use as ocular treatment (64).

An important consideration regarding most of the works with a pharmacological use of Nutlin-3 for *in vivo* ocular treatment is that they were performed by intravitreal injection, an invasive method unfortunately needed for many eyes' disease therapies. A delivery method through eye drops, as was proposed by Brennan more than a decade ago, would be certainly easier for approaching preclinical studies, and more desirable and better accepted by future patients.

Nutlin-3 might represent a new therapeutic tool to counteract SARS-CoV-2 infection of the eye

Following the here resumed evidence that external ocular tissues can be the portal of entry and site for spreading of SARS-CoV-2, and that p53 pathway can be activated in these tissues, it appears that a therapeutic intervention with Nutlin-3 can be feasible to counteract SARS-CoV-2 infection of the eye and merits to be pursued. In particular, Nutlin-3 subconjunctival formulations represent unique opportunities to efficiently deliver the drug topically, with a non-invasive method. Moreover, this approach could be further improved and targeted in the next future with the help of nano-formulations, such as lipid vesicles.

A proposal about a possible combination of molecular mechanisms acting on corneal infected epithelium after pharmacological ophthalmic treatment with Nutlin-3 is presented in Figure 1. Since Nutlin-3 and coronavirus PLPs are competitors on MDM2, acting in opposite direction on the fate of p53, the first stabilizing the protein and the second promoting its degradation, the comprehension of the downstream mechanisms mediated by p53, other than the well-known apoptosis induction, and able to modulate infectivity and ocular symptoms are highly wished. Among these, considering that the "cytokine storm" is an important clinical manifestation during the severe SARS-CoV-2 infections (35), one of the potential beneficial anti-viral mechanisms of promoting p53 activation could be to reduce the senescence-associated secretory phenotype and the interleukin-6 secretion. Indeed, SASP is negatively controlled by p53 (65), and it was demonstrated that SARS-Cov-2 infection can promote both IL-6 secretion in tears (39, 66), and inflammatory response in human corneal epithelial cells (67). Nutlin-3 treatment could then drive SASP inhibition and IL-6 secretion, through the inhibition of MDM2 and upregulation of p53. Moreover, since ACE2 expression is upregulated in inflamed eye's tissues (24, 68), reducing IL-6 could be a strategy to reduce local inflammation and to restore baseline receptor levels.



Other mechanisms could involve the transcriptional modulation of ACE2 and TMPRSS2, and the activation of the innate immune response. In the last context, it is known that p53 participates to the antiviral innate immune responses by inducing apoptosis of infected cells and by mediating type I interferon, IFN-I, synthesis/signaling (3, 28). On the other hand, it was demonstrated that SARS-CoV-2 can manipulate the host innate immune response blocking or inducing IFN-I pathway to its advantage, and it is sensitive to IFN treatment (63, 69). How p53, enhanced by Nutlin-3, could win the battle is a new challenge. Interestingly, topical administration of IFN is largely used in ophthalmic practice for treatment of several ocular surface disorders, and its therapeutic and side effects

are well known (70, 71). In the spectrum of the responses induced thanks to the activation of p53, topical use of Nutlin-3 could also induce type I interferon signaling (3, 28), having the limitation compared to recombinant interferon of a less direct and focused action, but the advantage of being broader, with the activation of the innate immune response along with other effects discussed above. Obviously, this specific ability remains to be preclinically characterized at the ocular level. In parallel, it should be interesting to evaluate the efficacy of IFN-I local therapy in the eye that, to our knowledge, has not been investigated against SARS-CoV-2 so far.

We believe that all these considerations pose the basis for a pharmacological approach against ocular SARS-CoV-2 infection

mediated by the control of p53 expression with Nutlin-3 as preferred candidate.

Author contributions

Conceptualization: GZ. Writing original draft preparation and writing review and editing: GZ, SA, SA-S, RV, and PS. All authors have read and agreed to the published version of the manuscript.

Funding

Funds for open access publication fees were received by RV as local funding from University of Ferrara (FAR and FIR programs).

References

- Sato Y, Tsurumi T. Genome guardian p53 and viral infections. *Rev Med Virol*. (2013) 23:213–20. doi: 10.1002/rmv.1738
- Tisato V, Voltan R, Gonelli A, Secchiero P, Zauli G. MDM2/X inhibitors under clinical evaluation: perspectives for the management of hematological malignancies and pediatric cancer. *J Hematol Oncol*. (2017) 10:133. doi: 10.1186/s13045-017-0500-5
- Yuan L, Chen Z, Song S, Wang S, Tian C, Xing G, et al. 53 degradation by a coronavirus papain-like protease suppresses type I interferon signaling. *J Biol Chem*. (2015) 290:3172–82. doi: 10.1074/jbc.M114.619890
- Dong S, Sun J, Mao Z, Wang L, Lu YL, Li J, et al. A guideline for homology modeling of the proteins from newly discovered betacoronavirus 2019 novel coronavirus, (2019-nCoV). *J Med Virol*. (2020) 92:1542–8. doi: 10.1002/jmv.25768
- Ma-Lauer Y, Carbajo-Lozoya J, Hein MY, Müller MA, Deng W, Lei J, et al. Down-regulates SARS coronavirus replication and is targeted by the SARS-unique domain and PLpro via E3 ubiquitin ligase RCHY1. *Proc Natl Acad Sci U S A*. (2016) 113:E5192–201. doi: 10.1073/pnas.1603435113
- Kumar A, Grams TR, Bloom DC, Toth Z. Signaling Pathway Reporter Screen with SARS-CoV-2 Proteins Identifies nsp5 as a Repressor of p53 Activity. *Viruses*. (2022) 14:1039. doi: 10.3390/v14051039
- Zauli G, Tisato V, Secchiero P. Rationale for considering oral idasanutlin as a therapeutic option for COVID-19 patients. *Front Pharmacol*. (2020) 11:1156. doi: 10.3389/fphar.2020.01156
- Harford JB, Kim SS, Pirollo KF, Chang EH. TP53. Gene therapy as a potential treatment for patients with COVID-19. *Viruses*. (2022) 14:739. doi: 10.3390/v14040739
- Lukassen S, Chua RL, Trefzer T, Kahn NC, Schneider MA, Muley T, et al. SARS-CoV-2 receptor ACE2 and TMPRSS2 are primarily expressed in bronchial transient secretory cells. *EMBO J*. (2020) 39:e105114. doi: 10.15252/embj.2020105114
- Zhao Y, Zhao Z, Wang Y, Zhou Y, Ma Y, Zuo W, et al. Single-Cell RNA expression profiling of ACE2, the Receptor of SARS-CoV-2. *Am J Respir Crit Care Med*. (2020) 202:756–9. doi: 10.1164/rccm.202001-0179LE
- Szczeniak M, Brydak-Godowska J. SARS-CoV-2 and the eyes: a review of the literature on transmission, detection, and ocular manifestations. *Med Sci Monit*. (2021) 27:e931863. doi: 10.12659/MSM.931863
- Yener AÜ. COVID-19 and the Eye: ocular manifestations, treatment and protection measures. *Ocul Immunol Inflamm*. (2021) 14:1–9. doi: 10.1080/09273948.2021.1977829
- Hill JM, Clement C, Arceneaux L, Lukiw WJ. Angiotensin converting enzyme 2 (ACE2) expression in the aged brain and visual system. *J Aging Sci*. (2021) 9:001. doi: 10.21203/rs.3.rs-296818/v1
- Sungnak W, Huang N, Bécavin C, Berg M, Queen R, Litvinukova M, et al. SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with innate immune genes. *Nat Med*. (2020) 26:681–7. doi: 10.1038/s41591-020-0868-6
- Kitazawa K, Deinhardt-Emmer S, Inomata T, Deshpande S, Sotozono C. The transmission of SARS-CoV-2 infection on the ocular surface and prevention strategies. *Cells*. (2021) 10:796. doi: 10.3390/cells10040796
- Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. (2020) 382:1708–20. doi: 10.1056/NEJMoa2002032
- Chen L, Deng C, Chen X, Zhang X, Chen B, Yu H, et al. Ocular manifestations and clinical characteristics of 535 cases of COVID-19 in Wuhan, China: a cross-sectional study. *Acta Ophthalmol*. (2020) 98:e951–9. doi: 10.1111/aos.14472
- Nasiri N, Sharifi H, Bazrafshan A, Noori A, Karamouzian M, Sharifi A, et al. Ocular Manifestations of COVID-19: a systematic review and meta-analysis. *J Ophthalmic Vis Res*. (2021) 16:103–12. doi: 10.18502/jovr.v16i1.8256
- Kaplan N, Gonzalez E, Peng H, Battle D, Lavker RM. Emerging importance of ACE2 in external stratified epithelial tissues. *Mol Cell Endocrinol*. (2021) 529:111260. doi: 10.1016/j.mce.2021.111260
- Zhou L, Xu Z, Castiglione GM, Soiberman US, Eberhart CG, Duh EJ, et al. ACE2 and TMPRSS2 are expressed on the human ocular surface, suggesting susceptibility to SARS-CoV-2 infection. *Ocul Surf*. (2020) 18:537–44. doi: 10.1016/j.jtos.2020.06.007
- Qu JY, Xie HT, Zhang MC. Evidence of SARS-CoV-2 transmission through the ocular route. *Clin Ophthalmol*. (2021) 15:687–96. doi: 10.2147/OPHTH.S295283
- Casagrande M, Fitzek A, Spitzer MS, Püschel K, Glatzel M, Krasemann S, et al. Presence of SARS-CoV-2 RNA in the Cornea of Viremic Patients With COVID-19. *JAMA Ophthalmol*. (2021) 139:383–8. doi: 10.1001/jamaophthalmol.2020.6339
- Singh S, Garcia G, Shah R, Kramerov AA, Wright RE, et al. SARS-CoV-2 and its beta variant of concern infect human conjunctival epithelial cells and induce differential antiviral innate immune response. *Ocul Surf*. (2021) 23:184–94. doi: 10.1016/j.jtos.2021.09.007
- Li X, Chan JF, Li KK, Tso EY, Yip CC, Sridhar S, et al. Detection of SARS-CoV-2 in conjunctival secretions from patients without ocular symptoms. *Infection*. (2021) 49:257–65. doi: 10.1007/s15010-020-01524-2
- Ma R, Gan L, Jiang S, Ding P, Chen D, Wu J, et al. High expression of SARS-CoV-2 entry factors in human conjunctival goblet cells. *Exp Eye Res*. (2021) 205:108501. doi: 10.1016/j.exer.2021.108501
- Eriksen AZ, Møller R, Makovoz B, Uhl SA, tenOever BR, Blenkinsop TA, et al. SARS-CoV-2 infects human adult donor eyes and hESC-derived ocular epithelium. *Cell Stem Cell*. 28, 1205–1220.e7. doi: 10.1016/j.stem.2021.04.028

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

27. Yoshinaga Y, Soma T, Azuma S, Maruyama K, Hashikawa Y, Katayama T et al. High expression of SARS-CoV2 viral entry-related proteins in human limbal stem cells. *Ocul. Surf.* (2021) 23:197–200. doi: 10.1016/j.jtos.2021.10.002
28. Munoz-Fontela C, Macip S, Martinez-Sobrido L, Brown L, Ashour J, Garcia-Sastre A, et al. Transcriptional role of p53 in interferon-mediated antiviral immunity. *J Exp Med.* (2008) 205:1929–38. doi: 10.1084/jem.2008.0383
29. Roehrich H, Yuan C, Hou JH. Immunohistochemical study of SARS-CoV-2 viral entry factors in the cornea and ocular surface. *Cornea.* (2020) 39:1556–62. doi: 10.1097/ICO.0000000000002509
30. Zhou L, Xu Z, Guerra J, Rosenberg AZ, Fenaroli P, Eberhart CG, et al. Expression of the SARS-CoV-2 Receptor ACE2 in Human Retina and Diabetes-Implications for Retinopathy. *Invest Ophthalmol Vis Sci.* (2021) 62:6. doi: 10.1167/iovs.62.7.6
31. Casagrande M, Fitzek A, Püschel K, Aleshcheva G, Schultheiss HP, Berneking L, et al. Detection of SARS-CoV-2 in human retinal biopsies of deceased COVID-19 patients. *Ocul Immunol Inflamm.* (2020) 28:721–5. doi: 10.1080/09273948.2020.1770301
32. Marinho PM, Marcos AAA, Romano AC, Nascimento H, Belfort RJr. Retinal findings in patients with COVID-19. *Lancet.* (2020) 395:1610. doi: 10.1016/S0140-6736(20)31014-X
33. Landecho MF, Yuste JR, Gándara E, Sunsendegui P, Quiroga J, Alcaide AB, et al. COVID-19 retinal microangiopathy as an in vivo biomarker of systemic vascular disease? *J. Intern Med.* (2021) 289:116–20. doi: 10.1111/joim.13156
34. Sharma S, Jagadeesh H, Saxena A, Chakravarthy H, Devanathan V. Central nervous system as a target of novel coronavirus infections: Potential routes of entry and pathogenic mechanisms. *J Biosci.* (2021) 46:106. doi: 10.1007/s12038-021-00232-9
35. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* (2020) 395:497–506. doi: 10.1016/S0140-6736(20)30183-5
36. Troisi M, Zannella C, Troisi S, Bernardo D, Galdiero M, et al. Ocular surface infection by SARS-CoV-2 in COVID-19 pneumonia patients admitted to sub-intensive unit: preliminary results. *Microorganisms.* (2022) 10:347. doi: 10.3390/microorganisms10020347
37. Azzolini C, Donati S, Premi E, Baj A, Siracusa C, Genoni A, et al. SARS-CoV-2 on ocular surfaces in a cohort of patients with COVID-19 from the Lombardy Region, Italy. *JAMA Ophthalmol.* (2021) 139:956–63. doi: 10.1001/jamaophthalmol.2020.5464
38. Arora R, Goel R, Kumar S, Chhabra M, Saxena S, Manchanda V, et al. Evaluation of SARS-CoV-2 in tears of patients with moderate to severe COVID-19. *Ophthalmology.* (2020) 128:494–503. doi: 10.1016/j.ophtha.2020.08.029
39. Gijs M, Veugen MJM, Wolffs PFG, Savelkoul PHM, Tas J, van Bussel BCT, et al. In-Depth Investigation of Conjunctival Swabs and Tear Fluid of Symptomatic COVID-19 Patients, an Observational Cohort Study. *Transl Vis Sci Technol.* (2021) 10:32. doi: 10.1167/tvst.10.12.32
40. Li S, Li D, Fang J, Liu Q, Cao W, Sun X, et al. SARS-CoV-2 receptor ACE2 is expressed in human conjunctival tissue, especially in diseased conjunctival tissue. *Ocul. Surf.* (2020) 19:249–51. doi: 10.1016/j.jtos.2020.09.010
41. Xia J, Tong J, Liu M, Shen Y, Guo D. Evaluation of coronavirus in tears and conjunctival secretions of patients with SARS-CoV-2 infection. *J Med Virol.* (2020) 92:589–94. doi: 10.1002/jmv.25725
42. Wu P, Duan F, Luo C, Liu Q, Qu X, Liang L, et al. Characteristics of ocular findings of patients with coronavirus disease 2019 (COVID-19) in Hubei province, China. *JAMA Ophthalmol.* (2020) 138:575–8. doi: 10.1001/jamaophthalmol.2020.1291
43. Liang L, Wu P. There may be virus in conjunctival secretion of patients with COVID-19. *Acta Ophthalmol.* (2020) 98:223. doi: 10.1111/aos.14413
44. Seah IYJ, Anderson DE, Kang AEZ, Wang L, Rao P, Young BE, et al. Assessing viral shedding and infectivity of tears in Coronavirus Disease 2019 (COVID-19) patients. *Ophthalmology.* (2020) 127:977–979. doi: 10.1016/j.ophtha.2020.03.026
45. Campione E, Lanna C, Cosio T, Rosa L, Conte MP, Iacovelli F, et al. Lactoferrin Against SARS-CoV-2: In Vitro and In Silico Evidences. *Front Pharmacol.* (2021) 17:666600. doi: 10.3389/fphar.2021.666600
46. Oh SM, Pyo CW, Kim Y, Choi SY. Neutrophil lactoferrin upregulates the human p53 gene through induction of NF-kappaB activation cascade. *Oncogene.* (2004) 23:8282–91. doi: 10.1038/sj.onc.1208021
47. Petronio Petronio G, Marco D, Costagliola R. Do ocular fluids represent a transmission route of SARS-CoV-2 infection? *Front Med.* (2021) 7:620412. doi: 10.3389/fmed.2020.620412
48. Ocansey S, Abu EK, Abraham CH, Owusu-Ansah A, Boadi-Kusi SB, Ilechie AA, et al. Ocular symptoms of SARS-CoV-2: indication of possible ocular transmission or viral shedding. *Ocul Immunol Inflamm.* (2020) 28:1269–79. doi: 10.1080/09273948.2020.1799035
49. Armstrong L, Collin J, Mostafa I, Queen R, Figueiredo FC, Lako M, et al. In the eye of the storm: SARS-CoV-2 infection and replication at the ocular surface? *Stem Cells Transl. Med.* (2021) 10:976–86. doi: 10.1002/sctm.20-0543
50. Spencer SKR, Francis IC, Coroneo MT. Spontaneous face- and eye-touching: Infection risk versus potential microbiome gain. *Ocu Surf.* (2021) 21, 64–65. doi: 10.1016/j.jtos.2021.04.008
51. Tendler Y, Panshin A. Features of p53 protein distribution in the corneal epithelium and corneal tear film. *Sci Rep.* (2020) 10:10051. doi: 10.1038/s41598-020-67206-z
52. Pokroy R, Tendler Y, Pollack A, and Zinder, O., and Weisinger, G. (2002). 53 expression in the normal murine eye. *Invest. Ophthalmol. Vis. Sci.* 43, 1736–1741
53. Cursiefen C. Immune privilege and angiogenic privilege of the cornea. *Chem Immunol Allergy.* (2007) 92:50–7. doi: 10.1159/000099253
54. Chen Y, Wang S, Alemi H, Dohlman T, Dana R. Immune regulation of the ocular surface. *Exp Eye Res.* (2022) 218:109007. doi: 10.1016/j.exer.2022.109007
55. Tendler Y, Pokroy R, Panshin A, Weisinger G. 53 protein subcellular localization and apoptosis in rodent corneal epithelium cell culture following ultraviolet irradiation. *Int J Mol Med.* (2013) 31:540–6. doi: 10.3892/ijmm.2013.1247
56. Pan S, Weng H, Hu G, Wang S, Zhao T, Yao X, et al. Lactoferrin may inhibit the development of cancer via its immunostimulatory and immunomodulatory activities. *Int J Oncol.* (2021) 59:85. doi: 10.3892/ijo.2021.5265
57. Alpgan O, Karakucuk S. Lactoferrin: the natural protector of the eye against coronavirus-19. *Ocul Immunol Inflamm.* (2021) 29:751–2. doi: 10.1080/09273948.2021.1954202
58. Vagge A, Senni C, Bernabei F, Pellegrini M, Scordia V, Traverso CE, et al. Therapeutic effects of lactoferrin in ocular diseases: from dry eye disease to infections. *Int J Mol Sci.* (2020) 21:6668. doi: 10.3390/ijms21186668
59. Cao D, Ng TK, Yip YWY, Young AL, Pang CP, Chu WK, et al. p53 inhibition by MDM2 in human pterygium. *Exp Eye Res.* (2018) 175:142–7. doi: 10.1016/j.exer.2018.06.021
60. Song D, Cismas S, Crudden C, Trocme E, Worrall C, Suleymanova N, et al. IGF-1R is a molecular determinant for response to p53 reactivation therapy in conjunctival melanoma. *Oncogene.* (2022) 41:600–11. doi: 10.1038/s41388-021-02111-x
61. Brennan RC, Federico S, Bradley C, Zhang J, Flores-Otero J, Wilson M, et al. Targeting the p53 pathway in retinoblastoma with subconjunctival Nutlin-3a. *Cancer Res.* (2011) 71:4205–13. doi: 10.1158/0008-5472.CAN-11-0058
62. Chae JB, Jang H, Son C, Park CW, Choi H, Jin S, et al. Targeting senescent retinal pigment epithelial cells facilitates retinal regeneration in mouse models of age-related macular degeneration. *Geroscience.* (2021) 43:2809–33. doi: 10.1007/s11357-021-00457-4
63. Lei H, Rheume MA, Cui J, Mukai S, Maberley D, Samad A, et al. A novel function of p53: a gatekeeper of retinal detachment. *Am J Pathol.* (2012) 181:866–74. doi: 10.1016/j.ajpath.2012.05.036
64. Jiang H, Luo J, Lei H. The roles of mouse double minute 2 (MDM2) oncoprotein in ocular diseases: a review. *Exp Eye Res.* (2022) 5:108910. doi: 10.1016/j.exer.2021.108910
65. Wiley CD, Schaum N, Alimirah F, Lopez-Dominguez JA, Orjalo AV, Scott G, et al. Small-molecule MDM2 antagonists attenuate the senescence-associated secretory phenotype. *Sci Rep.* (2018) 8:2410. doi: 10.1038/s41598-018-20000-4
66. Gulias-Cañizo R, Ruiz-Leyja ED, Sánchez-Huerta V, Ochoa-de la Paz L. SARS-CoV-2 and the eye: a relationship for a possible prognostic tool in COVID-19 patients. *Clin Ophthalmol.* (2021) 15 707–712. doi: 10.2147/OPHT.S284191
67. Zhu G, Lyu L, Yang H, Liu G, Yang S, Gu C, et al. SARS-CoV-2 spike protein-induced host inflammatory response signature in human corneal epithelial cells. *Mol Med Rep.* (2021) 24:584. doi: 10.3892/mmr.2021.12223

68. Jiang Z, Zhang H, Gao J, Yu H, Han R, Zhu L, et al. ACE2 expression is upregulated in inflammatory corneal epithelial cells and attenuated by resveratrol. *Invest Ophthalmol Vis Sci.* (2021) 62:25. doi: 10.1167/iovs.62.7.25
69. Lei X, Dong X, Ma R, Wang W, Xiao X, Tian Z, et al. Activation and evasion of type I interferon responses by SARS-CoV-2. *Nat Commun.* (2020) 11:3810. doi: 10.1038/s41467-020-17665-9
70. Mackensen F, Max R, Becker MD. Interferons and their potential in the treatment of ocular inflammation. *Clinical ophthalmology (Auckland, NZ).* (2009) 3:559–66. doi: 10.2147/OPTH.S3308
71. Lewczuk N, Zdebek A, Bogusławska J. Interferon Alpha 2a and 2b in Ophthalmology: a review. *J Interferon Cytokine Res.* (2019) 39:259–72. doi: 10.1089/jir.2018.0125