



Enantioselective Utilization of D-Amino Acids by Deep-Sea Microorganisms

Takaaki Kubota^{1*}, Tohru Kobayashi¹, Takuro Nunoura¹, Fumito Maruyama² and Shigeru Deguchi¹

¹ Research and Development Center for Marine Biosciences, Japan Agency for Marine-Earth Science and Technology, Yokosuka, Japan, ² Department of Microbiology, Graduate School of Medicine, Kyoto University, Kyoto, Japan

OPEN ACCESS

Edited by:

Hongyue Dang,
Xiamen University, China

Reviewed by:

Gerhard J. Josef Herndl,
University of Vienna, Austria
Uener Kolukisaoglu,
University of Tübingen, Germany

*Correspondence:

Takaaki Kubota
takaakikubota@jamstec.go.jp

Specialty section:

This article was submitted to
Aquatic Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 25 November 2015

Accepted: 29 March 2016

Published: 19 April 2016

Citation:

Kubota T, Kobayashi T, Nunoura T,
Maruyama F and Deguchi S (2016)
Enantioselective Utilization of D-Amino
Acids by Deep-Sea Microorganisms
Front. Microbiol. 7:511.
doi: 10.3389/fmicb.2016.00511

Microorganisms that utilize various D-amino acids (DAAs) were successfully isolated from deep-sea sediments. The isolates were phylogenetically assigned to Alphaproteobacteria, Gammaproteobacteria, and Bacilli. Some of the isolates exhibited high enantioselective degradation activities to various DAAs. In particular, the Alphaproteobacteria *Nautella* sp. strain A04V exhibited robust growth in minimal medium supplemented with D-Val as a sole carbon and nitrogen source, whereas its growth was poor on minimal medium supplemented with L-Val instead of D-Val. Its growth was facilitated most when racemic mixtures of valine were used. In contrast, the *Nautella* strains isolated from shallow-sea grew only with L-Val. No significant differences were found among the strains in the genome sequences including genes possibly related to DAA metabolisms.

Keywords: D-amino acid deaminase, D-amino acid utilization, D-valine, deep sea, marine bacteria

INTRODUCTION

Homochirality is one of the most important features of biological systems. Why and how life on earth chose L-amino acids (LAAs) instead of D-amino acids (DAAs) to build proteins remains open questions related to the chemical origins of life (Siegel, 1998; Blackmond, 2010; Hein and Blackmond, 2012). In the context of the homochirality of life, it was considered that organisms preferentially utilized LAAs over DAAs. However, recently, there have been a number of reports that the organisms utilize DAAs.

It is well-known that the peptidoglycan in bacterial cell walls comprise of DAAs (Schleifer and Kandler, 1972; Vollmer et al., 2008; Typas et al., 2011). The glycan strands are made of N-acetylglucosamine and N-acetylmuramic acid units that are connected by β -1,4-glycosidic linkages, and short peptide chains that cross-link the strands are enriched with D-Ala and D-Glu. Other DAAs such as D-Met, D-Val, D-Phe, and D-Trp are also found in the peptidoglycan (Lam et al., 2009; Cava et al., 2010). DAAs are also found in various antibiotic peptides that are produced by bacteria (Brückner and Fujii, 2010a,b; Velkov et al., 2010; Tempelaars et al., 2011; Doveston et al., 2012).

DAAs even mediate vital physiological functions in mammals. In the central nerve system, for example, D-Ser that is racemized from L-Ser by serine racemase acts as

Abbreviations: DAA, D-amino acid; LAA, L-amino acid; MBTH, 3-methyl-2-benzothiazolinone hydrazine; TCA, Trichloroacetic acid; DOM, dissolved organic matter.

a co-agonist for the *N*-methyl-D-aspartate receptor and controls synaptic plasticity and memory functions (Baumgart and Rodríguez-Crespo, 2008; Billard, 2012). Irukayama-Tomobe et al. (2009) reported that aromatic DAAs (D-Phe and D-Trp) act as bacteria-derived chemoattractants that elicit a chemotactic response in human neutrophils. They also suggested that D-Phe and D-Trp had lipid-lowering effects in human blood.

Identification of metabolic pathways relating to DAAs is crucial to understand their physiological functions. Such knowledge may provide novel insights into various pathophysiological conditions, thereby leading to development of new therapeutic strategies. However, our knowledge of the diversity of DAA metabolic pathways is still extremely limited.

The rates of DAA utilization and degradation are significantly lower than those of LAAs in soil and oceanic waters (Hopkins and Ferguson, 1994a; Hopkins et al., 1994b; O'Dowd et al., 1999; Amon et al., 2001; Calleja et al., 2013; Azúa et al., 2014). DAAs have been detected in diverse environments such as soil (Pollock et al., 1977), rivers (Stepanuskas et al., 2000; Dittmar et al., 2001; Tremblay and Benner, 2009), lakes (Kawasaki and Benner, 2006), and oceans (Lee and Bada, 1977; McCarthy et al., 1998; Dittmar et al., 2001; Kawasaki and Benner, 2006; Wedyan and Preston, 2008). Intriguingly, the abundance of DAAs among the total amino acids was higher in oceanic dissolved organic matter (DOM) than in terrestrial DOM (Dittmar et al., 2001; Kawasaki and Benner, 2006).

In oceanic environments, the presence of DAA-utilizing microbes was suggested by the activity measurements. In deep-sea water, the greater assimilation rate of D-Asp than that of L-Asp was observed (Perez et al., 2003). However, microbes that utilize DAAs as carbon and nitrogen sources have not been previously isolated from marine environments. Here, we report the first isolation of DAA-utilizing bacteria from deep-sea sediments. Characteristics of physiological and genomic traits of a representative strain are also given.

MATERIALS AND METHODS

Strains

Nautella italica LMG24365 was provided by the Belgian Co-ordinated Collections of Micro-organisms/Laboratory of Microbiology UGent (BCCM/LMG) Bacteria Collection (Ghent, Belgium). *N. italica* strain R11 was kindly provided by Dr. Torsten Thomas (University of New South Wales, Sydney, Australia).

Screening of DAA-Utilizing Microorganisms

Deep-sea sediments were collected from Sagami Bay by manned submersibles, *Shinkai 2000* and *Shinkai 6500*, and by a remotely operated vehicle, *ROV Hyper Dolphin*, during Japan Agency for Marine-Earth Science and Technology (JAMSTEC) expeditions (Cruise NT01-11, NT04-06, YK05-15, NT06-01, NT06-17, YK07-05, NT08-24) (Table S1). The samples were stored at -80°C .

To enrich DAA-utilizing bacteria in the sediments, two types of media, DBM and DAM, were prepared using artificial seawater (Daigo's artificial seawater SP for marine microalgae medium, Nihon Pharmaceutical Co. Ltd., Tokyo, Japan) as a basal medium.

DBM was used for screening of bacteria utilizing branched-chain DAA, and composed of artificial seawater (pH 8.0) supplemented with 0.005% (wt/v) D-valine, 0.005% D-leucine, 0.1% glucose, 0.001% yeast extract (Becton Dickinson, Franklin Lakes, New Jersey, USA), and 0.005% K_2HPO_4 . DAM was used for screening of bacteria utilizing aromatic DAAs, and composed of artificial seawater (pH 8.0) supplemented with 0.005% D-phenylalanine, 0.005% D-tryptophan, 0.1% glucose, 0.001% yeast extract, and 0.005% K_2HPO_4 . DAAs and LAAs were obtained from Wako Pure Chemical Industries (Osaka, Japan). When the media was prepared for isolating colonies on plate, 1.5% (wt/v) agar was added to DBM and DAM.

Slurry of the deep-sea sediments was inoculated into 2 mL of DBM or DAM in 5-mL test tubes. The cultures were incubated at 30°C for 5 days with shaking at 130 rpm. Afterwards, 0.02 mL of enrichment was inoculated into fresh medium, and incubated again under the same conditions. This procedure was repeated three times. Each enrichment cultures thus obtained were spread onto glucose-deficient DBM or DAM plates and incubated at 30°C . Morphologically distinct colonies were selected, transferred into 2 mL of the same liquid medium, and incubated at 30°C with shaking at 130 rpm. The resulting cultures were then spread onto solid media and incubated at 30°C . This purification procedure was repeated twice. The colonies in this secondary enrichment culture were isolated as DAA-utilizing bacteria.

Nucleotide Sequences of 16S rRNA Genes and Phylogenetic Analysis

A universal primer set of bacterial 16S rRNA gene was used for PCR amplification of the 16S rRNA genes from the isolates (Lane, 1991). A colony of each isolate was used as a template in the PCR reaction with LA Taq DNA polymerase (Takara Bio, Otsu, Japan) using a thermal cycler (model ABI 9700; Applied Biosystems, Foster City, CA, USA). The PCR products were directly sequenced using a DNA sequencer (ABI3130; Applied Biosystems) and an ABI Prism BigDye terminator sequencing kit (Applied Biosystems). 16S rRNA gene sequences are deposited in DDBJ/EMBL/GenBank database under accession numbers summarized in Table S2.

Preparation of Resting-Cell Suspensions

N. italica LMG24365, *N. italica* strain R11, and our isolates were pre-cultured in 20-mL test tubes containing 4 mL Difco Marine Broth 2216 (MB; Becton Dickinson) at 30°C with shaking at 120 rpm. Aliquots (0.5 mL) of the pre-culture were transferred to 50 mL MB in 200-mL Erlenmeyer flasks and incubated at 30°C with shaking at 180 rpm. Cells were harvested during the pre-stationary phase by centrifugation at $10,000 \times g$ for 15 min at 4°C , washed twice with 3% NaCl, and once with 50 mM HEPES-NaOH buffer (pH 7.5). The washed cells were suspended in 50 mM HEPES-NaOH buffer (pH 7.5) containing 10% dimethyl sulfoxide. The cell concentration was adjusted to 100 mg wet cell mL^{-1} and stored at -80°C .

Before activity measurements, the cell concentrations of the resting cell samples were measured turbidimetrically at 660 nm (OD_{660}) after diluting the culture broths with the appropriate medium. The optical density value was converted into the dry cell

weight (DCW) using a calibration equation that was established by plotting the weight of a known volume of desiccated cells. The desiccated cells were prepared by washing the wet-cell samples (usually from 45 mL culture broth) twice with pure water and drying them at 105°C for 12 h.

Assay of Amino Acid Degradation Activity

The amino acid degradation activity was measured based on the quantity of α -keto acid produced from each amino acid (AA) using the 3-methyl-2-benzothiazolinone hydrazone (MBTH) method (Soda, 1968). Various DAAs and LAAs (aliphatic AAs: Ala, Val, Leu, and Ile; aromatic AAs: Phe, and Trp; hydroxyl AAs: Ser and Thr; sulfur-containing AAs: Cys and Met; AA amides: Asn and Gln; imino acid: Pro; acidic AAs: Asp and Glu; and basic AAs: Lys, Arg, and His) were used as substrates.

The reaction mixture (50 μ L) was composed of 50 mM HEPES-NaOH buffer (pH 7.5), 50 mM DAA or LAA, and 0.5–6.75 μ g-DCW mL⁻¹ resting cells. The reaction was performed at 30°C for 60 min and terminated by heating at 96°C for 5 min. The reaction mixture was centrifuged at 10,000 \times g at room temperature for 15 min to separate the cells, and 20 μ L of supernatant was taken and mixed with 10 μ L 1 M sodium acetate (pH 5.0) and 10 μ L 25 mM MBTH. The mixture was incubated at 50°C for 30 min and then cooled to room temperature. The absorbance was measured at 320 nm against a blank, which contained all of the components except amino acids. α -Ketoglutaric acid was used as a calibration standard.

For determination of the kinetic parameters, the reaction was performed at 30°C for 30 min in 50 mM HEPES-NaOH buffer (pH 7.5) with 1, 5, 25, 50, 100, or 125 mM substrate, and 0.5–6.75 μ g-DCW mL⁻¹ resting-cells.

D- and L-Valine Degradation Studies

Nautella sp. strain A04V and *N. italica* LMG24365 were pre-cultured in 20-mL test tubes containing 5 mL MB for 12–15 h at 30°C with shaking at 120 rpm. Cells were harvested by centrifugation at 5,000 \times g for 5 min, and washed three times with sterile minimal medium composed of 2.6% (wt/v) NaCl, 0.08% KCl, 0.56% MgCl₂·6H₂O, 0.76% MgSO₄·7H₂O, 0.0005% FeSO₄, 0.154% CaCl₂·2H₂O, and 0.01% Na₂HPO₄ (pH 7.8). Cells were suspended in the same medium and the cell density was adjusted to OD₆₆₀ = 0.1. Aliquots (0.05 mL) of the cell suspension were inoculated into 5 mL minimal medium in L-shaped test tubes, which contained 6 mM D-, and L-valine mixtures (D:L= 0:6, 1:5, 2:4, 3:3, 4:2, 5:1, or 6:0) as the sole carbon and nitrogen source. For each strain, seven cultures were prepared in duplicate and incubated at 30°C for 250 h with shaking at 70 rpm. The OD₆₆₀ was recorded automatically every 30 min using an Advantec TVS062CA biophotorecorder (Advantec Toyo Co. Ltd., Tokyo, Japan). The morphologies of the cultured cells were observed using an Olympus BX51 microscope (Tokyo, Japan).

Analysis of L- and D-Valine Using HPLC

The valine concentration was measured using a Shimadzu HPLC Prominence system (Shimadzu Corporation, Kyoto, Japan).

Valine was derivatized to a fluorescent diastereomer, and detected by using a Shimadzu fluorimeter (RF 20Axs, Kyoto, Japan). The excitation and emission wavelengths for detection were set to 350 and 450 nm, respectively.

A 30 μ L aliquot of the culture medium was sampled during each growth phase of the cultures, and centrifuged at 10,000 \times g at 4°C for 15 min. The supernatant was diluted to the appropriate concentrations with distilled water. A 10 μ L aliquot was then mixed with 10 μ L 0.1 N HCl, 80 μ L 0.2 M borate-NaOH buffer (pH 10) containing 1.75 mg mL⁻¹ *N*-isobutyl-L-cysteine, and 1.25 mg mL⁻¹ *o*-phthalaldehyde. The mixture was filtered through 0.20 μ m PTFE cartridge filters (Millipore, MA, USA) and then injected into an RP-HPLC column (YMC-pack Pro C18, 4.6 mM i.d. \times 250 mM). Gradient elution was applied using the following eluents: A, 10 mM sodium-borate/10 mM sodium-phosphate buffer (pH 8.2); B, 40% methanol/40% acetonitrile/10% water. The gradient program was as follows: 35% B for 1 min; 35–60% B for 17 min; and 60% B for 3 min. The derivatized pure L- and D-valine were used as calibration standards.

Draft Genome Sequencing, Mapping, Assembly, and Annotation

Genomic DNA was extracted from strains A04V and LMG24365 using Qiagen DNeasy Tissue Kits (Qiagen, Valencia, CA, USA). Genomic DNA sequencing of the strains was performed at the Beijing Genome Institute (BGI; Shenjun, China) using the Illumina HiSeq 2000 platform (Illumina Inc., San Diego, CA, USA) with the single-end strategy for 90-bp reads. The quality of the reads was checked and trimmed (Phred quality score >15) with FASTQ tools using the online data analysis platform Galaxy (Goecks et al., 2010). The sequencing reads were aligned with the *N. italica* strain R11 genome reference using Bowtie2 software version 2.1.0 with the default settings. The unmapped reads of strain A04V in the reference genome were assembled into 2587 contigs using ABySS software version 1.3.2 (using kmer = 31). The protein-coding sequences (CDSs) were identified using GLIMMER software version 3.0.2 (Delcher et al., 2007) and those shorter than 50 codons were eliminated. Functional assignments of the predicted genes were made using BLASTP ($e \leq e^{-5}$) with the GenBank and Pfam databases (pfam-A, version 26.0; Punta et al., 2012). Draft genome sequences are deposited in DDBJ/EMBL/GenBank database under PRJDB4276 and PRJDB4510 (Table S2).

RESULTS AND DISCUSSION

Enrichment and Isolation of DAA-Utilizing Bacteria from Deep-Sea Sediments

To isolate DAA-utilizing bacteria, we used branched-chain DAAs (D-Val and D-Leu) and aromatic DAAs (D-Phe and D-Trp) as the substrates for enrichment. These DAAs have been detected as minor D-forms in the Sargasso Sea (D-Ile) (Lee and Bada, 1977), in the Equatorial Pacific (D-Leu) (Lee and Bada, 1977; McCarthy et al., 1998), and in the Gulf of Mexico (D-Phe) (McCarthy et al., 1998).




FIGURE 1 | Phylogenetic tree based on the 16S rRNA gene sequences of D-amino acid-degrading isolates from deep-sea sediments. *Thermotoga maritima* MSB-8 was used as an outgroup. The phylogenetic tree was constructed using the neighbor-joining method (Saitou and Nei, 1987). The accession numbers of the sequences are shown in parentheses. Numbers at each node indicate the bootstrap value based on 1000 replicates. Strains obtained in this study are shown in bold and the first letter of the strain number indicates the screening medium used: DAM (A) or DBM (B). Figures following the strain numbers are identifiers employed in this study.


Fifty six sediment samples from Sagami Bay, Japan (at water depths ranging from 800 to 1,500 m, Table S1) were inoculated into DBM and DAM, and incubated aerobically to obtain enrichment cultures of DAA-degrading bacteria using a total of 112 tubes. After enrichment and subsequent isolation of colonies, we obtained 17 and 11 isolates that produced morphologically distinct colonies on DBM and DAM agar plates, respectively. The purity of the isolates was confirmed by microscopic observation and by sequencing their 16S rRNA genes.

Partial nucleotide sequences (0.8–1.4 kb) of the 16S rRNA genes of the 28 isolates showed that 17 strains

belonged to Alphaproteobacteria, 5 strains belonged to Gammaproteobacteria, and 6 strains belonged to Bacilli (Figure 1). There were no significant relationships between the taxonomic diversity of the DAA-degrading bacteria and the two types of screening media used for their enrichment and isolation.

Determination of the DAA Degradation Activities of the Isolates

The DAA degradation activities of the isolates were examined using a spectrometric method with resting cells (Figure 2). The alphaproteobacterial strains, i.e., *Nautella* sp. strain A04V and




A04F (numbers 2 and 4 in Figure 2, respectively), and *Paracoccus* sp. strain A18Fa and A30V (numbers 13 and 14 in Figure 2, respectively), showed high DAA degradation activities compared with the others. Interestingly, the strains A04V, A04F, and A18Fa degraded various DAAs, but in most cases, did not degrade their L-counterparts in this reaction condition (Figure 3). The resting cells of the alphaproteobacterial isolates degraded not only the branched-chain and aromatic DAAs that were supplemented to the culture media for enrichment and isolation, but also D-Met and D-Pro as well as basic DAAs, i.e., D-Lys, D-Arg, and D-His. The isolates with no or weak DAA degradation activities showed poor growth on the glucose-deficient DBM or DAM. We also evaluated the amino acid racemase activities in resting cells according to the method described by Asano and Endo (1988), but did not observe significant activity except for alanine.

The presence of metabolic pathways to utilize DAAs has been reported in Gammaproteobacteria, i.e., *Pseudomonas fluorescens* (Tsukada, 1966), *Pseudomonas aeruginosa* (Marshall

and Sokatch, 1968), *Salmonella typhimurium* (Wild et al., 1974), and *Escherichia coli* (Raunio et al., 1973); Epsilonproteobacteria, i.e., *Helicobacter pylori* (Tanigawa et al., 2010); and Bacilli, i.e., *Ureibacillus thermosphaericus* (Akita et al., 2014). *P. aeruginosa* PAO1 is known to utilize six DAAs (D-Arg, D-Glu, D-Gln, D-Ala, D-Asn, and D-Val) as the sole carbon and nitrogen source in bacterial cultures, and harbor two amino acid dehydrogenases (LAA dehydrogenase and DAA dehydrogenase) for utilization of DAAs (Li et al., 2010). However, it remained unclear whether the pathway of DAA utilization is ubiquitously present among marine bacteria (Zhang et al., 2016). Our results suggest the presence of metabolic pathways to utilize DAAs in marine Alphaproteobacteria.

Growth Behavior of *Nautella* sp. Strain A04V

Among the four strains of *Nautella* that exhibited degradation activities to DAAs (Figure 2), strain A04V had the broadest DAA substrate range, i.e., D-Ala, D-Val, D-Leu, D-Ile, D-Asp,





D-Glu, and D-Phe. This strain grew on these DAAs and also utilized D-Ser, D-Thr, D-Cys, or D-Arg as the sole carbon and nitrogen source on solid media. Therefore, we used strain A04V to characterize marine DAA-degrading bacteria in greater detail. *Nautella* strain A04V was closely related to *N. italica* strain LMG24365 and strain R11 based on 16S rRNA gene similarity analysis (100% similarity).

During its growth with branched-chain AAs such as Val and Leu, strain A04V preferred the D-forms for growth rather than the L-forms (Figure 4). The preference for the D-forms of AAs was not observed when we tested other AAs such as Ala, Glu, and Phe as sole carbon and nitrogen sources (Figure S1). The optimum D-Val concentration for growth was 5–6 mM. Adverse effects of high concentrations of DAAs have been reported previously in a variety of bacteria (Teeri and Josselyn, 1953; Bopp, 1965; Champney and Jensen, 1970; Cosloy and McFall, 1973; Yabu and Huempfer, 1974; White, 1979). The growth of strain A04V was also completely inhibited in the presence of >10 mM D-Val. In contrast, high concentrations (<100 mM) of L-Val did not inhibit the growth of this strain at all. High concentrations of DAAs induce significant morphological changes for *Vibrio*

cholerae cells (Lam et al., 2009), but such effect was not observed for strain A04V even when it was incubated with a lethal level of D-Val (10 mM).

Moreover, strain A04V exhibited unique growth behavior in the presence of racemic Val, i.e., a mixture of L- and D-Val (the final total Val concentration was 6 mM), as the sole carbon and nitrogen source (Figure 5A). Growth was robust in the presence of both Val enantiomers, whereas growth was weak with each separate enantiomer of Val. In contrast, the type strain of *N. italica* (strain LMG24365), which was isolated from a shallow-sea environment (Vandecastelaere et al., 2009), grew only with L-Val, and not with D-Val or racemic mixtures (Figure 5B). As strain A04V and *N. italica* LMG24365 share 100% similarity in terms of its 16S rRNA genes, the capacity for utilizing D-Val as the sole carbon and nitrogen source for growth is not a common feature of this genus.

The rate of consumption of Val (D-, L-, and D, L-Val) during the growth of strain A04V was determined by HPLC (Figure 6). D-, L-, and D, L-Val were completely depleted from the medium in the early stationary phase. Intriguingly, D-Val was consumed faster than L-Val during cultivation when the L- or D-form



was supplemented as sole carbon and nitrogen sources, whereas L-Val and D-Val were consumed at the same rate when racemic mixtures were used (Figure 6).

In a soil microbial assemblage, LAAs were consumed faster than DAA except for alanine (Zhang and Sun, 2014). A lag time was observed to induce racemase for conversion of D-enantiomers to L-enantiomers except for alanine racemases that are synthesized constitutively in most microbes (Zhang and Sun, 2014). The marine isolate A04V was distinctly different. Although it consumed D- and L-alanine at equal rates (Figure S1), the consumption rates of D-enantiomers

of branched-chain AAs in strain A04V were faster than L-enantiomers.

DAA Degradation Activity of Strain A04V and Close Relatives

To elucidate the biochemical features of strain A04V, we compared the DAA degradation activities of *N. italica* strains A04V and LMG24365. The catalytic activities of the resting cells of strain A04V indicated that they could degrade various DAAs (Figure 7). DAA degradation activities were also detected in the resting cells of *N. italica* strains LMG24365, but they were




FIGURE 6 | Effects of valine concentrations in the medium on the growth of *Nautella* sp. strain A04V. The substrates used for cultivation were: 6 mM D-Val (A), 6 mM racemic Val at D:L = 5:1 mM (B), D:L = 4:2 mM (C), D:L = 3:3 mM (D), D:L = 2:4 mM (E), D:L = 1:5 mM (F), and 6 mM L-Val (G) as the sole carbon and nitrogen sources. The concentrations of D-Val (filled circles) and L-Val (open circles) were determined by HPLC. The growth of the strain was measured instrumentally by determining the optical density at 660 nm (OD₆₆₀) (open triangles).

extremely low compared with those of strain A04V (Figure 7). Both strains showed similar substrate specificities.

We also determined the kinetic parameters of the DAA degradation activities of these strains. The catalytic activity (V_{max}) of valine degradation of strain A04V was 20 times higher than that of LMG24365 (20.58 ± 2.88 , 1.11 ± 0.02), but both strains showed similar substrate affinity (K_m) (5.68 ± 0.57 for A04V, 8.77 ± 2.0 for LMG24365) (Table S3).

Enzymatic degradation of DAAs did not occur when the disrupted cells of *Nautella* strains, including the cell wall and membrane fraction, were used in the reaction, indicating that the enzymes involved in enantioselective degradation are membrane-bound proteins and/or located in the periplasmic space. Similar results were previously reported for bacterial DAA deaminases (Tsukada, 1966; Olsiewski et al., 1980). These enzymes catalyze deamination of various DAAs, but works




FIGURE 7 | Deamination of amino acids to α -keto acids catalyzed by resting cells of strains A04V, *N. italica* LMG24365, and R11. Reactions were performed at 30°C for 1 h. Double asterisks (**) represent amino acids with L-enantiomers that were preferentially deaminated and the single asterisk (*) denotes those that deamination occurred preferentially for D-enantiomers. The data represent the mean \pm SD (indicated by vertical bars) based on at least two independent experiments.

especially well for branched-chain, basic, and aromatic DAAs (Tsukada, 1966; Olsiewski et al., 1980; Satomura et al., 2002; Li et al., 2010). The DAA degradation activities of the A04V cells showed very similar substrate specificity to the known DAA deaminases (Figure 7). DAA aminotransferases are also involved in DAA metabolic pathway in several bacteria (Tanizawa et al., 1989; Lee et al., 2006). The DAA aminotransferases catalyze the transamination using the several DAAs as substrates, but they show no or weak activities for branched-chain and aromatic DAAs (Tanizawa et al., 1989; Lee et al., 2006). Accordingly, the DAA degradation activities of the *Nautella* strains are likely catalyzed by DAA deaminases.

Genomic Analysis of *Nautella* Strains

A complete genome sequence is only available for *N. italica* strain R11 in this genus (Fernandes et al., 2011), which was isolated as a pathogen causing a bleaching disease of red algae (Case et al., 2011). The strain showed DAA degradation activities similar to strain LMG24365 (Figure 7, Table S3), and grew with L-Val but not D-Val. Thus, we analyzed genome sequences of strains A04V and LMG24365, and performed comparative genome analysis of the three strains.

The genome sequences were analyzed using the Illumina HiSeq 2000 platform to identify the genomic backgrounds of DAA degradation in strain A04V. In total, we obtained 7,101,444 and 7,191,597 reads (90 bp) for strains A04V and LMG24365, respectively (Table S4). After removing low-quality reads, these reads were then mapped onto the genomic sequence of strain R11. The mapped reads shared 87% (6,190,244 out of 7,101,444 reads) of A04V and 90% (6,522,580 out of 7,191,597 reads) of LMG24365 in each of the total reads. The coverage rates were 94.9% (depth: 132.4 \times) and 96% (depth: 137.6 \times), respectively,

relative to the reference genome of strain R11. The average GC contents were 59%. The average nucleotide similarities with *N. italica* strain R11 were 97% (A04V) and 98% (LMG24365), respectively.

The genome sequences from strain R11 contained six predicted CDSs for putative DAA oxidases (or dehydrogenases) (pfam01266) and four CDSs for putative DAA transferases (pfam01063). CDSs were also identified in the draft genomes of strains A04V and LMG24365. The shared amino acid identities of the genes among the three strains were 99–100% and only non-conservative amino acid substitutions were observed (Table 1). Furthermore, we found no other candidate genes related to DAA degradation by BLAST and Pfam analyses of the mapped reads from strain A04V.

The unmapped reads for A04V and LMG24365 were assembled into 2,587 (N50 contig size: 613) and 716 (N50 contig size: 1,425) contigs, respectively. Analysis of the contig sequences of A04V detected 696 candidate peptide sequences, which contained 300 sequences similar to those of LMG24365. As a result, we found no candidate genes encoding DAA degradation enzymes among these sequences. According to the comparative genome analysis, we observed little difference between the genes involved with the known DAA metabolism processes in the three *Nautella* strains. Thus, we suggest that a few differences in terms of amino acid substitutions might explain the differences in DAA utilization among the three strains. However, according to our analysis, the amino acid substitutions in the six genes that encoded the putative DAA deaminase of strain A04V (shown in Table 1) had no significant relationships with the bacterial habitats based on analyses of the corresponding gene sequences in bacterial genome databases (including the sequences of the environmental sample).

TABLE 1 | Comparisons of the CDSs encoding known DAA deaminases among *Nautella* strains.

Pfam	Protein ID in strain R11	Annotation	Shared identity (%)		Amino acid substitution	
			A04V	LMG24365	A04V	LMG24365
DAO (Pfam01266: D-AMINO ACID OXIDASE)						
	RR11_23	Glycine/D-amino acid oxidase, deaminating	99	100	5	0
	RR11_898	D-Amino acid dehydrogenase	99	100	2	0
	RR11_1993	D-Amino acid dehydrogenase small subunit	99	100	4	0
	RR11_2500	Glycine/D-amino acid oxidase	99	100	4	0
	RR11_2832	Glycerol-3-phosphate dehydrogenase	99	100	2	0
	RR11_2957	Glycine/D-amino acid oxidase	99	100	3	0
AMINOTRAN_4 (Pfam01063: D-AMINO ACID TRANSFERASES)						
	RR11_862	D-Alanine aminotransferase	100	100	0	0
	RR11_1631	Branched-chain amino acid aminotransferase	100	100	0	0
	RR11_2182	Branched-chain amino acid aminotransferase	99	99	2	2
	RR11_3119	Aminotransferase, class IV	99	100	1	0

The accession numbers of the sequences are shown in Table S2.

Moreover, the six genes identified by the genomic analysis (shown in Table 1) as encoding putative DAA deaminases were cloned and expressed in *E. coli*. However, no significant activities were detected in the recombinant *E. coli* cells or the cell extracts (unpublished data), which suggests that the gene products require other components for activation such as membrane-associated molecules.

CONCLUSION

Microbiological and geochemical studies suggested the microbial production and consumption of DAAs in pelagic water ecosystems (Perez et al., 2003; Teira et al., 2006; Calleja et al., 2013; Zhang et al., 2016). In this geochemical cycle, importance of marine Alphaproteobacteria in the production of DAAs has already been pointed out (Pedersen et al., 2001; Lomstein et al., 2006; Kaiser and Benner, 2008; Kubota et al., 2016), however, phylogenetic and physiological features of DAA-utilizing marine bacteria are poorly investigated yet.

In this study, various DAA-utilizing bacteria were successfully isolated from deep-sea environments, and their catalytic activities with L- and D-AAAs were evaluated using resting cells. Especially, the resting cells of the several alphaproteobacterial isolates showed significantly higher DAA degradation activities than those of other isolates. The strain A04V exhibited robust growth on medium supplemented with D-Val, whereas its growth was poor on minimal medium supplemented with L-Val as a sole carbon and nitrogen source. In contrast, other *Nautella* strains (strain R11 and LMG24365) isolated from a shallow-sea

environment grew with L-Val but not D-Val. The catalytic activity of the DAA degradation in the strain A04V was 20 times greater than that of the other two *Nautella* strains. However, there are no significant differences in the gene sets involved with the DAA metabolisms among the three *Nautella* strains. These results suggest that minor genetic variance in the gene sets involved DAA metabolisms and/or gene regulation influence the variance of DAA utilization among the strains. Therefore, combination of reverse genetics and biochemical analyses is necessary to understand the metabolism and physiological role of DAAs in the marine DAA-utilizing bacteria.

AUTHOR CONTRIBUTIONS

TaK carried out conception and design of the study, analysis and interpretation of data, collection and assembly of data, drafting of the article, and critical revision of the article for important intellectual content. ToK, SD conceived of the study, and participated in its design and coordination and helped to draft the manuscript. TN, FM interpreted of data of the molecular genetic studies for the work and helped to drafted the manuscript. All authors read and approved the final manuscript.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2016.00511>

REFERENCES

Akita, H., Suzuki, H., Doi, K., and Ohshima, T. (2014). Efficient synthesis of D-branched-chain amino acids and their labeled compounds with stable isotopes using D-amino acid dehydrogenase. *Appl. Microbiol. Biotechnol.* 98, 1135–1143. doi: 10.1007/s00253-013-4902-1

Amon, R. M. W., Fitznar, H. P., and Benner, R. (2001). Linkages among the bioreactivity, chemical composition, and diagenetic state of marine dissolved organic matter. *Limnol. Oceanogr.* 46, 287–297. doi: 10.4319/lo.2001.46.2.0287

Asano, Y., and Endo, K. (1988). Amino acid racemase with broad substrate specificity, its properties and use in phenylalanine racemization. *Appl. Microbiol. Biotechnol.* 29, 523–527. doi: 10.1007/BF00260979

- Azúa, I., Goiriena, I., Baña, Z., Iriberry, J., and Unanue, M. (2014). Release and consumption of D-amino acids during growth of marine prokaryotes. *Microb. Ecol.* 67, 1–12. doi: 10.1007/s00248-013-0294-0
- Baumgart, F., and Rodríguez-Crespo, I. (2008). D-Amino acids in the brain: the biochemistry of brain serine racemase. *FEBS J.* 275, 3538–3545. doi: 10.1111/j.1742-4658.2008.06517.x
- Billard, J. M. (2012). D-Amino acids in brain neurotransmission and synaptic plasticity. *Amino Acids* 43, 1851–1860. doi: 10.1007/s00726-012-1346-3
- Blackmond, D. G. (2010). The origin of biological homochirality. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 366, 2878–2884. doi: 10.1098/rstb.2011.0130
- Bopp, M. (1965). Inhibition of *Agrobacterium tumefaciens* by D-amino acids. *Z. Naturforsch. B.* 20, 899–905.
- Brückner, H., and Fujii, N. (2010a). *D-Amino Acids in Chemistry, Life Sciences, and Biotechnology*. Weinheim: Wiley-VCH.
- Brückner, H., and Fujii, N. (2010b). Free and peptide-bound D-amino acids in chemistry and life sciences. *Chem. Biodivers* 7, 1333–1336. doi: 10.1002/cbdv.201000114
- Calleja, M. L., Batista, F. C., Peacock, M., Kudela, R., and McCarthy, M. D. (2013). Changes in compound specific $\delta^{15}\text{N}$ amino acid signatures and D/L ratios in marine dissolved organic matter induced by heterotrophic bacterial reworking. *Mar. Chem.* 149, 32–44. doi: 10.1016/j.marchem.2012.12.001
- Case, R. J., Longford, S. R., Campbell, A. H., Low, A., Tujula, N., Steinberg, P. D., et al. (2011). Temperature induced bacterial virulence and bleaching disease in a chemically defended marine macroalga. *Environ. Microbiol.* 13, 529–537. doi: 10.1111/j.1462-2920.2010.02356.x
- Cava, F., Lam, H., de Pedro, M. A., and Waldor, M. K. (2010). Emerging knowledge of regulatory roles of D-amino acids in bacteria. *Cell. Mol. Life Sci.* 68, 817–831. doi: 10.1007/s00018-010-0571-8
- Champney, W. S., and Jensen, R. A. (1970). Molecular events in the growth inhibition of *Bacillus subtilis* by D-tyrosine. *J. Bacteriol.* 104, 107–116.
- Cosloy, S. D., and McFall, E. (1973). Metabolism of D-serine in *Escherichia coli* K-12: mechanism of growth inhibition. *J. Bacteriol.* 114, 685–694.
- Delcher, A. L., Bratke, K. A., Powers, E. C., and Salzberg, S. L. (2007). Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* 23, 673–679. doi: 10.1093/bioinformatics/btm009
- Dittmar, T., Fitznar, H. P., and Kattner, G. (2001). Origin and biogeochemical cycling of organic nitrogen in the eastern Arctic Ocean as evident from D- and L-amino acid. *Geochim. Cosmochim. Acta.* 65, 4103–4114. doi: 10.1016/S0016-7037(01)00688-3
- Doveston, R. G., Steendam, R., Jones, S., and Taylor, R. J. (2012). Total synthesis of an oxepine natural product, (\pm)-janoxepin. *Org. Lett.* 14, 1122–1125. doi: 10.1021/ol300039x
- Fernandes, N., Case, R. J., Longford, S. R., Seyedsayamdost, M. R., Steinberg, P. D., Kjelleberg, S., et al. (2011). Genomes and virulence factors of novel bacterial pathogens causing bleaching disease in the marine red alga *Delisea pulchra*. *PLoS ONE* 6:e27387. doi: 10.1371/journal.pone.0027387
- Goecks, J., Nekrutenko, A., and Taylor, J. (2010). Galaxy: a comprehensive approach for supporting accessible, reproducible, and transparent computational research in the life sciences. *Genome Biol.* 11:R86. doi: 10.1186/gb-2010-11-8-r86
- Hein, J. E., and Blackmond, D. G. (2012). On the origin of single chirality of amino acids and sugars in biogenesis. *Acc. Chem. Res.* 45, 2045–2054. doi: 10.1021/ar200316n
- Hopkins, D. W., and Ferguson, K. E. (1994a). Substrate induced respiration in soil amended with different amino acid isomers. *Appl. Soil Ecol.* 1, 75–81. doi: 10.1016/0929-1393(94)90025-6
- Hopkins, D. W., Isabella, B. L., and Scott, S. E. (1994b). Relationship between microbial biomass and substrate induced respiration in soils amended with D- and L-isomers of amino acids. *Soil Biol. Biochem.* 26, 1623–1627. doi: 10.1016/0038-0717(94)90314-X
- Irukayama-Tomobe, Y., Tanaka, H., Yokomizo, T., Hashidate-Yoshida, T., Yanagisawa, M., and Sakurai, T. (2009). Aromatic D-amino acids act as chemoattractant factors for human leukocytes through a G protein-coupled receptor, GPR109B. *Proc. Natl. Acad. Sci. U.S.A.* 106, 3930–3934. doi: 10.1073/pnas.0811844106
- Kaiser, K., and Benner, R. (2008). Major bacterial contribution to the ocean reservoir of detrital organic carbon and nitrogen. *Limnol. Oceanogr.* 53, 99–112. doi: 10.4319/lo.2008.53.1.0099
- Kawasaki, N., and Benner, R. (2006). Bacterial release of dissolved organic matter during cell growth and decline: molecular origin and composition. *Limnol. Oceanogr.* 51, 2170–2180. doi: 10.4319/lo.2006.51.5.2170
- Kubota, T., Shimamura, S., Kobayashi, T., Nunoura, T., and Deguchi, S. (2016). Distribution of eukaryotic serine racemases in the bacterial domain and characterization of a representative protein in *Roseobacter litoralis* Och 149. *Microbiology* 162, 53–61. doi: 10.1099/mic.0.000200
- Lam, H., Oh, D. C., Cava, F., Takacs, C. N., Clardy, J., de Pedro, M. A., et al. (2009). D-Amino acids govern stationary phase cell wall remodeling in bacteria. *Science* 325, 1552–1555. doi: 10.1126/science.1178123
- Lane, D. J. (1991). “16S/23S rRNA sequencing,” in *Nucleic Acid Techniques in Bacterial Systematics*, eds E. Stackebrandt and M. Goodfellow (Chichester, UK: John Wiley and Sons), 115–175.
- Lee, C., and Bada, J. L. (1977). Dissolved amino acids in the equatorial Pacific, the Sargasso Sea, and Biscayne Bay. *Limnol. Oceanogr.* 22, 502–510. doi: 10.4319/lo.1977.22.3.0502
- Lee, S. G., Hong, S. P., Song, J. J., Kim, S. J., Kwak, M. S., and Sung, M. H. (2006). Functional and structural characterization of thermostable D-amino acid aminotransferases from *Geobacillus* spp. *Appl. Environ. Microbiol.* 72, 1588–1594. doi: 10.1128/AEM.72.2.1588-1594.2006
- Li, C., Yao, X., and Lu, C. D. (2010). Regulation of the dauBAR operon and characterization of D-amino acid dehydrogenase DauA in arginine and lysine catabolism of *Pseudomonas aeruginosa* PAO1. *Microbiology* 156, 60–71. doi: 10.1099/mic.0.033282-0
- Lomstein, B. A., Jørgensen, B. B., Schubert, C. J., and Niggemann, J. (2006). Amino acid biogeo- and stereochemistry in coastal Chilean sediments. *Geochim. Cosmochim. Acta.* 70, 2970–2989. doi: 10.1016/j.gca.2006.03.015
- Marshall, V. P., and Sokatch, J. R. (1968). Oxidation of D-amino acids by a particulate enzyme from *Pseudomonas aeruginosa*. *J. Bacteriol.* 95, 1419–1424.
- McCarthy, M. D., Hedges, J. I., and Benner, R. (1998). Major bacterial contribution to marine dissolved organic nitrogen. *Science* 281, 231–234. doi: 10.1126/science.281.5374.231
- O’Dowd, R. W., Barraclough, D., and Hopkins, D. W. (1999). Nitrogen and carbon mineralization in soil amended with D- and L-leucine. *Soil Biol. Biochem.* 31, 1573–1578.
- Olsiewski, P. J., Kaczorowski, G. J., and Walsh, C. (1980). Purification and properties of D-amino acid dehydrogenase, an inducible membrane-bound iron-sulfur flavoenzyme from *Escherichia coli* B. *J. Biol. Chem.* 255, 4487–4494.
- Pedersen, A. -G. U., Thomsen, T. R., Lomstein, B. A., and Jørgensen, N. O. G. (2001). Bacterial influence on amino acid enantiomerization in a coastal marine sediment. *Limnol. Oceanogr.* 46, 1358–1369. doi: 10.4319/lo.2001.46.6.1358
- Perez, M. T., Pausz, C., and Herndl, G. J. (2003). Major shift in bacterioplankton utilization of enantiomeric amino acids between surface waters and the ocean’s interior. *Limnol. Oceanogr.* 48, 755–763. doi: 10.4319/lo.2003.48.2.0755
- Pollock, G. E., Cheng, C. N., and Croni, S. E. (1977). Determination of the D and L isomers of some protein amino acids present in soils. *Anal. Chem.* 49, 2–7. doi: 10.1021/ac50009a008
- Punta, M., Coggill, P. C., Eberhardt, R. Y., Mistry, J., Tate, J., and Bournsnel, C., et al. (2012). The Pfam protein families database. *Nucleic Acids Res.* 40, D290–D301. doi: 10.1093/nar/gkr1065
- Raunio, R. P., Straus, L. P., and Jenkins, W. T. (1973). D-Alanine oxidase from *Escherichia coli*: participation in the oxidation of L-alanine. *J. Bacteriol.* 115, 567–573.
- Saitou, N., and Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4, 406–425.
- Satomura, T., Kawakami, R., Sakuraba, H., and Ohshima, T. (2002). Dye-linked D-proline dehydrogenase from hyperthermophilic archaeon *Pyrobaculum islandicum* is a novel FAD-dependent amino acid dehydrogenase. *J. Biol. Chem.* 277, 12861–12867. doi: 10.1074/jbc.M112272200
- Schleifer, K. H., and Kandler, O. (1972). Peptidoglycan types of bacterial cell walls and their taxonomic implications. *Bacteriol. Rev.* 36, 407–477.
- Siegel, J. S. (1998). Homochiral imperative of molecular evolution. *Chirality* 10, 24–27. doi: 10.1002/chir.5
- Soda, K. (1968). Microdetermination of D-amino acids and D-amino acid oxidase activity with 3, methyl-2-benzothiazolone hydrazone hydrochloride. *Anal. Biochem.* 25, 228–235. doi: 10.1016/0003-2697(68)90095-X

- Stepanuskas, R., Laudon, H., and Jørgensen, N. O. G. (2000). High DON bioavailability in boreal streams during a spring flood. *Limnol. Oceanogr.* 45, 1298–1307. doi: 10.4319/lo.2000.45.6.1298
- Tanigawa, M., Shinohara, T., Saito, M., Nishimura, K., Hasegawa, Y., Wakabayashi, S., et al. (2010). D-Amino acid dehydrogenase from *Helicobacter pylori* NCTC 11637. *Amino Acids* 38, 247–255. doi: 10.1007/s00726-009-0240-0
- Tanizawa, K., Masu, Y., Asano, S., Tanaka, H., and Soda, K. (1989). Thermostable D-amino acid aminotransferase from a thermophilic *Bacillus* species. Purification, characterization, and active site sequence determination. *J. Biol. Chem.* 264, 2445–2449.
- Teeri, A., and Josselyn, D. (1953). Effect of excess amino acids on growth of certain *Lactobacilli*. *J. Bacteriol.* 66, 72–73.
- Teira, E., van Aken, H., Veth, C., and Herndl, G. J. (2006). Archaeal uptake of enantiomeric amino acids in the meso- and bathypelagic waters of the North Atlantic. *Limnol. Oceanogr.* 51, 60–69. doi: 10.4319/lo.2006.51.1.0060
- Tempelaars, M. H., Rodrigues, S., and Abee, T. (2011). Comparative analysis of antimicrobial activities of valinomycin and cereulide, the *Bacillus cereus* emetic toxin. *Appl. Environ. Microbiol.* 77, 2755–2762. doi: 10.1128/AEM.02671-10
- Tremblay, L., and Benner, R. (2009). Organic matter diagenesis and bacterial contributions to detrital carbon and nitrogen in the Amazon River system. *Limnol. Oceanogr.* 54, 681–691. doi: 10.4319/lo.2009.54.3.0681
- Tsukada, K. (1966). D-Amino acid dehydrogenases of *Pseudomonas fluorescens*. *J. Biol. Chem.* 241, 4522–4528.
- Typas, A., Banzhaf, M., Gross, C. A., and Vollmer, W. (2011). From the regulation of peptidoglycan synthesis to bacterial growth and morphology. *Nat. Rev. Microbiol.* 10, 123–136. doi: 10.1038/nrmicro2677
- Vandecastelaere, I., Nercessian, O., Segart, E., Achouak, W., Mollica, A., Faimali, M., et al. (2009). *Nautella italica* gen. nov., sp. nov., isolated from a marine electroactive biofilm. *Int. J. Syst. Evol. Microbiol.* 59, 811–817. doi: 10.1099/ijs.0.002683-0
- Velkov, T., Thompson, P. E., and Nation, R. L., and Li, J. (2010). Structure–activity relationships of polymyxin antibiotics. *J. Med. Chem.* 53, 1898–1916. doi: 10.1021/jm900999h
- Vollmer, W., Blanot, D., and de Pedro, M. A. (2008). Peptidoglycan structure and architecture. *FEMS Microbiol. Rev.* 32, 149–167. doi: 10.1111/j.1574-6976.2007.00094.x
- Wedyan, M. A., and Preston, M. R. (2008). The coupling of surface seawater organic nitrogen and the marine aerosol as inferred from enantiomer-specific amino acid analysis. *Atm. Env.* 42, 8698–8705. doi: 10.1016/j.atmosenv.2008.04.038
- White, P. J. (1979). Inhibition by D-glutamate as the cause of diphasic growth of *Bacillus megaterium* NCIB 7581 with glycerol plus DL-glutamic acid. *Microbiology* 114, 149–158. doi: 10.1099/00221287-114-1-149
- Wild, J., Walczac, W., Krajewska-Grynkiewicz, K., and Klopotoski, T. (1974). D-Amino acid dehydrogenase: the enzyme of the first step of D-histidine and D-methionine racemisation in *Salmonella typhimurium*. *Mol. Gen. Genet.* 128, 131–146. doi: 10.1007/BF02654486
- Yabu, K., and Huempfer, H. R. (1974). Inhibition of growth of *Mycobacterium smegmatis* and of cell wall synthesis by D-serine. *Antimicrob. Agents Chemother.* 6, 1–10. doi: 10.1128/AAC.6.1.1
- Zhang, G., and Sun, H. J. (2014). Racemization in reverse: evidence that D-amino acid toxicity on Earth is controlled by bacteria with racemases. *PLoS ONE* 9:e92101. doi: 10.1371/journal.pone.0092101
- Zhang, Z., Zheng, Q., and Jiao, N. (2016). Microbial D-amino acids and marine carbon storage. *Sci. China Earth Sci.* 59, 17–24. doi: 10.1007/s11430-015-5155-x

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Kubota, Kobayashi, Nunoura, Maruyama and Deguchi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.