



Characterization of Shiga Toxin Subtypes and Virulence Genes in Porcine Shiga Toxin-Producing *Escherichia coli*

Gian Marco Baranzoni¹, Pina M. Fratamico^{1*}, Jayanthi Gangiredla², Isha Patel², Lori K. Bagi¹, Sabine Delannoy³, Patrick Fach³, Federica Boccia⁴, Aniello Anastasio⁴ and Tiziana Pepe⁴

¹ Eastern Regional Research Center, United States Department of Agriculture – Agricultural Research Service, Wyndmoor, PA, USA, ² Center of Food Safety and Applied Nutrition, U.S. Food and Drug Administration, Laurel, MD, USA, ³ Food Safety Laboratory, University of Paris-Est, Anses, Maisons-Alfort, France, ⁴ Department of Veterinary Medicine and Animal Production, University of Naples Federico II, Naples, Italy

OPEN ACCESS

Edited by:

Dustin Brisson,
University of Pennsylvania, USA

Reviewed by:

Jorge Blanco,
University of Santiago
de Compostela, Spain
Seamus Fanning,
University College Dublin, Ireland

*Correspondence:

Pina M. Fratamico
pina.fratamico@ars.usda.gov

Specialty section:

This article was submitted to
Evolutionary and Genomic
Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 26 February 2016

Accepted: 07 April 2016

Published: 21 April 2016

Citation:

Baranzoni GM, Fratamico PM, Gangiredla J, Patel I, Bagi LK, Delannoy S, Fach P, Boccia F, Anastasio A and Pepe T (2016) Characterization of Shiga Toxin Subtypes and Virulence Genes in Porcine Shiga Toxin-Producing *Escherichia coli*. *Front. Microbiol.* 7:574. doi: 10.3389/fmicb.2016.00574

Similar to ruminants, swine have been shown to be a reservoir for Shiga toxin-producing *Escherichia coli* (STEC), and pork products have been linked with outbreaks associated with STEC O157 and O111:H-. STEC strains, isolated in a previous study from fecal samples of late-finisher pigs, belonged to a total of 56 serotypes, including O15:H27, O91:H14, and other serogroups previously associated with human illness. The isolates were tested by polymerase chain reaction (PCR) and a high-throughput real-time PCR system to determine the Shiga toxin (Stx) subtype and virulence-associated and putative virulence-associated genes they carried. Select STEC strains were further analyzed using a Minimal Signature *E. coli* Array Strip. As expected, *stx*_{2e} (81%) was the most common Stx variant, followed by *stx*_{1a} (14%), *stx*_{2d} (3%), and *stx*_{1c} (1%). The STEC serogroups that carried *stx*_{2d} were O15:H27, O159:H16 and O159:H-. Similar to *stx*_{2a} and *stx*_{2c}, the *stx*_{2d} variant is associated with development of hemorrhagic colitis and hemolytic uremic syndrome, and reports on the presence of this variant in STEC strains isolated from swine are lacking. Moreover, the genes encoding heat stable toxin (*est*1a) and enteroaggregative *E. coli* heat stable enterotoxin-1 (*ast*A) were commonly found in 50 and 44% of isolates, respectively. The hemolysin genes, *hly*A and *ehx*A, were both detected in 7% of the swine STEC strains. Although the *eae* gene was not found, other genes involved in host cell adhesion, including *lpf*A_{O113} and *paa* were detected in more than 50% of swine STEC strains, and a number of strains also carried *iha*, *lpf*A_{O26}, *lpf*A_{O157}, *fed*A, *orf*A, and *orf*B. The present work provides new insights on the distribution of virulence factors among swine STEC strains and shows that swine may carry Stx1a-, Stx2e-, or Stx2d-producing *E. coli* with virulence gene profiles associated with human infections.

Keywords: *Escherichia coli*, STEC, swine, Shiga toxins variants, virulence genes

INTRODUCTION

Shiga Toxin-producing *Escherichia coli* (STEC) are food-borne pathogens responsible for outbreaks and serious illness including hemorrhagic colitis (HC) and hemolytic uremic syndrome (HUS). STEC O157:H7 is the serotype that has most often been associated with outbreaks and severe forms of diarrhea; however, recently a number of non-O157 STEC serogroups that cause similar illnesses have emerged (Gould et al., 2013). Cattle and other ruminants are important reservoirs of STEC; infection is asymptomatic, and they can carry the pathogens for long periods of time. Similarly, healthy swine may shed STEC, as demonstrated by several studies in which STEC were detected and isolated from swine fecal samples (Tseng et al., 2014b). Many of the investigations focused on serotype O157:H7; however, some studies also tested for non-O157 STEC serogroups and identified serogroups previously associated with human cases of illness (Fratamico et al., 2004; Kaufmann et al., 2006; Tseng et al., 2014b). The possibility that swine can transmit pathogenic STEC to humans is supported by a few outbreaks linked to the consumption of pork products contaminated with STEC O157:H7, O157:NM, and O111:H- (Tseng et al., 2014b).

Shiga toxins (Stx) are divided in two major antigenic forms: Stx1 and Stx2. Variants for Stx1 and Stx2 are grouped in three (Stx1a, Stx1c, Stx1d) and seven (Stx2a, Stx2b, Stx2c, Stx2d, Stx2e, Stx2f, and Stx2g) subtypes, respectively (Scheutz et al., 2012). Although Stx1a has been linked to human illness, STEC that produce subtypes Stx2a, Stx2c, and Stx2d are more often associated with the development of HC and HUS (Friedrich et al., 2002; Melton-Celsa, 2014). *In vitro* studies in two different cell lines showed that Stx2a and Stx2d were more potent than Stx2b and Stx2c. These results were also confirmed by experimentation in mice showing a significantly higher potency of Stx2a and Stx2d than Stx1, Stx2b, and Stx2c (Fuller et al., 2011). Stx variants are not homogeneously distributed among the STEC population and certain variants are frequently detected in association with different animals (Martin and Beutin, 2011; Hofer et al., 2012; Fuente et al., 2015). Swine STEC strains commonly produce Stx2e (Fratamico et al., 2004; Meng et al., 2014; Tseng et al., 2015), which may cause edema disease in weaned pigs, often leading to ataxia and death. Stx2e-producing *Escherichia coli*, do not represent a particular threat for humans (Friedrich et al., 2002; Tseng et al., 2014b). Nevertheless, STEC carrying the *stx2e* gene have been isolated from human cases with mild diarrhea (Muniesa et al., 2000; Friedrich et al., 2002; Beutin et al., 2004; Sonntag et al., 2005) and from two patients with HUS (Thomas et al., 1994; Fasel et al., 2014). The severe outcome of the first HUS case was probably due to a co-infection with another STEC strain (Thomas et al., 1994), while the second patient with HUS was described as having a very weak immune system (Fasel et al., 2014). Besides Stx2e, there is a lack of information on the presence of other Stx subtypes in STEC strains isolated from swine.

The production of Stx is necessary to provoke HUS; however, other virulence factors are also important in causing illness. These include genes involved in cell adhesion, proteases, and

toxins, as well as other putative virulence factors. The presence of specific combinations of virulence factors may determine the risk of developing severe symptoms. The *eae* gene, found on the locus of enterocyte effacement (LEE), encodes intimin, which is an adhesin involved in gut colonization. LEE-positive STEC are expected to provoke HUS or HC more frequently than LEE-negative STEC (Ethelberg et al., 2004; Toma et al., 2004; Luna-Gierke et al., 2014). Nevertheless, cases of HUS provoked by LEE-negative STEC have been reported (Karmali et al., 1985; Paton et al., 1999; Bielaszewska et al., 2009), including a large outbreak in 2011 in Europe caused by an enteroaggregative *E. coli* that acquired the *stx2a* gene, and it possessed a combination of virulence genes increasing its virulence (Boisen et al., 2015). This suggests that LEE is not essential in the development of severe symptoms, and other genes involved in adherence may also be important. Many adherence gene candidates, including *eibG*, *lpfA*, *saa*, and *sab* have been identified in STEC (Croxen et al., 2013). Nevertheless, mechanisms for attachment of LEE-negative STEC to the intestinal epithelium have not been studied as extensively as attachment of LEE-positive STEC.

In 2000, one objective of the U.S. Department of Agriculture's Animal and Plant Health Inspection Service National Animal Health Monitoring System (NAHMS) Swine 2000 study was to determine the prevalence of STEC in swine. Fecal samples were from states with the highest production of swine in the U.S. (U.S. Department of Agriculture, 2001). As a result of this work, 219 STEC isolates were recovered and characterized (Fratamico et al., 2004, 2008). Since this work was conducted, the knowledge of the importance of non-O157 STEC in human illness has increased, and there is a need to develop a model for molecular risk assessment associated with STEC. Knowledge of the virulence gene combinations that distinguish highly pathogenic *E. coli* from less virulent strains remains unclear, particularly for LEE-negative STEC (Beutin and Fach, 2014). Additionally, new virulence-associated and putative virulence-associated factors are being identified (Coombes et al., 2008; Brandt et al., 2011; Bugarel et al., 2011). The aim of the present study was to characterize STEC recovered from swine, belonging to a variety of serotypes to determine their Stx subtype and virulence gene profiles to understand their virulence potential.

MATERIALS AND METHODS

Bacterial Strains

Swine STEC strains were isolated and serotyped during the NAHMS swine 2000 study (NAHMS 2000) as described by Fratamico et al. (2004). Briefly, fresh swine feces were recovered from the pen floor of swine operations from the main pork-producing states in U.S. A total of 687 swine fecal samples were enriched using tryptic soy broth (TSB) and screened for the presence of *stx1* and *stx2* by polymerase chain reaction (PCR). Positive samples were plated onto Luria-Bertani agar, and *stx1*- and *stx2*-positive colonies were detected following DNA hybridization and confirmed by PCR. Two hundred and nineteen

STEC strains were serotyped and frozen in TSB with 20% of glycerol. From this collection, 181 STEC strains were used in this study and maintained on tryptic soy agar plates or TSB as working stock cultures.

Besides the NAHMS swine isolates, three STEC O91 strains from our collection were also used for comparison. STEC O91:H14 (strains 2.4111 and 2.4114) were isolated from ground beef while STEC O91:H21 (strain B2F1) was isolated from a case of HUS (Ito et al., 1990).

Identification of Shiga-toxin Subtypes

DNA extraction and PCR assays to identify *stx* subtypes and *stx* partial sequences were performed according to Scheutz et al. (2012) using a ProFlex PCR system (Thermo Fisher, Waltham, MA, USA) with slight modifications. TaqMan Environmental Master Mix 2.0 (Thermo Fisher) was used, and the annealing temperature was raised to 65°C when cross-reaction was observed, as suggested by the authors (Scheutz et al., 2012). Gel electrophoresis was performed using 1.5% UltraPure Agarose (Invitrogen, Carlsbad, CA, USA) gel with 0.5X GelRed (Phenix Research Products, Candler, NC, USA) in 1X Tris-acetate-EDTA buffer at 100 V for 1 h. One microliter of amplified DNA was analyzed by agarose gel electrophoresis and visualized using an AlphaImager gel documentation system (Alpha Innotech, San Leandro, CA, USA).

Polymerase chain reaction products for sequencing were cleaned with Agencourt AMPure XP (Beckman Coulter, Brea, CA, USA), and 1.2 µl were amplified in a reaction consisting of 7 µl of 2.5X buffer, 1 µl of 3.2 µM primer *stx2-F4* or *stx2-R1* (Scheutz et al., 2012), 1 µl of Big Dye Terminator (Applied Biosystems), and 10 µl of nuclease-free water. Thermocycling conditions consisted of 30 cycles of 95°C for 10 s, 55°C for 5 s and 60°C for 4 min. The sequencing reaction products were then purified and sequenced using Agencourt CleanSEQ (Beckman Coulter) and 3730 DNA Analyzer (Applied Biosystems), respectively. The sequences were manually curated using Sequencher v5.2.3 (Gene Code Corporation, Ann Arbor, MI, USA), run in VirulenceFinder 1.5 (Joensen et al., 2014), and blasted against the NCBI database¹. The nucleotide sequences were deposited in the GenBank nucleotide sequence database under the following accession numbers: strain 306, KU682619; strain 308, KU682620; strain 326, KU682621; strain 341, KU682622; strain 360, KU682623, and strain 500, KU682624.

High-throughput Real-time PCR Assay and Testing for Hemolysis

DNA was extracted from the swine isolates using the PrepMan Ultra Sample Preparation Reagent (Thermo Fisher) according to the manufacturer's instructions. The high-throughput real-time PCR (hrPCR) assay was carried out using the BioMark real-time PCR system (Fluidigm, San Francisco, CA, USA), targeting 67 virulence-associated and putative virulence-associated genes, 14 O-group-associated genes (O26, O45, O55, O91, O103, O104, O111, O113, O118, O121, O128, O145, O146, and O157) and 11

H-group-associated genes (H2, H4, H7, H8, H11, H16, H19, H21, H25, H28, and H32). Primers were designed in several studies (Perelle et al., 2003; Fratamico et al., 2008; Bugarel et al., 2010, 2011; Delannoy et al., 2013) and summarized by Tseng et al. (2014a). Reagents for DNA amplification and thermal cycling conditions were previously reported (Tseng et al., 2014a). Swine STEC strains positive to *ehxA* and *hlyA* genes were tested for hemolysis by plating onto SHIBAM agar (Hardy Diagnostic, Santa Maria, CA, USA).

FDA Minimal Signature *E. coli* Array

Swine Stx2d-producing *E. coli* and non-Stx2e STEC belonging to a serotype associated with human disease were further analyzed using the Minimal Signature *E. coli* Array Strip (FDA-ECID; Affymetrix, Santa Clara, CA, USA). Genomic DNA was isolated and concentrated using the DNeasy Tissue Kit (QIAGEN Inc., Valencia, CA, USA) and SC100 Speedvac Concentrator (Savant Instruments, Inc. Holbrook, NY, USA), respectively. Two micrograms of DNA were tested using the FDA-ECID array as described in detail by Lacher et al. (2014). Robust multi-array average summarized probe intensity data were analyzed using R-Bioconductor software v3.1.2 and affy package with parameters defined by Lacher et al. (2014). The Hierarchical clustering was done using overview function in MADE4 package that uses average linkage cluster analysis with a correlation metric distance (Culhane et al., 2005; Culhane and Thioulouse, 2006).

RESULTS


Swine STEC Serotypes

All of the strains had been previously serotyped at the *E. coli* Reference Center at the Pennsylvania State University (University Park, PA, USA). In addition, many O-group- and H-group-specific targets were included in the hrPCR assay. Several discrepancies were found and serotypes that did not match with the traditional serotyping are indicated in bold in **Figure 1**. Selected swine STEC strains were also analyzed using the FDA-ECID microarray, and the resulting serotypes were in agreement with the hrPCR. Moreover, the grouping within the phylogenetic tree was consistent with the serotypes proposed by the FDA-ECID microarray (**Table 1**).

Shiga-toxin Subtype Characterization

The swine STEC strains were analyzed by singleplex and multiplex PCR assays to determine their Shiga-toxin subtype. *Stx*-encoding genes were carried by 177/181 (99.8%) of the tested isolates. Four strains previously identified as STEC likely lost the *Stx* genes due to loss of *Stx*-encoding phages, as has been shown by other investigators (Joris et al., 2011) since PCR results were negative for any of the subtypes. *stx1* or *stx2* genes were carried by 25 and 151 strains, respectively. *Stx* subtype analysis revealed that the 25 *stx1*-positive strains carried the *stx1a* subtype. Among the 151 *stx2*-positive strains, 146 and 5 isolates carried *stx2e* and *stx2d*, respectively. STEC strain 308 was the only isolate that carried both *stx1* and *stx2*, subtypes *stx1c*

¹<http://blast.ncbi.nlm.nih.gov/Blast.cgi>



and *stx2d*, respectively. Stx subtypes divided by STEC serotype are reported in **Figure 1**. Strains carrying Stx subtypes *stx1d*, *stx2a*, *stx2b*, *stx2c*, *stx2f*, and *stx2g* were not identified. Selected swine strains were analyzed using the FDA-ECID microarray and results of Stx subtypes are reported in **Table 1**. Nucleotide sequencing of *stx2* was carried out from STEC strains that were *stx2d* positive by PCR. The STEC strain 308 *stx2* sequence showed 100% identity to a portion of *stx2d* subunit B (AF479829) using VirulenceFinder. When blasted against the NCBI database, the STEC strain 308 *stx2* partial sequence matched EF441621 (*stx2d*) with 100% identity and no gaps. The *stx2* partial sequences of STEC strains 306, 326, 341, 360, and 500 were identical. VirulenceFinder results showed 100% identity with a portion of *stx2d*, subunit B (DQ059012). The most similar sequence present in the NCBI database was KC339670 (*stx2e*) with an identity on 99%.

Distribution of Virulence-associated Genes among the Swine STEC Collection

Genomic DNA extracted from the swine STEC strains was analyzed using hrPCR. Genes encoding the enteroaggregative *E. coli* heat-stable enterotoxin 1 (*astA*) and the heat-stable enterotoxin (*estIa*) were detected in 79 (44%) and 91 (50%) of the isolates, respectively. Toxins and cytotoxic factors encoded by *cdtI*, *cdtIII*, *elt*, *ent/espL2*, *cnf2*, and *subAB* were not detected. Regarding cytolysins, enterohemolysin (*ehxA*) and α -hemolysin (*hlyA*) encoding genes were found non-simultaneously in 13 (7%) of the swine STEC strains each. These strains were also hemolytic when plated onto SHIBAM plates.

None of the isolated swine STEC strains carried the intimin-encoding gene, *eae*, effector genes involved in the type III secretion system (*espK*, *espM1*, *espM2*, *espN*, *espO1-1*, *espV*, *espX7*, *nleA*, *nleB*, *nleE*, *nleF*, *nleG5*, *nleG6-2*, and *nleH1-2*) or the type II secretion system effector (*etpD*). Other genes encoding factors involved in adhesion and colonization to the host intestine were also investigated. Among these, the most prevalent were *lpfA*_{O113}, *paa*, *ihA*, and *lpfA*_{O26} present in 116 (64%), 98 (54%), 41 (23%), and 33 (18%) isolates, respectively. Genes *orfA*, *orfB*, and *fedA* were detected in less than 8% of the isolates. While *bfp*, *efa1*, *fasA*, *fimF41a*, *saa*, and *toxB* were not found, one swine STEC strain carried *lpfA*_{O157}. Three autotransporter protein genes *ehaA*, *espP*, and *sab* were found in 60 (33%) 13 (7%), and 15 (8%) isolates.

Other gene targets were also included in the high-throughput real-time PCR assay. Positive results were obtained for the *ecs1763*, *terE*, *katP*, and *ureD* genes in 56 (31%), 31 (17%), 27 (15%), and 23 (13%) of isolates, respectively. While less than 8% were positive for *pagC*, *eibG*, *irp2*, *fyuA*, *ecf1*, *ecf2*, *ecf3*, *ecf4*, and Z2099. None of the swine STEC strains carried *ecs1822*, *epeA*, *sfp*, *stcE*, Z2096, or Z2098.

DISCUSSION

It is well-known that swine shed a variety of STEC serogroups, which may be carried along the food production chain. Most of the STEC isolated from these animals have adapted to the swine host and seem to have low potential to infect humans. Nevertheless, outbreaks associated with pork products have occurred (Meng et al., 2014; Tseng et al., 2014b). The sampling area covered by the NAHMS swine 2000 study was large, covering all the main pork-producing States (Fratamico et al., 2004). A subset of 181 STEC strains were analyzed and their pathogenic potential was assessed by detection of virulence and putative virulence factors.

The *stx* subtypes carried by the swine STEC were identified, and the majority of the isolates carried *stx2e* (81%), which was consistent with the data reported by Fratamico et al. (2004). The second most prevalent subtype was *stx1a* (14%), followed by *stx2d* (3%), and *stx1c* (1%). Stx2d is a potent toxin, and infection with strains carrying this subtype can lead to severe symptoms such as HC and HUS in humans (Melton-Celsa, 2014). Besides the Stx genes, the thermostable enterotoxin genes, *astA* and/or *estIa*, genes were found in ~71%

TABLE 1 | Serotypes and phylogenetic tree of select swine STEC strains analyzed using the FDA-ECID array.

Height	FDA-ECID serotype	Serotype (traditional serotyping)	Strain	Isolation (State, Date)	stx ₁ subtype	stx ₂ subtype
0.10 0.00						
	O159:H16	O159:H-	360	NC, 11/28/2000	neg	2e
	O159:H16	O159:H-	306	IO, 10/2/2000	neg	2i
	O159:H16	O159:H4	500	IO, 10/2/2000	neg	2i
	O159:H-	O159:H4	326	KS, 2/6/2001	neg	2i
	O159:H16	OX10:H-	341	MN, 9/19/2000	neg	2i
	O-:H38	O91:H44	448	NE, 10/16/2000	neg	2e
	O-:H38	O91:H44	477	NE, 10/16/2000	neg	2e
	O15:H27	O15:H-	308	IO, 10/2/2000	1c	2acd
	O152:H19	O20:H19	339	MI, 9/20/2000	1a	neg
	O-:H19 ^a	O163:H-	318	IO, 10/17/2000	1a	neg
	O-:H19 ^a	O163:H-	313	IO, 2/5/2001	1a	neg
	O-:H19 ^a	O163:H41 and 51	316	IO, 2/5/2001	1a	neg
	O-:H19 ^a	O163:H-	303	IO, 2/5/2001	1a	neg
	O-:H19 ^a	O163:H-	332	IO, 3/20/2001	1a	neg
	O91:H14	O91:H-	300	IO, 9/18/2000	1a	neg
	O91:H14	O91:H-	337	IO, 10/30/2000	1a	neg
	O91:H14	O91:H-	336	IO, 10/30/2000	1a	neg
	O91:H14	O91:H12	311	SD, 10/2/2000	1a	neg
	O91:H14	O91:H12	312	SD, 10/2/2000	1a	neg
	O91:H14	O91:H-	309	SD, 10/2/2000	1a	neg
	O91:H14	O91:H-	424	IO, 10/30/2000	1a	neg
	O91:H14	O91:H-	307	SD, 10/2/2000	1a	neg
	O91:H14	A:H-	319	IO, 10/16/2000	1a	neg
	O91:H14	O91:H-	315	IO, 10/16/2000	1a	neg
	O91:H14	O91:H-	317	NE, 10/16/2000	1a	neg
	O91:H14 ^b	O91:H14	2.4114	NA	1a	neg
	O91:H14	O91:H14	347	IO, 10/16/2000	1a	neg
	O91:H14 ^b	O91:H14	2.4111	NA	1a	neg
	O91:H14	O91:H14	346	WI, 10/16/2000	1a	neg
	O159:H21	O159:H34	387	IO, 10/2/2000	neg	neg
	O91:H21 ^b	O91:H21	B2F1	NA	neg	2acd

A, autoagglutination; O-, O non-typeable; H-, H non-typeable; A, neg, negative results; NA, not available.

^aThe FDA-ECID does not include probes for O163 markers.

^bSwine STEC isolates were analyzed together with two strains of *E. coli* O91:H14 and *E. coli* O91:H21 for comparison.

of the isolates. Thermostable enterotoxins are usually carried by enterotoxigenic *E. coli*, which are the major pathogens responsible for traveler's diarrhea. Twenty-two percent of the swine STEC strains were positive for both genes. The exotoxins HlyA (α -hemolysin) and EhxA (enterohemolysin) produce pores in the cytoplasmic membranes of eukaryotic cells causing their death. Their role in STEC pathogenesis is still not clear; HlyA may increase the virulence of extraintestinal pathogenic *E. coli* and, in the case of EhxA, a correlation between *ehxA*-positive STEC and development of severe symptoms in humans has been observed (Karch and Bielaszewska, 2001; Mainil, 2013). Thirteen isolates carried the *hlyA* gene. Nine of them belonged to serotypes O121:H- or O121:H10, presenting a virulence gene profile typical of strains associated with edema disease in swine due to the presence of *stx*_{2e}, *hlyA* and *fedA* (Tseng et al., 2014b). The *ehxA* gene is commonly found in STEC.

From 40 to 77% of strains collected from patients, food, and cattle carry this gene (Karch and Bielaszewska, 2001; Slanec et al., 2009; Bosilevac and Koohmaraie, 2011; Feng, 2014). Swine isolates appear to carry *ehxA* less frequently (Meng et al., 2014; Tseng et al., 2014a), and this observation is in agreement with our study where only 7% of the isolates was *ehxA* positive.

All of the swine STEC strains were LEE-negative. Although the adhesion mechanisms of LEE-negative STEC are not well characterized, several factors have been described to play an important role in adhesion to the intestinal epithelium. The long polar fimbriae gene *lpfA*_{O113} was identified in STEC O113:H21 (Doughty et al., 2002). These investigators demonstrated that the removal of *lpfA*_{O113} reduces the bacterial capacity to adhere to epithelial cells. Similar *lpfA* genes were found in *E. coli* O157 and O26 (Hayashi et al., 2001; Toma et al.,

2004). Another bacterial adherence-conferring gene is the iron-regulated gene A homolog adhesin *iha*. Similarly to *lpfA*_{O113}, the *iha* gene is commonly found in STEC strains associated with human cases of HUS (Newton et al., 2009; Galli et al., 2010). Nevertheless, non-pathogenic *E. coli* can also carry *lpfA*_{O113} and *iha*, suggesting that the presence of these genes is insufficient to establish an infection (Toma et al., 2004). Over 80% of the strains analyzed in this study carried *lpfA*_{O26}, *lpfA*_{O113}, or *lpfA*_{O157}; while *iha* was found in almost one quarter of swine isolates. *iha*-positive STEC were also described in a longitudinal study of two Midwestern U.S. pork production sites (Tseng et al., 2014a, 2015). On the contrary, none of the swine STEC strains collected in another interesting study in China carried *iha* (Meng et al., 2014). The second most prevalent adhesion factor found in this dataset was the porcine attaching and effacing-associated adhesin, *paa*, which is associated with neonatal post-weaning diarrhea in pigs (An et al., 1999). In addition, a few strains carried *orfA* and *orfB*, which encode for adhesins involved in diffuse adherence (Charbonneau et al., 2006).

Autotransporter proteins have a peculiar structure that allows them to move through the membrane system and execute their function outside the bacterial cell. The genes *ehaA* and *sab* were discovered in O157:H7 strain EDL933 and LEE-negative O113:H21, respectively. They encode for two different autotransporter proteins that contribute to adhesion and biofilm formation (Wells et al., 2008; Herold et al., 2009). Together with LEE genes, *iha* and *ehaA* are highly expressed in the intestines of pigs presenting attaching and effacing lesions (Liu et al., 2015). While the *ehaA* gene was present in over 30% of the swine isolates, *sab* was carried by 13 STEC strains only belonging to O-group O91.

As stated above, 12 to 18% of the isolates were positive for *katP*, *ureD*, and *terE*. The genes *katP* and *ureD* encode for a catalase/peroxidase and urease transporter, respectively. Their role in *E. coli* pathogenesis is unclear; however, they appear to be prevalent in diarrheagenic *E. coli* (Dorothea et al., 2006; Delannoy et al., 2013). The gene *terE* is a component of the *ter* cluster, which confers tellurite resistance (Orth et al., 2007). The *ecs1763* and *ecs1822* genes have been proposed to be novel markers for enterohemorrhagic *E. coli*. Their function is unknown, and they were shown to be shared by a clonal group of enterohemorrhagic *E. coli* that includes O26, O111, and O118 (Abu-Ali et al., 2009). Tseng et al. (2014a) observed that *ecs1763* is frequently found in swine STEC, which was confirmed by the present study where 31% of the isolates carried *ecs1763*. *ecs1822* was absent in all the tested strains.

Traditional serotyping of *E. coli* is time consuming, and cross-reactions among antisera often occur. Based on the hrPCR and FDA-ECID results, 71 strains present in this collection belonged to serotypes O8:H9, O8:H19, O8:H-, O15:H27, O20:H-, O91:H14, O101:H-, O121:H-, O145:H25, O159:H21, and O163:H19 that were previously isolated from human patients (Blanco et al., 1992; Beutin and Fach, 2014). Serotypes O8:H19, O15:H27, O145:H25, and O163:H19 have also been associated with cases of HUS (Prager et al., 2005; Bielaszewska et al., 2006; Galli et al., 2010). All of the strains belonging to the

serotypes O8:H9, O8:H19, O8:H-, O20:H-, O101:H-, O121:H-, O145:H25, and O159:H21 analyzed in this study carried *stx*_{2e}, which is a subtype that is generally not associated with STEC that cause serious human illness. Human infections linked to Stx2e-producing *E. coli* generally cause asymptomatic infections or mild diarrhea (Tseng et al., 2014b). The work of Sonntag et al. (2005) reported that human Stx2e-producing *E. coli* carry different virulence factors compared to swine Stx2e-producing *E. coli* associated with edema disease. They also detected *fyuA* and *irp2* genes in five strains isolated from humans. These genes are included in the high-pathogenicity island (HPI), which is involved in the iron metabolism of *Yersinia*. Mouse models showed that the HPI increases *E. coli* virulence in extraintestinal infections (Schubert et al., 2002). The hrPCR results revealed that some swine STEC strains belonged to the same serotypes as human Stx2e-producing *E. coli* (O8:H19 and O8:H-) reported by Sonntag et al. (2005). STEC O8:H19 and STEC O8:H- also carried markers for the HPI. Moreover, their virulence gene profiles included adhesins (*lpfA*_{O26}, *lpfA*_{O113}, *paa*) and enterotoxins (*astA* and *estIa*), which suggest that they can potentially provoke mild diarrhea in humans. The HPI genes *fyuA* and *irp2* were also found in Stx2e-producing *E. coli* belonging to serotypes O5:H4 and O8:H4 (Table 1).

Shiga toxin-producing *Escherichia coli* strain 308 was re-typed as O15:H27 using the FDA-ECID array and was found to have the same *stx*_{2d} sequence as *E. coli* O15:H27 (strain 88-1509) in the STEC isolate database at Michigan State University². *E. coli* strain 88-1509 was collected in 1988 from a human case of HC and HUS in Canada. Other strains belonging to serotype O15:H27 have been isolated from human and cattle feces, and from meat sources (Piérard et al., 1997; Woodward et al., 2002; Bosilevac et al., 2007; Galli et al., 2010). The LEE-negative swine STEC O15:H27 has a virulence gene profile consisting of *stx*_{1c}, *stx*_{2d}, *ehaA*, *espP*, *fyuA*, *ihA*, *irp2*, *lpfA*_{O113}, and Z2099. The relevance of some of these genes was mentioned above. *E. coli* secreted protein P (EspP) is an autotransporter protein with serine protease activity, and is used by the bacteria to impair the complement response of the host (Orth et al., 2010). Recently, In et al. (2013) reported that EspP boosts macropinocytosis in the intestinal epithelium increasing Stx uptake. The open reading frame Z2099 is highly prevalent in typical and emerging enterohemorrhagic *E. coli*, while it is significantly less prevalent in non-pathogenic *E. coli* (Delannoy et al., 2013).

Six of the swine STEC strains carried *stx*_{2d} according to PCR and VirulenceFinder results, and they belonged to serotypes O159:H-, O159:H4, and OX10:H-. DebRoy et al. (2016) reported that serological cross-reactions between the O159 and OX10 O-groups often occur and that the nucleotide sequences of O159 and OX10 O-antigen gene clusters are almost identical. Based on the FDA-ECID analysis, the strains 306, 360, 341, and 500 were re-typed as O159:H16; while the strain 326 was re-typed as O159:H- (Table 1). STEC belonging to O-group O159 rarely infect humans (Brooks et al., 2005; Gould et al., 2013). STEC O159:H16 and O159:H- have been isolated only

²<http://shigatox.net/new/database.html>

from swine samples, such as feces and carcasses (DesRosiers et al., 2001; Kaufmann et al., 2006; Meng et al., 2014). Stx subtype analysis of these strains often gives ambiguous results (Kaufmann et al., 2006; Meng et al., 2014). In this work, STEC O159:H16 and O159:H- were positive for *stx_{2d}* when tested by PCR; however, they were positive for *stx_{2e}* or *stx_{2i}* using the FDA-ECID array. Note that probes of the FDA-ECID array corresponding to *stx_{2i}* were designed using the Stx sequences AM904726 and FN252457 (Patel et al., 2016) that belong to the *stx_{2e}* subtype according to Scheutz et al. (2012). The product obtained from partial sequencing of *stx₂* was 99% identical to the sequence KC339670 when blasted against the NCBI database. KC339670 is a complete *stx₂* sequence belonging to a STEC O159:H16 strain isolated from swine in China. After a neighbor-joining cluster analysis of the sequence, Meng et al. (2014) concluded that KC339670 represented a new variant of *stx_{2e}*. Further investigations using cell lines and animal models are needed to understand the virulence potential of this Stx2 variant. Another STEC O159 was detected in this collection. It belonged to the H21 H-group and was positioned distantly from the clade of O159:H16 and O159:H- (Table 1). This strain was positive for *stx_{2e}* only by PCR. *E. coli* belonging to serotype O159:H21 was isolated in 1983 during a small outbreak of diarrhea involving newborn children in Spain (Blanco et al., 1992), and no other infections associated with serotype O159:H21 have been reported.

Locus of enterocyte effacement-negative STEC belonging to O-group O91 are frequently associated with adult human infections with symptoms ranging from mild diarrhea to HC and HUS. The main serotypes are O91:H14 and O91:H21, and the latter is usually linked with development of severe symptoms (Bielaszewska et al., 2009). Human STEC O91:H14 and O91:H21 isolates carried mainly *stx₁* and *stx_{2d}*, respectively (Prager et al., 2005; Bielaszewska et al., 2009; Galli et al., 2010). These STEC have been isolated from food samples derived from bovine, swine, and ovine origin, and from both domestic and wild animals (Martin and Beutin, 2011; Ju et al., 2012). From the NAHMS swine 2000 study, 15 strains belonging to serotypes O91:H12, O91:H14, O91:H44, and O91:H- were isolated from fresh fecal samples collected from four different states (Fratamico et al., 2004). Eight of these strains were re-typed as O91:H14, while STEC O91:H44 strains 448 and 477 did not belong to the O91 O-group based on FDA-ECID and hrPCR results (Figure 1; Table 1). STEC strain 319 that carried an identical virulence gene profile to other O91:H14 strains was also re-typed as O91:H14 by FDA-ECID array (Table 1). According to the phylogenetic tree in Table 1, the clade of STEC O91:H14 strains is well separated from the STEC O91:H21 strain B2F1 isolated from a human case of HUS. Interestingly, two O91:H14 strains were more closely related to two STEC O91:H14 strains isolated from ground beef samples than the other swine STEC O91:H14 strains. Despite the fact that one strain was *katP*-negative, all 13 STEC O91:H14 strains presented a conserved virulence gene profile (*ehaA*, *ehxA*, *eibG*, *espP*, *ihaA*, *katP*, *lpfA_{O26}*, *lpfA_{O113}*, *pagC*, *sab*, and *stx_{1a}*), which is very similar to profiles of strains from human clinical samples (Prager et al., 2005; Bielaszewska et al., 2009). Similar to *ihaA*, *lpfA_{O26}*, and *lpfA_{O113}*, the proteins encoded by the genes

eibG and *sab* are involved in host gut colonization. The *E. coli* immunoglobulin-binding protein encoded by *eibG* binds human immunoglobulin G and immunoglobulin A, and contributes to epithelial host cell adhesion (Lu et al., 2006), and *sab* is a gene encoding for an autotransporter protein involved in biofilm formation and found in a pathogenic LEE-negative STEC (Herold et al., 2009). Lastly, the *pagC* gene encodes for an outer membrane protein present in different Enterobacteriaceae that contributes to serum resistance (Nishio et al., 2005).

Human infections caused by STEC O163:H19 are rare (Brooks et al., 2005; Gould et al., 2013). However, Stx2-producing *E. coli* O163:H19 provoked sporadic cases of HUS (Prager et al., 2005) and have been found associated with cattle and produce (Woodward et al., 2002; Galli et al., 2010; Feng, 2014). In this work, five strains of STEC O163:H- or O163:H41/H51 were re-typed as O163:H19. They all carried *stx_{1a}* similar to the Stx1-producing *E. coli* O163:H19 strain isolated from swine by DesRosiers et al. (2001).

STEC O20:H19 is associated with human cases of HUS (Galli et al., 2010), and one strain belonging to this serotype was isolated in the NAHMS study (Fratamico et al., 2004). However, this same strain was re-analyzed using the FDA-ECID array, and it was re-typed as O152:H19, which is not known to be a human pathogen.

CONCLUSION

Using state-of-the-art DNA-based techniques, this study provides new insights on the distribution of virulence factors in a heterogeneous collection of STEC isolated from the major pork-producing states of the U.S. Stx2e-producing *E. coli* known to provoke mild diarrhea in humans carried different virulence factors than Stx2e-producing *E. coli* associated with edema disease in pigs; this finding suggests that Stx2e-producing *E. coli* that cause human illnesses may not have a swine origin (Sonntag et al., 2005). In our work, STEC strains carrying *stx_{2e}* belonging to the same serotype and having similar virulence gene profiles as Stx2e-producing *E. coli* isolated from humans were identified. Additionally, the majority of Stx2e-producing *E. coli* carried thermostable enterotoxin genes usually found in enterotoxigenic *E. coli*.

This work suggests that STEC, including serotypes O15:H27 and O91:H14 that have been associated with human illness and are found in multiple hosts or environments, could also be carried by swine. Interestingly, a strain of O15:H27 found to carry *stx_{2d}* and other virulence genes may have the potential to produce severe symptoms in humans. Moreover, STEC O91:H14 strains presented a virulence gene profile very similar to profiles found in human isolates.

AUTHOR CONTRIBUTIONS

GMB and PF. designed research; GMB, LKB, SD, PF, FB, AA, and TP performed research; GMB, JG, and IP analyzed data; GMB and PF wrote the paper.

ACKNOWLEDGMENTS

This research was supported in part by an appointment to the Agricultural Research services (ARS) Research Participation Program administered by the Oak Ridge Institute for Science and Education (ORISE) through an interagency agreement between the U.S. Department of Energy (DOE) and the

USDA. ORISE is managed by ORAU under DOE contract number DE-AC05-06OR23100. The hrPCR development was partially financed by the French joint ministerial program of R&D against CBRNE risks. All opinions expressed in this manuscript are the authors' and do not necessarily reflect the policies and views of USDA, ARS, DOE, or ORAU/ORISE.

REFERENCES

- Abu-Ali, G. S., Lacher, D. W., Wick, L. M., Qi, W., and Whittam, T. S. (2009). Genomic diversity of pathogenic *Escherichia coli* of the EHEC 2 clonal complex. *BMC Genom.* 10:296. doi: 10.1186/1471-2164-10-296
- An, H., Fairbrother, J. M., Desautels, C., and Harel, J. (1999). "Distribution of a novel locus called Paa (porcine attaching and effacing associated) among Enteric *Escherichia coli*," in *Mechanisms in the Pathogenesis of Enteric Diseases 2 Advances in Experimental Medicine and Biology*, eds P. S. Paul and D. H. Francis (Boston, MA: Springer), 179–184. doi: 10.1007/978-1-4615-4143-1
- Beutin, L., and Fach, P. (2014). Detection of Shiga toxin-producing *Escherichia coli* from nonhuman sources and strain typing. *Microbiol. Spectr.* 2:EHEC-0001-2013. doi: 10.1128/microbiolspec.EHEC-0001-2013
- Beutin, L., Krause, G., Zimmermann, S., Kaulfuss, S., and Gleier, K. (2004). Characterization of Shiga toxin-producing *Escherichia coli* strains isolated from human patients in Germany over a 3-year period. *J. Clin. Microbiol.* 42, 1099–1108. doi: 10.1128/JCM.42.3.1099
- Bielaszewska, M., Friedrich, A. W., Aldick, T., Schürk-Bulgrin, R., and Karch, H. (2006). Shiga toxin activatable by intestinal mucus in *Escherichia coli* isolated from humans: predictor for a severe clinical outcome. *Clin. Infect. Dis.* 43, 1160–1167. doi: 10.1086/508195
- Bielaszewska, M., Stoewe, F., Fruth, A., Zhang, W., Prager, R., Brockmeyer, J., et al. (2009). Shiga toxin, cytolethal distending toxin, and hemolysin repertoires in clinical *Escherichia coli* O91 isolates. *J. Clin. Microbiol.* 47, 2061–2066. doi: 10.1128/JCM.00201-09
- Blanco, J., González, E. A., Espinosa, P., Blanco, M., Garabal, J. I., and Alonso, M. P. (1992). Enterotoxigenic and necrotizing *Escherichia coli* in human diarrhoea in Spain. *Eur. J. Epidemiol.* 8, 548–552. doi: 10.1007/BF00146375
- Boisen, N., Melton-Celsa, A. R., Scheutz, F., O'Brien, A. D., and Nataro, J. P. (2015). Shiga toxin 2a and enteroaggregative *Escherichia coli* – A deadly combination. *Gut Microbes* 6, 272–278. doi: 10.1080/19490976.2015.1054591
- Bosilevac, J. M., Guerini, M. N., Brichta-Harhay, D. M., Arthur, T. M., and Koohmaraie, M. (2007). Microbiological characterization of imported and domestic boneless beef trim used for ground beef. *J. Food Prot.* 70, 440–449.
- Bosilevac, J. M., and Koohmaraie, M. (2011). Prevalence and characterization of non-O157 shiga toxin-producing *Escherichia coli* isolates from commercial ground beef in the United States. *Appl. Environ. Microbiol.* 77, 2103–2112. doi: 10.1128/AEM.02833-10
- Brandt, S. M., King, N., Cornelius, A. J., Premaratne, A., Besser, T. E., and On, S. L. W. (2011). Molecular risk assessment and epidemiological typing of Shiga toxin-producing *Escherichia coli* by using a novel PCR binary typing system. *Appl. Environ. Microbiol.* 77, 2458–2470. doi: 10.1128/AEM.02322-10
- Brooks, J. T., Sowers, E. G., Wells, J. G., Greene, K. D., Griffin, P. M., Hoekstra, R. M., et al. (2005). Non-O157 Shiga toxin – producing *Escherichia coli* infections in the United States, 1983 – 2002. *J. Infect. Dis.* 192, 1422–1424. doi: 10.1086/466536
- Bugarel, M., Beutin, L., and Fach, P. (2010). Low-density microarray targeting non-locus of enterocyte effacement effectors (nle genes) and major virulence factors of Shiga toxin-producing *Escherichia coli* (STEC): a new approach for molecular risk assessment of STEC isolates. *Appl. Environ. Microbiol.* 76, 203–211. doi: 10.1128/AEM.01921-09
- Bugarel, M., Martin, A., Fach, P., and Beutin, L. (2011). Virulence gene profiling of enterohemorrhagic (EHEC) and enteropathogenic (EPEC) *Escherichia coli* strains: a basis for molecular risk assessment of typical and atypical EPEC strains. *BMC Microbiol.* 11:142. doi: 10.1186/1471-2180-11-142
- Charbonneau, M. È., Berthiaume, F., and Mourez, M. (2006). Proteolytic processing is not essential for multiple functions of the *Escherichia coli* autotransporter adhesin involved in diffuse adherence (AIDA-I). *J. Bacteriol.* 188, 8504–8512. doi: 10.1128/JB.00864-06
- Coombes, B. K., Wickham, M. E., Mascarenhas, M., Gruenheid, S., Finlay, B. B., and Karmali, M. A. (2008). Molecular analysis as an aid to assess the public health risk of non-O157 Shiga toxin-producing *Escherichia coli* strains. *Appl. Environ. Microbiol.* 74, 2153–2160. doi: 10.1128/AEM.02566-07
- Croxen, M. A., Law, R. J., Scholz, R., Keeney, K. M., Wlodarska, M., and Finlay, B. B. (2013). Recent advances in understanding enteric pathogenic *Escherichia coli*. *Clin. Microbiol. Rev.* 26, 822–880. doi: 10.1128/CMR.00022-13
- Culhane, A. C., and Thioulouse, J. (2006). A multivariate approach to integrating datasets using made4 and ade4. *Newsl. R Proj.* 6, 54–58.
- Culhane, A. C., Thioulouse, J., Perrière, G., and Higgins, D. G. (2005). MADE4: An R package for multivariate analysis of gene expression data. *Bioinformatics* 21, 2789–2790. doi: 10.1093/bioinformatics/bti394
- DeRoy, C., Fratamico, P. M., Yan, X., Baranzoni, G., Liu, Y., Needleman, D. S., et al. (2016). Comparison of O-antigen gene clusters of all O-serogroups of *Escherichia coli* and proposal for adopting a new nomenclature for O-typing. *PLoS ONE* 11:e0147434. doi: 10.1371/journal.pone.0147434
- Delannoy, S., Beutin, L., and Fach, P. (2013). Discrimination of enterohemorrhagic *Escherichia coli* (EHEC) from non-EHEC strains based on detection of various combinations of type III effector genes. *J. Clin. Microbiol.* 51, 3257–3262. doi: 10.1128/JCM.01471-13
- DesRosiers, A., Fairbrother, J. M., Johnson, R. P., Desautels, C., Letellier, A., and Quessy, S. (2001). Phenotypic and genotypic characterization of *Escherichia coli* verotoxin-producing isolates from humans and pigs. *J. Food Prot.* 64, 1904–1911. doi: 10.1128/JCM.01095-08
- Dorothea, O., Grif, K., Dierich, M. P., and Wu, R. (2006). Prevalence, structure and expression of urease genes in Shiga toxin-producing *Escherichia coli* from humans and the environment. *Int. J. Hyg. Environ. Health* 209, 513–520. doi: 10.1016/j.ijheh.2006.06.003
- Doughty, S., Sloan, J., Bennett-Wood, V., Robertson, M., Robins-Browne, R. M., and Hartland, E. L. (2002). Identification of a novel fimbrial gene cluster related to long polar fimbriae in locus of enterocyte effacement-negative strains of enterohemorrhagic *Escherichia coli*. *Infect. Immun.* 70, 6761–6769. doi: 10.1128/IAI.70.12.6761-6769.2002
- Ethelberg, S., Olsen, K. E. P., Scheutz, F., Jensen, C., Schiellerup, P., Engberg, J., et al. (2004). Virulence factors for hemolytic uremic syndrome, Denmark. *Emerg. Infect. Dis.* 10, 842–847. doi: 10.3201/eid1005.030576
- Fasel, D., Mellmann, A., Cernela, N., Hächler, H., Fruth, A., Khanna, N., et al. (2014). Hemolytic uremic syndrome in a 65 year-old male linked to a very unusual type of stx2e and eae harboring O51:H49 Shiga-toxin producing *Escherichia coli*. *J. Clin. Microbiol.* 52, 1301–1303. doi: 10.1128/JCM.03459-13
- Feng, P. (2014). Shiga Toxin-Producing *Escherichia coli* (STEC) in fresh produce — A food safety dilemma. *Microbiol. Spectr.* 2:EHEC-0010-2013. doi: 10.1128/microbiolspec.EHEC-0010
- Fratamico, P. M., Bagi, L. K., Bush, E. J., and Solow, B. T. (2004). Prevalence and characterization of Shiga toxin-producing *Escherichia coli* in swine feces recovered in the National Animal Health Monitoring system's swine 2000 study. *Appl. Environ. Microbiol.* 70, 7173–7178. doi: 10.1128/AEM.70.12.7173-7178.2004
- Fratamico, P. M., Bhagwat, A. A., Injaian, L., and Fedorka-Cray, P. J. (2008). Characterization of Shiga toxin-producing *Escherichia coli* strains isolated from swine feces. *Foodborne Pathog. Dis.* 5, 827–838. doi: 10.1089/fpd.2008.0147
- Friedrich, A. W., Bielaszewska, M., Zhang, W., Pulz, M., Kuczius, T., Ammon, A., et al. (2002). *Escherichia coli* harboring Shiga toxin 2 gene variants: frequency and association with clinical symptoms. *J. Infect. Dis.* 185, 74–84. doi: 10.1086/338115

- Fuente, R. D. La Horcajo, P., and Dom, G. (2015). Association of vt1c with verotoxin-producing *Escherichia coli* from goats and sheep. *J. Vet. Diagn. Invest.* 334, 332–334.
- Fuller, C. A., Pellino, C. A., Flagler, M. J., Strasser, J. E., and Weiss, A. A. (2011). Shiga toxin subtypes display dramatic differences in potency. *Infect. Immun.* 79, 1329–1337. doi: 10.1128/IAI.01182-10
- Galli, L., Miliwebsky, E., Irino, K., Leotta, G., and Rivas, M. (2010). Virulence profile comparison between LEE-negative Shiga toxin-producing *Escherichia coli* (STEC) strains isolated from cattle and humans. *Vet. Microbiol.* 143, 307–313. doi: 10.1016/j.vetmic.2009.11.028
- Gould, L. H., Mody, R. K., Ong, K. L., Clogher, P., Cronquist, A. B., Garman, K. N., et al. (2013). Increased recognition of non-O157 Shiga toxin-producing *Escherichia coli* infections in the United States during 2000–2010: epidemiologic features and comparison with *E. coli* O157 infections. *Foodborne Pathog. Dis.* 10, 453–460. doi: 10.1089/fpd.2012.1401
- Hayashi, T., Makino, K., Ohnishi, M., Kurokawa, K., Ishii, K., Yokoyama, K., et al. (2001). Complete genome sequence of enterohemorrhagic *Escherichia coli* O157:H7 and genomic comparison with a laboratory strain K-12. *DNA Res.* 8, 11–22. doi: 10.1093/dnares/8.1.47
- Herold, S., Paton, J. C., and Paton, A. W. (2009). Sab, a novel autotransporter of locus of enterocyte effacement-negative Shiga-toxigenic *Escherichia coli* O113:H21, contributes to adherence and biofilm formation. *Infect. Immun.* 77, 3234–3243. doi: 10.1128/IAI.00031-09
- Hofer, E., Cernela, N., and Stephan, R. (2012). Shiga toxin subtypes associated with shiga toxin-producing *Escherichia coli* strains isolated from red deer, roe deer, chamois, and ibex. *Foodborne Pathog. Dis.* 9, 792–795. doi: 10.1089/fpd.2012.1156
- In, J., Lukyanenko, V., Foulke-Abel, J., Hubbard, A. L., Delannoy, M., Hansen, A.-M., et al. (2013). Serine protease EspP from enterohemorrhagic *Escherichia coli* is sufficient to induce Shiga toxin macropinocytosis in intestinal epithelium. *PLoS ONE* 8:e69196. doi: 10.1371/journal.pone.0069196
- Ito, H., Terai, A., Kurazono, H., Takeda, Y., and Nishibuchi, M. (1990). Cloning and nucleotide sequencing of Vero toxin 2 variant genes from *Escherichia coli* O91:H21 isolated from a patient with the hemolytic uremic syndrome. *Microb. Pathog.* 8, 47–60. doi: 10.1016/0882-4010(90)90007-D
- Joensen, K. G., Scheutz, F., Lund, O., Hasman, H., Kaas, R. S., Nielsen, E. M., et al. (2014). Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J. Clin. Microbiol.* 52, 1501–1510. doi: 10.1128/JCM.03617-13
- Joris, M. -A., Verstraete, K., De Reu, K., and De Zutter, L. (2011). Loss of vtx genes after the first subcultivation step of verocytotoxigenic *Escherichia coli* O157 and Non-O157 during isolation from naturally contaminated fecal samples. *Toxins (Basel)* 3, 672–677. doi: 10.3390/toxins3060672
- Ju, W., Shen, J., Li, Y., Toro, M. A., Zhao, S., Ayers, S., et al. (2012). Non-O157 Shiga toxin-producing *Escherichia coli* in retail ground beef and pork in the Washington D.C. area. *Food Microbiol.* 32, 371–377. doi: 10.1016/j.fm.2012.07.017
- Karch, H., and Bielaszewska, M. (2001). Sorbitol-fermenting Shiga toxin-producing *Escherichia coli* O157:H(-) strains: epidemiology, phenotypic and molecular characteristics, and microbiological diagnosis. *J. Clin. Microbiol.* 39, 2043–2049. doi: 10.1128/JCM.39.6.2043-2049.2001
- Karmali, M. A., Petric, M., Lim, C., Fleming, P. C., Arbus, G. S., and Lior, H. (1985). The association between idiopathic hemolytic uremic syndrome and infection by verotoxin-producing *Escherichia coli*. *J. Infect. Dis.* 151, 775–782. doi: 10.1093/infdis/151.5.775
- Kaufmann, M., Zweifel, C., Blanco, M., Blanco, J. E., Blanco, J. E., Beutin, L., et al. (2006). *Escherichia coli* O157 and non-O157 Shiga toxin-producing *Escherichia coli* in fecal samples of finished pigs at slaughter in Switzerland. *J. Food Prot.* 69, 260–266.
- Lacher, D. W., Gangiredla, J., Jackson, S. A., Elkins, C. A., and Feng, P. C. H. (2014). Novel microarray design for molecular serotyping of Shiga toxin-producing *Escherichia coli* isolated from fresh produce. *Appl. Environ. Microbiol.* 80, 4677–4682. doi: 10.1128/AEM.01049-14
- Liu, B., Yin, X., Yu, H., Feng, Y., Ying, X., Gong, J., et al. (2015). Alteration of the microbiota and virulence gene expression in *E. coli* O157:H7 in pig ligated intestine with and without AE lesions. *PLoS ONE* 10:e0130272. doi: 10.1371/journal.pone.0130272
- Lu, Y., Iyoda, S., Satou, H., Satou, H., Itoh, K., Saitoh, T., et al. (2006). A new immunoglobulin-binding protein, EibG, is responsible for the chain-like adhesion phenotype of locus of enterocyte effacement-negative, Shiga toxin-producing *Escherichia coli*. *Infect. Immun.* 74, 5747–5755. doi: 10.1128/IAI.00724-06
- Luna-Gierke, R. E., Griffin, P. M., Gould, L. H., Herman, K., Bopp, C. A., Strockbine, N., et al. (2014). Outbreaks of non-O157 Shiga toxin-producing *Escherichia coli* infection: USA. *Epidemiol. Infect.* 142, 2270–2280. doi: 10.1017/S0950268813003233
- Mainil, J. (2013). *Escherichia coli* virulence factors. *Vet. Immunol. Immunopathol.* 152, 2–12. doi: 10.1016/j.vetimm.2012.09.032
- Martin, A., and Beutin, L. (2011). Characteristics of Shiga toxin-producing *Escherichia coli* from meat and milk products of different origins and association with food producing animals as main contamination sources. *Int. J. Food Microbiol.* 146, 99–104. doi: 10.1016/j.ijfoodmicro.2011.01.041
- Melton-Celsa, A. R. (2014). Shiga toxin (Stx) classification, structure, and function. *Microbiol. Spectr.* 2, 1–13. doi: 10.1128/microbiolspec.EHEC-0024-2013
- Meng, Q., Bai, X., Zhao, A., Lan, R., Du, H., Wang, T., et al. (2014). Characterization of Shiga toxin-producing *Escherichia coli* isolated from healthy pigs in China. *BMC Microbiol.* 14:5. doi: 10.1186/1471-2180-14-5
- Muniesa, M., Recktenwald, J., Bielaszewska, M., Karch, H., and Schmidt, H. (2000). Characterization of a Shiga toxin 2e-converting bacteriophage from an *Escherichia coli* strain of human origin. *Infect. Immun.* 68, 4850–4855. doi: 10.1128/IAI.68.9.4850-4855.2000
- Newton, H. J., Sloan, J., Bulach, D. M., Seemann, T., Allison, C. C., Tauschek, M., et al. (2009). Shiga toxin-producing *Escherichia coli* strains negative for locus of enterocyte effacement. *Emerg. Infect. Dis.* 15, 372–380. doi: 10.3201/eid1502.080631
- Nishio, M., Okada, N., Miki, T., Haneda, T., and Danbara, H. (2005). Identification of the outer-membrane protein PagC required for the serum resistance phenotype in *Salmonella enterica* serovar Choleraesuis. *Microbiology* 151, 863–873. doi: 10.1099/mic.0.27654-0
- Orth, D., Ehrlenbach, S., Brockmeyer, J., Khan, A. B., Huber, G., Karch, H., et al. (2010). EspP, a serine protease of enterohemorrhagic *Escherichia coli*, impairs complement activation by cleaving complement factors C3/C3b and C5. *Infect. Immun.* 78, 4294–4301. doi: 10.1128/IAI.00488-10
- Orth, D., Grif, K., Dierich, M. P., and Würzner, R. (2007). Variability in tellurite resistance and the ter gene cluster among Shiga toxin-producing *Escherichia coli* isolated from humans, animals and food. *Res. Microbiol.* 158, 105–111. doi: 10.1016/j.resmic.2006.10.007
- Patel, I. R., Gangiredla, J., Lacher, D. W., Mammel, M. K., Jackson, S. A., Lampel, K. A., et al. (2016). FDA-*Escherichia coli* identification (FDA-ECID) microarray: a pan-genome molecular toolbox for serotyping, virulence profiling, molecular epidemiology, and phylogeny. *Appl. Environ. Microbiol.* (in press). doi: 10.1128/AEM.04077-15
- Paton, A. W., Woodrow, M. C., Doyle, R. M., Lanser, J. A., and Paton, J. C. (1999). Molecular characterization of a Shiga toxigenic *Escherichia coli* O113:H21 strain lacking eae responsible for a cluster of cases of hemolytic-uremic syndrome. *J. Clin. Microbiol.* 37, 3357–3361.
- Perelle, S., Dilasser, F., Grout, J., and Fach, P. (2003). Development of a 5'-nuclease PCR assay for detecting Shiga toxin-producing *Escherichia coli* O145 based on the identification of an "O-island 29" homologue. *J. Appl. Microbiol.* 94, 587–594. doi: 10.1046/j.1365-2672.2003.01872.x
- Piérard, D., Stevens, D., Moriau, L., Lior, H., and Lauwers, S. (1997). Isolation and virulence factors of verocytotoxin-producing *Escherichia coli* in human stool samples. *Clin. Microbiol. Infect.* 3, 531–540. doi: 10.1111/j.1469-0691.1997.tb00303.x
- Prager, R., Annemüller, S., and Tschäpe, H. (2005). Diversity of virulence patterns among Shiga toxin-producing *Escherichia coli* from human clinical cases - need for more detailed diagnostics. *Int. J. Med. Microbiol.* 295, 29–38. doi: 10.1016/j.ijmm.2004.12.009
- Scheutz, F., Teel, L. D., Beutin, L., Piérard, D., Buvens, G., Karch, H., et al. (2012). Multicenter evaluation of a sequence-based protocol for subtyping Shiga toxins and standardizing Stx nomenclature. *J. Clin. Microbiol.* 50, 2951–2963. doi: 10.1128/JCM.00860-12

- Schubert, S., Picard, B., Gouriou, S., Heesemann, J., and Denamur, E. (2002). Yersinia high-pathogenicity island contributes to virulence in *Escherichia coli* causing extraintestinal infections. *Infect. Immun.* 70, 5335–5337. doi: 10.1128/IAI.70.9.5335
- Slanec, T., Fruth, A., Creuzburg, K., and Schmidt, H. (2009). Molecular analysis of virulence profiles and Shiga toxin genes in food-borne Shiga toxin-producing *Escherichia coli*. *Appl. Environ. Microbiol.* 75, 6187–6197. doi: 10.1128/AEM.00874-09
- Sontag, A. K., Bielaszewska, M., Mellmann, A., Dierksen, N., Schierack, P., Wieler, L. H., et al. (2005). Shiga toxin 2e-producing *Escherichia coli* isolates from humans and pigs differ in their virulence profiles and interactions with intestinal epithelial cells. *Appl. Environ. Microbiol.* 71, 8855–8863. doi: 10.1128/AEM.71.12.8855-8863.2005
- Thomas, A., Cheasty, T., Chart, H., and Rowe, B. (1994). Isolation of Vero cytotoxin-producing *Escherichia coli* serotypes O9ab:H- and O101:H-carrying VT2 variant gene sequences from a patient with haemolytic uraemic syndrome. *Eur. J. Clin. Microbiol. Infect. Dis.* 13, 1074–1076. doi: 10.1007/BF02111832
- Toma, C., Martínez Espinosa, E., Song, T., Miliwebsky, E., Chinen, I., Iyoda, S., et al. (2004). Distribution of putative adhesins in different seropathotypes of Shiga toxin-producing *Escherichia coli*. *J. Clin. Microbiol.* 42, 4937–4946. doi: 10.1128/JCM.42.11.4937-4946.2004
- Tseng, M., Fratamico, P. M., Bagi, L., Delannoy, S., Fach, P., Manning, S. D., et al. (2014a). Diverse virulence gene content of Shiga toxin-producing *Escherichia coli* from finishing swine. *Appl. Environ. Microbiol.* 80, 6395–6402. doi: 10.1128/AEM.01761-14
- Tseng, M., Fratamico, P. M., Bagi, L., Manzinger, D., and Funk, J. A. (2015). Shiga toxin-producing *E. coli* (STEC) in swine: prevalence over the finishing period and characteristics of the STEC isolates. *Epidemiol. Infect.* 143, 505–514. doi: 10.1017/S0950268814001095
- Tseng, M., Fratamico, P. M., Manning, S. D., and Funk, J. A. (2014b). Shiga toxin-producing *Escherichia coli* in swine: the public health perspective. *Anim. Health Res. Rev.* 15, 1–13. doi: 10.1017/S1466252313000170
- U.S. Department of Agriculture. (2001). Part I. *Reference of Swine Health and Health Management in the United States, 2000*. Fort Collins, CO: National Animal Health Monitoring System. Publication no. 338.0801.
- Wells, T. J., Sherlock, O., Rivas, L., Mahajan, A., Beatson, S. A., Torpdahl, M., et al. (2008). EhaA is a novel autotransporter protein of enterohemorrhagic *Escherichia coli* O157:H7 that contributes to adhesion and biofilm formation. *Environ. Microbiol.* 10, 589–604. doi: 10.1111/j.1462-2920.2007.01479.x
- Woodward, D. L., Clark, C. G., Caldeira, R. A., Ahmed, R., and Rodgers, F. G. (2002). Verotoxigenic *Escherichia coli* (VTEC): A major public health threat in Canada. *Can. J. Infect. Dis.* 13, 321–330. doi: 10.1155/2002/383840

Disclaimer: Mention of trade names or commercial products is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Baranzoni, Fratamico, Gangiredla, Patel, Bagi, Delannoy, Fach, Boccia, Anastasio and Pepe. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.