



Photoregulation in a Kleptochloroplastic Dinoflagellate, *Dinophysis acuta*

Per J. Hansen^{1*}, Karin Ojamäe², Terje Berge^{1,3}, Erik C. L. Trampe¹, Lasse T. Nielsen³, Inga Lips² and Michael Kühl^{1,4}

¹ Marine Biological Section, Department of Biology, University of Copenhagen, Helsingør, Denmark, ² Marine Systems Institute, Tallinn University of Technology, Tallinn, Estonia, ³ Centre for Ocean Life, DTU Aqua National Institute for Aquatic Resources, Technical University of Denmark, Charlottenlund, Denmark, ⁴ Plant Functional Biology and Climate Change Cluster, University of Technology Sydney, Sydney, NSW, Australia

Some phagotrophic organisms can retain chloroplasts of their photosynthetic prey as so-called kleptochloroplasts and maintain their function for shorter or longer periods of time. Here we show for the first time that the dinoflagellate *Dinophysis acuta* takes control over “third-hand” chloroplasts obtained from its ciliate prey *Mesodinium* spp. that originally ingested the cryptophyte chloroplasts. With its kleptochloroplasts, *D. acuta* can synthesize photosynthetic as well as photoprotective pigments under long-term starvation in the light. Variable chlorophyll fluorescence measurements showed that the kleptochloroplasts were fully functional during 1 month of prey starvation, while the chlorophyll *a*-specific inorganic carbon uptake decreased within days of prey starvation under an irradiance of 100 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. While *D. acuta* cells can regulate their pigmentation and function of kleptochloroplasts they apparently lose the ability to maintain high inorganic carbon fixation rates.

Keywords: acquired phototrophy, *Dinophysis*, kleptochloroplasts, photoregulation, photosynthesis

OPEN ACCESS

Edited by:

Senjie Lin,
University of Connecticut, USA

Reviewed by:

Kathleen Scott,
University of South Florida, USA
Holly V. Moeller,
Woods Hole Oceanographic
Institution, USA

*Correspondence:

Per J. Hansen
pjhansen@bio.ku.dk

Specialty section:

This article was submitted to
Aquatic Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 07 March 2016

Accepted: 09 May 2016

Published: 30 May 2016

Citation:

Hansen PJ, Ojamäe K, Berge T,
Trampe ECL, Nielsen LT, Lips I
and Kühl M (2016) Photoregulation
in a Kleptochloroplastic
Dinoflagellate, *Dinophysis acuta*.
Front. Microbiol. 7:785.
doi: 10.3389/fmicb.2016.00785

INTRODUCTION

Some free-living phagotrophic protists sequester chloroplasts from their algal prey and utilize them for shorter or longer time; a life style that is common among ciliates, dinoflagellates and radiolarians (e.g., Stoecker et al., 2009; Johnson, 2011). In many such species, however, other prey cell organelles are retained as well. Red tide ciliates, *Mesodinium* spp., ingest certain cryptophyte prey species and sequester not only the chloroplasts (Johnson et al., 2006; Moeller et al., 2011; Hansen et al., 2012), but also a number of other prey cell organelles, such as mitochondria, the prey nucleus, and the nucleomorph (a reduced former nucleus of an earlier endosymbiont found in cryptophytes). These red-pigmented *Mesodinium* spp. can keep the ingested prey organelles functionally active for several months gaining enough photosynthate for survival in periods of low prey abundance (Johnson and Stoecker, 2005; Hansen and Fenchel, 2006; Johnson et al., 2006, 2007; Smith and Hansen, 2007). In fact, the ingestion of only a single cryptophyte prey cell per day (~1% of daily carbon needs) is sufficient to maintain maximum growth rate (Smith and Hansen, 2007). The red-pigmented *Mesodinium* spp. can control the division of its ingested prey organelles (Hansen and Fenchel, 2006), but this ability seems to get lost after a few cell divisions in prey-starved cultures probably due to loss of prey nuclei material (Johnson et al., 2007; Moeller et al., 2011). Red-pigmented *Mesodinium* spp. display photoacclimation, i.e., increases in cellular photosynthetic pigments at low irradiance and a change in photosynthesis vs. irradiance response curves (Johnson et al., 2006; Moeller et al., 2011).

In other protist species, only prey chloroplasts are sequestered and typically only remain functional in the predator cell for hours up to a few days. However, in some cases such kleptochloroplasts are kept active over several weeks, allowing kleptochloroplastic predators to survive periods of prey starvation a lot better than purely heterotrophic protists (Stoecker et al., 2009). One conspicuous case is found among the dinoflagellate genus *Dinophysis*, which are associated with diarrhetic shellfish poisoning worldwide. Species within this genus have chloroplasts of cryptophyte origin and the first culture of a *Dinophysis* spp. was established by feeding it a red-pigmented *Mesodinium* spp. suggesting that its cryptophyte chloroplasts were supplied from the ciliate prey (Park et al., 2006). This was later confirmed by Kim et al. (2012) and *Dinophysis* spp. can thus employ “third-hand” chloroplasts. A total of eight species have now been cultured all relying on red-pigmented *Mesodinium* spp. as prey (Hansen et al., 2013). *Dinophysis* spp. use a peduncle to suck out the contents of the ciliate prey. *Dinophysis* spp. do not ingest cryptophytes directly and thus rely entirely on *Mesodinium* spp. for the supply of chloroplasts and food. There are no direct evidence suggesting that *Dinophysis* spp. may feed other protists, detritus, or bacteria (Poulsen et al., 2011; Hansen et al., 2013).

It has recently been shown that *Dinophysis acuta* and *D. caudata*, grown together with red *Mesodinium* spp. under low irradiance have an elevated cellular Chl *a* content compared to cells grown at high irradiance (Rial et al., 2013). *Dinophysis* cells grown under low irradiance contain more and larger kleptochloroplasts as compared to cells grown at high irradiance (Nielsen et al., 2012, 2013). This suggests that *Dinophysis* spp. may have the capacity to photoregulate, but how is this possible when neither prey nuclei nor nucleomorphs are retained by *Dinophysis* spp. remains unknown.

The apparent regulation of kleptochloroplasts in *Dinophysis* spp. could work in three ways, alone or in combination: (i) Behavioral regulation: *Dinophysis* spp. may increase the number of chloroplasts incorporated via ingestion when grown under low irradiance and may fuse ingested chloroplasts leading to the observed increase in chloroplast size and cellular Chl *a* content; (ii) Photoregulation: *Dinophysis* spp. may be able to produce Chl *a* and other photosynthetic pigments, while the photosynthetic light response curves are unaffected; (iii) Photoacclimation: If *Dinophysis* spp. perform photoacclimation, an increase in cellular Chl *a* concentration as well as a change in the photosynthetic light response curve is expected. This would indicate full control over the acquired chloroplasts. In the latter two cases, the genes involved in either of such regulations must have been transferred to the host cell. A few chloroplast housekeeping genes have been found in the genome of *Dinophysis acuminata*, but it is unlikely that they alone allow for either photoregulation or photoacclimation (Wisecaver and Hackett, 2010).

In the present study, we investigated the photoregulation potential of *D. acuta* cells deprived of prey while being subjected to low irradiance to test whether photoregulation or photoacclimation occurs in sequestered chloroplasts of cryptophyte origin. For this, we employed a suite of different experimental techniques quantifying cell devisions,

photosynthetic performance, inorganic carbon (C_i) uptake, and respiration of *D. acuta* in cultures starved at the initiation of the experiments and incubated at different irradiance levels. We hypothesize that *D. acuta* is only capable of the first of the three options listed above, behavioral regulation, and thus will display no, or limited, control over its kleptochloroplasts during starvation. This would include the lack of ability to synthesize Chl *a* and limited adaptation of photosynthetic light curve response parameters. Thus, we expect to find an exponential decline in cellular Chl *a* content as well as declining values of C_i assimilation rate and photosynthetic performance, as cells divide during prey starvation.

MATERIALS AND METHODS

Organisms and Culture Conditions

Cultures of the cryptophyte *Teleaulax amphioxeia* (K-1837; SCCAP) and the ciliate *Mesodinium rubrum* (MBL-DK2009) were established from water samples from Helsingør Harbor in 2009. Cultures of *M. rubrum* were fed *T. amphioxeia* at a predator:prey ratio of 1:10 twice a week. Cultures of *D. acuta* were established in June 2010 from the North Sea (Nielsen et al., 2013). *M. rubrum* was added as prey organism twice per week at a predator:prey ratio of ~1:10 to allow mixotrophic growth. Only *M. rubrum* cultures that had completely removed their cryptophyte prey were used to feed *D. acuta*. All three species were grown in *f*/2 medium (Guillard, 1983) based on autoclaved seawater with a salinity of 32, a dissolved inorganic carbon (DIC) concentration of 2.3 ± 0.1 mmol l^{-1} and a pH of 8.0 ± 0.05 . pH was monitored directly in the flasks with a SenTix41® pH electrode (WTW, Germany) connected to a pH meter (WTW, pH 3210, Germany), and calibrated by measurements in pH 7 and pH 10 standard buffers (WTW, Technischer, NIST, buffers). All organisms were grown at $15.0 \pm 1.0^\circ C$ in a temperature regulated room, under a photon irradiance of $100 \mu mol photons m^{-2} s^{-1}$ (PAR, 400–700 nm), as controlled by a timer to a light:dark cycle of 16:8 h, unless otherwise stated. Culture flasks were placed on a glass table with light coming from below. Light was provided by cool white fluorescent tubes (OSRAM 58W, 840) and photon irradiance was measured (in air) at the level of incubation flasks using a light meter equipped with a spherical quantum sensor (ULM & US-SQS/L, Walz GmbH, Germany). All cultures were xenic.

Experimental Design

Experiment 1

This experiment was designed to study photoregulation in a recently prey starved culture of *D. acuta*. A culture of *D. acuta* was maintained in 750 ml tissue culture flasks filled with 500 ml culture medium under a photon irradiance of $100 \mu mol photons m^{-2} s^{-1}$ with prey for at least 2 weeks prior to the experiment and was eventually allowed to deplete its ciliate prey (below detection limit which was a few cells ml^{-1}). At the initiation of the experiment, subsamples of the culture were split in two and poured into 270 ml tissue culture flasks to capacity at a *D. acuta* cell density of $200 cells ml^{-1}$ (in triplicate). One set

of flasks were maintained at $100 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (I_{100}), while the other set of flasks was shifted to a photon irradiance of $15 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (I_{15}). Subsamples were withdrawn on day 3, 6, 9, 13, 16, and 29 (I_{100}), and day 4, 7, 10, 14, 17, and 30 (I_{15}) for measurements of cell concentration, ^{14}C fixation, algal pigment concentration using both fluorometry and HPLC techniques, photosynthetic capacity using variable chlorophyll fluorimetry, and O_2 optode-based respiration measurements (see descriptions of these techniques below).

pH was checked on each sampling occasion to avoid physiological effects of elevated pH in laboratory cultures (Hansen, 2002). Any cryptophytes left in the incubation flasks at the start of the experiment grew to concentrations where they could affect pH, especially at I_{100} . As the *Dinophysis* culture increased in cell concentration they also affected the pH of the culture medium. Therefore, we removed most of the culture medium once each week during the experiment, using gentle inverse filtration, and applied fresh $f/2$ culture medium to ensure optimal growth conditions (i.e., nutrients, pH, etc.). A plastic tube with plankton gauze (mesh size $20 \mu\text{m}$) attached to the tip was lowered directly into the experimental flasks and $\sim 90\%$ “old” growth medium was removed gently using a pipette and replaced with fresh growth medium. This was repeated three times each time.

Experiment 2

This experiment was carried out to measure Chl *a*, and physiological rates in well fed cultures of *D. acuta* at an irradiance of $100 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$. Cultures were maintained in 750 ml tissue culture flasks filled with 500 ml culture medium prior to the experiment. At the initiation of the experiment, 270 ml tissue culture flasks were filled to capacity with an initial concentration of 400 *D. acuta* cells ml^{-1} and 1000 *M. rubrum* cells ml^{-1} (in triplicate). Samples were withdrawn three times over 6 days for measurements of cell concentration, ^{14}C fixation, photosynthetic capacity using variable chlorophyll fluorimetry, and O_2 optode-based respiration measurements (see descriptions of these techniques below). Fresh growth medium was added at each sampling to replace the water volume removed. *Dinophysis* cells were picked individually using a drawn Pasteur glass pipette and transferred to clean growth medium twice to remove all *M. rubrum* cells before measurements.

Cell Enumeration

Cells were fixed in Lugol’s solution (1% final conc.) and were enumerated using a Sedgewick Rafter cell and an inverted microscope (Olympus®, CK-40). A minimum of 200 *Dinophysis* cells were counted each time and all samples were checked for possible leftover of ciliate prey cells.

Photosynthetic Pigments

Fluorimetric Measurements of Algal Pigments

Cellular chlorophyll concentration was measured by extracting chlorophyll of single cells according to Skovgaard et al. (2000). One ml of 96% ethanol was added to the borosilicate measuring vials. A total of 80 individual *Dinophysis* cells were pipetted directly into the ethanol and placed in the dark. This process took place in a temperature-regulated room kept at experimental

temperature. The chlorophyll *a* concentration ($\mu\text{g Chl } a \text{ l}^{-1}$) was measured in the extract after 20 min using a bench-top fluorometer (Trilogy, Turner designs, USA) equipped with the manufacture’s module for Chl *a* determination. The fluorometer was calibrated against a pure Chl *a* standard ($2.13 \text{ mg Chl } a \text{ l}^{-1}$) of cyanobacterial origin (DHI, Hørsholm, Denmark).

Additionally, we measured the *in vivo* chlorophyll, and phycoerythrin fluorescence using the appropriate filter set modules for the fluorimeter. For this, 80 individual *Dinophysis* cells were isolated and transferred into plastic measuring cuvettes containing 1 ml of fresh medium. The cells were kept in suspension by carefully blowing air with a pipette immediately before the measurements. Due to high variation between single measurements, the *in vivo* fluorescence was measured every second and averaged over a period of 10 s.

Photopigment Analysis

We analyzed the pigment composition of *D. acuta* with high-pressure liquid chromatography (HPLC) using a slightly modified method from Frigaard et al. (1996) measuring the major light harvesting pigment Chl *a*, the photoprotective pigment alloxanthin, and the ratio between the two in *D. acuta* throughout the starvation experiments. For pigment conservation prior to HPLC analysis, 200 *D. acuta* cells were picked with a micropipette and transferred twice into fresh growth medium, before they were spun down at 10 g for 5 min. The supernatant was removed, and the pellet was frozen at -80°C . Before HPLC analysis, each pellet containing 200 *D. acuta* cells was resuspended for pigment extraction in $125 \mu\text{l}$ of an acetone-methanol (7:2, vol/vol) mixture and sonicated using an ultrasonicator (Misonix 4000, Qsonica LLC., Newtown, CT, USA) under dim light for 20 s. The sonicated cells were extracted in darkness on ice for 2 min. The extracts were briefly centrifuged to pellet cell debris, and the supernatants were mixed with $15 \mu\text{L}$ ammoniumacetate (1 M) in 0.3 ml HPLC vials. A $100 \mu\text{L}$ sample of the mixture was then immediately injected into the HPLC.

Pigment extracts were separated and analyzed by a diode array detector connected to the HPLC system (HPLC-DAD & Agilent 1260 Infinity, Agilent Technologies, Santa Clara, CA, USA) fitted with a Nova-pak C18 column (dimensions: $3.9 \times 300 \text{ mm}$), detecting specific absorption wavelengths of compounds. The extracts were run at a column temperature of 30°C for 69 min. Injected extracts moved with a flow-rate of 1.0 ml min^{-1} in solvent A (methanol:acetonitrile:water, 42:33:25, vol/vol/vol), and solvent B (methanol:acetonitrile:ethyl acetate, 50:20:30, vol/vol/vol). In the separation process, the mobile phase changed linearly from 30% solvent B at the time of injection to 100% at 52 min, staying at 100% for 15 min and then falling back to 30% within 2 min. Chl *a* and alloxanthin were identified manually from HPLC chromatograms, and pigment ratios were calculated from the derived integrated peak areas using the manufacturers software (OpenLAB CDS ChemStation Edition, Agilent Technologies, Santa Clara, CA, USA).

Variable Chlorophyll Fluorescence Analysis

We assessed the photosynthetic capacity of *D. acuta*, mainly by means of evaluating the dark adapted maximum PSII quantum

yield, F_v/F_m as measured by the pulse saturation method with pulse-amplitude modulated (PAM) fluorimeters (Schreiber et al., 1994). Individual cells were imaged with a variable chlorophyll fluorescence imaging system (RGB Microscopy PAM, Walz GmbH, Germany; Trampe et al., 2011), while bulk culture samples were investigated in a cuvette-based chlorophyll fluorimeter (MULTI-COLOR-PAM; Walz GmbH, Germany). Detailed descriptions of the two types of PAM measurements, and definitions of the parameters used, are provided in the Supplementary Material, Section Material and Methods.

Sample preparation – Imaging PAM

1.5 ml subsamples of *D. acuta* culture were transferred to 1.5 ml tubes and centrifuged at 43 g for 2 min. The 1.4 ml supernatant was removed, and the pellet was resuspended in the remaining 0.1 ml yielding a concentrated sample facilitating a low search time for cells of interest at high magnification under the microscope. Control experiments showed no significant effect on cell structure or photosynthetic capacity by the treatment (data not shown). A droplet (16.5 μ L) of concentrated sample was transferred to a special treated microscope slide displaying a permanently positive charge (Superfrost Ultra Plus, Thermo Scientific, Gerhard Menzel GmbH, Braunschweig, Germany) resulting in good cell adhesion by electrostatic attraction, limiting motion of cells during imaging. Variable chlorophyll fluorescence imaging measurements rely on recording of consecutive images when calculating photosynthetic parameters, and it is thus crucial that there is no cell movement during the saturation pulse. The sample was sealed with a cover glass using petroleum jelly (VaselineTM) at the periphery to avoid evaporation, and the slide was placed at 12°C in a thermostated slide holder (see details in Trampe et al., 2011).

Sample preparation – MULTI-COLOR PAM

For each sample, 1000 (Experiment 1) or 300 (Experiment 2) cells were isolated with a drawn out Pasteur pipette and transferred several times through sterile-filtered seawater before final transfer into quartz cuvettes and the volume was adjusted to 1 ml using 30 psu sterile-filtered f/2 medium. Samples were stirred with a magnetic stirrer during measurements.

Inorganic Carbon Uptake and Respiration

¹⁴C Incorporation

Subsamples (1–2 ml) were removed from each culture flask for photosynthesis measurements. A total of 80 *D. acuta* cells were picked from the subsamples and washed in f/2 medium by micromanipulation using a stereoscope to remove all prey. A total of 40 *D. acuta* cells were transferred to each of two 20 ml glass scintillation vials containing 2 ml f/2 medium, and 20 μ l of a NaH¹⁴CO₃ stock solution (specific activity = 100 μ Ci ml⁻¹; Carbon-14 Centralen, Denmark) was added. One vial was incubated for ~3 h under the treatment irradiance, while the other was kept in complete darkness (by wrapping in several layers of aluminum foil). After incubation, a 100 μ L sub-sample was transferred to a new vial containing 200 μ l phenylethylamine for determination of specific activity (see Skovgaard et al., 2000

for a detailed method description). The remaining 1.9 ml were acidified with 2 ml 10% glacial acetic acid in methanol, and evaporated overnight at 60°C to remove all inorganic carbon. The residue was then re-dissolved in 2 ml Milli-Q water. All vials were treated with 10 ml Packard Insta-Gel Plus scintillation cocktail, and disintegrations per minute were measured using a Packard 1500 Tri-Carb liquid scintillation analyzer with automatic quench correction. The photosynthetic activity (PA, pg C cell⁻¹ h⁻¹) per cell was calculated as follows:

$$PA = \frac{DPM \times [DIC]}{^{14}C \times h \times N}$$

where DPM is disintegrations min⁻¹ (in 1.9ml) in the light corrected for dark values, DIC is the concentration of DIC (pg C ml⁻¹), ¹⁴C is the specific activity (disintegrations min⁻¹ ml⁻¹), *h* is the incubation time, and *N* is the number of cells in the vial (1.9 ml). DIC concentrations were measured on 1 ml subsamples using an infrared gas analyzer (ADC 225 Mk3 Gas analyzer, Analytic Development Co. Ltd., Hoddesdon, England) as described in detail elsewhere (Nielsen et al., 2007). Glass vials with screw caps were used for DIC samples allowing no headspace, and the samples were analyzed within a few hours.

Respiration Measurements

Respiration rates of *D. acuta* were measured in 1.8 ml glass vials (*n* = 4) equipped with calibrated optical O₂ sensor spots with optical isolation (Pyroscience GmbH, Germany). The O₂-dependent luminescence of each sensor spot was monitored contactless across the vial wall using an optical fiber cable fixed at one end to the glass vial using a solid plexi-glass adaptor and connected at the other end to a fiber-optic O₂ meter (FireSting, PyroScience, Germany). The sensor spot readout was calibrated from readings in anoxic and fully aerated medium. A total of 400 *Dinophysis* cells were isolated by micropipetting and washed 3 times in sterile filtered (pore size 0.22 μ m) f/2 culture medium prior to addition to each of the 4 glass vials containing 1 ml sterile f/2 medium. A glass bead was added to each of the vials that were filled to capacity with sterile filtered medium and carefully sealed avoiding any air bubbles inside the closed vials. The vials were mounted on a modified whirley mixer run at low speed to homogenize internal O₂ gradients in the sample (Trampe et al., 2015). The measurements were done in darkness in a temperature regulated room (15°C), and O₂ was measured with the built-in temperature correction of the O₂ meter. The O₂ concentration in the glass vials was measured for 1–3 h, and the linear decrease in O₂-concentration (μ mol l⁻¹ s⁻¹) was converted into cellular respiration rate (pg C cell⁻¹ day⁻¹) by assuming a respiratory quotient of 1.

RESULTS

Changes in Photosynthetic Pigments in *D. acuta* Starved of Prey

No ciliate prey cells were observed during enumeration of *Dinophysis* cells for the entire duration the prey starvation experiment. *D. acuta* cultures subjected to prey starvation at high photon irradiance ($I_{100} = 100 \mu$ mol photons m⁻² s⁻¹) increased

from 200 to 1400 cells ml^{-1} during 1 month of incubation, equivalent to an average growth of 2.8 cell divisions in total (Experiment 1, **Figure 1A**). The subculture that was exposed to low photon irradiance ($I_{15} = 15 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) increased in cell concentration from 200 to 1000 cells ml^{-1} and thus had <2.3 cell divisions during the 1 month of incubation (**Figure 1A**). The cellular Chl *a* concentration in *D. acuta* incubated at I_{100} decreased exponentially from the initial ~ 53 – $16 \text{ pg Chl } a \text{ cell}^{-1}$ (**Figure 1B**), leading to an overall increase in Chl *a* concentration from 11 to a maximum of $26 \text{ ng Chl } a \text{ ml}^{-1}$ at Day 16 (**Figure 1C**). Cells incubated at I_{15} , however, maintained their cellular Chl *a* concentration during the 1 month long incubation, even though cells divided 2.3 times. This led to an increase in the amount of Chl *a* in these incubations from 11 to $46 \text{ ng Chl } a \text{ ml}^{-1}$ at Day 30, indicating a higher net production of Chl *a* production in prey-starved *Dinophysis* cells grown at I_{15} , compared to those at I_{100} (**Figure 1C**). For comparison, well fed cultures of *D. acuta* grown at I_{100} had a cellular Chl *a* concentration of $\sim 40 \text{ pg Chl } a \text{ cell}^{-1}$ (Experiment 2).


Cultures exposed to I_{100} exhibited a slight increase in the alloxanthin:Chl *a* ratio from 0.86 to ~ 1.2 over the first 10 days. Hereafter, the ratio remained constant indicating that Chl *a* was initially lost at a faster rate than alloxanthin (**Figure 2A**). With cultures exposed to I_{15} , a significant decrease in the alloxanthin:Chl *a* ratio from 0.86 to 0.7 was observed over the first 2 weeks indicating dilution due to growth, which was followed by an increase of the alloxanthin:Chl *a* ratio reaching ~ 1 at the end of the experiment. *In vivo* measurements of Chl *a* and phycoerythrin showed that the changes in phycoerythrin content under the two experimental irradiance regimes matched those found in Chl *a* (**Figure 2B**).

Single-Cell Variable Chlorophyll Fluorescence Imaging


In well fed cultures of *D. acuta*, incubated at an irradiance of $100 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, microscopy revealed a broad distribution of cellular Chl *a* with hotspots clearly defining 4–5 chloroplast centers with high fluorescence. Maximum PSII quantum yields (F_v/F_m) of ~ 0.5 in these centers indicated a healthy and functioning photosynthetic apparatus (**Figure 3A**). During the starvation experiment at I_{100} , we found a continuously declining Chl *a* coverage in the cells with incubation time (**Figures 3B–D**). After a slight increase in F_v/F_m over the first 2 days, F_v/F_m decreased with the declining chlorophyll coverage (**Figure 3B**; Supplementary Figure S1). At I_{15} we observed a rapid increase in F_v/F_m to ~ 0.6 over the first 3 days, which remained high even after 10 days, while the Chl *a* coverage had decreased a bit (**Figures 3E,F**; Supplementary Figure S1). After 30 days, we still observed an even coverage, almost similar to the well-fed cells, with similar condensation in four chloroplast centers, and with a sustained high level of F_v/F_m (**Figure 3G**).

Bulk Measurements of Photochemical and Non-photochemical Quenching

Measurements on bulk cell samples yielded more insight into the photochemical, and non-photochemical quenching processes



in starved *D. acuta* cultures. The maximum PSII quantum yield (F_v/F_m) was on average higher for cells at I_{15} (0.74 ± 0.04 ; average \pm SE, $n = 18$) than I_{100} (0.65 ± 0.05 , $n = 21$; **Figure 4A**). In both treatments, F_v/F_m showed an initial increase during the 1 week, but only cells at I_{15} were able to maintain a high F_v/F_m of ~ 0.76 , indicating better photosynthetic capacity than cells at I_{100} . The change in F_v/F_m resulted primarily from the change in



the fraction of “open” PSII reaction centers, qP (Supplementary Figure S2A). An increase in qP to 0.94 ± 0.02 ($n = 15$) at I_{15} indicates a decrease in the proportion of “closed” PSII centers due to lower excitation pressure, than compared to cells at I_{100} , which had a lower qP of 0.67 ± 0.02 ($n = 21$) throughout the starvation experiment.

Absorbed light energy partitioning profiles for *D. acuta* (Supplement Figures S2A,B) show a photoregulative change in response to changes in photon irradiance. The use of excitation energy for photosynthesis (ϕ_{II}) in cultures kept at I_{100} was lowered with respect to F_v/F_m by partial closure of PSII centers (Supplementary Figure S2A), and non-photochemical energy losses (ϕ_{NPQ} and ϕ_{NO}) induced by illumination. Only 38% of the absorbed light energy was used for photosynthetic reactions (Supplementary Figure S2B), whereas cells kept at I_{15} used 70% of the absorbed light for photosynthesis (Supplementary Figure S2C).

The maximum relative electron transport rate ($rETR_{max}$), and the initial slope of the light response curves (α rETR) showed no changes between the treatments (Figures 4B,C). For cultures incubated at I_{100} , the average $rETR_{max}$ and α rETR values were 76.3 ± 8.14 and 0.25 ± 0.01 ($n = 21$), respectively. Cultures at I_{15} showed average $rETR_{max}$ and α rETR values of 88.2 ± 2.58 and 0.31 ± 0.02 ($n = 18$), respectively, (Figures 4B,C).

Inorganic Carbon Uptake and Respiration before and during Starvation


Inorganic carbon uptake was 138 ± 6 pg C cell⁻¹ h⁻¹ (average \pm SE) in well fed cultures of *D. acuta* at I_{100} , equivalent to 2.21 ± 0.94 ng C cell⁻¹ day⁻¹ (light:dark cycle of 16:8 h), leading to a Chl *a* specific rate of 3.5 ± 0.5 pg C pg Chl *a*⁻¹. The inorganic carbon uptake of *D. acuta* cells subjected to starvation at I_{100} decreased exponentially from 177 ± 14 to 23.7 ± 4.2 pg C cell⁻¹ h⁻¹ at Day 16 (equivalent to 2.83 ± 0.23 and 0.38 ng C cell⁻¹ day⁻¹, respectively, Figure 5A), i.e., a 6.1-fold decrease in photosynthetic carbon fixation. This translated into a significant decline in the Chl *a*-specific carbon uptake from the initial 3.4 to 1.0 pg C pg Chl *a*⁻¹ h⁻¹ at Day 16 (one-way ANOVA, $P < 0.05$; Figure 5B). The following 2 weeks, the inorganic carbon uptake rate dropped further to 16.8 ± 1.1 pg C cell⁻¹ h⁻¹ ($=0.27 \pm 0.02$ ng C cell⁻¹ day⁻¹), but the Chl *a*-specific carbon uptake stayed constant (1.07 pg C pg Chl *a*⁻¹ h⁻¹).

Under low irradiance (I_{15}), inorganic carbon uptake remained constant for the first 17 days at 33.1 ± 2.8 pg C cell⁻¹ h⁻¹ ($=0.53 \pm 0.03$ ng C cell⁻¹ day⁻¹), and dropped only marginally to 20.7 ± 1.9 pg C cell⁻¹ h⁻¹ ($=0.33 \pm 0.03$ ng C cell⁻¹ day⁻¹) after 1 month of incubation. Thus, the Chl *a*-specific inorganic carbon uptake rates of the starved low irradiance *D. acuta* stayed constant at ~ 0.6 – 0.7 pg C μ g Chl *a*⁻¹ h⁻¹ during the entire experiment.

Well-fed cells of *D. acuta* had a respiration rate of 86.5 ± 0.16 pg C cell⁻¹ h⁻¹ at I_{100} , equivalent to 2.08 ± 0.38 ng C cell⁻¹ day⁻¹ (average \pm SE, $n = 9$). No difference in respiration rate was observed as a function of prey starvation after 1 day of starvation in either of the light treatments, thus data were merged for each irradiance (One way-ANOVA, $P > 0.05$). Data on respiration of prey starved cells are only available until day 16 and 17 for the high and low irradiances, respectively. Starved cells had approximately halved their respiration rates [42.9 ± 7.1 ($n = 11$) and 36.2 ± 6.6 ($n = 12$) pg C cell⁻¹ h⁻¹ at I_{100} and I_{15} , respectively]; respiration rates at the two irradiances were not statistically different (Two way-ANOVA, $P > 0.05$).

DISCUSSION

The present study documents for the first time that kleptochloroplasts (without prey nuclear material) taken up by *D. acuta* exhibit photoregulation, where photosynthetic pigments are produced to improve growth under both high and low irradiances. However, we saw no changes in photosynthesis vs. irradiance parameters like α rETR or max rETR during 1 month of starvation, similar to what is usually found in regular



algae displaying photoacclimation (e.g., MacIntyre et al., 2001). Thus, we found photoregulation but not true photoacclimation in *D. acuta*.

Under low irradiance, photoregulation enabled *D. acuta* to maintain their cellular Chl *a* and phycoerythrin contents even though cells divided several times. We also found that the concentration of the alloxanthin increased relative to Chl *a* in cells exposed to high irradiance, while this pigment was initially reduced in cells incubated under low irradiance. While the role of alloxanthin as a light harvesting or photoprotective pigment in cryptophytes is not well established, the few available reports suggest that the alloxanthin:Chl *a* ratio in cryptophytes is elevated at high irradiances indicating that it may have photoprotective properties (Schlüter et al., 2000; Laviale and Neveux, 2011) and alloxanthin may thus play a similar role in *D. acuta*.

The literature provides no evidence for division of kleptochloroplasts in *Dinophysis* spp., suggesting that the cellular number of chloroplasts will decrease during cell division (Minnhagen et al., 2008). Hence, the ability of *Dinophysis* to maintain cellular photosynthetic pigment concentrations may potentially lead to enlargement (rather than division) of individual chloroplasts (Nielsen et al., 2013). Enlargement of kleptochloroplasts has recently been shown in other dinoflagellates such as *Nusuttodinium aeruginosum* and *N. myriopyrenoides*, where a 20-fold enlargement of the chloroplasts was observed within 120 min after ingestion in *N. aeruginosum* (Yamaguchi et al., 2011; Onuma and Horiguchi, 2015). Up to 10-fold enlargement of ingested chloroplasts has also been observed in the katablepharid *Hatena arenicola*

(Yamaguchi et al., 2011). However, in all these cases, prey nuclei and nucleomorphs were retained by the host cell, which is not the case in *Dinophysis* spp.

Variable chlorophyll fluorescence data suggested that chloroplasts in *D. acuta* were competent with maximum PSII quantum yields (F_v/F_m) between 0.5 and 0.8 throughout the duration of the experiment (30 days). These values are similar to what is usually found in well-functioning microalgae (Falkowski and Raven, 2007). Similar observations have been obtained on *D. caudata* cells starved for up to 85 days (Park et al., 2008), where F_v/F_m began to decline after 24 days of starvation, and reached 0.4 after 45 days of starvation, i.e., at the same time as cell divisions stopped. Thereafter, F_v/F_m decreased steadily over the next days and reached zero after 85 days of starvation. We saw no indications of photodamage. Photodamage would be characterized by a decrease in ϕ_{NPQ} , which is one of the most important safety valves for the regulation of light harvest (Müller et al., 2001). Also, if the cells were photodamaged we would have expected to see a consecutive decrease of qP (Mackey et al., 2008), which gives information if photosynthetic efficiency has been altered by a changed proportion of functional reaction centers. In our case, ϕ_{NPQ} was relatively stable ($\sim 20\%$) and qP was constant in high light (60–70%) throughout the experiment, so the cells were unlikely to experience photodamage.

The prey species of *Dinophysis* spp., i.e., the red *Mesodinium* spp. retain ingested cryptophyte nuclei for up to 100 days and these have been shown to be transcriptionally active (Johnson et al., 2007; Lasek-Nesselquist et al., 2015). However, it has also been shown that the expression of a cryptophyte nuclear-encoded




FIGURE 4 | Variable chlorophyll fluorescence measurements on *D. acuta* culture samples incubated at I_{100} (o) and I_{15} (●) as a function of time after initiation of prey starvation. The culture had been growing at I_{100} prior to the initiation of the experiment. (A) Maximum PSII quantum yield (F_v/F_m). (B) Maximum relative electron transport rates (rETRmax) and (C) initial slopes of the light response curves (α rETR), i.e., rETR vs irradiance. Data points represent the mean value parameter estimates obtained from the rapid light response curves. Error bars show standard error of the mean ($n = 3$).

gene for the plastid-targeted protein, *LHCC10*, involved in plastid function, declined in *Mesodinium* cells as the sequestered nuclei disappeared from the population (Johnson et al., 2007). In the red *Mesodinium* spp., F_v/F_m values of 0.54–0.66 have been observed when grown for up to a month without prey (Johnson et al., 2006; Kim et al., 2014).

Another kleptochloroplastic dinoflagellate, *Amylax triacantha*, also preys on red *Mesodinium* spp. like *Dinophysis* spp. (Kim et al., 2014). Besides the cryptophyte chloroplasts,




FIGURE 5 | Changes in (A) Cellular inorganic carbon uptake ($\text{pg C cell}^{-1} \text{h}^{-1}$), and (B) Chl *a*-specific inorganic carbon uptake ($\text{pg C pg Chl } a^{-1} \text{h}^{-1}$) in *D. acuta* during the prey starvation experiment at two irradiances, I_{100} (o) and I_{15} (●), respectively. Data points represent treatment means and error bars indicate standard errors ($n = 3$). Arrows indicate values from well fed cultures for comparison.

A. triacantha also retains cryptophyte mitochondria, cryptophyte nuclei and nucleomorphs. However, unlike in *M. rubrum*, the ingested cryptophytes are kept as individual packages. In *A. triacantha* cultures subjected to prey starvation, F_v/F_m remains at 0.6 for the first 5–8 days, whereafter it declines over time to reach F_v/F_m values of ~ 0.05 after 25 days of prey starvation in conjunction with the loss of cryptophyte nuclear material, i.e., a much faster decline in photosynthetic capacity compared to the red *Mesodinium* spp.

Available evidence thus suggests that the cryptophyte chloroplasts are not independent entities, which will function on their own inside a host cell. In the red *Mesodinium* spp. and in *A. triacantha* the photosynthetic performance of kleptochloroplasts is coupled to the sequestration of prey nuclei and nucleomorph material. This is not the case in *Dinophysis* spp. Thus, it seems that the function of chloroplasts in *Dinophysis* may depend upon genes, which in the past have been transferred from prey nuclei and nucleomorphs to the dinoflagellate genome.

Very little is known about the transfer of genes from the cryptophyte genome to the genome of *Dinophysis* spp., and only one species, *D. acuminata*, has been investigated (Wisecaver and

Hackett, 2010). Five proteins, complete with plastid-targeting peptides that may function in photosystem stabilization and metabolite transport, have been found encoded in the nuclear genome of *D. acuminata* (Wisecaver and Hackett, 2010). It seems unlikely that those genes alone allow for the extensive regulation of photosynthetic and photoprotective pigments that we observed in *D. acuta*, and more genes controlling photoregulation in *D. acuta* and other *Dinophysis* spp. remain to be identified.

The Chl *a*-specific inorganic carbon uptake dropped dramatically in cells grown at high irradiance indicating that, although *D. acuta* was able to produce Chl *a*, this did not translate into increased inorganic carbon uptake. There may be several reasons for this observation: First, since *Dinophysis* spp. only sequester the chloroplasts from their prey, any loss of genes involved in the regulation of RuBisCO that are located in the cryptophyte nuclei material are lost and this will strongly affect inorganic carbon fixation. Such genes may come from the prey nucleus, but they may also come from the nucleomorph. The Rubisco regulatory protein, CbbX, for instance, is encoded by the cryptophyte nucleomorph in cryptophytes (Maier et al., 2000).

Second, RuBisCo is located in the single terminally placed pyrenoid, contained by every chloroplast in most algae (Holdsworth, 1971; Giordano et al., 2005; Garcia-Cuetos et al., 2010). As *D. acuta* cannot divide the acquired chloroplasts, it will also be unable to divide its pyrenoids, which seems to be a key missing control factor, lacked by these protists to fully regulate and divide the sequestered “third hand” chloroplasts.

In purely phototrophic dinoflagellates, respiration rates typically account for ~20 and ~40% of gross photosynthesis in the exponential and stationary growth phase, respectively (Geider and Osborne, 1989; Geider, 1992; López-Sandoval et al., 2014). In the kleptochloroplastidic *D. acuta*, respiration rates accounted for ~50% of gross photosynthesis in well fed cells, and that percentage increased when the cells were subjected to prey starvation. Under high irradiance, *D. acuta* cells became full of storage material (unpublished observations, probably lipids and starch; Durand-Clement et al., 1988) upon initiation of prey starvation, in accordance with earlier observations in *D. caudata* (Park et al., 2008). Accordingly, we found excess inorganic carbon uptake during the first ~9 days under high irradiance, whereafter respiration rates started to slightly exceed rates of inorganic carbon uptake. Under low irradiance, respiration rates slightly exceeded the inorganic carbon uptake at all times. The build up of carbon storage upon onset of prey starvation and maintenance of fully active kleptochloroplasts thus allows *D. acuta* to survive for extended periods of time (months; Park et al., 2008; Nielsen et al., 2012, 2013). This is an important trait of these kleptochloroplastidic dinoflagellates, and explains how they can survive in a fluctuating environment and still depend upon *M. rubrum* as a single type of prey.

CONCLUSION

We found strong evidence of photoregulation via production of photosynthetic and photoprotective pigments in the klepto-

chloroplastidic dinoflagellate, *D. acuta*, which only retains the chloroplasts from its prey. No direct evidence of changes in photosynthesis vs. irradiance parameters and thus no signs of photoacclimation were observed in *D. acuta* incubated at different irradiances, and thus no signs of photoacclimation were found. A decrease in Chl *a*-specific inorganic carbon uptake in prey-starved cells indicated a dilution of carbon fixing units (Rubisco) among daughter cells in combination with the production of Chl *a*. However, in their natural environment *D. acuta* and the >100 species that belong to the genus *Dinophysis* may also indirectly photoregulate via increased retention of kleptochloroplasts, when prey cells are available. It is also possible that *Dinophysis* cells may indirectly achieve “photoacclimation” by ingestion of photoacclimated prey cells and maintaining a higher number of chloroplasts. Our data point to a hitherto unstudied role of gene transfer from prey to *D. acuta* that may enable it to regulate the function of its kleptochloroplast, although the exact genetic and biochemical mechanisms remain to be identified.

AUTHOR CONTRIBUTIONS

All authors were involved in the design and planning of the experiments. PH prepared the cultures for experimentation and was in charge of sampling scheme, carried out all ¹⁴C measurements and data analysis, except for the last sampling where TB carried out the measurements. TB did all respiration measurements and data analysis, isolation of cells for all pigment analyses and carried out fluorometric measurements and connected data analysis of phytopigments. ET carried out all Imaging PAM and HPLC measurements of phytopigments and the connected data analysis. KO carried out all MultiColorPAM measurements and the connected data analysis. PH wrote the manuscript with inputs from all of the co-authors.

FUNDING

This work was supported by the Villum Foundation via the “Centre for Ocean Life” (PH, TB, and LN), a grant from the Danish Research Council for Independent Research, grant no 4181-00484 (PH), VKR Institutional funding IUT (19-6) for the Estonian Ministry of Education and Research and by the Estonian Science Foundation, grant no 8930 (KO), an EliteForsk Travel Award from the Danish Ministry of Higher Education and Science, no 1323-00065B (ET), and a Sapere Aude Advanced Grant from the Danish Research Council for Independent Research of Natural Science (MK), grant no 3071-00007B.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2016.00785>

REFERENCES

- Durand-Clement, M., Clement, J.-C., Moreau, A., Jeanne, N., and Puiseux-Dao, S. (1988). New ecological and ultrastructural data on the dinoflagellate *Dinophysis* sp. from the French coast. *Mar. Biol.* 97, 37–44. doi: 10.1007/BF00391243
- Falkowski, P. G., and Raven, J. A. (2007). “The photosynthetic light reactions,” in *Aquatic Photosynthesis*, 2nd Edn, eds P. G. Falkowski and J. A. Raven (Oxfordshire: Princeton University Press), 81–117.
- Frigaard, N. U., Larsen, K. L., and Cox, R. P. (1996). Spectrochromatography of photosynthetic pigments as a fingerprinting technique for microbial phototrophs. *FEMS Microb. Ecol.* 20, 69–77. doi: 10.1111/j.1574-6941.1996.tb00306.x
- García-Cuetos, L., Moestrup, Ø, Hansen, P. J., and Daugbjerg, N. (2010). The toxic dinoflagellate *Dinophysis acuminata* harbors permanent chloroplasts of cryptomonad origin, not kleptochloroplasts. *Harmful Algae* 9, 25–38. doi: 10.1016/j.hal.2009.07.002
- Geider, R. J. (1992). “Respiration: taxation without representation?,” in *Primary Productivity and Biogeochemical Cycles in the Sea*, eds P. G. Falkowski and A. D. Woodhead (London: Plenum Press), 333–360.
- Geider, R. J., and Osborne, B. A. (1989). Respiration and microalgal growth: a review of the quantitative relationship between dark respiration and growth. *New Phytol.* 112, 327–341. doi: 10.1111/j.1469-8137.1989.tb00321.x
- Giordano, M., Beardall, J., and Raven, J. A. (2005). CO₂ concentrating mechanisms in algae: mechanisms, environmental modulation, and evolution. *Annu. Rev. Plant Biol.* 56, 99–131. doi: 10.1146/annurev.arplant.56.032604.144052
- Guillard, R. R. L. (1983). “Culture of the phytoplankton for feeding invertebrate animals,” in *Culture of Marine Invertebrates*, ed. C. J. Berg (Stroudsburg, PA: Hutchinson Ross), 123–128.
- Hansen, P. J. (2002). The effect of high pH on the growth and survival of marine phytoplankton: implications for species succession. *Aquat. Microb. Ecol.* 28, 279–288. doi: 10.3354/ame028279
- Hansen, P. J., and Fenchel, T. (2006). *Mesodinium rubrum* harbours a permanent endosymbiont. *Mar. Biol. Res.* 2, 169–177. doi: 10.1080/17451000600719577
- Hansen, P. J., Moldrup, M., Tarangkoon, W., García-Cuetos, L., and Moestrup, Ø. (2012). Direct evidence for symbiont sequestration in the marine red tide ciliate *Mesodinium rubrum*. *Aquat. Microb. Ecol.* 66, 63–75. doi: 10.3354/ame01559
- Hansen, P. J., Nielsen, L. T., Johnson, M. D., Berge, T., and Flynn, K. J. (2013). Acquired phototrophy in *Mesodinium* and *Dinophysis* – a review of cellular organization, prey selectivity, nutrient uptake and bioenergetics. *Harmful Algae* 28, 126–139. doi: 10.1016/j.hal.2013.06.004
- Holdsworth, R. H. (1971). The isolation and partial characterization of the pyrenoid protein of *Eremosphaera viridis*. *J. Cell Biol.* 51, 499–513. doi: 10.1083/jcb.51.2.499
- Johnson, M. D. (2011). The acquisition of phototrophy: adaptive strategies of hosting endosymbionts and organelles. *Photosynth. Res.* 107, 117–132. doi: 10.1007/s11220-010-9546-8
- Johnson, M. D., Oldach, D., Delwiche, C. F., and Stoecker, D. K. (2007). Retention of transcriptionally active cryptophyte nuclei by the ciliate *Myrionecta rubra*. *Nature* 445, 426–428. doi: 10.1038/nature05496
- Johnson, M. D., and Stoecker, D. K. (2005). Role of feeding in growth and photophysiology of *Myrionecta rubra*. *Aquat. Microb. Ecol.* 39, 303–312. doi: 10.3354/ame039303
- Johnson, M. D., Tengs, T., Oldach, D., and Stoecker, D. K. (2006). Sequestration, performance, and functional control of cryptophyte plastids in the ciliate *Myrionecta rubra* (Ciliophora). *J. Phycol.* 42, 1235–1246. doi: 10.1111/j.1529-8817.2006.00275.x
- Kim, M., Kim, K. Y., Nam, S. W., Shin, W., Yih, W., and Park, M. G. (2014). The effect of starvation on plastid number and photosynthetic performance in the kleptoplastidic dinoflagellate *Amylax triacantha*. *J. Eukaryot. Microb.* 61, 354–363. doi: 10.1111/jeu.12115
- Kim, M., Nam, S. W., Shin, W., Coats, D. W., and Park, M. G. (2012). *Dinophysis caudata* (Dinophyceae) sequesters and retains plastids from the mixotrophic ciliate prey *Mesodinium rubrum*. *J. Phycol.* 48, 569–579. doi: 10.1111/j.1529-8817.2012.01150.x
- Lasek-Nesselquist, E., Wisecaver, J. H., Hackett, J. D., and Johnson, M. D. (2015). Insights into transcriptional changes that accompany organelle sequestration from the stolen nucleus of *Mesodinium rubrum*. *BMC Genomics* 16:805. doi: 10.1186/s12864-015-2052-9
- Laviale, M., and Neveux, J. (2011). Relationships between pigment ratios and growth irradiance in 11 marine phytoplankton species. *Mar. Ecol. Prog. Ser.* 425, 63–77. doi: 10.3354/meps09013
- López-Sandoval, D. C., Rodríguez-Ramos, T., Cermeño, P., Sobrino, C., and Marañón, E. (2014). Photosynthesis and respiration in marine phytoplankton: relationship with cell size, taxonomic affiliation, and growth phase. *J. Exp. Mar. Biol. Ecol.* 457, 151–159. doi: 10.1016/j.jembe.2014.04.013
- MacIntyre, H. L., Kana, T. M., Anning, T., and Geider, R. J. (2001). Photoacclimation of photosynthesis irradiance response curves and photosynthetic pigments in microalgae and cyanobacteria. *J. Phycol.* 38, 17–38. doi: 10.1046/j.1529-8817.2002.00094.x
- Mackey, K. R. M., Paytan, A., Grossman, A. R., and Bailey, S. (2008). A photosynthetic strategy for coping in a high-light, low-nutrient environment. *Limnol. Oceanogr.* 53, 900–913. doi: 10.4319/lo.2008.53.3.0900
- Maier, U. G., Fraunholz, M., Zauner, S., Penny, S., and Douglas, S. (2000). A nucleomorph-encoded CbbX and the phylogeny of RuBisCo regulators. *Mol. Biol. Evol.* 17, 576–583. doi: 10.1093/oxfordjournals.molbev.a026337
- Minnhagen, S., Carvalho, W. F., Salomon, P. S., and Janson, S. (2008). Chloroplast DNA content in *Dinophysis* (Dinophyceae) from different cell cycle stages is consistent with kleptoplasty. *Environ. Microbiol.* 10, 2411–2417. doi: 10.1111/j.1462-2920.2008.01666.x
- Moeller, H. V., Johnson, M. D., and Falkowski, P. G. (2011). Photoacclimation in the phototrophic marine ciliate *Mesodinium rubrum* (Ciliophora). *J. Phycol.* 47, 324–332. doi: 10.1111/j.1529-8817.2010.00954.x
- Müller, P., Li, X.-P., and Niyogi, K. N. (2001). Non-photochemical quenching. A response to excess light energy. *Plant Physiol.* 125, 1558–1566. doi: 10.1104/pp.125.4.1558
- Nielsen, L. T., Krock, B., and Hansen, P. J. (2012). Influence of irradiance and food availability on DSP toxin production, growth and photosynthesis in *Dinophysis acuminata*. *Mar. Ecol. Prog. Ser.* 471, 34–50. doi: 10.3354/meps10027
- Nielsen, L. T., Krock, B., and Hansen, P. J. (2013). Production and excretion of okadaic acid, pectenotoxin-2 and a novel dinophysistoxin from the DSP-causing marine dinoflagellate *Dinophysis acuta* – effects of light, food availability and growth phase. *Harmful Algae* 23, 34–45. doi: 10.1016/j.hal.2012.12.004
- Nielsen, L. T., Lundholm, N., and Hansen, P. J. (2007). Does irradiance influence the tolerance of marine phytoplankton to high pH? *Mar. Biol. Res.* 3, 446–453. doi: 10.1080/17451000701711820
- Onuma, R., and Horiguchi, T. (2015). Kleptochloroplast enlargement, karyoklepty and the distribution of the cryptomonad nucleus in *Nusuttodinium* (= *Gymnodinium*) aeruginosum. *Protist* 166, 177–195. doi: 10.1016/j.protis.2015.01.004
- Park, M. G., Kim, S., Kim, H. S., Myung, G., and Kang, Y. G. (2006). First successful culture of the marine dinoflagellate *Dinophysis acuminata*. *Aquat. Microb. Ecol.* 45, 101–106. doi: 10.3354/ame045101
- Park, M. G., Park, J. S., and Kim, M. (2008). Plastid dynamics during survival of *Dinophysis caudata* without its ciliate prey. *J. Phycol.* 44, 1154–1163. doi: 10.1111/j.1529-8817.2008.00579.x
- Poulsen, L. K., Moldrup, M., Berge, T., and Hansen, P. J. (2011). Feeding on fecal pellets – a new trophic role for dinoflagellates as detritivores. *Mar. Ecol. Prog. Ser.* 441, 65–77. doi: 10.3354/meps09357
- Rial, P., Garrido, J. L., Jaén, D., and Rodríguez, F. (2013). Pigment composition in three *Dinophysis* species (Dinophyceae) and the associated cultures of *Mesodinium rubrum* and *Tealeaulax amphioxeia*. *J. Plankton Res.* 35, 433–437. doi: 10.1093/plankt/fbs099
- Schlüter, L., Møhlenberg, F., Havskum, H., and Larsen, S. (2000). The use of phytoplankton pigments for identifying and quantifying phytoplankton groups in coastal areas: testing the influence of light and nutrients on pigment/chlorophyll a ratios. *Mar. Ecol. Prog. Ser.* 192, 49–63. doi: 10.3354/meps192049
- Schreiber, U., Bilger, W., and Neubauer, C. (1994). “Chlorophyll fluorescence as a non-intrusive indicator for rapid assessment of in vivo photosynthesis,” in *Ecological Studies*, Vol. 100, eds E.-D. Schulze and M. M. Caldwell (Heidelberg: Springer), 49–70.

- Skovgaard, A., Hansen, P. J., and Stoecker, D. K. (2000). Physiology of the mixotrophic dinoflagellate *Fragilidium subglobosum*. I. Effects of phagotrophy and irradiance on photosynthesis and carbon content. *Mar. Ecol. Prog. Ser.* 201, 129–136. doi: 10.3354/meps201129
- Smith, M., and Hansen, P. J. (2007). Interaction between *Mesodinium rubrum* and its prey: importance of prey concentration, irradiance and pH. *Mar. Ecol. Prog. Ser.* 338, 61–70. doi: 10.3354/meps338061
- Stoecker, D. K., Johnson, M. D., de Vargas, C., and Not, F. (2009). Acquired phototrophy in aquatic protists. *Aquat. Microb. Ecol.* 57, 279–310. doi: 10.3354/ame01340
- Trampe, E., Hansen, P. J., and Kühl, M. (2015). A comparison of photosynthesis measurements by O₂ evolution, ¹⁴C assimilation and variable chlorophyll fluorescence during light acclimatization of the diatom *Coscinodiscus granii*. *Algae* 30, 103–119. doi: 10.4490/algae.2015.30.2.103
- Trampe, E., Kolbowski, J., Schreiber, U., and Kühl, M. (2011). Rapid assessment of different oxygenic phototrophs and single cell photosynthesis with multicolour variable chlorophyll fluorescence imaging. *Mar. Biol.* 158, 1667–1675. doi: 10.1007/s00227-011-1663-1
- Wisecaver, J. H., and Hackett, J. D. (2010). Transcriptome analysis reveals nuclear-encoded proteins for the maintenance of temporary plastids in the dinoflagellate *Dinophysis acuminata*. *BMC Genomics* 11:360. doi: 10.1186/1471-2164-11-366
- Yamaguchi, H., Nakayama, T., Kai, A., and Inouye, I. (2011). Taxonomy and phylogeny of a new kleptoplastidal dinoflagellate *Gymnodinium myriopyrenoides* sp. nov. (Gymnodiniales, Dinophyceae) and its cryptophyte symbiont. *Protist* 162, 650–677. doi: 10.1016/j.protis.2011.01.002

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Hansen, Ojamäe, Berge, Trampe, Nielsen, Lips and Kühl. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.