



Cell-Free Spent Media Obtained from *Bifidobacterium bifidum* and *Bifidobacterium crudilactis* Grown in Media Supplemented with 3'-Sialyllactose Modulate Virulence Gene Expression in *Escherichia coli* O157:H7 and *Salmonella* Typhimurium

OPEN ACCESS

Edited by:

Francesca Turroni,
University College Cork, Ireland

Reviewed by:

Alberto Amaretti,
University of Modena and Reggio
Emilia, Italy
Frédéric Borges,
École Nationale Supérieure
d'Agronomie et des Industries
Alimentaires - Université de Lorraine,
France

Pauline Bondue¹, Sébastien Crèvecoeur¹, François Brose¹, Georges Daube¹,
Marie-Christine Seghaye², Mansel W. Griffiths³, Gisèle LaPointe³ and
Véronique Delcenserie^{1*}

¹ Department of Food Science, Fundamental and Applied Research for Animal and Health, Faculty of Veterinary Medicine, University of Liège, Liège, Belgium, ² Department of Pediatrics, University Hospital Liège-Notre-Dame des Bruyères, Belgium, ³ Canadian Research Institute for Food Safety, University of Guelph, Guelph, Canada

*Correspondence:

Véronique Delcenserie
veronique.delcenserie@ulg.ac.be

Specialty section:

This article was submitted to
Microbial Symbioses,
a section of the journal
Frontiers in Microbiology

Received: 29 April 2016

Accepted: 01 September 2016

Published: 22 September 2016

Citation:

Bondue P, Crèvecoeur S, Brose F,
Daube G, Seghaye M-C, Griffiths MW,
LaPointe G and Delcenserie V (2016)
Cell-Free Spent Media Obtained from
Bifidobacterium bifidum and
Bifidobacterium crudilactis Grown in
Media Supplemented with
3'-Sialyllactose Modulate Virulence
Gene Expression in *Escherichia coli*
O157:H7 and *Salmonella*
Typhimurium.
Front. Microbiol. 7:1460.
doi: 10.3389/fmicb.2016.01460

Complex oligosaccharides from human milk (HMO) possess an antimicrobial activity and can promote the growth of bifidobacteria such as *Bifidobacterium bifidum* and *Bifidobacterium longum* subsp. *infantis*. In addition, fermentation of carbohydrates by bifidobacteria can result in the production of metabolites presenting an antivirulence effect on several pathogenic bacteria. Whey is rich in complex bovine milk oligosaccharides (BMO) structurally similar to HMO and *B. crudilactis*, a species of bovine origin, is able to metabolize some of those complex carbohydrates. This study focused on the ability of *B. bifidum* and *B. crudilactis* to grow in a culture medium supplemented in 3'-sialyllactose (3'SL) as the main source of carbon, a major BMO encountered in cow milk. Next, the effects of cell-free spent media (CFSM) were tested against virulence expression of *Escherichia coli* O157:H7 and *Salmonella enterica* serovar Typhimurium. Both strains were able to grow in presence of 3'SL, but *B. crudilactis* showed the best growth (7.92 ± 0.3 log cfu/ml) compared to *B. bifidum* (6.84 ± 0.9 log cfu/ml). Then, CFSM were tested for their effects on virulence gene expression by *ler* and *hilA* promoter activity of luminescent mutants of *E. coli* and *S. Typhimurium*, respectively, and on wild type strains of *E. coli* O157:H7 and *S. Typhimurium* using RT-qPCR. All CFSM resulted in significant under expression of the *ler* and *hilA* genes for the luminescent mutants and *ler* (ratios of -15.4 and -8.1 respectively) and *qseA* (ratios of -2.1 and -3.1) for the wild type strain of *E. coli* O157:H7. The 3'SL, a major BMO, combined with some

bifidobacteria strains of bovine or human origin could therefore be an interesting synbiotic to maintain or restore the intestinal health of young children. These effects observed *in vitro* will be further investigated regarding the overall phenotype of pathogenic agents and the exact nature of the active molecules.

Keywords: *Bifidobacterium bifidum*, *Bifidobacterium crudilactis*, bovine milk oligosaccharide, *Escherichia coli* enterohemorrhagic O157:H7, *Salmonella enterica* serovar Typhimurium, virulence expression, 3'-sialyllactose, whey

INTRODUCTION

Due to the influence on the microbiota of carbohydrate source present in food, breast-fed children are generally in better health than children fed with formula (Arrieta et al., 2014; Smilowitz et al., 2014; Scott et al., 2015). Human milk oligosaccharides (HMO) are complex oligosaccharides found in human milk. Their concentration can reach 15 g/l and more than 500 structures have been identified (Pacheco et al., 2015). These HMO, produced in mammary glands, cannot be metabolized either by the host or most bacteria, while bifidobacteria species have enzymatic activity able to degrade specific α - and β -bonds (Garrido et al., 2013). These bacteria such as *Bifidobacterium bifidum* subsp. *infantis* or *Bifidobacterium bifidum* are mainly found in the feces of breast-fed children. Indeed, the fecal microbiota of breast-fed child contain more than 75% bifidobacteria (Di Gioia et al., 2014). In addition to facilitating the growth of beneficial bacteria such as bifidobacteria, HMO can bind with pathogenic virus or bacteria, limiting adhesion to intestinal epithelium and therefore decreasing pathogens colonization by promoting natural excretion (Smilowitz et al., 2014). Also, other metabolites produced by lactic acid bacteria and bifidobacteria are able to inhibit virulence gene expression of *Escherichia coli* O157:H7 (Medellin-Pena and Griffiths, 2009; Zeinhom et al., 2012), *Salmonella enterica* serovar Typhimurium SA 941256 (Medellin-Pena et al., 2007; Bayoumi and Griffiths, 2012; S. Typhimurium) and *Campylobacter jejuni* (Mundi et al., 2013).

Five monosaccharides can be found in different HMO structures: glucose (Glc), galactose (Gal), N-acetylglucosamine (GlcNAc), fucose (Fuc), and sialic acid, also called N-acetylneuraminic (NeuAc). High quantities of lacto-N-biose type I (LNB: Gal β 1-3GlcNAc) and fucosylated HMO are an important characteristic of human milk (Chichlowski et al., 2011; Garrido et al., 2013; Dotz et al., 2014; Smilowitz et al., 2014). *B. infantis* is able to fully degrade HMO intracellularly, contrary to *B. longum*, *B. breve*, and *B. bifidum* (Sela and Mills, 2010; Underwood et al., 2015). It possesses genes encoding specific transporters and four enzymes necessary for HMO degradation (α -fucosidase, α -sialidase, β -galactosidase, and β -N-hexosaminidase) (Sela et al.,

2008; Sela, 2011; Smilowitz et al., 2014). HMO degradation by *B. bifidum* occurs outside the cells. Indeed, *B. bifidum* possesses a lacto-N-biosidase, which cleaves LNB from HMO. LNB is internalized using a specific transport system and is then degraded using LNB-phosphorylase (Sela, 2011; Smilowitz et al., 2014). Therefore, *B. infantis* and *B. bifidum*, despite their different gene clusters, both grow very well in the presence of HMO as sole source of carbon (Asakuma et al., 2011; Barile and Rastall, 2013).

Common oligosaccharides used in infant formula are galacto-oligosaccharides (GOS) and fructo-oligosaccharides (FOS), including inulin. GOS are composed of Glc and Gal, while FOS are composed of fructose and Glc. Their structures are very simple and linear. They are also bifidogenic, but because of their simple structure, they can also be consumed by other members of the intestinal microbiota such as *Bacteroides* spp. or *Clostridium* spp. (Chichlowski et al., 2011; Scholtens, 2014). This is probably why the fecal microbiota of formula-fed children contains only 30% bifidobacteria (Di Gioia et al., 2014) and the species that are present are different from those observed in the feces of breast-fed children. The predominant bifidobacteria in formula-fed children are those encountered in adult feces, such as *Bifidobacterium longum* subsp. *longum* and *Bifidobacterium adolescentis*, which present a less diverse enzymatic arsenal (De Vuyst et al., 2013).

Bovine milk oligosaccharides (BMO) can have similar composition and branching as HMO, so they could share some common properties. In addition, *B. infantis* or *B. bifidum* can grow in the presence of these carbohydrates (Sela, 2011; Milani et al., 2014). More than 60 BMO have been identified so far (Pacheco et al., 2015) and whey, a by-product of the dairy industry, is an important low cost source of BMO (Barile et al., 2009; Zivkovic and Barile, 2011). Furthermore, *Lactobacillus acidophilus* La-5 grown in medium supplemented with dairy ingredients such as whey, presented a protective effect in EHEC-infected mice (Zeinhom et al., 2012). However, BMO concentration in bovine milk is lower than HMO concentration in human milk (Barile et al., 2009; Tao et al., 2009; Kelly et al., 2013). Also, fucosylation occurs at very low frequency compared to sialylation, which is contrary to human milk (Tao et al., 2008). Even if the degree of polymerization in BMO is lower than in HMO, they are also protected by α - and β -bonds which are less accessible to other bacteria (Chichlowski et al., 2011). One of the most important BMO found in cow milk is 3'-sialyllactose (3'SL: NeuAc α 2-3Gal β 1-4Glc) and its concentration in colostrum can reach 0.85 mg/ml (Nakamura et al., 2003; Urashima et al., 2013). The majority of bifidobacteria grows only in anaerobic conditions, an environment very difficult to reproduce on an industrial scale. In addition, they have to survive

Abbreviations: 3'SL, 3'-sialyllactose; AE, attaching and effacing; BHI, brain heart infusion; BMO, bovine milk oligosaccharide; CFSM, cell-free spent medium; EHEC, enterohaemorrhagic *Escherichia coli*; FOS, fructo-oligosaccharide; Fuc, fucose; Gal, galactose; Glc, glucose; GlcNAc, N-acetylglucosamine; GOS, galacto-oligosaccharide; HMO, human milk oligosaccharide; LacNAc, N-acetyllactosamine; LB, Luria Bertani; LNB, lacto-N-biose; LNT, lacto-N-neotetraose; LNT, lacto-N-tetraose; MRS, De Man, Rogosa and Sharpe medium; NeuAc, N-acetylneuraminic acid or sialic acid; NeuGc, N-glycolylneuraminic acid; OD, optical density; T3SS, type III secretion system.

et al., 2003). *B. bifidum* BBA1 and *B. crudilactis* FR/62/B/3 were grown in three independent experiments under the same anaerobic conditions as previously at 37°C for 48 h. Five log/ml of bifidobacteria from a fresh 48 h culture of bifidobacteria were inoculated into the fresh media (1% v/v). The concentration of 5 log/ml was confirmed by plating several dilutions of bifidobacteria at day 0 post inoculation. Bacterial growth was determined by viable counts after 48 h incubation. Cell free spent media (CFSM) were obtained after two centrifugation steps at 5000 x (Eppendorf Centrifuge 5804, Hamburg, Germany) for 10 min. Supernatants were then sterilized by filtration (Minisart® 0.45 µm and 0.2 µm, Sartorius, Vilvoorde, Belgium). Next, CFSM were freeze-dried (Virtis Benchtop 3.3 EL, SP Scientific, Suffolk, United-Kingdom) and rehydrated with sterile distilled water to obtain a 10x concentration. The same treatment was applied to non-fermented culture media (controls). The pH of rehydrated CFSM was adjusted to 7 using 1 M NaOH.

Measurement of LEE and *hilA* Promoter Activity

Both *E. coli* reporter strains were grown overnight in BHI broth supplemented with ampicillin (BHI-Amp). Each overnight culture was diluted 1:100 with fresh BHI-Amp broth supplemented (test samples) or not (control samples) with 10% concentrated CFSM obtained from fermented MRS2-3'SL. Two hundred microliters of each sample were distributed into triplicate wells of a sterile, opaque 96-well microliter plate (Corning 3610, Fisher Scientific, Ottawa, Ontario, Canada) and incubated at 30°C. Luminescence was measured every hour for 24 h, with a Victor multilabel counter (Wallac, PerkinElmer Life Sciences Canada, Woodbridge, Ontario, Canada). Luminescence was expressed in counts per second. Optical density (OD) was determined using a Genesys 20 spectrophotometer (Thermo scientific, Erembodegem, Belgium) adjusted to 600 nm.

Contact between Concentrated CFSM and Wild Type Pathogenic Strains

E. coli O157:H7 ATCC 43890 was grown in LB agar and a single colony was taken from the plate and incubated overnight in LB broth at 37°C with aeration. The same procedure was applied for *S. Typhimurium* ATCC 14028 using BHI medium. The cultures were homogenized and 50 µl were diluted in 4.5 ml of LB broth for *E. coli* and BHI broth for *S. Typhimurium*. Then, 450 µl of each concentrated CFSM was added to the bacterial suspensions. Triplicate cultures were incubated at 37°C for 4 h on a shaker at 150 rpm. Bacterial growth was determined by OD measurement at 600 nm. *E. coli* O157:H7 and *S. Typhimurium* were grown in LB and BHI broth alone, respectively, as controls.

Gene Expression Analysis by RT-qPCR

The method was adapted with some modifications from Tellez et al. (2012), Mith et al. (2014) and Guri et al. (2016). After 4 h of incubation (Delcenserie et al., 2012), cells were collected by centrifugation at 5000 × g for 10 min at room temperature (Eppendorf Centrifuge 5804, Hamburg, Germany). The pellet was suspended in 100 µl TE buffer (10 mM Tris and 1 mM EDTA) containing 1% lysozyme (Roche, Mannheim, Germany).

Samples were stored at −20°C overnight. On the next day, RNA was extracted using the RNeasy® Mini Kit (Qiagen, Antwerp, Belgium). DNA contamination was eliminated from each sample using the DNase I Recombinant RNase-free Kit (Roche Diagnostics GmbH, Mannheim, Germany). To inactivate the DNase, samples were heated at 75°C for 10 min. The quantity of RNA was determined by measuring the absorbance at 260 nm using a Nanodrop 2000 Spectrophotometer (Thermo Scientific, USA). The purity and quality of RNA were verified by measuring the ratio of absorbance (260 nm/280 nm) and by using agarose gel electrophoresis (Eurogentec, Seraing, Belgium). The concentration of RNA used for reverse transcription was normalized to 100 ng/µl for *E. coli* and to 50 ng/µl for *S. Typhimurium*. Next, the RNA was subjected to reverse transcription polymerase chain reaction (RT-PCR) using a high-capacity cDNA Reverse Transcription Kit (Applied Biosystems, Ghent, Belgium). Briefly, 1 µg of RNA was reverse transcribed with 0.8 µl of desoxyribonucleoside triphosphate (dNTP; 100 mM), 1 µl of Multiscribe® Reverse Transcriptase (50 U/µl), 2 µl of 10X RT Random Primers and 2 µl of 10X RT Buffer in an adjusted total volume of 20 µl. For each sample, a no-RT control was included to confirm the absence of DNA contamination. Synthesis of cDNA was performed in a Mastercycler Gradient Thermocycler (Flexigene, Cambridge, UK) under the following conditions: 25°C for 10 min, 37°C for 120 min, 85°C for 5 min and a cooling step at 4°C. Then, cDNA was stored at −20°C.

To study the effects of bioactive molecules present in culture supernatant on gene expression of *E. coli* O157:H7 ATCC 43890, the expression of genes *ler* (involved in attaching effacing lesions), *fliC* (involved in mobility), *stxB2* (encoding subunit B of Shiga-toxin 2), *luxS* (major regulator of quorum sensing and producing AI-2), and *qseA* (involved in quorum sensing and regulator of LEE expression) was determined using qPCR. The same method was used to study effects on *S. Typhimurium* virulence gene expression of *hilA* (invasion protein regulator), *ssrB2* (Type III secretion system regulator), and *sopD* (secreted effector protein). Quantitative PCR amplification was conducted using the GoTaq® qPCR Master Mix (Promega, Leiden, Netherlands) and using the ABI 7300 Real Time PCR System (Applied Biosystems, Singapore) for *E. coli* or the Light Cycler 480 (Roche Diagnostics, Mannheim, Germany) for *S. Typhimurium*. The primers were synthesized by Eurogentec (Liège, Belgium) and have been used in previous studies (Table 2). The RT-qPCR was performed in a total volume of 20 µl, containing 10 µl of GoTaq® Master Mix, 5.75 µl of molecular grade water, 1 µl of forward primer (10 µM), 1 µl of reverse primer (10 µM), 0.25 µl of carboxy-X-rhodamine (30 µM), and 2 µl of diluted cDNA. The qPCR conditions for *E. coli* were: initial denaturation at 95°C for 3 min; denaturation, annealing and elongation repeated 45 times: 95°C for 15 s, 58°C for 30 s and 72°C for 45 s; melting curve program: 60–95°C with a heating rate of 0.1°C/s. The qPCR conditions for *S. Typhimurium* were: denaturation at 95°C for 10 min; 40 cycles of amplification and quantification: 95°C for 30 s, 56°C for 30 s and 72°C for 30 s; melting curve program: 60–95°C with a heating rate of 0.1°C/s. The annealing temperature, optimized at 56°C, was determined experimentally. Each specific amplicon was validated for the

presence of a single melting temperature peak and a single band of expected size on a 2% agarose gel after electrophoresis. Cycle threshold (C_t) values were determined using the ABI 7300 System SDS Software for *E. coli* and the Light Cycler Software 480 version 1.5 for *S. Typhimurium*. Four housekeeping genes were tested for *E. coli*: *gnd* (6-phosphogluconate dehydrogenase), *gst* (glutathione S-transferase), 16S gene (ribosomal RNA) and *recA* (recombinase A). Three housekeeping genes were tested for *S. Typhimurium*: *gmk* (guanylate kinase), *rpoD* (sigma factor) and 16S gene (ribosomal RNA gene). Because *recA* and *gmk* were the most stable under different treatments, they were selected for normalizing transcript expression levels. The experiments were replicated three independent times. To determine relative changes in gene expression, the formula described by Pfaffl (Pfaffl, 2001) was used: ratio = $(E_{\text{target}})^{\Delta C_t \text{ target}(\text{control-sample})} / (E_{\text{reference}})^{\Delta C_t \text{ reference}(\text{control-sample})}$,

where E is the efficiency of the qPCR, calculated according to the equation: $E = 10^{(-1/\text{slope})}$.

Statistical Analysis

The data are means ± standard error of three replicates. A Student's *t*-test was used to assess the statistical significance of the differences between test and control groups, where $p \leq 0.05$ was considered as significant.

RESULTS

Growth of *Bifidobacterium bifidum* and *Bifidobacterium crudilactis*

B. crudilactis FR/62/B/3 showed increase in viable counts in MRS2, MRS2-Wh and MRS2-3'SL compared to MRS2 G, but the highest counts were observed on MRS2-Wh (8.9 ± 0.6 log cfu/ml, Table 3). The same trend was observed for *B. bifidum* BBA1 with slightly lower counts (Table 3) compared to *B. crudilactis* FR/62/B/3. The highest counts were also observed for MRS2-Wh (8.1 ± 0.3 log cfu/ml).

Activity of CFSM from MRS2-3'SL Fermented by Bifidobacteria on Bioluminescent Reporter Gene Expression

Luminescence expression of the plasmids LEE::*luxCDABE* and *hila::luxCDABE* reached its maximum at 4 h for the *E. coli* mutant and at 13 h for the *S. Typhimurium* mutant, respectively. In the presence of supernatants from fermented MRS2-3'SL medium, bioluminescence induction decreased for both mutants (Figures 1, 2) showing a decrease in promoter expression of *ler* and *hila*. These results, statistically significant for both strains, were more pronounced for *hila* gene expression of *S. Typhimurium* (Figure 2).

Effect of CFSM on *E. coli* O157:H7 Virulence Gene Expression

After incubation of 4 h, the different CFSM had no negative impact on growth. The average OD of *E. coli* O157:H7 at 600 nm after 4 h exposure was around 1.286 ± 0.119 . From every tested housekeeping gene, *recA* was the most stable and was chosen to normalize the results according to the efficiency of each pair of primers (virulence genes *ler*, *fliC*, *stxB2*, *luxS*, and *qseA*) monitored using qPCR (Pfaffl, 2001; Tellez et al., 2012) (Table 4). The fermented MRS2 G, MRS2, and MRS2-Wh media did not show significant impact on gene expression

TABLE 2 | Primers of virulence genes used for qPCR (F: forward; R: reverse).

| Primer | Sequence of PCR primers (5'-3') ^a | References |
|---|--|----------------------------|
| <i>E. coli</i> Housekeeping and Virulence Genes | | |
| <i>gnd</i> | F: 5'-GGTAATACCTTCTCCAGGACACC-3' | Rashid et al., 2006 |
| | R: 5'-TAGTGCGCCCTCCTCACC-3' | |
| <i>gst</i> | F: 5'-CTTTGCCGTTAAACCCTAAGGG-3' | Pfaffl, 2001 |
| | R: 5'-GCTGCAATGTGCTCTAACCC-3' | |
| <i>recA</i> | F: 5'-CAATATCCCCACTGCTGCC-3' | Takle et al., 2007 |
| | R: 5'-CACCTAGGCGACGATCCCT-3' | |
| 16S | F: 5'-GGTGAGCTGGTTGATCTGGG-3' | Takle et al., 2007 |
| | R: 5'-GCATTGCTTTACCTGACC-3' | |
| <i>ler</i> | F: 5'-TTTCTTTCAGTGTCTTCA-3' | Medellin-Pena et al., 2007 |
| | R: 5'-TGCGGAGATTATTATTATGA-3' | |
| <i>fliC</i> | F: 5'-TACCATCGAAAAGCAACTCC-3' | Medellin-Pena et al., 2007 |
| | R: 5'-GTCGGCAACGTTAGTGATACC-3' | |
| <i>luxS</i> | F: 5'-GATCATACCCGGATGGAAG-3' | Medellin-Pena et al., 2007 |
| | R: 5'-AGAATGCTACGCGCAATATC-3' | |
| <i>stxB2</i> | F: 5'-AGATGTTTATGGCGTTTGA-3' | Medellin-Pena et al., 2007 |
| | R: 5'-TTAACTGCACCTCAGCAAA-3' | |
| <i>qseA</i> | F: 5'-CGCGGATCCCCTTGGCACAGTTTGTACA-3' | Medellin-Pena et al., 2007 |
| | R: 5'-CGCGGATCCCCTTGGCACAGTTTGTACA-3' | |
| <i>S. Typhimurium</i> Housekeeping and Virulence Genes | | |
| <i>gmk</i> | F: 5'-TTGGCAGGGAGGCGTTT-3' | Rashid et al., 2006 |
| | R: 5'-GCGCGAAGTGCCGTAGTAAT-3' | |
| <i>rpoD</i> | F: 5'-ACATGGGTATTACAGTAATGGAAGA-3' | Botteldoorn et al., 2006 |
| | R: 5'-CGGTGGGTATTACAGTAATGGAAGA-3' | |
| 16S | F: 5'-AGGCCTTCGGGTTGTAAGT-3' | Xu et al., 2010 |
| | R: 5'-GTTAGCCGGTGCTTCTCTG-3' | |
| <i>hila</i> | F: 5'-TGTCGGAAGATAAAGAGCAT-3' | Guri et al., 2016 |
| | R: 5'-AAGGAAGTATCGCCAATGTA-3' | |
| <i>sopD</i> | F: 5'-ATTAATGCCGGTAACTTTGA-3' | Guri et al., 2016 |
| | R: 5'-CTCTGAAAACGGTGAATAGC-3' | |
| <i>ssrB2</i> | F: 5'-TGGTTTACACAGCATACCAA-3' | Guri et al., 2016 |
| | R: 5'-GGTCAATGTAACGCTTGT-3' | |

TABLE 3 | Counts of *B. bifidum* and *B. crudilactis* after 48 h of incubation in MRS2 G, MRS2, MRS2-Wh, and MRS2-3'SL media.

| | Final concentrations after 48 h incubation (log cfu/ml) | |
|-----------|---|-----------------------|
| | <i>B. bifidum</i> | <i>B. crudilactis</i> |
| MRS2 G | 6.9 ± 0.3 | 5.5 ± 0.5 |
| MRS2 | 7.3 ± 0.8 | 7.8 ± 1.4 |
| MRS2-Wh | 8.1 ± 0.3 | 8.9 ± 0.6 |
| MRS2-3'SL | 6.8 ± 0.9 | 7.9 ± 0.3 |




FIGURE 1 | Effect of CFMS from MRS2-3'SL medium fermented with *B. bifidum* and *B. crudilactis* on *E. coli* O157:H7 (*stx*⁻, *LEE1:lux*) ATCC 43888 expression. The *E. coli* control is *E. coli* grown on BHI only. The MRS2-3'SL control is *E. coli* grown on BHI and CFMS from MRS2-3'SL medium unfermented. Data are the means ± the standard deviations derived from triplicate and expressed as relative light units (RLU) defined as counts per seconds, adjusted to OD600 (RLU/OD600) and where OD is fixed at 0.806 and was taken after 4 h of incubation. LUM: luminescence; OD: optical density. ****P* ≤ 0.005.




FIGURE 2 | Effect of CFMS from MRS2-3'SL medium fermented with *B. bifidum* and *B. crudilactis* on *S. Typhimurium* (*hilA:lux*) SA 941256 expression. The *Salmonella* control is *Salmonella* grown on BHI only. The MRS2-3'SL control is *S. Typhimurium* grown on BHI and CFMS from MRS2-3'SL medium unfermented. Data are the means ± the standard deviations derived from triplicate and expressed as relative light units (RLU) defined as counts per seconds, adjusted to OD600 (RLU/OD600) and where OD is fixed at 0.909 and was taken after 13 h of incubation. LUM: luminescence; OD: optical density. ****P* ≤ 0.005.

(Figures 3A–C) compared to non-fermented control media, meaning that CFMS had no effects. However, significant down-regulation of virulence genes of *E. coli* O157:H7 was observed in the presence of fermented MRS2-3'SL medium (Figure 3D). The medium fermented by *B. bifidum* BBA1 and *B. crudilactis* FR/62/B/3 induced a down-regulation of the *E. coli* *ler* gene (ratios of −15.4; *P* ≤ 0.01 and −8.1; *P* ≤ 0.05, respectively). A down-regulation of the *qseA* gene was also observed (ratios of −2.1; *P* ≤ 0.01 and −3.1; *P* ≤ 0.05, respectively). A non-significant trend for up-regulation of the *fliC* gene (ratios of 25.8 and +20.8, respectively) was noted while a non-significant trend for down-regulation of the *stxB2* gene (ratios of −4.6 and −4.2, respectively) was observed. In the case of fermentation by *B. crudilactis* FR/62/B/3, a slight non-significant trend for down-regulation of the *luxS* gene was observed (ratio of −2.1).

Effect of CFMS on *S. Typhimurium* Virulence Gene Expression

After incubation of 4 h, the OD measurements showed no negative impact on growth. The average OD of *S. Typhimurium* at 600 nm and after 4 h of exposure was 0.862 ± 0.078. From every tested housekeeping gene, *gmk* was the most stable and was chosen to normalize and adjust the results according to the efficiency of each pair of primers (virulence genes *hilA*, *ssrB2*, and *sopD*) monitored using qPCR (Guri et al., 2016, Table 5). The CFMS of MRS2-3'SL medium fermented by *B. bifidum* induced a slight down-regulation of the *ssrB2* gene (−2.1; *P* ≤ 0.05). The same trend was observed for the genes *hilA* and *sopD* (−2.5 and −1.9, respectively). MRS2-3'SL CFMS fermented by *B. crudilactis* FR/62/B/3 did not show any significant effect on virulence gene expression (Figure 4B). A significant increase of *sopD* expression is observed but too light to be biologically meaningful (1.1; *P* ≤

TABLE 4 | Effect of CFSM on expression (cycle threshold values ± standard error) of virulence genes of enterohaemorrhagic *E. coli* O157:H7 after 4 h incubation.

| Gene | PCR efficiency ^a (%) | LB control ^b | Test supernatants from unfermented media ^c | | | CFSM from fermented media by <i>B. bifidum</i> ^d | | | CFSM from fermented media by <i>B. crudilactis</i> ^e | | | | | |
|--------------|---------------------------------|-------------------------|---|------------|------------|---|------------|------------|---|------------|------------|------------|------------|------------|
| | | | MRS2 G | MRS2 | MRS2-Wh | MRS2 G | MRS2 | MRS2-Wh | MRS2 G | MRS2 | MRS2-Wh | MRS2-3'SL | | |
| <i>recA</i> | 100 | 20 | 21.3 ± 0.6 | 18.2 ± 0.5 | 21.0 ± 0.7 | 20.3 ± 0.3 | 21.2 ± 0.4 | 21.4 ± 1.2 | 23.6 ± 1.9 | 19.9 ± 0.9 | 20.9 ± 0.5 | 23.7 ± 0.2 | 21.7 ± 0.3 | 19.9 ± 1.0 |
| <i>ler</i> | 93 | 18.9 | 25.2 ± 0.4 | 20.3 ± 1.3 | 26.4 ± 1.6 | 19.9 ± 0.8 | 25.6 ± 1.2 | 25.4 ± 0.3 | 28.4 ± 1.6 | 23.8 ± 0.4 | 24.2 ± 1.4 | 25.6 ± 1.0 | 27.8 ± 2.0 | 22.8 ± 2.0 |
| <i>flhC</i> | 97 | 20.8 | 18.2 ± 4.0 | 22.8 ± 1.4 | 24.1 ± 1.1 | 23.4 ± 2.0 | 19.1 ± 0.3 | 24.9 ± 4.2 | 26.0 ± 4.9 | 18.7 ± 1.5 | 19.8 ± 1.6 | 21.7 ± 3.7 | 25.3 ± 3.1 | 19.3 ± 0.6 |
| <i>luxS</i> | 91 | 23.6 | 26.8 ± 0.9 | 20.7 ± 1.6 | 30.1 ± 1.9 | 25.0 ± 1.3 | 27.7 ± 1.0 | 28.3 ± 0.9 | 32.6 ± 4.7 | 25.4 ± 1.5 | 27.4 ± 0.3 | 26.8 ± 3.1 | 29.8 ± 3.3 | 25.8 ± 2.2 |
| <i>stxB2</i> | 95 | 26.1 | 28.8 ± 1.0 | 25.7 ± 1.0 | 33.4 ± 2.4 | 26.5 ± 2.6 | 29.3 ± 3.0 | 30.4 ± 1.5 | 35.3 ± 1.0 | 27.6 ± 0.5 | 29.7 ± 2.5 | 30.1 ± 2.3 | 33.8 ± 3.8 | 27.8 ± 2.0 |
| <i>qseA</i> | 91 | 22.5 | 24.1 ± 0.4 | 26.3 ± 0.4 | 24.4 ± 0.8 | 21.9 ± 0.5 | 24.7 ± 1.4 | 25.3 ± 2.0 | 27.0 ± 2.1 | 22.7 ± 1.2 | 24.1 ± 0.9 | 23.5 ± 3.0 | 25.2 ± 1.0 | 23.3 ± 1.0 |

^aPCR efficiency: $E = [(10^{-(1/\text{slope})})/2] \times 100\%$.

^b*E. coli* O157:H7 grown in LB broth for 4 h.

^c*E. coli* O157:H7 grown in LB broth supplemented with unfermented culture media CFSM for 4 h.

^d*E. coli* O157:H7 grown in LB broth supplemented with fermented concentrated culture media CFSM from *B. bifidum* for 4 h.


^e*E. coli* O157:H7 grown in LB broth supplemented with fermented concentrated culture media CFSM from *B. crudilactis* for 4 h.

0.05), and the same but non-significant trend is observed with *hila* gene (1.6). Interestingly, a down-regulation of virulence genes was observed with CFSM obtained from MRS2 without glucose. In the case of fermentation by *B. crudilactis* FR/62/B/3, the genes *hila*, *ssrB2* and *sopD* were down regulated (−8.3, −10.9, and −6.2, respectively; $P \leq 0.05$). The same but non-significant trend was observed for *B. bifidum* BBA1 (−8.5, −8.0, and −2.6, respectively, **Figure 4A**).

DISCUSSION

B. crudilactis FR/62/B/3 presented the best growth potential compared to *B. bifidum* BBA1, particularly with whey or 3'SL instead of glucose. This could be explained by the fact that *B. crudilactis* FR/62/B/3 was originally isolated from raw cow milk and raw milk cheese. This species possesses the genetic machinery suggesting some ability to synthesize specific enzymes for BMO degradation, as highlighted by the presence of genes coding for β-galactosidase and α- or β-glucosidase, genes also present in other bacteria such as *B. bifidum*, *B. longum* subsp. *infantis*, *B. mongoliense*, *B. biavatii*, *B. kashiwanohense*, and *B. stellenboschense* (Delcenserie et al., 2007; Milani et al., 2014, 2015; Bondue and Delcenserie, 2015). *B. bifidum* BBA1 grew as well on medium containing 3'SL as the main carbohydrate source. This species possesses genes encoding some of the enzymes cleaving BMO bonds, so their expression would lead to growth. However, the growth results of bifidobacteria on this media were similar to those observed with MRS2 G (control). This could mean that those genes may not be expressed efficiently under the conditions tested, or another function is necessary. A next step would be to measure the expression of genes involved in carbohydrate metabolism such as genes coding for β-galactosidase and α- or β-glucosidase to validate the expression of those genes in the presence of BMO. Another hypothesis could be attributed to the presence of residual glucose from MRS culture allowing bifidobacteria to grow in medium exempt of carbohydrate. Indeed, the glucose present in initial MRS medium was in excess (Tanimomo et al., 2016), but a negligible part of it (maximum 1 mg) has been transferred with the inoculum. Another source of glucose could be the presence of residual carbohydrate in the yeast extracts or peptone extracts used in MRS media. The positive effects of media supplemented with milk products on growth of probiotics has been demonstrated previously (Champagne et al., 2014). This is confirmed in the present study as the best levels of growth were reached on MRS2-Wh media for both strains studied. In addition to BMO, whey is rich in lactose (Food and Agriculture Organization/Organisation Mondiale de la Santé, 1998), a carbohydrate source that is easily consumed by bifidobacteria (Delcenserie et al., 2007).

As demonstrated previously, the CFSM obtained from some lactic acid bacteria or bifidobacteria can induce a decrease in virulence gene expression of pathogenic bacteria such as *C. jejuni* (Mundi et al., 2013). Likewise, *B. bifidum* ATCC 29521, and *L. acidophilus* La-5 CFSM were able to produce metabolites inhibiting virulence gene expression of *E. coli* O157:H7 (Medellin-Pena et al., 2007) and *S. Typhimurium*



(Bayoumi and Griffiths, 2012). In our study, CFSM collected from MRS2-3'SL medium fermented by *B. bifidum* BBA1 and *B. crudilactis* FR/62/B/3 down-regulated most of the virulence genes tested in *E. coli* O157:H7, except the *fliC* gene, which tended to be up-regulated. This is not surprising according to the fact that *fliC* gene is not coded by the LEE operon and therefore not necessarily regulated as other virulence genes involved in T3SS and situated within the LEE operon (Falcao et al., 2004). No significant effect has been observed with CFSM from MRS2 and MRS2-Wh medium. In addition, CFSM obtained from bifidobacteria grown in media enriched in 3'SL were able to affect virulence gene expression of *E. coli* O157:H7 without having any impact on its growth, at least during the first 4 h of incubation.

In those media, higher in carbohydrates, more fermentation products such as lactate or acetate are synthesized and could have an inhibiting effect on pathogenic bacteria, as well as acidifying the media. However, all CFSM were neutralized before testing them against *E. coli* or *Salmonella*, meaning that the pH did not exert any effect on *E. coli* O157:H7 growth. Furthermore, under neutral pH, the organic acids were under

dissociated form and should not present any bactericidal or bacteriostatic action, contrary to un-dissociated forms (Momose and Hirayama, 2008).

The genes involved in virulence expression such as *ler* but also *fliC* genes are regulated by *luxS*, involved in quorum sensing. However, nutrients can interfere with quorum sensing mechanisms (Henke and Bassler, 2004; Kaper et al., 2004; Nakanishi et al., 2006; Mellies et al., 2007) and induce a decrease in virulence gene expression through a decrease in *luxS* expression. Delcenserie et al. (2012) previously demonstrated the effects of glucose in down-regulating virulence gene expression of *E. coli* O157:H7. The present study brought out similar observations with lactose instead of glucose (data not shown). The *ler* gene was the most affected by the presence of those carbohydrates and the effect was dose-dependent. Media used as controls and containing glucose or lactose (MRS2 and MRS2-Wh) down-regulated this gene but no effect was observed with medium containing mainly 3'SL as a source of carbohydrate (MRS2-3'SL).

To be able to metabolize 3'SL, *B. bifidum* and *B. crudilactis* have to secrete sialidases through which NeuAc (sialic acid) can

TABLE 5 | Effect of CFMSM on expression (cycle threshold values ± standard error) of virulence gene expression of *S. Typhimurium* after 4 h incubation.

| Gene | PCR efficiency ^a (%) | BHI control ^b | Test supernatants from unfermented media ^c | | CFMSM from fermented media by <i>B. bifidum</i> ^d | | CFMSM from fermented media by <i>B. crudilactis</i> ^e | |
|--------------|---------------------------------|--------------------------|---|------------|--|------------|--|------------|
| | | | MRS2 G | MRS2-3'SL | MRS2 G | MRS2-3'SL | MRS2 G | MRS2-3'SL |
| <i>gmk</i> | 100 | 23 | 23.3 ± 0.3 | 24.4 ± 1.2 | 23.4 ± 0.4 | 24.4 ± 1.6 | 21.8 ± 0.2 | 25.0 ± 1.6 |
| <i>hilA</i> | 91 | 31.5 | 28.6 ± 0.2 | 31.2 ± 0.7 | 31.4 ± 2.8 | 32.5 ± 1.2 | 30.2 ± 0.4 | 31.3 ± 2.0 |
| <i>ssrB2</i> | 115 | 31.2 | 28.7 ± 1.1 | 30.8 ± 0.1 | 30.9 ± 1.6 | 31.6 ± 1.6 | 30.2 ± 0.6 | 32 ± 2.0 |
| <i>sopD</i> | 91 | 30.1 | 27.7 ± 0.5 | 30.2 ± 1.0 | 29.3 ± 1.6 | 31.2 ± 1.5 | 28.9 ± 0.6 | 30.9 ± 2.1 |


^aPCR efficiency: $E = [(10^{-1/\text{slope}})/2] \times 100\%$.

^b*S. Typhimurium* grown in BHI broth for 4 h.

^c*S. Typhimurium* grown in BHI broth supplemented with concentrated supernatants from culture media unfermented for 4 h.

^d*S. Typhimurium* grown in BHI broth supplemented with concentrated supernatants from culture media fermented *B. bifidum* for 4 h.

^e*S. Typhimurium* grown in BHI broth supplemented with concentrated supernatants from culture media fermented *B. crudilactis* for 4 h.



be produced. *B. bifidum* does not use this sialic acid, which is available for other bacteria such as *B. breve* (Milani et al., 2015). Therefore, if 3'SL is metabolized by *B. bifidum*, free sialic acid was probably present in fermented CFMSM from 3'SL medium. Usually, pathogenic bacteria are able to bind this free sialic acid to their cell surface and use it to improve their resistance to the host's innate immune response, or can consume it as a nutrient (Vimr et al., 2004; Severi et al., 2007). NeuAc also exerts a major role in *Salmonella enterica* subsp. *enterica* serovar Typhi adhesion to intestinal epithelium (Sakarya et al., 2010). This means that in theory, sialic acid could have an impact on *S. Typhimurium* and *E. coli* growth, but no effect on growth as measured by OD has been observed. In addition, NeuAc had probably no impact on virulence gene expression in our study, when supplied in the medium.

The non-significant trend for up-regulation of *fliC* observed in our study should be clarified. Indeed, this trend seems higher when *E. coli* O157:H7 was exposed to CFMSM from fermented 3'SL. A hypothesis could be that the presence of residual complex carbohydrates affects gene expression of *fliC*. The presence or absence of some nutrients, or stress, could play a role in virulence expression, including *fliC* (Mei et al., 2015). Several studies

investigated the effects of some stress (oxidative stress, heat shock, long storage) on down-regulation of *fliC* gene while other virulence genes were upregulated (Carey et al., 2009; Mei et al., 2015; Singh and Jiang, 2015). The experimental protocol of this study submitted *E. coli* O157:H7 to some stress, which may influence virulence gene expression. Genes involved in general stress (*uspA* and *rpoS*), in starvation (*phoA* and *dpS*), in cold shock (*cspA*, *cspC*, and *cspE*) and in acid resistance (*gadW*) have been investigated and the results have shown that the different treatments did not affect expression of stress-related genes (data not shown). This leads us to suppose that virulence factor expression has not been influenced by the experimental conditions.

The results observed using RT-qPCR with CFMSM from fermented MRS2-3'SL on *S. Typhimurium* virulence gene expression did not confirm the results observed with luminescent reporter strains. However, CFMSM obtained after fermentation of MRS2 without any glucose fermented by *B. crudilactis* FR/62/B/3 down-regulated several virulence genes. Regarding *B. bifidum* BBA1 CFMSM, a trend to down-regulation was observed as well. These down-regulations could be caused by other non-carbohydrate metabolites produced by bifidobacteria

and these bioactive molecules could originate from the degradation of proteins. Most known bioactive molecules, such as subpeptin JM4-A and subpeptin JM4-B are antimicrobial peptides synthesized by *Bacillus subtilis* and active against *Salmonella*, *Staphylococcus aureus*, and *Bacillus cereus* (Sumi et al., 2015). Nisin, a bacteriocin well known, is produced by *Lactococcus lactis* and has a negative impact on *Listeria* or *Clostridium* (Ebbensgaard et al., 2015). The results of our study suggest that the CFMS activity is not due to an antimicrobial effect but due to an antivirulent effect.

In conclusion, this study provides the information that CFMS obtained from MRS2-3'SL medium fermented by *B. bifidum* BBA1 and *B. crudilactis* FR/62/B/3 down-regulated LEE1 expression of the luminescent *E. coli* reporter strain and *hilA* expression of luminescent *S. Typhimurium* reporter strain. These results agree with the decreasing virulence gene expression of *ler* and *qseA* for *E. coli*, but not for *S. Typhimurium*. The contact between *S. Typhimurium* and CFMS from fermented MRS without glucose showed down-regulation of genes *hilA*, *ssrB2*, and *sopD*. According to this *in vitro* study, the antivirulent metabolites issuing from fermentation by bifidobacteria could have a negative impact on T3SS of both pathogens, decreasing expression of genes mainly implicated in this virulence mechanism (*ler* and *qseA* genes for *E. coli* O157:H7; *hilA*, *ssrB2*, and *sopD* genes for *S. Typhimurium*). The potential upregulation of *fliC* in *E. coli* O157:H7 could increase the motility as well as biofilm formation. A phenotypic analysis

of the pathogens under the experimental conditions could bring more insights about its virulence pattern. Information is lacking about the nature of the active molecules, but the activity of those CFMS might be due to small peptides or proteins with low molecular weight and resistant to pH modification and heat, or products obtained from carbohydrate metabolism. Size exclusion chromatography could contribute to separating and isolating these bioactive molecules in order to identify them. In the future, the effects of these metabolites will be investigated in a human gastrointestinal model to study the impact on microbiota to mimic *in vivo* conditions.

AUTHOR CONTRIBUTIONS

PB did the experiments, interpreted the results and wrote the manuscript. SC and FB participated to the experiments. MS, GD, GL, and MG were involved in the design of the study and provided help for interpretation of the results. VD participated to the design of the study, interpretation of the results and writing of the manuscript.

ACKNOWLEDGMENTS

This study was funded by the “Fond Opportunité” from Ulg. The author wishes to acknowledge Nicolas Korda, Emilie Arnould, staff from the Bruyères hospital (Liège, Belgium) and Herve society (Liège, Belgium) for their valuable help on this project.

REFERENCES

- Arrieta, M. C., Stiensma, L. T., Amenogbe, N., Brown, E. M., and Finlay, B. (2014). The intestinal microbiome in early life: health and disease. *Front. Immunol.* 5:427. doi: 10.3389/fimmu.2014.00427
- Asakuma, S., Hatakeyama, E., Urashima, T., Yoshida, E., Katayama, T., Yamamoto, K., et al. (2011). Physiology of the consumption of human milk oligosaccharides by infant-gut associated bifidobacteria. *J. Biol. Chem.* 40, 34583–34592. doi: 10.1074/jbc.M111.248138
- Barile, D., and Rastall, R. A. (2013). Human milk and related oligosaccharides as prebiotics. *Curr. Opin. Biotechnol.* 24, 214–219. doi: 10.1016/j.copbio.2013.01.008
- Barile, D., Tao, N., Lebrilla, C. B., Coisson, J. D., Arlorio, M., and German, J. B. (2009). Permeate from cheese whey ultrafiltration is a source of milk oligosaccharides. *Int. Dairy J.* 19, 524–530. doi: 10.1016/j.idairyj.2009.03.008
- Bayoumi, M. A., and Griffiths, M. W. (2010). Probiotics down-regulate genes in *Salmonella enterica* serovar Typhimurium pathogenicity islands 1 and 2. *J. Food Prot.* 73, 452–460.
- Bayoumi, M. A., and Griffiths, M. W. (2012). *In vitro* inhibition of expression of virulence genes responsible for colonization and systemic spread of enteric pathogens using *Bifidobacterium bifidum* secreted molecules. *Int. J. Food Microbiol.* 156, 255–263. doi: 10.1016/j.ijfoodmicro.2012.03.034
- Bondue, P., and Delcenserie, V. (2015). Genome of bifidobacteria and carbohydrate metabolism. *Korean J. Food Sci. Anim. Resour.* 35, 1–9. doi: 10.5851/kosfa.2015.35.1.1
- Botteldoorn, N., Van Coillie, E., Grijspeerd, K., Werbrouck, H., Haesebrouck, F., Donné, E., et al. (2006). Real-time reverse transcription PCR for the quantification of the *mntH* expression of *Salmonella enterica* as a function of growth phase and phagosome-like conditions. *J. Microbiol. Methods* 66, 125–135. doi: 10.1016/j.mimet.2005.11.003
- Carey, C. M., Kostrzynska, M., and Thompson, S. (2009). *Escherichia coli* O157:H7 stress and virulence gene expression on romaine lettuce using comparative real-time PCR. *J. Microbiol. Methods* 2, 235–242. doi: 10.1016/j.mimet.2009.02.010
- Champagne, C. P., Raymond, Y., Pouliot, Y., Gauthier, S. F., and Lessard, M. (2014). Effect of bovine colostrum, cheese whey, and spray-dried porcine plasma on the *in vitro* growth of probiotic bacteria and *Escherichia coli*. *Can. J. Microbiol.* 60, 287–295. doi: 10.1139/cjm-2014-0130
- Chichlowski, M., German, J. B., Lebrilla, C. B., and Mills, D. A. (2011). The influence of milk oligosaccharides on microbiota of infants: opportunities for formulas. *Annu. Rev. Food Sci. Technol.* 2, 331–351. doi: 10.1146/annurev-food-022510-133743
- Daube, G., Delcenserie, V., and Gavini, F. (2006). *Probiotic Bifidobacterial Species*. PCT/EP2006/061247, US 20080274085, WO 2006/122850, 31-03-2006. European Patent Office, European Union.
- Delcenserie, V., Gavini, F., Beerens, H., Tresse, O., Franssen, C., and Daube, G. (2007). Description of a new species, *Bifidobacterium crudilactis* sp. nov., isolated from raw milk and raw milk cheeses. *Syst. Appl. Microbiol.* 30, 381–389. doi: 10.1016/j.syapm.2007.01.004
- Delcenserie, V., Lapointe, G., Charaslertrangsri, T., Rabalski, A., and Griffiths, M. W. (2012). Glucose decreases virulence gene expression of *Escherichia coli* O17:H7. *J. Food Prot.* 75, 748–752. doi: 10.4315/0362-028X.JFP-11-384
- Delcenserie, V., Taminiau, B., Gavini, F., de Schaetzen, M. A., Cleenwerck, I., Theves, M., et al. (2013). Detection and characterization of *Bifidobacterium crudilactis* and *B. mongoliense* able to grow during the manufacturing process of French raw milk cheeses. *BMC Microbiol.* 13:239. doi: 10.1186/1471-2180-13-239
- De Vuyst, L., Moens, F., Selak, M., Riviere, A., and Leroy, F. (2013). Summer Meeting 2013: growth and physiology of bifidobacteria. *J. Appl. Microbiol.* 116, 477–491. doi: 10.1111/jam.12415
- Di Gioia, D., Aloisio, I., Mazzola, G., and Biavati, B. (2014). Bifidobacteria: their impact on gut microbiota composition and their applications as probiotics

- in infants. *Appl. Microbiol. Biotechnol.* 98, 563–577. doi: 10.1007/s00253-013-5405-9
- Dotz, V., Rudloff, S., Meyer, C., Lochnit, G., and Kunz, C. (2014). Metabolic fate of neutral human milk oligosaccharides in exclusively breastfed infants. *Mol. Nutr. Food Res.* 59, 355–364. doi: 10.1002/mnfr.201400160
- Ebbensgaard, A., Mordhorst, H., Overgaard, M. T., Nielsen, C. G., Aarestrup, F. M., and Hansen, E. B. (2015). Comparative evaluation of antimicrobial activity of different antimicrobial peptides against a range of pathogenic bacteria. *PLoS ONE* 10:e0144611. doi: 10.1371/journal.pone.0144611
- Falcao, J. P., Falcao, D. P., and Gomes, T. A. T. (2004). Ice as a vehicle for diarrheagenic *Escherichia coli*. *Int. J. Food Microbiol.* 91, 99–103. doi: 10.1016/S0168-1605(03)00327-1
- Food and Agriculture Organization/Organisation Mondiale de la Santé (FAO/OMS) (1998). *Le lait et les Produits Laitiers dans la Nutrition Humaine*. Available online at: <http://www.fao.org/docrep/t4280f/t4280f0h.htm> (Accessed June 16, 2016).
- Garrido, D., Dallas, D. C., and Mills, D. A. (2013). Consumption of human milk glycoconjugates by infant-associated bifidobacteria: mechanisms and implications. *Microbiology* 159, 649–664. doi: 10.1099/mic.0.064113-0
- Guri, A., Paligot, M., Crèvecoeur, S., Piedboeuf, B., Claes, J., Daube, G., et al. (2016). *In vitro* screening of mare's milk antimicrobial effect and antiproliferative activity. *FEMS Microbiol. Lett.* 363, 1–7. doi: 10.1093/femsle/fnv234
- Henke, J. B., and Bassler, B. (2004). Quorum sensing regulates type III secretion in *Vibrio Harveyi* and *Vibrio parahaemolyticus*. *J. Bacteriol.* 186, 3794–3805. doi: 10.1128/JB.186.12.3794-3805.2004
- Kaper, J. B., Nataro, J. P., and Mobley, H. L. (2004). Pathogenic *Escherichia coli*. *Nat. Rev. Microbiol.* 2, 123–140. doi: 10.1038/nrmicro818
- Kelly, V., Davis, S., Berry, S., Melis, J., Spelman, R., Snell, R., et al. (2013). Rapid, quantitative analysis of 3'- and 6'-sialyllactose in milk by flow-injection analysis-mass spectrometry: screening of milks for naturally elevated sialyllactose concentration. *J. Dairy Sci.* 96, 7684–7691. doi: 10.3168/jds.2013-6972
- Medellin-Pena, M. J., and Griffiths, M. W. (2009). Effect of molecules secreted by *Lactobacillus acidophilus* strain La-5 on *Escherichia coli* O157:H7 colonization. *Appl. Environ. Microbiol.* 75, 1165–1172. doi: 10.1128/AEM.01651-08
- Medellin-Pena, M. J., Wang, H., Johnson, R., Anand, S., and Griffiths, M. W. (2007). Probiotics affect virulence-related gene expression in *Escherichia coli* O157:H7. *Appl. Environ. Microbiol.* 73, 4259–4267. doi: 10.1128/AEM.00159-07
- Mei, G. Y., Tang, J., Carey, C., Bach, S., and Kostrzynska, M. (2015). The effect of oxidative stress on gene expression of Shiga toxin-producing *Escherichia coli* (STEC) O157:H7 and non-O157 serotypes. *Int. J. Food Microbiol.* 215, 7–15. doi: 10.1016/j.ijfoodmicro.2015.07.029
- Mellies, J. L., Barron, A. M. S., and Carmona, A. M. (2007). Enteropathogenic and enterohemorrhagic *Escherichia coli* virulence gene regulation. *Infect. Immun.* 75, 4199–4210. doi: 10.1128/IAI.01927-06
- Milani, C., Lugli, G. A., Duranti, S., Turrioni, F., Bottacini, F., Mangifesta, M., et al. (2014). Genomic encyclopedia of type strains of the genus bifidobacterium. *Appl. Environ. Microbiol.* 80, 6290–6302. doi: 10.1128/AEM.02308-14
- Milani, C., Lugli, G. A., Duranti, S., Turrioni, F., Mancabelli, L., Ferrario, C., et al. (2015). Bifidobacteria exhibit social behavior through carbohydrate resource sharing in the gut. *Sci. Rep.* 5:15782. doi: 10.1038/srep15782
- Mith, H., Clinquart, A., Zhiri, A., Daube, G., and Delcenserie, V. (2014). The impact of oregano (*Origanum heracleoticum*) essential oil and carvacrol on virulence gene transcription by *Escherichia coli* O157:H7. *FEMS Microbiol. Lett.* 362, 1–7. doi: 10.1093/femsle/fnu021
- Momose, Y., and Hirayama, K. (2008). Effect of organic acids on inhibition of *Escherichia coli* O157:H7 colonization in gnotobiotic mice associated with infant intestinal microbiota. *Anton. Leeuw. Int. J. G.* 93, 141–149. doi: 10.1007/s10482-007-9188-9
- Mundi, A., Delcenserie, V., Amiri-Jami, M., Moorhead, S., and Griffiths, M. W. (2013). Cell-free preparations of *Lactobacillus acidophilus* strain La-5 and *Bifidobacterium longum* strain NCC2705 affect virulence gene expression in *Campylobacter jejuni*. *J. Food Prot.* 76, 1740–1746. doi: 10.4315/0362-028X.JFP-13-084
- Nakamura, T., Kawase, H., Kimura, K., Watanabe, Y., Ohtani, M., Arai, I., et al. (2003). Concentrations of sialyloligosaccharides in bovine colostrum and milk during the prepartum and early lactation. *J. Dairy Sci.* 86, 1315–1320. doi: 10.3168/jds.S0022-0302(03)73715-1
- Nakanishi, N., Abe, H., Ogura, Y., Hayashi, T., Tashiro, K., Kuhara, S., et al. (2006). ppGpp with DksA controls gene expression in the locus of enterocyte effacement (LEE) pathogenicity island of enterohaemorrhagic *Escherichia coli* through activation of two virulence regulatory genes. *Mol. Microbiol.* 61, 194–205. doi: 10.1111/j.1365-2958.2006.05217.x
- Pacheco, A. R., Barile, D., Underwood, M. A., and Mills, D. A. (2015). The impact of the milk glyco-biome on the neonate gut microbiota. *Annu. Rev. Anim. Biosci.* 3, 419–445. doi: 10.1146/annurev-animal-022114-111112
- Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* 29, 2002–2007. doi: 10.1093/nar/29.9.e45
- Rashid, R. A., Tabata, T. A., Oatley, M. J., Besser, T. E., Tarr, P. I., and Moseley, S. L. (2006). Expression of putative virulence factors of *Escherichia coli* O157:H7 differs in bovine and human infections. *Infect. Immun.* 74, 4142–4148. doi: 10.1128/IAI.00299-06
- Sakarya, S., Göktürk, C., Öztürk, T., and Ertugrul, M. B. (2010). Sialic acid is required for nonspecific adherence of *Salmonella enterica* ssp. *enterica* serovar Typhi on Caco-2 cells. *FEMS Immunol. Med. Microbiol.* 58, 330–335. doi: 10.1111/j.1574-695X.2010.00650.x
- Scholten, P. A. (2014). Stool characteristics of infants receiving short-chain galacto-oligosaccharides and long-chain fructo-oligosaccharides: a review. *World, J. Gastroenterol.* 20:13446. doi: 10.3748/wjg.v20.i37.13446
- Scott, K. P., Antoine, J., Midtvedt, T., and Van Hemert, S. (2015). Manipulating the gut microbiota to maintain health and treat disease. *Microb. Ecol. Health Dis.* 1, 1–10. doi: 10.3402/mehd.v26.25877
- Sela, D. A. (2011). Bifidobacterial utilization of human milk oligosaccharides. *Int. J. Food Microbiol.* 149, 58–64. doi: 10.1016/j.ijfoodmicro.2011.01.025
- Sela, D. A., Chapman, J., Adeuya, A., Kim, J. H., Chen, F., Whitehead, T. R., et al. (2008). The genome sequence of *Bifidobacterium longum* subsp. *infantis* reveals adaptations for milk utilization within the infant microbiome. *Proc. Natl. Acad. Sci. U.S.A.* 105, 18964–18969. doi: 10.1073/pnas.0809584105
- Sela, D. A., and Mills, D. A. (2010). Nursing our microbiota: molecular linkages between bifidobacteria and milk oligosaccharides. *Trends Microbiol.* 18, 298–307. doi: 10.1016/j.tim.2010.03.008
- Severi, E., Hood, D. W., and Thomas, G. H. (2007). Sialic acid utilization by bacterial pathogens. *Microbiology* 153, 2817–2822. doi: 10.1099/mic.0.2007/009480-0
- Singh, R., and Jiang, X. (2015). Expression of stress and virulence genes in *Escherichia coli* O157:H7 heat shocked in fresh dairy compost. *J. Food Prot.* 78, 31–41. doi: 10.4315/0362-028X.JFP-13-529
- Smilowitz, J. T., Lebrilla, C. B., Mills, D. A., German, J. B., and Freeman, S. L. (2014). Breast milk oligosaccharides: structure-function relationships in the neonate. *Annu. Rev. Nutr.* 34, 1–27. doi: 10.1146/annurev-nutr-071813-105721
- Sumi, C. D., Yang, B. W., Yeo, I. C., and Hahm, Y. T. (2015). Antimicrobial peptides of the genus *Bacillus*: a new era for antibiotics. *Can. J. Microbiol.* 61, 93–103. doi: 10.1139/cjm-2014-0613
- Takle, G. W., Toth, I. K., and Brurberg, M. B. (2007). Evaluation of reference genes for real-time RT-PCR expression studies in the plant pathogen *Pectobacterium atrosepticum*. *BMC Plant Biol.* 7:50. doi: 10.1186/1471-2229-7-50
- Tanimou, J., Delcenserie, V., Taminiau, B., Daube, G., Saint-Hubert, C., and Durieux, A. (2016). Growth and freeze-drying optimization of *Bifidobacterium crudilactis*. *Food Nutr. Sci.* 7, 616–626. doi: 10.4236/fns.2016.77063
- Tao, N., DePeters, E. J., Freeman, S., German, J. B., Grimm, R., and Lebrilla, C. B. (2008). Bovine milk glycome. *J. Dairy Sci.* 91, 3768–3778. doi: 10.3168/jds.2008-1305
- Tao, N., DePeters, E. J., German, J. B., Grimm, R., and Lebrilla, C. B. (2009). Variations in bovine milk oligosaccharides during early and middle lactation stages analyzed by high-performance liquid chromatography-chip/mass spectrometry. *J. Dairy Sci.* 92, 2991–3001. doi: 10.3168/jds.2008-1642
- Tellez, A., Corredig, M., Guri, A., Zanabria, R., Griffiths, M. W., and Delcenserie, V. (2012). Bovine milk fat globule membrane affects virulence expression in *Escherichia coli* O157:H7. *J. Dairy Sci.* 95, 6313–6319. doi: 10.3168/jds.2012-5560
- Underwood, M. A., German, J. B., Lebrilla, C. B., and Mills, D. A. (2015). *Bifidobacterium longum* subspecies *infantis*: champion colonizer in the infant gut. *Pediatr. Res.* 77, 229–235. doi: 10.1038/pr.2014.156

- Urashima, T., Taufik, E., Fukuda, K., and Asakuma, S. (2013). Recent advances in studies on milk oligosaccharides of cows and other domestic farm animals. *Biosci. Biotechnol. Biochem.* 77, 455–466. doi: 10.1271/bbb.120810
- Vimr, E. R., Kalivoda, K. A., Deszo, E. L., and Steenbergen, S. M. (2004). Diversity of microbial sialic acid metabolism. *Microbiol. Mol. Biol. Rev.* 68, 132–153. doi: 10.1128/MMBR.68.1.132-153.2004
- Wood, K. T., Gonzalez Barrios, A. F., Herzberg, M., and Lee, J. (2006). Motility influences biofilm architecture in *Escherichia coli*. *Appl. Microbiol. Biotechnol.* 72, 361–367. doi: 10.1007/s00253-005-0263-8
- Xu, H., Lee, H. Y., and Ahn, J. (2010). Growth and virulence properties of biofilm-forming *Salmonella enterica* serovar Typhimurium under different acidic conditions. *Appl. Environ. Microbiol.* 76, 7910–7917. doi: 10.1128/AEM.01508-10
- Zeinhom, M., Tellez, A. M., Delcenserie, V., El-Kholy, A. M., El-Shinawy, and Griffiths, M. W. (2012). Yoghurt containing bioactive molecules produced by *Lactobacillus acidophilus* La-5 exerts a protective effect against enterohaemorrhagic *Escherichia coli* (EHEC) in mice. *J. Food Prot.* 10, 1796–1805. doi: 10.4315/0362-028X.JFP-11-508
- Zivkovic, A. M., and Barile, D. (2011). Bovine milk as a source of functional oligosaccharides for improving human health. *Adv. Nutr.* 2, 284–289. doi: 10.3945/an.111.000455

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Bondue, Crèvecoeur, Brose, Daube, Seghayé, Griffiths, LaPointe and Delcenserie. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.