



How to Control HTLV-1-Associated Diseases: Preventing *de Novo* Cellular Infection Using Antiviral Therapy

Amandine Pasquier^{1,2,3}, Sandrine Alais^{1,2}, Loic Roux^{4†}, Maria-Isabel Thoulouze⁵, Karine Alvarez⁴, Chloé Journo^{1,2}, H el ene Dutartre^{1,2‡} and Renaud Mahieux^{1,2*‡}

¹ International Center for Research in Infectiology, Retroviral Oncogenesis Laboratory, INSERM U1111 – Universit e Claude Bernard Lyon 1, CNRS, UMR5308, Ecole Normale Sup erieure de Lyon, Universit e Lyon, Lyon, France, ² Equipe Labellis ee Ligue Nationale Contre le Cancer, Paris, France, ³ Ecole Pratique des Hautes Etudes, Paris, France, ⁴ CNRS UMR 7257, Architecture et Fonction des Macromol cules Biologiques, Aix-Marseille Universit e, Marseille, France, ⁵ "Biofilm and Viral Transmission" Team, Structural Virology Unit, Department of Virology, CNRS UMR 3569, Institut Pasteur, Paris, France

OPEN ACCESS

Edited by:

Umberto Bertazzoni,
University of Verona, Italy

Reviewed by:

Beatrice Macchi,
Universit a degli Studi di Roma Tor
Vergata, Italy
Johan Van Weyenbergh,
KU Leuven, Belgium

*Correspondence:

Renaud Mahieux
renaud.mahieux@ens-lyon.fr

† Present address:

Loic Roux,
RNA Therapeutics Institute,
Worcester, MA, United States

‡ These authors have contributed
equally to this work.

Specialty section:

This article was submitted to
Virology,
a section of the journal
Frontiers in Microbiology

Received: 20 December 2017

Accepted: 07 February 2018

Published: 13 March 2018

Citation:

Pasquier A, Alais S, Roux L,
Thoulouze M-I, Alvarez K, Journo C,
Dutartre H and Mahieux R (2018)
How to Control HTLV-1-Associated
Diseases: Preventing *de Novo* Cellular
Infection Using Antiviral Therapy.
Front. Microbiol. 9:278.
doi: 10.3389/fmicb.2018.00278

Five to ten million individuals are infected by Human T-cell Leukemia Virus type 1 (HTLV-1). HTLV-1 is transmitted through prolonged breast-feeding, by sexual contacts and by transmission of infected T lymphocytes through blood transfusion. One to ten percent of infected carriers will develop a severe HTLV-1-associated disease: Adult-T-cell leukemia/lymphoma (ATLL), or a neurological disorder named Tropical Spastic Paraparesis/HTLV-1 Associated Myelopathy (TSP/HAM). *In vivo*, HTLV-1 is mostly detected in CD4⁺ T-cells, and to a lesser extent in CD8⁺ T cells and dendritic cells. There is a strong correlation between HTLV-1 proviral load (PVL) and clinical status of infected individuals. Thus, reducing PVL could be part of a strategy to prevent or treat HTLV-1-associated diseases among carriers. Treatment of ATLL patients using conventional chemotherapy has very limited benefit. Some chronic and acute ATLL patients are, however, efficiently treated with a combination of interferon α and zidovudine (IFN- α /AZT), to which arsenic trioxide is added in some cases. On the other hand, no efficient treatment for TSP/HAM patients has been described yet. It is therefore crucial to develop therapies that could either prevent the occurrence of HTLV-1-associated diseases or at least block the evolution of the disease in the early stages. *In vivo*, reverse transcriptase (RT) activity is low in infected cells, which is correlated with a clonal mode of viral replication. This renders infected cells resistant to nucleoside RT inhibitors such as AZT. However, histone deacetylase inhibitors (HDACi) associated to AZT efficiently induces viral expression and prevent *de novo* cellular infection. In asymptomatic STLV-1 infected non-human primates, HDACi/AZT combination allows a strong decrease in the PVL. Unfortunately, rebound in the PVL occurs when the treatment is stopped, highlighting the need for better antiviral compounds. Here, we review previously used strategies targeting HTLV-1 replication. We also tested a series of HIV-1 RT inhibitors in an *in vitro* anti-HTLV-1 screen, and report that bis-POM-PMEA

(adefovir dipivoxil) and bis-POC-PMPA (tenofovir disoproxil) are much more efficient compared to AZT to decrease HTLV-1 cell-to-cell transmission *in vitro*. Our results suggest that revisiting already established antiviral drugs is an interesting approach to discover new anti-HTLV-1 drugs.

Keywords: HTLV-1, antiviral therapy, proviral load, acyclic nucleoside phosphonates and thiophosphonates, AZT, prodrugs, cell-cell transmission

INTRODUCTION

Five to ten million individuals are infected worldwide by the oncogenic Human T-cell Leukemia Virus type 1 (HTLV-1) (Gessain and Cassar, 2012). This retrovirus is mainly present in Japan, Sub-Saharan Africa, the Caribbean region and Brazil. HTLV-1 is transmitted from mother-to-child through prolonged breast-feeding, by sexual contacts mostly from man to woman and by transmission of infected T lymphocytes through blood transfusion (Willems et al., 2017). HTLV-1 is mostly detected in CD4⁺ T-lymphocytes, and to a lesser extent CD8⁺ T-cells and dendritic cells *in vivo*. Interestingly, the differences in the number of HTLV-1-infected cells among distinct cell types are not due to the ability of HTLV-1 to infect these cells, but rather to its ability to persist and eventually transform the infected cells. Indeed, *in vitro*, human monocyte-derived dendritic cells are more susceptible to HTLV-1 infection than autologous CD4⁺ T-cells (Alais et al., 2015). In animal models, although it infects CD8⁺ T-cells and dendritic cells in the early phase of infection, HTLV-1 does not transform CD8⁺ T-cells (Rahman et al., 2010; Valeri et al., 2010; Kannian et al., 2013). Nevertheless, there is a strong correlation between HTLV-1 proviral load (PVL, i.e., the number of single integrated copies of the viral genome in cells), and the number and abundance of HTLV-1 infected CD4⁺ T-cells clones (Melamed et al., 2015), and the clinical status of the individuals (Gillet et al., 2013; Cook et al., 2014). A biomarker that would predict which carrier will develop an HTLV-1-associated pathology has not been described so far.

Difficulties to Treat HTLV-1 Patients: Two Associated Diseases and Treatments


One to ten percent of infected carriers develop a severe HTLV-1-associated disease during their life: Adult-T-cell leukemia/lymphoma (ATLL), a CD4⁺ T lymphoproliferation of very poor prognosis with a mean survival time of 6 months in the acute form, or a progressive neurological disorder named Tropical Spastic Paraparesis/HTLV-1-Associated Myelopathy (TSP/HAM) (Yamano and Sato, 2012; Bangham et al., 2015). About 1000 cases of ATLL are diagnosed each year in Japan (Iwanaga et al., 2012). In the Martinique Island (French West Indies), a 14-year follow-up study allowed the detection of 123 TSP/HAM cases for only 400,000 inhabitants (Olindo et al., 2006). Because HTLV-1-infected cells are resistant to apoptosis-inducing agents, treatment of ATLL patients using conventional chemotherapy has very limited benefit (Utsunomiya et al., 2015). In Japan, when possible, ATLL patients undergo allogeneic hematopoietic stem cell transplantation (Utsunomiya et al., 2001). In Europe and United States, chronic and acute

ATLL patients are treated with an anti-viral combination of recombinant interferon alpha (IFN- α) and AZT (zidovudine), combined in some cases with arsenic trioxide (El-Sabban et al., 2000; Nasr et al., 2011). It is worth noting that these results were convincingly confirmed by a world-wide meta analysis (Bazarbachi et al., 2010). However, this treatment is efficient only in a subset of patients, despite the efficacy of arsenic trioxide for inducing the death of HTLV-1-infected cells *in vitro* (El-Sabban et al., 2000; Mahieux and Hermine, 2005). While one report described the use of cyclosporin A for treating TSP/HAM patients with some benefits (Martin et al., 2012), treatment of these patients is in fact still an issue. Other drugs have been tried in clinic, although they generally have a very limited effect. As an example, a trial involving IFN- α had a modest but significant effect (Izumo et al., 1996). Different open trials [summarized by (Nakagawa et al., 1996)] have shown clinical benefit for glucocorticoids, followed by IFN- α , azathioprine and high-dose vitamin C. On the other hand, antiviral effects and/or a decrease in PVL, as well as immunomodulatory effects have been demonstrated for IFN- α (Saito et al., 2004; Rafatpanah et al., 2012), IFN- β (Oh et al., 2005), vitamin C (Moens et al., 2012), cyclosporine (Martin et al., 2012), danazol (Harrington et al., 1991), HDAC inhibitors (Lezin et al., 2007) in HAM/TSP *ex vivo* or *in vivo*.

Viral Amplification Through Clonal Expansion in Addition to *de Novo* Infection: A Difference With HIV-1 Propagation

Contrary to other viruses, HTLV-1 cannot be transmitted efficiently through cell-free viral particles. Using a cell-free experimental system, it was shown that compared to HIV-1, HTLV-1 had a low infectivity (at least 1000-fold lower luciferase activity) and that this was linked to some properties of the viral core and to post-entry processes that are still unclear (Derse et al., 2001). In contrast, HTLV-1 is efficiently transmitted following contacts between an infected donor cell and an uninfected target cell through the establishment of viral synapses and the transfer of viral biofilm (**Figure 1**, left part) (Igakura et al., 2003; Pais-Correia et al., 2010; Thoulouze and Alcover, 2011; Alais et al., 2015). Of note, cell-associated viral transmission of HIV-1 through nanotubes, filopodes or viral synapses is also more efficient than the cell-free infection protocol that is commonly used in *in vitro* experiments (Jolly and Sattentau, 2005; Sherer et al., 2007; Sowinski et al., 2008; Rudnicka et al., 2009; Zhong et al., 2013).

HTLV-1 infection then leads to the clonal expansion of infected cells (**Figure 1**, right part) (Wattel et al., 1995; Bangham



et al., 2014; Turpin et al., 2017; Watanabe, 2017). Because the reverse transcriptase (RT) is not involved in replication by clonal expansion, this phenomenon may explain the very low genetic variability of the virus despite the low fidelity of its RT (Mansky, 2000). Interestingly, HIV-1 clonal expansion also occurs (Maldarelli et al., 2014; Boritz et al., 2016) and has been suggested to allow the virus to overcome antibody neutralization and surface retention by the tetherin restriction factor (Zhong et al., 2013), although the consequences of such a phenomenon remain to be fully understood.

Targeting Viral Replication With the Use of Antivirals: The HIV Example

Nowadays, 25 antiretroviral agents classified in six classes have been approved to treat HIV infections (Cihlar and Fordyce, 2016). The antiretroviral therapy (cART) involves combinations of drugs to achieve maximal response and is generally composed of two nucleoside/nucleotide reverse transcriptase inhibitors (NRTIs) and a third active antiviral from a different class (NNRTI: non-nucleosidic RT inhibitor, INSTI: integrase inhibitor, PI: protease inhibitor, EI: entry inhibitor).

Nucleotide reverse transcriptase inhibitors were the first class of compounds to be used in HIV therapy, by the

approval of zidovudine (AZT) in 1987 (Yarchoan et al., 1986), initially discovered as potent anti-cancer agent (Furmanski et al., 1980). NRTIs are compounds that become active after being phosphorylated into their triphosphate forms, in a process that involves three distinct phosphorylation steps catalyzed by cellular kinases. While they are generally poor substrates for cellular polymerases, triphosphorylated NRTIs compete with natural triphosphate nucleotides for incorporation into growing viral DNA by HIV RT (Furman et al., 1986), resulting in DNA chain termination by blocking further DNA extension (Mitsuya and Broder, 1987).

Historically, AZT and ddI (didanosine) were first shown to block HIV replication in T cells (Mitsuya et al., 1985) and subsequently were shown to suppress HIV replication in monocytes and macrophages *in vitro* (Perno et al., 1988). Other nucleoside analogs without oxacyclopentane sugar moiety, as well as 2',3'-dideoxynucleosides such as d4T (Balzarini et al., 1987), or carbocyclic nucleosides such as Abacavir (ABC) (Daluge et al., 1997), were then identified to be active against HIV, resulting in the emergence of structure activity relationship (SAR) knowledge (Balzarini et al., 1987). However, nucleoside analogs were not equivalent in either activity or toxicity profiles when used as therapeutic drugs *in vitro* and *in vivo* (Mitsuya et al., 1990), and in particular in their capacity to be activated through

phosphorylation in cells. Indeed, during this indispensable process, the first phosphorylation step is often rate-limiting and thus an important bottleneck that limits NRTI efficacy in cells.

To circumvent this drawback, acyclic nucleoside phosphonates (ANPs) have revolutionized the antiviral drug field (Holy, 2003; De Clercq and Holy, 2005). ANPs are nucleotide analogs with a phosphonate group attached to the nucleoside moiety, which require only two phosphorylation steps to be converted into their fully phosphorylated active form (Sharma et al., 2004). The anti-HIV properties of Tenofovir (PMPA, 9-[(R)-2-(Phosphonomethoxy)Propyl]Adenine) were first described in Balzarini et al. (1993) and 8 years later, the oral prodrug, namely Tenofovir disoproxil (TDF) was licensed for clinical use for the treatment of HIV infection. As such, TDF has remained a cornerstone for 14 years and has been replaced recently by a new oral prodrug of the same compound, namely Tenofovir Alafenamide (TAF), which allows a better pharmacokinetic profile in patient fluids (Ray et al., 2016).

Undesirable emergence of viral resistance, toxicity, drug-related adverse effects and drug-drug interactions associated with treatment using NRTIs have challenged the antiviral therapy. Over the last 30 years, it has led to the discovery and approval of several new compounds (Cihlar and Fordyce, 2016) with new ones still in ongoing development (Sax et al., 2015). This intense antiviral research provided the clinic with different anti-HIV active drugs that are now used in combination and have markedly decreased mortality and morbidity from HIV-1 infections in the developed world (Palella et al., 1998). This illustrates that the effort to provide the clinic with new drug regimens results in successful therapies that benefit to patients.

In 2017, the NRTIs approved by the FDA for the treatment of HIV-1 include Retrovir® (Zidovudine, AZT, 1987), Videx® (didanosine, ddI, 1991), Zerit® (stavudine, d4T, 1994), Epivir® (lamivudine, 3TC, 1995), Combivir® (AZT + 3TC, 1997), Ziagen® (abacavir sulfate, ABC, 1998), Trizivir® (AZT + 3TC + ABC, 2000), Viread® (tenofovir disoproxil fumarate, TDF, 2001), Emtriva® (emtricitabine, FTC, 2003), Epzicom® (3TC + ABC, 2004), Truvada® (TDF + FTC, 2004), Atripla® (TDF + FTC + Efavirenz (NNRTI), 2006), Complera® [TDF + FTC + Elvitegravir (INSTI) + booster, 2011], Triumeq® [3TC + ABC + dolutegravir (INSTI), 2014], Genvoya® [FTC + Tenofovir Alafenamide (TAF) + Elvitegravir (INSTI) + booster, 2015], Odefsey® [FTC + TAF + rilpivirine (NNRTI), 2016], Descovy® (FTC + TAF, 2016). Hivid® (zalcitabine, ddC, 1992) was discontinued due to genotoxicity. This antiviral success prompted clinician and researchers to test whether some of the anti-HIV drugs, and especially the acyclic phosphonates ones, could also be active against infections in which viral replication use a viral DNA polymerase enzyme, such as HBV, HSV, EBV, or CMV (De Clercq and Holy, 2005). Interestingly, although the acyclic phosphonate-including class has a broad-spectrum efficacy against several divergent DNA virus families, each acyclic phosphonate within the class has its own specificity. For example, HPMPC is able to inhibit the replication of *Papovaviridae*, *Adenoviridae* and some viruses from the *Herpesviridae* family, while PMEA and PMPA are able to inhibit *Hepadnaviridae* and *Retroviridae*, and PMEA and

HPMPC are able to inhibit the replication of several, but not all *Herpesviridae* (De Clercq and Holy, 2005). Thus, some inhibitors first described as active against HIV may have a broader antiviral spectrum, even on viruses belonging to different classes, strongly supporting the concept of re-positioning of existing drugs.

HIV-NRTI Repositioning on HTLV-1

Because replication of both HIV-1 and HTLV-1 retroviruses (whether it is a frequent step or not, as discussed above) requires RNA reverse transcription performed by phylogenetically related RTs, it is tempting to hypothesize that some anti-HIV-1 drugs targeting HIV-1 RT might also be efficient against HTLV-1 RT. Indeed, using cellular assays of viral transmission, it was shown that AZT, 3TC, phosphonated carbocyclic 2'-Oxa-3'-Aza-nucleoside and Tenofovir each reduced HTLV-1 transmission through cell-cell contact (see **Table 1**) (Macchi et al., 1997; Zhang et al., 2001; Balestrieri et al., 2002, 2005, 2008; Chiacchio et al., 2005, 2007; Pais-Correia et al., 2010). Furthermore, *in vitro* RT assays using HTLV-1 virions confirmed that these NRTIs target the enzymatic RT activity of HTLV-1 (Garcia-Lerma et al., 2001; Balestrieri et al., 2005, 2008; Macchi et al., 2011). The first inhibitory values were obtained using a test that is not quantitative but allows only to determine the drug efficiency (Yamamoto et al., 1996). However, a clinical trial involving AZT as a putative HTLV-1 RT inhibitor, showed only a very limited efficacy in reducing HTLV-1 PVL and no immunological or clinical responses could be demonstrated (Taylor et al., 2006). Furthermore, other data suggested that AZT could in fact inhibit telomerase and lead to cell death, irrespective of the infected status of treated cells (Datta et al., 2006).

A number of studies were designed to decipher why nucleoside analogs, contrary to HIV-1, are so inefficient against HTLV-1. One obvious reason, already mentioned, is the pseudo-latency of the virus (Taylor and Matsuoka, 2005; Taylor et al., 2006). Indeed, in chronically infected asymptomatic carriers, HTLV-1 does not frequently replicate by reverse transcription but rather replicates through clonal expansion of the infected cells (Wattell et al., 1995; Gillet et al., 2011, 2017). Another explanation lies in the mode of viral cell-to-cell transmission. As discussed above, HTLV-1 transmission through cell-free viral particles does not seem efficient. In contrast, cell-to-cell transmission mostly occurs through cell-cell contacts between an infected cell and an uninfected target (Derse et al., 2001; Ceccaldi et al., 2006; Mazurov et al., 2010; Alais et al., 2015). HTLV-1 transmission after cell-to-cell contacts is thought to lead to a high multiplicity of infection and to drug resistance, as was previously described in the case of HIV (Sigal et al., 2011; Zhong et al., 2013). Interestingly, while AZT is indisputably recognized as an efficient HIV-1 replication inhibitor, reported IC₅₀ are consistently higher when inhibition assays are performed using cell-to-cell infection settings instead of cell-free virus (Sigal et al., 2011; Agosto et al., 2014), suggesting that cell-to-cell transmitted viruses have an inherent resistance to reverse transcription inhibitors, which may not be linked to the molecular enzymatic resistance of the RT. Thus, efficacy of

TABLE 1 | Summary of reported inhibition tests using NRTI in cell-to-cell transmission (A) or *in vitro* RT assays (B).

(A) Cell-to-cell transmission.							
	Hill et al., 2003	Zhang et al., 2001	Macchi et al., 1997	Balestrieri et al., 2008	Balestrieri et al., 2008		
	Single-cycle infection (VLP and cell-to-cell transmission)	Cell-to-cell transmission with MT2 (donors) and cord blood PBMCs (targets)	Cell-to-cell transmission with MT2 (donors) and blood PBMCs (targets)				
	24 h p.i. single cycle infection (IC₅₀)	4 days p.i. viral RNA (IC₅₀)	4–7 days viral RNA (IC₅₀)	4 weeks (p19 EC₅₀)	4–7 days viral RNA (EC₁₀₀)		
Tenofovir	5 nM	<i>Not tested</i>	<i>Not tested</i>	0.73 μM	2.3 μM		
AZT	0.11 μM	0.3 ± 0.15 μM DNA PCR and 0.01 ± 0.00 μM RT-PCR	0.01 ± 0.0 μM RNA dot blot and 0.03 ± 0.01 μM DNA dot blot	ND	0.7 μM		
ddC	0.27 μM	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>		
Abacavir	4.6 μM	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>		
d4T	14.5 μM	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>		
3TC	22 μM	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>		
(B) In vitro RT activity.							
	Garcia-Lerma et al., 2001	Balestrieri et al., 2008			Balestrieri et al., 2005		Macchi et al., 2011
	Indirect RT activity assay on virions from infected cells. The amount of RT in the viral lysates was indirectly assessed through the antigen capture assay by measuring the amount of p19 in the supernatant from infected cells. The RT added to the assay was equivalent to a standardized amount of 32 pg of p19. RNA template = RNA extracted from transfected cells ectopically expressing the glycoprotein D of herpes simplex virus type 1 (HSV-1gD, 1 μg). The assayed compounds phosphorylated by crude extract were added at concentration ranging from 0.1 to 1000 nM; 1 mM dNTP, and DNA-PCR was carried out for 35 cycles (Yamamoto et al., 1996).	Indirect RT activity assay on virions from infected cells (RT added was equivalent to a standardized amount of 32 pg of p19), using RNA template corresponding to total RNA extracted from HSV-D glycoprotein transfected cells (HSV-1gD 1 μg). Inhibitors were phosphorylated by crude extract (0.1–1000 nM) and assays were carried out with 1 mM dNTP, and 35 cycles DNA-PCR.					
	RT from MT-2; Hut102 and patients' isolates (EC₁₀₀)	MT2 (EC₁₀₀)	C91 (EC₁₀₀)	Hut102 (EC₁₀₀)	MT2 (EC₁₀₀)	MT2 (EC₁₀₀)	Patients' isolates (EC₁₀₀)
Tenofovir	<i>Not tested</i>	0.4 nM	1 nM	10 nM	1 nM	0.4 nM	10–1000 nM (TDF)
AZT	0.12–0.38 μM	0.4 nM	<i>Not tested</i>	<i>Not tested</i>	10 nM	0.4 nM	4–8 nM
3TC	>10 μM, i.e., no inhibition	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>	<i>Not tested</i>

IC₅₀ is inhibitory concentration 50%, EC₅₀ is effective concentration 50%. EC₁₀₀ is effective concentration 100%.

AZT or other NRTIs against HTLV-1 might be further limited by the almost exclusive cell-to-cell transmission mode of the virus.

Several parameters might further account for the discrepancies between results obtained *in vitro* and *in vivo*. First, the efficacy of AZT against HTLV-1 as measured by the IC₅₀ might be dependent on the experimental settings. This is illustrated by a study that reported a 1000-fold lower efficacy of compounds when used in cell-to-cell transmission assays (inhibition range of 1 μM) compared to *in vitro* RT

inhibition assays performed on virions from supernatant of infected cells (inhibition range of around 1 nM) [Table 1, (Balestrieri et al., 2008)]. Thus, IC₅₀ determined *in vitro* or in cellular assays may not be relevant to what occurs *in vivo*. Furthermore, sequence variation within the RT gene, in addition to its influence on replication efficiency (Mitchell et al., 2007), might also be responsible for different sensitivity of RT from virions produced by different HTLV-1 infected cells lines (Balestrieri et al., 2008) or RT derived from patient isolates (Macchi et al., 2011) to NRTIs (Table 1). Even if inhibition

is measured in cell–cell transmission assays using the same infected donor cells, the use of different target cells might explain the variability in NRTI efficacy. For example, AZT efficacy was measured on HTLV-1 transmission after co-culture of chronically infected MT2 cells with primary cord PBMCs (Zhang et al., 2001) or adult blood PBMCs (Macchi et al., 1997; Balestrieri et al., 2008) (Table 1). In these assays, AZT was estimated to inhibit HTLV-1 transmission after 4–7 days of culture, as measured by viral DNA or viral RNA detected in primary PBMCs. AZT IC₅₀ values were 0.03 ± 0.01 and $0.01 \pm 0.00 \mu\text{M}$ for DNA and RNA, respectively, in PBMCs and 0.30 ± 0.15 and $0.01 \pm 0.00 \mu\text{M}$ for DNA and RNA in cord blood, respectively (Table 1). The lower susceptibility to AZT inhibition of PBMCs from adult blood highlights that besides the direct effect of AZT on RT activity in virions released from MT2 infected cells (Table 1), other mechanisms exist in the target cells that modulate the efficacy of inhibition. This could include drug uptake and intracellular phosphorylation that have been shown to be cell-type dependent (Deville-Bonne et al., 2010; Roux et al., 2013), or RT activity in the target cell that could differ depending on the cellular environment. Very little is known regarding RT activity in cells during the early steps of the viral cycle. Yet, because cell-to-cell infection is much more efficient than cell-free infection, it is possible that either a higher number of virions is delivered to the target cell during cell-to-cell contact or that contact-dependent signals are required to enhance virus replication. Such signals could also modulate the sensitivity of target cells to RT inhibition, in a target cell-specific manner.

Of note, although co-culture assays probably mimic HTLV-1 transmission *in vivo*, inhibition is measured after several days of culture. This long-term culture of infected cells could select for T-cells susceptible to infection and able to survive after HTLV-1 infection, due to the immortalization potency of the virus. Furthermore, several rounds of infection could occur that may not be suitable for quantitative analysis of infection and replication. Alternative experimental systems have been developed to measure HTLV-1 spread after a single-cycle replication (Derse et al., 2001) and to measure viral inhibition using NRTI (Hill et al., 2003) (Table 1). This elegant system [reviewed in (Aloia et al., 2013)] is based on virus-like particles (VLPs) that contain HTLV-1 RT and that deliver a surrogate retroviral genome designed to express only a reporter gene product in infected cells. These VLPs are generated by co-transfecting cells (usually adherent cell lines such as HEK293T) with a packaging plasmid (allowing the expression of HTLV-1 structural, enzymatic and regulatory proteins under the control of a CMV promoter), a transfer reporter vector and an envelope-encoding plasmid [reviewed in (Derse et al., 2001)]. In these settings, the inhibition properties of several nucleoside analogs were determined (Table 1) and results confirmed AZT IC₅₀ to be around $0.1 \mu\text{M}$ (Hill et al., 2003). It is worth noting that efficiencies of all compounds tested against HTLV-1 were at least 100-fold lower than that tested against HIV-1, further suggesting that although both RTs have similar sequence and enzymatic organization, the catalytic properties of the

two RTs are different and/or differently regulated by the cellular environment of the infected cells. Moreover, although useful to monitor single-cycle infections, this system uses VLPs that may be different from WT virions and more importantly, it uses transfected HEK293T donor cells that are very different from the natural infected T-cells expected to transfer the virus *in vivo*. Knowing that cell–cell contacts may regulate HTLV-1 RT activity, the nature of the infected cell used to transfer the virus is expected to modulate the infectivity of the transferred virions as well as the efficacy of NRTIs.

Strategies to Control HTLV-1 PVL *in Vivo*: Replication Reactivation and Use of NRTIs

As discussed above, HTLV-1 pseudo-latency renders this retrovirus usually insensitive to RT inhibitors. Interestingly, viral expression and RT-dependent *de novo* cellular infection can be induced using histone deacetylase inhibitors (HDACi) such as valproic acid (VPA) (Lezin et al., 2007; Olindo et al., 2011). Importantly, long-term treatment of HTLV-1-infected patients with VPA is proven to be safe (Lezin et al., 2007; Olindo et al., 2011). In an earlier work conducted in non-human primates naturally infected with Simian T-Lymphotropic Virus type 1 (STLV-1), the simian equivalent of HTLV-1, we tested a combination of VPA, used to increase viral replication, and AZT, used to inhibit *de novo* cellular infection by targeting the RT activity (Afonso et al., 2010). Because the PVL is higher in TSP/HAM patients than in asymptomatic carriers and can therefore be considered as a prognostic marker (Yakova et al., 2005), we selected asymptomatic STLV-1-infected animals with a high PVL (Afonso et al., 2010). Our results showed for the first time that the AZT/VPA combination leads to a very strong and significant PVL decrease in all tested animals. However, STLV-1 PVL rebounded after the treatment was stopped, thus demonstrating that the virus had not been cleared during the treatment, a disappointing result reminiscent of what is described for HIV-1. As for HIV-1, this rebound of HTLV-1 viral load suggests (i) the existence of a putative viral reservoir inaccessible to the drugs, or (ii) that AZT is not efficient enough to block *de novo* HTLV-1 infection that occurs at a level below the detection threshold, or (iii) that viral escape mutants are selected during treatment, leading to resistance to AZT, although this is unlikely given that resistance mutation were not reported in human patients undergoing AZT. Nevertheless, these results demonstrate that preventing *de novo* infection is an option for controlling the virus and foster the use of more efficient drugs that would better control HTLV-1 dissemination within infected individuals.

Here we show a series of results obtained with already tested drugs as well as with newly compounds. We report for the first time that bis-POM-PMEA (adefovir dipivoxil) and bis-POC-PMPA (tenofovir disoproxil) are much more efficient compared to AZT to decrease HTLV-1 cell-to-cell transmission *in vitro*.

MATERIALS AND METHODS

Cells

Jurkat, Jurkat-LTR-Luc (Pais-Correia et al., 2010), and HTLV-1-infected T-cell line (C91PL) were maintained in RPMI 1640 medium supplemented with 10% fetal calf serum and penicillin-streptomycin (100 µg/ml, Gibco Life Technology). All cells were grown at 37°C in 5% CO₂. Jurkat-LTR-Luc cells, stably transfected with a plasmid encoding luciferase, under the control of the HTLV-1 long terminal repeat (LTR) promoter are maintained under hygromycin (450 µg/ml, Sigma) selection.

Drugs

AZT, d4T, 3TC, tenofovir alafenamide, tenofovir disoproxil and tenofovir disoproxil fumarate were purchased from Selleckchem. PMPA, PMEAs were provided by Dr. Alvarez (AFMB, Marseille). S-PMPA, S-PMEA, bis-POC-S-PMPA, bis-POM-PMEA, and bis-POM-S-PMEA were synthesized and provided by Dr Alvarez (AFMB, Marseille). All drugs were resuspended in DMSO or water according to the furnisher, aliquoted and stored at -80°C until used. Aliquots were used only once to avoid drug degradation under thawing-freezing cycles.

Inhibition of Viral Transmission Test

Jurkat reporter cells (JK-LTR-Luc, 50,000 cells) were seeded in 96-well plates and treated for 18 h in presence or absence of the different drugs (1–50 µM, diluted in medium). Then, cells were washed once and HTLV-1-infected cells (C91PL, 10,000 cells) or control cells (Jurkat, 10,000 cells) were added together with or without an additional 1/2 dose of drugs, and the co-culture was resumed for 24 h. Cell viability was measured by trypan blue exclusion counting. The mortality index was determined as the percentage of dead cells in each condition normalized to the percentage of dead cells in the control condition set to 1. A mortality index higher than 1.5 signs drug toxicity. Reporter activities were assayed using the luciferase reporter assay system (Promega). Luciferase activity was normalized according to the amount of total proteins in the co-culture determined by Bradford (Bio-Rad). Results were then expressed as a percentage of control in the absence of drugs.

Statistical Analysis

One-way analysis of variance (ANOVA) with Bonferroni's *post hoc* multiple comparison test was used to determine statistically significant differences. Differences were considered significant if the *p*-value was < 0.05.

RESULTS

New Strategies for the Repositioning of Anti-HIV Inhibitors as Efficient Anti-HTLV-1 Drugs


High-throughput screening efforts against HIV-1 followed by structure-based drug design have allowed the discovery of several drugs that are more potent than AZT and related

nucleoside analogs, and are still ongoing (Taylor et al., 2016). Highly active non-nucleoside inhibitors eliciting hardly any resistant mutants have been designed when purified target enzymes (RT, protease, integrase) were made available, providing essential structural and functional models. In strong contrast, discovery of anti-HTLV-1 specific drugs is currently significantly impeded by the unavailability of purified viral enzymes. In addition, and as stated above, because the virus is almost exclusively transmitted through cell-cell contacts, cell infection cannot be easily induced *in vitro* using cell-free particles as a source of infectious viruses (Alais et al., 2015). This further challenges the development of drug screening protocols. Since HIV-1 and HTLV-1 use very similar mechanisms of reverse transcription, we hypothesized that anti-HIV-1 NRTIs might be efficient against HTLV-1 RT. NNRTIs, which are highly specific to HIV-1 RT, were thus excluded from our screen.

We thus set up an *in vitro* cell co-culture assay that allows for the monitoring of HTLV-1 *de novo* infection in LTR-luciferase reporter Jurkat T-cells using HTLV-1 chronically infected C91PL T-cells as a source of virus (Alais et al., 2017) (Figure 2A). This system was used to test the efficacy of several NRTIs known to inhibit HIV-1 RT (Figure 2B). In addition, we selected pro-drugs of these NRTIs and evaluated their efficacy and toxicity. AZT (1) was used as a positive control. 2',3'-dideoxynucleosides such as 3TC (2) and d4T (3) were also used as controls (Figure 2B). Because highly variable data are reported in the literature on the effectiveness of these compounds (Table 1), we also included them to calibrate the stringency of our assay. While the ANP Tenofovir (PMPA, 5) has been tested against HTLV-1 previously, its congener Adefovir (PMEA, 4) has never been evaluated on HTLV-1 so far, although its efficiency to inhibit HIV and HBV infections is described (De Clercq and Holy, 2005). Both compounds were included in our screen. We also included new ANP analogs, named thiophosphonates, S-PMEA (6) and S-PMPA (7), which have shown potent activity against HIV and HBV infections (Barral et al., 2011; Roux et al., 2013; Priet et al., 2015).

As discussed above, ANPs are NRTIs that contain a stable phosphonate group. After penetration into the target cells, ANPs are phosphorylated into their diphosphate forms which serve as substrates for RT and thus exert the antiviral effect. Because ANPs are negatively charged at physiological pH, they are less efficient in passing through cell membranes than neutral, lipophilic species. To address this issue, these compounds can be prepared as prodrugs. The activity of ANPs is dramatically increased by the use of their biolabile prodrugs: Adefovir dipivoxil (Hepsera®) was licensed for the treatment of chronic HBV infections during several years (Merle and Trepo, 2001) and Tenofovir is commercialized as two prodrugs, which are much more potent than their parent counterparts.


We therefore also evaluated Adefovir dipivoxil (bis-POM-PMEA, 8), its thiophosphonate equivalent bis-POM-S-PMEA (9), Tenofovir disoproxil (bis-POC-PMPA, 10), Tenofovir disoproxil fumarate (bis-POC-PMPA, 11), Tenofovir alafenamide (12) and the thiophosphonate congener bis-POC-S-PMPA (13).



First, we tested whether these compounds had any impact on target cell viability (**Figure 3**). The concentration of these drugs varied between 1 and 50 μM . The viability of treated cells was represented relative to that of the untreated cells, set to 1. An index > 1 means that the mortality of the treated cells is higher than that of the untreated cells. We considered that cell viability was impaired if the index was > 1.5 and thus excluded all the drug concentrations leading to a mortality index > 1.5 . None of the drugs, i.e., AZT (1), 3TC (2), d4T (3) (**Figure 3A**) as well as the PMEa (4–6) and PMPA (5–7) families had any significant effect on cell viability, i.e., their mortality index was lower than 1.5 (**Figure 3B**). As already reported (Roux et al., 2013), Bis-POM-PMEa (8) and Bis-POM-S-PMEa (9) were toxic when used at 10 μM (**Figure 3C**), as well as BIS-POC-PMPA (10) when used at 50 μM (**Figure 3C**). Thus, these concentrations were excluded for the subsequent evaluation of their ability to prevent HTLV-1 transmission to reporter T-cells after co-culture with infected donor cells (**Figure 4**). LTR-luciferase Jurkat reporter T-cells were used as target cells (Alais et al., 2017). Given that Tax

is absent from the viral particle, transcription from the viral Tax-dependent LTR requires productive infection and Tax expression in the target cells, which can occur after contact with the infected donor cells through the establishment of viral synapses or transfer of viral biofilm (Pais-Correia et al., 2010). Interestingly, this experimental system does not rely on cell-free infection and is therefore closer to *in vivo* viral transmission.

Consistent with previous reports, AZT (1) had a significant but limited effect on HTLV-1 transmission, even at the high concentration of 50 μM (inhibition of 10–20% **Figures 4A,B**), while 3TC (2) had no effect, independently of the concentration used (**Figure 4A**). Surprisingly, AZT (1), had the same significant inhibitory activity for all the doses tested, without any dose-dependent effect (**Figure 4B**). d4T (3) also had no effect (**Figure 4A**), suggesting that previously published cell-free experiments using virion-associated enzymatic RT assays might not be relevant for determining that a drug has an anti-HTLV-1 effect (Garcia-Lerma et al., 2001). We then tested phosphonate PMEa (4) and PMPA (5) (**Figure 4B**) and compared their effect



to that of AZT. Contrary to AZT, PMEAs (4) had a statistically significant dose-dependent inhibitory activity (**Figure 4B**). When used at 50 μM , PMEAs was statistically more efficient than AZT (70% vs. 30% of viral transmission decrease, respectively, $p < 0.0001$). Of note, PMEAs active concentration on HTLV-1 was much higher than the dose reported to inhibit HIV-1 cell-free transmission, which is around 5 μM (Barral et al., 2011). Importantly, it is worth noting that when HIV-1 inhibition assays were performed using cell-to-cell infection instead of infection with cell-free virus, reported IC_{50} were also consistently higher (Sigal et al., 2011; Agosto et al., 2014). PMPA (5) did not show any inhibitory effect compared to control, AZT (1) or PMEAs (4), independently of the concentration used (**Figure 4B**). This is somehow different from a previous report which showed that overnight Tenofovir treatment (1 μM) before co-culture had some effect on viral transmission from chronically-infected MT2 cells to PBMCs, while the same product at 0.01 μM had no effect (Balestrieri et al., 2005). We also tested S-PMEAs (6) and S-PMPA (7) (**Figure 4B**), which showed a limited effect for (6) at the highest concentrations, yet not significantly better than (1) and significantly worse than (4). Thus, modifying PMEAs into the thio-derivative S-PMEAs (6) results in a strong decrease in HTLV-1 transmission inhibition [compare (4) vs. (6)], consistent with the results on HIV replication for which IC_{50} is threefold higher than that of the parental derivative (Barral et al.,

2011). Interestingly, modifying PMPA into its thio-derivative (7) resulted in a limited but significant better inhibition than 5, at 1 and 10 μM (25% of inhibition $p < 0.01$), an effect that is lost when the concentration is increased to 50 μM (**Figure 4B**), probably due to the higher toxicity of 7 at 50 μM (**Figure 3B**).

We then tested ANP pro-drugs since they were reported to display an active concentration that was better than that of their parent compound due to the higher ability in passing through cell membranes (**Figure 4C**). bis-POM-PMEAs (8) showed a statistically better, although modest, inhibition of LTR activation at 1 μM than the parent compound PMEAs (4) tested at the same concentration (compare **Figure 4C**). Although S-PMEAs (6) had a significant yet modest inhibitory effect at 1 μM (20% of inhibition $p < 0.001$, **Figure 4C**), its bis-POM modification (9) results in a loss of this effect when used at the same concentration. This suggests that although the cell-membrane penetration is increased, it does not translate into a better efficacy of the compound. This could be explained by the higher toxicity of the pro-drug (6) at 1 μM (**Figure 3C**), which within the 42 h of treatment could impair the metabolism of the pro-drug after its penetration into the cells. Higher concentration could not be used because of compound 9 toxicity (see **Figure 3C**).

For the PMPA compound family, no effect was observed when pro-drugs (10, 11, or 12) were used at 1 μM . This is consistent with the absence of inhibition observed with the parent PMPA (5)




FIGURE 4 | Inhibition efficacy of drugs on HTLV-1 cell-to-cell transmission. **(A–E)** Luciferase activity was normalized to protein concentration (in μg , quantified by Bradford assay) and expressed as percent of untreated (control) cells set as 100%. Drug concentrations and drug names are indicated. Results were grouped based on the drug family: **(A)** AZT (1), 3TC (2) and d4T (3) nucleoside analogs, **(B)** AZT (1) and acyclic nucleoside phosphonates, **(C)** PMEA and S-PMEA and their prodrugs, **(D)** PMPA and a series of PMPA pro-drugs, **(E)** S-PMPA and its pro-drug. Statistical differences were determined using one-way ANOVA (ns, not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, $n = 2\text{--}4$ independent experiments with triplicate samples).

compound (**Figure 4D**). However, when used at $10\ \mu\text{M}$, prodrugs (10–12) had a higher effect than PMPA (5) itself and induced a significant reduction (50%, $p < 0.0001$) of HTLV-1 replication.

Finally, bis-POC-S-PMPA (13) did not show any improvement over S-PMPA (7) and even had a statistically significant positive effect on viral transmission (**Figure 4E**). This could be explained by the lower de-protection rate of (13) in cell, thus leading to a lower concentration of active S-PMPA within cells (Roux et al., 2013). Surprisingly, the pro-drug effect was lower than expected from the results observed in HIV tests in which IC_{50} are decreased by at least 120-fold (Roux et al., 2013) by bis-POC or bis-POM modifications. This suggests that in our settings, the rate of pro-drugs metabolism and their subsequent use as substrates by HTLV-1 RT might be very low, although they could increase over time, leading to a potential better inhibition.

Altogether, our results show that PMPA (5) is not active when HTLV-1 infection occurs through cell–cell contacts, although (5) inhibits HTLV-1 RT in *in vitro* enzymatic assays (see **Table 1**). However, we demonstrate that several pro-drugs of PMPA are efficient (10, 11, 12), confirming earlier studies (Macchi et al., 2011) and suggesting that they could be used in clinical trials. More importantly, our results extend the inhibitory potential of phosphonate derivatives by showing that PMEA (4) is more efficient than PMPA (5) to block HTLV-1

viral transmission, without any associated-toxicity. In addition, PMEA pro-drug (8) is also efficient, and at a 10-fold lower concentration than all PMPA pro-drugs. However, its associated-toxicity might compromise its use for clinical trial. Finally, the thio-modification of PMEA (6) does not increase PMEA (4) inhibitory potential, while PMPA thio-modification (7) increases PMPA (5) efficiency, although this effect is lost for the pro-drug (13).

DISCUSSION

HTLV-1-induced diseases, i.e., ATLL and TSP/HAM, are of bad prognosis. Despite the early discovery of HTLV-1 more than 30 years ago, and despite the outstanding genetic stability of the virus (which could be a topic of interest for the vaccine industry), specific anti-HTLV-1 drugs have not been developed yet. In addition, and contrary to the HIV-1 field, asymptomatic HTLV-1-infected individuals are not treated, although a high PVL is established to be a valuable risk factor that could be used as a biomarker to identify individuals at risk for developing HTLV-1-associated diseases. Rather, patients are treated once they develop the disease, which is probably too late. Clonal populations of infected T-cells are detected *in vivo*, suggesting that the virus

mostly replicates through clonal expansion (Bangham et al., 2014), although viral expression, potentially leading to viral production, may occur upon metabolic changes occurring in protected body areas (Kulkarni et al., 2017). In addition, HTLV-1 does not replicate only through clonal expansion given that AZT as an *in vivo* effect in asymptomatic individuals (Afonso et al., 2010).

Anti-HIV-1 drugs were first tested using cell-free infection systems in which it is easy to quantify viral production. This is the reason why reverse-transcriptase inhibitors have also been tested using HTLV-1 purified virions (Garcia-Lerma et al., 2001; Balestrieri et al., 2005, 2008), or after co-culture between MT2 infected cell line and PBMCs from normal blood donor (Macchi et al., 2011). This is, however, an issue for translating results into the clinic. Indeed, when it occurs, HTLV-1 transmission goes through cell-cell contacts rather than through cell-free viral particles, a transmission mode that was proven to increase viral resistance to drugs in the HIV-1 situation. Previously described co-culture assays for drug screening used MT2 cells as donor cells (Macchi et al., 1997; Zhang et al., 2001; Balestrieri et al., 2008). Of note, this cell line was recently demonstrated to produce poorly infectious viral particles, even when these cells were used for viral transfer after cell-to-cell contacts (Alais et al., 2015). We have developed an easy-to-use assay relying on the co-culture between HTLV-1-infected C91PL T-cells and LTR-luciferase reporter Jurkat T-cells (Pais-Correia et al., 2010). Here, we demonstrate that this assay is useful to screen a series of drugs, in experimental settings that most closely model the *in vivo* viral transmission mode. Our results using stavudine d4T and lamivudine 3TC are consistent with the literature and show that, for some unexplained reasons, these drugs have no effect on HTLV-1 RT. More interestingly, we show here for the first time that Adefovir (PMEA) as well as its prodrug (Adefovir dipivoxil) work much better than AZT. In the case of Tenofovir (PMPA), the drug has no effect, however, the prodrug (Tenofovir disoproxil) has a significant inhibitory effect when used at 10 μ M.

This encourages us to validate its inhibitory potential *in vivo*, in naturally infected and currently asymptomatic STLV-1 infected non-human primates. The ongoing pre-clinical test combines Tenofovir disoproxil fumarate (Viread) to valproate, in a protocol similar to that used in the Afonso study (Afonso et al., 2010), in which the combination of AZT with valproate led to a strong PVL decrease as long as the treatment did go on. From these results and the results of the present study, it can therefore be expected to get better results with Tenofovir disoproxil fumarate (Viread). This could help HTLV-1 infected patients' treatment in the future.

AUTHOR CONTRIBUTIONS

AP and SA performed the experiments. LR and KA synthesized all phosphonates derivatives except the bis-POC-PMPA fumarate and PMPA Alafenamide which were purchased from Selleckchem. M-IT contributed with technical developments for some experiments and discussion. RM, HD, and KA designed the experiments. RM, CJ, HD, and KA wrote the article.

FUNDING

AP was supported by INSERM. RM and CJ are supported by Ecole Normale Supérieure de Lyon. RM is part of the French Laboratory of Excellence project ECOFECT (ANR-11-LABX-0048). SA and HD are supported by INSERM. KA is supported by CNRS and Aix-Marseille Université. LR was supported by Sidaction (Ph.D. grant). M-IT research was supported by AXA Research fund as well as by institutional grants from Institut Pasteur and the Centre National de la Recherche Scientifique (CNRS). The authors acknowledge the support of ARC, La Ligue Contre le Cancer (Equipe Labellisée EL2013-3 Mahieux) and ANRS.

REFERENCES

- Afonso, P. V., Mekaouche, M., Mortreux, F., Toulza, F., Moriceau, A., Wattel, E., et al. (2010). Highly active antiretroviral treatment against STLV-1 infection combining reverse transcriptase and HDAC inhibitors. *Blood* 116, 3802–3808. doi: 10.1182/blood-2010-02-270751
- Agosto, L. M., Zhong, P., Munro, J., and Mothes, W. (2014). Highly active antiretroviral therapies are effective against HIV-1 cell-to-cell transmission. *PLoS Pathog.* 10:e1003982. doi: 10.1371/journal.ppat.1003982
- Alais, S., Dutartre, H., and Mahieux, R. (2017). Quantitative analysis of human T-lymphotropic virus type 1 (HTLV-1) infection using co-culture with Jurkat LTR-luciferase or Jurkat LTR-GFP reporter cells. *Methods Mol. Biol.* 1582, 47–55. doi: 10.1007/978-1-4939-6872-5_4
- Alais, S., Mahieux, R., and Dutartre, H. (2015). Viral source-independent high susceptibility of dendritic cells to human T-cell leukemia virus type 1 infection compared to that of T lymphocytes. *J. Virol.* 89, 10580–10590. doi: 10.1128/JVI.01799-15
- Aloia, A. L., Duffy, L., Pak, V., Lee, K. E., Sanchez-Martinez, S., Derse, D., et al. (2013). A reporter system for replication-competent gammaretroviruses: the inGluc-MLV-DERSE assay. *Gene Ther.* 20, 169–176. doi: 10.1038/gt.2012.18
- Balestrieri, E., Forte, G., Matteucci, C., Mastino, A., and Macchi, B. (2002). Effect of lamivudine on transmission of human T-cell lymphotropic virus type 1 to adult peripheral blood mononuclear cells in vitro. *Antimicrob. Agents Chemother.* 46, 3080–3083. doi: 10.1128/AAC.46.9.3080-3083.2002
- Balestrieri, E., Matteucci, C., Ascolani, A., Piperno, A., Romeo, R., Romeo, G., et al. (2008). Effect of phosphonated carbocyclic 2'-oxa-3'-aza-nucleoside on human T-cell leukemia virus type 1 infection in vitro. *Antimicrob. Agents Chemother.* 52, 54–64. doi: 10.1128/AAC.00470-07
- Balestrieri, E., Sciortino, M. T., Mastino, A., and Macchi, B. (2005). Protective effect of the acyclic nucleoside phosphonate tenofovir toward human T-cell leukemia/lymphotropic virus type 1 infection of human peripheral blood mononuclear cells in vitro. *Antiviral Res.* 68, 154–162. doi: 10.1016/j.antiviral.2005.09.001
- Balzarini, J., Holy, A., Jindrich, J., Naesens, L., Snoeck, R., Schols, D., et al. (1993). Differential antiherpesvirus and antiretrovirus effects of the (S) and (R) enantiomers of acyclic nucleoside phosphonates: potent and selective in vitro and in vivo antiretrovirus activities of (R)-9-(2-phosphonomethoxypropyl)-2,6-diaminopurine. *Antimicrob. Agents Chemother.* 37, 332–338. doi: 10.1128/AAC.37.2.332
- Balzarini, J., Kang, G. J., Dalal, M., Herdewijn, P., De Clercq, E., Broder, S., et al. (1987). The anti-HTLV-III (anti-HIV) and cytotoxic activity of 2',3'-didehydro-2',3'-dideoxyribonucleosides: a comparison with their parental 2',3'-dideoxyribonucleosides. *Mol. Pharmacol.* 32, 162–167.

- Bangham, C. R., Araujo, A., Yamano, Y., and Taylor, G. P. (2015). HTLV-1-associated myelopathy/tropical spastic paraparesis. *Nat. Rev. Dis. Primers* 1:15012. doi: 10.1038/nrdp.2015.12
- Bangham, C. R., Cook, L. B., and Melamed, A. (2014). HTLV-1 clonality in adult T-cell leukaemia and non-malignant HTLV-1 infection. *Semin. Cancer Biol.* 26, 89–98. doi: 10.1016/j.semcancer.2013.11.003
- Barral, K., Weck, C., Payrot, N., Roux, L., Durafour, C., Zoulim, F., et al. (2011). Acyclic nucleoside thiophosphonates as potent inhibitors of HIV and HBV replication. *Eur. J. Med. Chem.* 46, 4281–4288. doi: 10.1016/j.ejmech.2011.06.034
- Bazarbachi, A., Plumelle, Y., Carlos Ramos, J., Tortevoe, P., Otrock, Z., Taylor, G., et al. (2010). Meta-analysis on the use of zidovudine and interferon- α in adult T-cell leukemia/lymphoma showing improved survival in the leukemic subtypes. *J. Clin. Oncol.* 28, 4177–4183. doi: 10.1200/JCO.2010.28.0669
- Boritz, E. A., Darko, S., Swaszek, L., Wolf, G., Wells, D., Wu, X., et al. (2016). Multiple origins of virus persistence during natural control of HIV infection. *Cell* 166, 1004–1015. doi: 10.1016/j.cell.2016.06.039
- Ceccaldi, P. E., Delebecque, F., Prevost, M. C., Moris, A., Abastado, J. P., Gessain, A., et al. (2006). DC-SIGN facilitates fusion of dendritic cells with human T-cell leukemia virus type 1-infected cells. *J. Virol.* 80, 4771–4780. doi: 10.1128/JVI.80.10.4771-4780.2006
- Chiacchio, U., Balestrieri, E., Macchi, B., Iannazzo, D., Piperno, A., Rescifina, A., et al. (2005). Synthesis of phosphonated carbocyclic 2'-oxa-3'-aza-nucleosides: novel inhibitors of reverse transcriptase. *J. Med. Chem.* 48, 1389–1394. doi: 10.1021/jm049399i
- Chiacchio, U., Rescifina, A., Iannazzo, D., Piperno, A., Romeo, R., Borrello, L., et al. (2007). Phosphonated carbocyclic 2'-oxa-3'-azanucleosides as new antiretroviral agents. *J. Med. Chem.* 50, 3747–3750. doi: 10.1021/jm070285r
- Cihlar, T., and Fordyce, M. (2016). Current status and prospects of HIV treatment. *Curr. Opin. Virol.* 18, 50–56. doi: 10.1016/j.coviro.2016.03.004
- Cook, L. B., Melamed, A., Niederer, H., Valganon, M., Laydon, D., Foroni, L., et al. (2014). The role of HTLV-1 clonality, proviral structure, and genomic integration site in adult T-cell leukemia/lymphoma. *Blood* 123, 3925–3931. doi: 10.1182/blood-2014-02-553602
- Daluge, S. M., Good, S. S., Faletto, M. B., Miller, W. H., St. Clair, M. H., Boone, L. R., et al. (1997). 1592U89, a novel carbocyclic nucleoside analog with potent, selective anti-human immunodeficiency virus activity. *Antimicrob. Agents Chemother.* 41, 1082–1093.
- Datta, A., Bellon, M., Sinha-Datta, U., Bazarbachi, A., Lepelletier, Y., Canioni, D., et al. (2006). Persistent inhibition of telomerase reprograms adult T-cell leukemia to p53-dependent senescence. *Blood* 108, 1021–1029. doi: 10.1182/blood-2006-01-0067
- De Clercq, E., and Holy, A. (2005). Acyclic nucleoside phosphonates: a key class of antiviral drugs. *Nat. Rev. Drug Discov.* 4, 928–940. doi: 10.1038/nrd1877
- Derse, D., Hill, S. A., Lloyd, P. A., Chung, H., and Morse, B. A. (2001). Examining human T-lymphotropic virus type 1 infection and replication by cell-free infection with recombinant virus vectors. *J. Virol.* 75, 8461–8468. doi: 10.1128/JVI.75.18.8461-8468.2001
- Deville-Bonne, D., El Amri, C., Meyer, P., Chen, Y., Agrofoglio, L. A., and Janin, J. (2010). Human and viral nucleoside/nucleotide kinases involved in antiviral drug activation: structural and catalytic properties. *Antiviral Res.* 86, 101–120. doi: 10.1016/j.antiviral.2010.02.001
- El-Sabban, M. E., Nasr, R., Dbaibo, G., Hermine, O., Abboushi, N., Quignon, F., et al. (2000). Arsenic-interferon- α -triggered apoptosis in HTLV-1 transformed cells is associated with tax down-regulation and reversal of NF- κ B activation. *Blood* 96, 2849–2855.
- Furman, P. A., Fyfe, J. A., St. Clair, M. H., Weinhold, K., Rideout, J. L., Freeman, G. A., et al. (1986). Phosphorylation of 3'-azido-3'-deoxythymidine and selective interaction of the 5'-triphosphate with human immunodeficiency virus reverse transcriptase. *Proc. Natl. Acad. Sci. U.S.A.* 83, 8333–8337. doi: 10.1073/pnas.83.21.8333
- Furmanski, P., Bourguignon, G. J., Bolles, C. S., Corombos, J. D., and Das, M. R. (1980). Inhibition by 2',3'-dideoxythymidine of retroviral infection of mouse and human cells. *Cancer Lett.* 8, 307–315. doi: 10.1016/0304-3835(80)90146-9
- Garcia-Lerma, J. G., Nidtha, S., and Heneine, W. (2001). Susceptibility of human T cell leukemia virus type 1 to reverse-transcriptase inhibitors: evidence for resistance to lamivudine. *J. Infect. Dis.* 184, 507–510. doi: 10.1086/322785
- Gessain, A., and Cassar, O. (2012). Epidemiological aspects and world distribution of HTLV-1 infection. *Front. Microbiol.* 3:388. doi: 10.3389/fmicb.2012.00388
- Gillet, N. A., Cook, L., Laydon, D. J., Hlela, C., Verdonck, K., Alvarez, C., et al. (2013). Strongyloidiasis and infective dermatitis alter human T lymphotropic virus-1 clonality in vivo. *PLoS Pathog.* 9:e1003263. doi: 10.1371/journal.ppat.1003263
- Gillet, N. A., Malani, N., Melamed, A., Gormley, N., Carter, R., Bentley, D., et al. (2011). The host genomic environment of the provirus determines the abundance of HTLV-1-infected T-cell clones. *Blood* 117, 3113–3122. doi: 10.1182/blood-2010-10-312926
- Gillet, N. A., Melamed, A., and Bangham, C. R. (2017). High-throughput mapping and clonal quantification of retroviral integration sites. *Methods Mol. Biol.* 1582, 127–141. doi: 10.1007/978-1-4939-6872-5_10
- Harrington, W. J. Jr., Sheremata, W. A., Snodgrass, S. R., Emerson, S., Phillips, S., and Berger, J. R. (1991). Tropical spastic paraparesis/HTLV-1-associated myelopathy (TSP/HAM): treatment with an anabolic steroid danazol. *AIDS Res. Hum. Retroviruses* 7, 1031–1034. doi: 10.1089/aid.1991.7.1031
- Hill, S. A., Lloyd, P. A., McDonald, S., Wykoff, J., and Derse, D. (2003). Susceptibility of human T cell leukemia virus type I to nucleoside reverse transcriptase inhibitors. *J. Infect. Dis.* 188, 424–427. doi: 10.1086/376531
- Holy, A. (2003). Phosphonomethoxyalkyl analogs of nucleotides. *Curr. Pharm. Des.* 9, 2567–2592. doi: 10.2174/1381612033453668
- Igakura, T., Stinchcombe, J. C., Goon, P. K., Taylor, G. P., Weber, J. N., Griffiths, G. M., et al. (2003). Spread of HTLV-1 between lymphocytes by virus-induced polarization of the cytoskeleton. *Science* 299, 1713–1716. doi: 10.1126/science.1080115
- Iwanaga, M., Watanabe, T., and Yamaguchi, K. (2012). Adult T-cell leukemia: a review of epidemiological evidence. *Front. Microbiol.* 3:322. doi: 10.3389/fmicb.2012.00322
- Izumo, S., Goto, I., Itoyama, Y., Okajima, T., Watanabe, S., Kuroda, Y., et al. (1996). Interferon- α is effective in HTLV-1-associated myelopathy: a multicenter, randomized, double-blind, controlled trial. *Neurology* 46, 1016–1021. doi: 10.1212/WNL.46.4.1016
- Jolly, C., and Sattentau, Q. J. (2005). Human immunodeficiency virus type 1 virological synapse formation in T cells requires lipid raft integrity. *J. Virol.* 79, 12088–12094. doi: 10.1128/JVI.79.18.12088-12094.2005
- Kannian, P., Fernandez, S., Jones, K. S., and Green, P. L. (2013). Human T lymphotropic virus type 1 SU residue 195 plays a role in determining the preferential CD4⁺ T cell immortalization/transformation tropism. *J. Virol.* 87, 9344–9352. doi: 10.1128/JVI.01079-13
- Kulkarni, A., Mateus, M., Thinnis, C. C., McCullagh, J. S., Schofield, C. J., Taylor, G. P., et al. (2017). Glucose metabolism and oxygen availability govern reactivation of the latent human retrovirus HTLV-1. *Cell Chem. Biol.* 24, 1377.e3–1387.e3. doi: 10.1016/j.chembiol.2017.08.016
- Lezin, A., Gillet, N., Olindo, S., Signate, A., Grandvaux, N., Verlaeten, O., et al. (2007). Histone deacetylase mediated transcriptional activation reduces proviral loads in HTLV-1 associated myelopathy/tropical spastic paraparesis patients. *Blood* 110, 3722–3728. doi: 10.1182/blood-2007-04-085076
- Macchi, B., Balestrieri, E., Ascolani, A., Hilburn, S., Martin, F., Mastino, A., et al. (2011). Susceptibility of primary HTLV-1 isolates from patients with HTLV-1-associated myelopathy to reverse transcriptase inhibitors. *Viruses* 3, 469–483. doi: 10.3390/v3050469
- Macchi, B., Faraoni, I., Zhang, J., Grelli, S., Favalli, C., Mastino, A., et al. (1997). AZT inhibits the transmission of human T cell leukaemia/lymphoma virus type I to adult peripheral blood mononuclear cells in vitro. *J. Gen. Virol.* 78(Pt 5), 1007–1016. doi: 10.1099/0022-1317-78-5-1007
- Mahieux, R., and Hermine, O. (2005). In vivo and in vitro treatment of HTLV-1 and HTLV-2 infected cells with arsenic trioxide and interferon- α . *Leuk. Lymphoma* 46, 347–355. doi: 10.1080/10428190400019966
- Maldarelli, F., Wu, X., Su, L., Simonetti, F. R., Shao, W., Hill, S., et al. (2014). HIV latency. Specific HIV integration sites are linked to clonal expansion and persistence of infected cells. *Science* 345, 179–183. doi: 10.1126/science.1254194
- Mansky, L. M. (2000). In vivo analysis of human T-cell leukemia virus type 1 reverse transcription accuracy. *J. Virol.* 74, 9525–9531. doi: 10.1128/JVI.74.20.9525-9531.2000
- Martin, F., Castro, H., Gabriel, C., Adonis, A., Fedina, A., Harrison, L., et al. (2012). Ciclosporin A proof of concept study in patients with active, progressive

- HTLV-1 associated myelopathy/tropical spastic paraparesis. *PLoS Negl. Trop. Dis.* 6:e1675. doi: 10.1371/journal.pntd.0001675
- Mazurov, D., Ilinskaya, A., Heidecker, G., Lloyd, P., and Derse, D. (2010). Quantitative comparison of HTLV-1 and HIV-1 cell-to-cell infection with new replication dependent vectors. *PLoS Pathog.* 6:e1000788. doi: 10.1371/journal.ppat.1000788
- Melamed, A., Laydon, D. J., Al Khatib, H., Rowan, A. G., Taylor, G. P., and Bangham, C. R. (2015). HTLV-1 drives vigorous clonal expansion of infected CD8⁺ T cells in natural infection. *Retrovirology* 12:91. doi: 10.1186/s12977-015-0221-1
- Merle, P., and Trepo, C. (2001). Therapeutic management of hepatitis B-related cirrhosis. *J. Viral Hepat.* 8, 391–399. doi: 10.1046/j.1365-2893.2001.00299.x
- Mitchell, M. S., Bodine, E. T., Hill, S., Princler, G., Lloyd, P., Mitsuya, H., et al. (2007). Phenotypic and genotypic comparisons of human T-cell leukemia virus type 1 reverse transcriptases from infected T-cell lines and patient samples. *J. Virol.* 81, 4422–4428. doi: 10.1128/JVI.02660-06
- Mitsuya, H., and Broder, S. (1987). Strategies for antiviral therapy in AIDS. *Nature* 325, 773–778. doi: 10.1038/325773a0
- Mitsuya, H., Weinhold, K. J., Furman, P. A., St Clair, M. H., Lehrman, S. N., Gallo, R. C., et al. (1985). 3'-Azido-3'-deoxythymidine (BW A509U): an antiviral agent that inhibits the infectivity and cytopathic effect of human T-lymphotropic virus type III/lymphadenopathy-associated virus in vitro. *Proc. Natl. Acad. Sci. U.S.A.* 82, 7096–7100. doi: 10.1073/pnas.82.20.7096
- Mitsuya, H., Yarchoan, R., and Broder, S. (1990). Molecular targets for AIDS therapy. *Science* 249, 1533–1544. doi: 10.1126/science.1699273
- Moens, B., Decanine, D., Menezes, S. M., Khouri, R., Silva-Santos, G., Lopez, G., et al. (2012). Ascorbic acid has superior ex vivo antiproliferative, cell death-inducing and immunomodulatory effects over IFN-alpha in HTLV-1-associated myelopathy. *PLoS Negl. Trop. Dis.* 6:e1729. doi: 10.1371/journal.pntd.0001729
- Nakagawa, M., Nakahara, K., Maruyama, Y., Kawabata, M., Higuchi, I., Kubota, H., et al. (1996). Therapeutic trials in 200 patients with HTLV-I-associated myelopathy/ tropical spastic paraparesis. *J. Neurovirol.* 2, 345–355. doi: 10.3109/13550289609146899
- Nasr, R., El Hajj, H., Kfoury, Y., de The, H., Hermine, O., and Bazarbachi, A. (2011). Controversies in targeted therapy of adult T cell leukemia/lymphoma: ON target or OFF target effects? *Viruses* 3, 750–769. doi: 10.3390/v3060750
- Oh, U., Yamano, Y., Mora, C. A., Ohayon, J., Bagnato, F., Butman, J. A., et al. (2005). Interferon-beta1a therapy in human T-lymphotropic virus type I-associated neurologic disease. *Ann. Neurol.* 57, 526–534. doi: 10.1002/ana.20429
- Olindo, S., Belrose, G., Gillet, N., Rodriguez, S., Boxus, M., Verlaeten, O., et al. (2011). Safety of long-term treatment of HAM/TSP patients with valproic acid. *Blood* 118, 6306–6309. doi: 10.1182/blood-2011-04-349910
- Olindo, S., Cabre, P., Lezin, A., Merle, H., Saint-Vil, M., Signate, A., et al. (2006). Natural history of human T-lymphotropic virus 1-associated myelopathy: a 14-year follow-up study. *Arch. Neurol.* 63, 1560–1566. doi: 10.1001/archneur.63.11.1560
- Pais-Correia, A. M., Sachse, M., Guadagnini, S., Robbiati, V., Lasserre, R., Gessain, A., et al. (2010). Biofilm-like extracellular viral assemblies mediate HTLV-1 cell-to-cell transmission at virological synapses. *Nat. Med.* 16, 83–89. doi: 10.1038/nm.2065
- Parella, F. J. Jr., Delaney, K. M., Moorman, A. C., Loveless, M. O., Fuhrer, J., Satten, G. A., et al. (1998). Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. HIV outpatient study investigators. *N. Engl. J. Med.* 338, 853–860. doi: 10.1056/NEJM199803263381301
- Perno, C. F., Yarchoan, R., Cooney, D. A., Hartman, N. R., Gartner, S., Popovic, M., et al. (1988). Inhibition of human immunodeficiency virus (HIV-1/HTLV-IIIb/L) replication in fresh and cultured human peripheral blood monocytes/macrophages by azidothymidine and related 2',3'-dideoxynucleosides. *J. Exp. Med.* 168, 1111–1125. doi: 10.1084/jem.168.3.1111
- Priet, S., Roux, L., Saez-Ayala, M., Ferron, F., Canard, B., and Alvarez, K. (2015). Enzymatic synthesis of acyclic nucleoside thiophosphonate diphosphates: effect of the alpha-phosphorus configuration on HIV-1 RT activity. *Antiviral Res.* 117, 122–131. doi: 10.1016/j.antiviral.2015.03.003
- Rafatpanah, H., Rezaee, A., Etemadi, M. M., Hosseini, R. F., Khorram, B., Afshar, L., et al. (2012). The impact of interferon-alpha treatment on clinical and immunovirological aspects of HTLV-1-associated myelopathy in northeast of Iran. *J. Neuroimmunol.* 250, 87–93. doi: 10.1016/j.jneuroim.2012.05.004
- Rahman, S., Manuel, S. L., Khan, Z. K., Wigdahl, B., Acheampong, E., Tangy, F., et al. (2010). Depletion of dendritic cells enhances susceptibility to cell-free infection of human T cell leukemia virus type 1 in CD11c-diphtheria toxin receptor transgenic mice. *J. Immunol.* 184, 5553–5561. doi: 10.4049/jimmunol.0903226
- Ray, A. S., Fordyce, M. W., and Hitchcock, M. J. (2016). Tenofovir alafenamide: a novel prodrug of tenofovir for the treatment of human immunodeficiency virus. *Antiviral Res.* 125, 63–70. doi: 10.1016/j.antiviral.2015.11.009
- Roux, L., Priet, S., Payrot, N., Weck, C., Fournier, M., Zoulim, F., et al. (2013). Ester prodrugs of acyclic nucleoside thiophosphonates compared to phosphonates: synthesis, antiviral activity and decomposition study. *Eur. J. Med. Chem.* 63, 869–881. doi: 10.1016/j.ejmech.2013.02.039
- Rudnicka, D., Feldmann, J., Porrot, F., Wietgreffe, S., Guadagnini, S., Prevost, M. C., et al. (2009). Simultaneous cell-to-cell transmission of human immunodeficiency virus to multiple targets through polysynapses. *J. Virol.* 83, 6234–6246. doi: 10.1128/JVI.00282-09
- Saito, M., Nakagawa, M., Kaseda, S., Matsuzaki, T., Jonosono, M., Eiraku, N., et al. (2004). Decreased human T lymphotropic virus type I (HTLV-I) provirus load and alteration in T cell phenotype after interferon-alpha therapy for HTLV-I-associated myelopathy/tropical spastic paraparesis. *J. Infect. Dis.* 189, 29–40. doi: 10.1086/380101
- Sax, P. E., Wohl, D., Yin, M. T., Post, F., DeJesus, E., Saag, M., et al. (2015). Tenofovir alafenamide versus tenofovir disoproxil fumarate, coformulated with elvitegravir, cobicistat, and emtricitabine, for initial treatment of HIV-1 infection: two randomised, double-blind, phase 3, non-inferiority trials. *Lancet* 385, 2606–2615. doi: 10.1016/S0140-6736(15)60616-X
- Sharma, P. L., Nurpeisov, V., Hernandez-Santiago, B., Beltran, T., and Schinazi, R. F. (2004). Nucleoside inhibitors of human immunodeficiency virus type 1 reverse transcriptase. *Curr. Top. Med. Chem.* 4, 895–919. doi: 10.2174/1568026043388844
- Sherer, N. M., Lehmann, M. J., Jimenez-Soto, L. F., Horensavitz, C., Pypaert, M., and Mothes, W. (2007). Retroviruses can establish filopodial bridges for efficient cell-to-cell transmission. *Nat. Cell Biol.* 9, 310–315. doi: 10.1038/ncb1544
- Sigal, A., Kim, J. T., Balazs, A. B., Dekel, E., Mayo, A., Milo, R., et al. (2011). Cell-to-cell spread of HIV permits ongoing replication despite antiretroviral therapy. *Nature* 477, 95–98. doi: 10.1038/nature10347
- Sowinski, S., Jolly, C., Berninghausen, O., Purbhoo, M. A., Chauveau, A., Kohler, K., et al. (2008). Membrane nanotubes physically connect T cells over long distances presenting a novel route for HIV-1 transmission. *Nat. Cell Biol.* 10, 211–219. doi: 10.1038/ncb1682
- Taylor, B. S., Olender, S. A., Tieu, H. V., and Wilkin, T. J. (2016). CROI 2016: advances in antiretroviral therapy. *Top. Antivir. Med.* 24, 59–81.
- Taylor, G. P., Goon, P., Furukawa, Y., Green, H., Barfield, A., Mosley, A., et al. (2006). Zidovudine plus lamivudine in human T-lymphotropic virus type-I-associated myelopathy: a randomised trial. *Retrovirology* 3:63. doi: 10.1186/1742-4690-3-63
- Taylor, G. P., and Matsuoka, M. (2005). Natural history of adult T-cell leukemia/lymphoma and approaches to therapy. *Oncogene* 24, 6047–6057. doi: 10.1038/sj.onc.1208979
- Thoulouze, M. I., and Alcover, A. (2011). Can viruses form biofilms? *Trends Microbiol.* 19, 257–262. doi: 10.1016/j.tim.2011.03.002
- Turpin, J., Alais, S., Marçais, A., Bruneau, J., Melamed, A., Gadot, N., et al. (2017). Whole body clonality analysis in an aggressive STLV-1 associated leukemia (ATLL) reveals an unexpected clonal complexity. *Cancer Lett.* 389, 78–85. doi: 10.1016/j.canlet.2016.12.022
- Utsunomiya, A., Choi, I., Chihara, D., and Seto, M. (2015). Recent advances in the treatment of adult T-cell leukemia-lymphomas. *Cancer Sci.* 106, 344–351. doi: 10.1111/cas.12617
- Utsunomiya, A., Miyazaki, Y., Takatsuka, Y., Hanada, S., Uozumi, K., Yashiki, S., et al. (2001). Improved outcome of adult T cell leukemia/lymphoma with allogeneic hematopoietic stem cell transplantation. *Bone Marrow Transplant.* 27, 15–20. doi: 10.1038/sj.bmt.1702731
- Valeri, V. W., Hryniewicz, A., Andresen, V., Jones, K., Fenizia, C., Bialuk, I., et al. (2010). Requirement of the human T-cell leukemia virus p12 and p30 products for infectivity of human dendritic cells and macaques but not rabbits. *Blood* 116, 3809–3817. doi: 10.1182/blood-2010-05-284141

- Watanabe, T. (2017). Adult T-cell leukemia: molecular basis for clonal expansion and transformation of HTLV-1-infected T cells. *Blood* 129, 1071–1081. doi: 10.1182/blood-2016-09-692574
- Wattel, E., Vartanian, J. P., Pannetier, C., and Wain-Hobson, S. (1995). Clonal expansion of human T-cell leukemia virus type I-infected cells in asymptomatic and symptomatic carriers without malignancy. *J. Virol.* 69, 2863–2868.
- Willems, L., Hasegawa, H., Accolla, R., Bangham, C., Bazarbachi, A., Bertazzoni, U., et al. (2017). Reducing the global burden of HTLV-1 infection: an agenda for research and action. *Antiviral Res.* 137, 41–48. doi: 10.1016/j.antiviral.2016.10.015
- Yakova, M., Lezin, A., Dantin, F., Lagathu, G., Olindo, S., Jean-Baptiste, G., et al. (2005). Increased proviral load in HTLV-1-infected patients with rheumatoid arthritis or connective tissue disease. *Retrovirology* 2:4. doi: 10.1186/1742-4690-2-4
- Yamamoto, S., Folks, T. M., and Heneine, W. (1996). Highly sensitive qualitative and quantitative detection of reverse transcriptase activity: optimization, validation, and comparative analysis with other detection systems. *J. Virol. Methods* 61, 135–143. doi: 10.1016/0166-0934(96)02078-2
- Yamano, Y., and Sato, T. (2012). Clinical pathophysiology of human T-lymphotropic virus-type 1-associated myelopathy/tropical spastic paraparesis. *Front. Microbiol.* 3:389. doi: 10.3389/fmicb.2012.00389
- Yarchoan, R., Klecker, R. W., Weinhold, K. J., Markham, P. D., Lyerly, H. K., Durack, D. T., et al. (1986). Administration of 3'-azido-3'-deoxythymidine, an inhibitor of HTLV-III/LAV replication, to patients with AIDS or AIDS-related complex. *Lancet* 1, 575–580. doi: 10.1016/S0140-6736(86)92808-4
- Zhang, J., Balestrieri, E., Grelli, S., Matteucci, C., Pagnini, V., D'Agostini, C., et al. (2001). Efficacy of 3'-azido 3'-deoxythymidine (AZT) in preventing HTLV-1 transmission to human cord blood mononuclear cells. *Virus Res.* 78, 67–78. doi: 10.1016/S0168-1702(01)00285-4
- Zhong, P., Agosto, L. M., Ilinskaya, A., Dorjbal, B., Truong, R., Derse, D., et al. (2013). Cell-to-cell transmission can overcome multiple donor and target cell barriers imposed on cell-free HIV. *PLoS One* 8:e53138. doi: 10.1371/journal.pone.0053138

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Pasquier, Alais, Roux, Thoulouze, Alvarez, Journou, Dutartre and Mahieux. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.