



Immunoproteomic Approach of Extracellular Antigens From *Paracoccidioides* Species Reveals Exclusive B-Cell Epitopes

André Luís Elias Moreira¹, Milton Adriano Pelli Oliveira², Lana O'Hara Souza Silva¹, Moisés Morais Inácio¹, Alexandre Melo Bailão¹, Juliana Alves Parente-Rocha¹, Vanessa Rafaela Milhomem Cruz-Leite¹, Juliano Domiraci Pაცეზ¹, Célia Maria de Almeida Soares¹, Simone Schneider Weber^{3,4} and Clayton Luiz Borges^{1*}

¹ Laboratório de Biologia Molecular, Instituto de Ciências Biológicas, Universidade Federal de Goiás, Goiânia, Brazil,

² Laboratório de Citocinas, Instituto de Patologia Tropical e Saúde Pública, Universidade Federal de Goiás, Goiânia, Brazil,

³ Laboratório de Biociência, Faculdade de Ciências Farmacêuticas, Alimentos e Nutrição, Universidade Federal de Mato Grosso do Sul, Campo Grande, Brazil, ⁴ Instituto de Ciências Exatas e Tecnologia, Universidade Federal do Amazonas, Itacoatiara, Brazil

OPEN ACCESS

Edited by:

José Roberto Mineo,
Federal University of Uberlândia, Brazil

Reviewed by:

Haroldo Oliveira,
Carlos Chagas Institute (ICC), Brazil
Marisa Mariel Fernandez,
Institute of Studies on Humoral
Immunity (IDEHU), Argentina

*Correspondence:

Clayton Luiz Borges
clayton@ufg.br;
clbluiz2@gmail.com

Specialty section:

This article was submitted to
Microbial Immunology,
a section of the journal
Frontiers in Microbiology

Received: 20 August 2019

Accepted: 09 December 2019


Published: 28 January 2020

Citation:

Moreira ALE, Oliveira MAP,
Silva LOS, Inácio MM, Bailão AM,
Parente-Rocha JA, Cruz-Leite VRM,
Pაცეზ JD, de Almeida Soares CM,
Weber SS and Borges CL (2020)
Immunoproteomic Approach
of Extracellular Antigens From
Paracoccidioides Species Reveals
Exclusive B-Cell Epitopes.
Front. Microbiol. 10:2968.
doi: 10.3389/fmicb.2019.02968

Fungi of the *Paracoccidioides* genus are the etiological agents of paracoccidioidomycosis (PCM), a systemic mycosis restricted to the countries of Latin America. Currently, the *Paracoccidioides* complex is represented by *Paracoccidioides lutzii*, *Paracoccidioides americana*, *Paracoccidioides brasiliensis*, *Paracoccidioides restrepiensis*, and *Paracoccidioides venezuelensis*. Even with advances in techniques used for diagnosing fungal diseases, high rates of false-positive results for PCM are still presented. Additionally, there is no efficient antigen that can be used to follow up the efficiency of patient treatment. The immunoproteomic is considered a powerful tool for the identification of antigens. In addition, antigens are molecules recognized by the immune system, which make them excellent targets for diagnostic testing of diseases caused by microorganisms. In this vein, we investigated which antigens are secreted by species representing *Paracoccidioides* complex to increase the spectrum of molecules that could be used for future diagnostic tests, patient follow-up, or PCM therapy. To identify the profile of antigens secreted by *Paracoccidioides* spp., immunoproteomic approaches were used combining immunoprecipitation, followed by antigen identification by nanoUPLC-MS^E-based proteomics. Consequently, it was possible to verify differences in the exoantigen profiles present among the studied species. Through a mass spectrometry approach, it was possible to identify 79 exoantigens in *Paracoccidioides* species. Using bioinformatics tools, two unique exoantigens in *P. lutzii* species were identified, as well as 44 epitopes exclusive to the *Paracoccidioides* complex and 12 unique antigenic sequences that can differentiate between *Paracoccidioides* species. Therefore, these results demonstrate that *Paracoccidioides* species have a range of B-cell epitopes exclusive to the complex as well as specific to each *Paracoccidioides* species. In addition, these analyses allowed us the identification of excellent biomarker candidates for epidemiology screening, diagnosis, patient follow-up, as well as new candidates for PCM therapy.

Keywords: *Paracoccidioides* spp., antigens secreted, epitopes, diagnostic, mass spectrometry



can be analyzed by optical microscopy. For a long time, immunodiagnosis was not recommended for specific diagnosis of the disease as it presented low specificity, being more commonly indicated for patient follow-up (Shikanai-Yasuda et al., 2006). Despite that, in the *Paracoccidioides*'s literature, there is a robust number of studies that have contributed to the improvement of PCM immunodiagnosis, with the development of serological tests that present high sensitivity and specificity (that reached 90–100%), as revised by Zancoppe-Oliveira et al. (2014). Even with these advances, the latest “Brazilian Guidelines for the Clinical Management of Paracoccidioidomycosis” still points to the fact that patients with histoplasmosis, aspergillosis, and leishmaniasis may have false-positive results for PCM. Additionally, determining which species of *Paracoccidioides* is the causative agent of the disease is still a restricted approach to some research centers, which limits the expansion of disease epidemiological data (Shikanai-Yasuda et al., 2017). Altogether, these facts point to the need for determination of specific *Paracoccidioides* antigens, for which the immunoproteomics approach is of great value.

The immunoproteomic is considered a powerful tool for the identification of antigens and has been used successfully to verify the response of antibodies to fungal antigens, enabling the identification of possible biomarkers, reactivity profile analysis, as well as characterizing molecules with potential for diagnosis, as demonstrated for several fungal pathogens such as *Aspergillus fumigatus* (Gautam et al., 2007; Virginio et al., 2014), *Candida albicans* (Bar et al., 2012; Lee et al., 2014; Pitarch et al., 2016), *Rhizopus oryzae* (Sircar et al., 2012), *Cryptococcus gattii* (Jobbins et al., 2010), *Coccidioides posadasii* (Tarcha et al., 2006), and *Sporothrix schenckii* (Rodrigues et al., 2015).

To determine specific antigens of *Paracoccidioides* spp., we employed the immunoproteomics approach for identification and characterization of secreted antigens (exoantigens) by representative species of the *Paracoccidioides* complex. With this, it was possible to verify differences in the exoantigen profiles among the species used in this study. Through mass spectrometry tools, it was possible to identify 79 antigens secreted by isolates of *Paracoccidioides*. In addition, bioinformatics tools were used to analyze which antigens are secreted by alternative or classic pathways, as well as which antigenic epitopes are present in these molecules. In addition, we identified which molecules are shared among the isolates under study and which molecules are unique to each representative of the *Paracoccidioides* complex. These results demonstrate that *Paracoccidioides* spp. have several proteins that can be recognized by the human immune system. In addition, these analyses allowed us the identification of new possible molecules for PCM diagnosis, patient follow-ups, and therapy.

MATERIALS AND METHODS

Culture and Maintenance of Microorganisms

Isolates from the *Paracoccidioides* complex, *Pb01* (*P. lutzii*), *Pb02* (*P. americana*), *Pb18* (*P. brasiliensis*), and *PbEpm83*

(*P. restrepiensis*), were used in this study. The *P. venezuelensis* species was not used during the study because it belongs to a monophyletic population restricted to Venezuela and cases of the disease caused by this species in other regions of Latin America have not been described (Theodoro et al., 2012; Turissini et al., 2017). The yeast phase was maintained *in vitro* at 36°C in a Fava Netto's semisolid medium for 72 h (Fava-Netto, 1955). The components of the Fava Netto's culture medium were 0.5% (w/v) yeast extract, 1% (w/v) peptone, 0.5% (w/v) meat extract, 0.3% (w/v) proteose-peptone, 0.5% (w/v) NaCl, 4% (w/v) glucose, 1% (w/v) agar, pH 7.2.

Production of Extracellular Extracts

Yeasts of each *Paracoccidioides* spp. were transferred to the Fava Netto's liquid medium and maintained at 36°C under agitation at 120 rpm for 96 h. After this period, cells were washed in phosphate buffered saline (PBS) and counted in Neubauer's chamber, and the viability was measured using Trypan blue dye. For inoculum, 5×10^6 viable cells/ml were used in liquid medium McVeigh-Morton (MMcM) (Restrepo and Jimenez, 1980) and incubated at 120 rpm for 24 h at 36°C. After incubation, the cells were centrifuged at $2,000 \times g$ for 20 min. The supernatant obtained after centrifugation was subjected to vacuum filtration on 0.22- μ m membranes (Millipore, United States). The filtered samples were concentrated using a 10-kDa ultrafiltration system (Amicon®, Millipore, United States) and then washed three times with PBS (Weber et al., 2012). The quantification of extracellular extracts was determined by the Bradford method (Bradford, 1976).

Cell Lysis Detection Assay Using Polymerase Chain Reaction (PCR)

The genomic DNA of the isolates was obtained according to the protocol described by Sambrook and Russell (2001). For the PCR reactions, the supernatant of sample secretomes (2 μ l) and the genomic DNA obtained were used. The reactions were performed in 40 cycles of 94°C for 30 s, 55°C for 30 s, and 72°C for 2 min. The 681 base pair PCR products were generated using oligonucleotides (sense 5'-GACATGCGTGATATCGACTTG-3' and antisense 5'-GTGCGCCATGCCATTCT-3') for the formamidase coding gene (GenBank accession number AY163575). PCR amplicons were detected by 1% (w/v) agarose gel electrophoresis using the GelRed intercalating agent (Biotium™, United States). To verify the sensitivity of PCR, a genomic DNA curve containing five dilutions (50 ng, 5 ng, 50 pg, 5 pg, and 1 pg) was constructed for each *Paracoccidioides* species, as described by Weber et al. (2012).

One-Dimensional Gel Electrophoresis (1-D SDS-PAGE)

The integrity of the extracellular extracts was verified by 1-D SDS-PAGE (Laemmli, 1970). Thirty micrograms of extracellular extracts from each isolate were prepared with sample buffer [0.2 M Tris-HCl, pH 6.8, 40% (w/v) SDS, 2% (w/v) β -mercaptoethanol, and bromophenol blue traces] and heated in the thermoblock at 100°C for 10 min. Subsequently, the samples

were submitted to 12% 1-D SDS-PAGE. As reference, the low-molecular-weight marker (GE Healthcare, United Kingdom) was used. Soon after this step, the gels were stained by Coomassie Blue. ImageQuant 300 (GE Healthcare, United Kingdom) was used to obtain the images.

Production of Polyclonal Antibodies

Balb/c mice from 8 to 12 weeks old were provided by bioterium of the Instituto de Patologia Tropical e Saúde Pública, da Universidade Federal de Goiás (IPTSP-UFG). The animals were handled according to Conselho Nacional de Controle de Experimentação Animal (CONCEA-MCTIC) and submitted to the Ethics Committee of Comissão de Ética no Uso de Animais (CEUA-UFG) under the registry number 030/2016. Four groups containing five animals each were immunized with extracellular extract samples to induce the production of polyclonal antibodies. Separately, each group was immunized using proteins secreted by *P. lutzii*, *P. americana*, *P. brasiliensis*, and *P. restrepiensis*. For the immunizations, 50 µg of extracellular extracts were used in three doses with intervals of 15 days using the complete Freund's adjuvant (Sigma-Aldrich, United States) for the first dose and the incomplete Freund's adjuvant (Sigma-Aldrich, United States) for the second and third immunizations. The negative control was obtained from immunized mice only with Freund's adjuvant as previously described. After the immunizations, the animals were euthanized, and the whole blood of the animals was collected. Whole blood was incubated at 37°C for 10 min, 4°C for 10 min, and centrifuged at 500 × g for 10 min. After centrifugation, the immunized sera containing the polyclonal antibodies were collected.

Immunoblotting Analysis

Immunoblotting was performed as described by de Oliveira et al. (2018) with some modifications. The extracellular extract samples (30 µg) were submitted to 1-D SDS-PAGE at 12% with subsequent transfer to nitrocellulose membranes. The primary antibodies (1:500) were incubated for 2 h, and the secondary antibody (1:20,000) (alkaline phosphatase-labeled mouse anti-IgG; Sigma-Aldrich, United Kingdom) was incubated for 2 h. The membranes were then washed twice with PBS and once with alkaline phosphatase buffer for 15 min. The chromogenic substrate solution for alkaline phosphatase containing BCIP (5-bromo-4-chloro-3-indolyl phosphate) and NBT (tetrazolium-nitroblue chloride) (Sigma-Aldrich, United Kingdom) was used.

Coupling of IgG to Protein G-Sepharose and Immunoprecipitation

Protein G-Sepharose® 4B resin (Invitrogen, United States) was used for affinity chromatography. The resin was washed three times with PBS and 1.5 ml of immunized or control serum at a concentration of 1.5 µg/µl. The resin was subsequently added to the affinity columns, incubated at 4°C for 30 min to promote the binding of the antibodies to protein G, which is covalently linked to Sepharose beads. After ligation, the columns

were washed three times with PBS. Then, 1.5 mg extracellular extracts from each *Paracoccidioides* species were added to the columns, maintained for 16 h at 4°C for the antigen-antibody binding. Afterward, the columns were washed three times using PBS, then the elution solution buffer [glycine 0.1 M, 0.02% sodium azide (w/v), pH 2.6] was added and incubated for 10 min. Subsequently, the supernatants containing antigens were obtained. The experimental procedure of immunoprecipitation is summarized in **Supplementary Figure S1**.

Tryptic Digestion of Exoantigens

Antigens of the *Paracoccidioides* species obtained by immunoprecipitation were subjected to tryptic digestion. Briefly, 10 µl of 50 mM ammonium bicarbonate pH 8.5 were added to 50 µg of the protein extracts and subsequently subjected to tryptic digestion as previously described (Murad and Rech, 2012; Bailao et al., 2014; de Curcio et al., 2017; de Oliveira et al., 2018). Then, 25 µl of 0.2% (w/v) RapiGest SF™ surfactant (Waters Corporation, United Kingdom) was added to the protein extracts and vortexed with subsequent incubation at 80°C for 15 min. After this incubation, 2.5 µl of 100 mM dithiothreitol (GE Healthcare, United States) was added and incubated at 60°C for 30 min. Then, 2.5 µl of 300 mM iodoacetamide (GE Healthcare, United States) was added and maintained at room temperature for 30 min. Subsequently, 10 µl of 50 mM trypsin (Promega, United States) solution was added and incubated for 16 h at 37°C. To precipitate the RapiGest SF™, 10 µl of 5% trifluoroacetic acid (Sigma-Aldrich, United Kingdom) was added and incubated for 90 min at 37°C. The samples were centrifuged at 20,000 × g for 30 min at 6°C, and the supernatants were collected. The supernatants were dried by using speed vacuum (Eppendorf, GER). All peptides obtained were resuspended in 80 µl of a solution containing 20 mM ammonium formate and Fosforilase MassPREP™ Digestion Standard (100 fmol/µl for *PbEpm83*, 400 fmol/µl for *Pb01*, 100 fmol/µl for *Pb02*, and 100 fmol/µl for *Pb18*) as the endogenous standard.

NanoUPLC-MS^E Analysis

After tryptic digestion of the purified exoantigens, the samples were submitted to nanoUPLC-MS^E. For the chromatographic analyses of peptides, the nanoACQUITY™ M-Class system (Waters Corporation, United Kingdom) was used. For the first dimension, the fragmentation of peptides was performed through a 5-µm UPLC M-Class Peptide BEH C18, 130 Å (300 µm × 50 mm; Waters Corporation, Milford, MA, United States). The fragmented peptides were then submitted to five different acetonitrile/0.1% (v/v) formic acid concentrations (F1, 11.4%; F2, 14.7%; F3, 17.4%; F4, 20.7%; and F5, 50%). Each fraction eluted was trapped in a 5-µm Acquity UPLC M-Class Symmetry C18 Trap Column, 100 Å (180 µm × 20 mm; Waters Corporation, Milford, MA, United States). For the second dimension, for the separation of peptides, an Acquity UPLC M-Class HSS T3 1.8 µm (75 µm × 150 mm) Analytical Column was used. For mass calibration, a solution of 200 fmol/µl of the precursor ion [Glu1]-Fibronopeptide B human (m/z 785.8426) (GFP) (Sigma-Aldrich, St. Louis, MO, United States) was used with a constant flow of 0.5 µl/min, and it was measured every

30 s. The eluted peptides were analyzed by Synapt G1 HDMSTM (Waters Corporation, United Kingdom) mass spectrometer. This device is equipped with nano-electrospray ion source and two mass analyzers, one quadrupole and one time-of-flight (Q-TOF) called nanoESI-Q-TOF (Waters Corporation, United Kingdom). This equipment operates in MS^E mode, alternating at 6 V (low power) and 40 V (high power) in each acquisition mode (0.4 s). Three biological replicates were performed for *P. lutzii* (Pb01), *P. americana* (Pb02), *P. brasiliensis* (Pb18), and *P. restrepiensis* (PbEmp83) samples.

Immunoproteomic NanoUPLC-MS^E Data Processing

After nanoUPLC-MS^E acquisition, the data processing was performed by using Protein Lynx Global Server version 3.0.2 software (Waters Corporation, United Kingdom). The obtained spectra were compared with sequences deposited in the database of Pb01 (*P. lutzii*), Pb03 (*P. americana*), Pb18 (*P. brasiliensis*), and PbCnh (*P. restrepiensis*)¹ to identify the peptides.

Bioinformatics Analyses

After data processing and protein identification, the identified proteins were subjected to *in silico* analyses. The identified proteins were functionally categorized based on UniProt² database. To predict the subcellular localization of the secreted antigens, WoLF PSORT software³ was employed. ExPasy software⁴ was used for isoelectric point prediction. For the prediction of proteins secreted by classical and non-classical pathways, SignalP 4.1⁵ and SecretomeP 2.0⁶ were, respectively, used. Regarding SignalP and SecretomeP software, values greater than or equal to 0.5 were considered indicative of secretion. For prediction of linear B-cell epitopes, we used the software BCPREDS⁷ and ABCpred⁸.

To analyze and generate the Venn diagram, we used OrthoVenn software⁹ and Draw Venn Diagram¹⁰. BLASTp (Basic Local Alignment Search Tool) software¹¹ and ClustalX 2.0 and R software¹² were used for the homology analysis of the identified exoantigens and heat map generation, respectively. ProtScale software¹³ was used for prediction of hydrophilic peptides. The three-dimensional (3-D) modeling of the peptides was performed using I-Tasser¹⁴ and the figures were visualized by using PyMol version 2.3 software¹⁵. To verify energy minimization and total

structural refinement of 3-D models, the GalaxyWEB¹⁶ and ModRefiner¹⁷ servers were used.

RESULTS

Monitoring of *Paracoccidioides* Species Cell Integrity Based on PCR Analysis

To verify if cell lysis would influence the profile of proteins secreted by the *P. lutzii* (Pb01), *P. americana* (Pb02), *P. brasiliensis* (Pb18), and *P. restrepiensis* (PbEpm83) species, the PCR technique was used, as described by Weber et al. (2012), with some modifications.

The formamidase gene of *Paracoccidioides* species was analyzed. The sensitivity of the technique was attested to by constructing a standard curve using genomic DNA (samples from 50-ng to 1-pg dilutions) and primers to formamidase gene, generating a 681 base pair amplicon, which was evaluated by agarose gel electrophoresis.

These results demonstrate that this assay was able to amplify extremely low amounts of genomic DNA (1 pg); nevertheless, no amplifications were detected in the supernatant of secretome samples of the isolates under study (Supplementary Figure S2). It was possible to observe that the extracellular extracts did not present PCR-detectable contamination and, accordingly, there are no detectable contaminants from the intracellular compartment since the gene that encodes the formamidase was not amplified in the samples analyzed during the PCR tests (Supplementary Figure S2). This demonstrates that even in the case of any cell lysis, it was undetectable by PCR and thus cannot influence the profile of samples during the proteomics analyses.

Profile of Extracellular Extracts Evaluated by 1-D SDS-PAGE

Extracellular extracts of *P. lutzii*, *P. americana*, *P. brasiliensis*, and *P. restrepiensis* were analyzed by 12% 1-D SDS-PAGE. After separation, the proteins of isolates depicted a molecular mass distribution between 103 and 16 kDa (Supplementary Figure S3).

High abundance of extracellular protein species by *P. lutzii*, *P. Americana*, and *P. restrepiensis* species was found in the 45-kDa range, demonstrating that the profile of the secreted proteins between the analyzed species presented some similarities (Supplementary Figure S3). Among these, the one with the highest abundance of proteins in the 45-kDa range was the *P. americana* species. The *P. brasiliensis* species did not present high expression of these proteins in the 45-kDa range (Supplementary Figure S3).

Exoantigen Profiles Assayed by Immunoblotting

Analyses were performed with the objective of obtaining the profile of exoantigens of *Paracoccidioides* species. For

¹ http://www.broadinstitute.org/annotation/genome/paracoccidioides_brasiliensis/Multihome.html

² <http://www.uniprot.org/>

³ <http://wolfsort.hgc.jp/>

⁴ https://web.expasy.org/compute_pi/

⁵ <http://www.cbs.dtu.dk/services/SignalP/>

⁶ <http://www.cbs.dtu.dk/services/SecretomeP/>

⁷ <http://omictools.com/bcpreds-tool>

⁸ <http://www.imtech.res.in/raghava/abcpred/>

⁹ www.bioinfogenome.net/OrthoVenn/

¹⁰ <http://bioinformatics.psb.ugent.be/webtools/Venn/>

¹¹ <http://blast.ncbi.nlm.nih.gov/>

¹² <http://cran.r-project.org>

¹³ <https://web.expasy.org/protscale/>

¹⁴ <http://zhanglab.ccmb.med.umich.edu/I-TASSER/>

¹⁵ <http://pymol.org/2/>

¹⁶ <http://galaxy.seoklab.org/>

¹⁷ <http://zhanglab.ccmb.med.umich.edu/ModRefiner/>

this, the extracellular extracts of the *P. lutzii*, *P. americana*, *P. brasiliensis*, and *P. restrepiensis* species were used. Initially, extracellular extracts were subjected to 1-D SDS-PAGE with subsequent transfer to nitrocellulose membranes. Subsequently, immunoblotting assays were performed using antibodies obtained from control and immunized mice (Figure 1).

These results demonstrated that immunoblottings probed with antibodies obtained from immunized mice only with Freund's adjuvant (controls) did not show any reactivity with the proteins of *Paracoccidioides* species. On the other hand, the serum obtained from mice immunized with secreted protein extracts from *P. lutzii*, *P. americana*, *P. brasiliensis*, and *P. restrepiensis* depicts reactivity against *Paracoccidioides* exoproteomes (Figure 1). In addition, by means of these data, it was possible to observe differences in antigenic profiles among the species representative of the *Paracoccidioides* complex. The antigenic profile of *P. lutzii* showed a wide range of antigenic proteins between 105 and 19 kDa. For the *P. americana* species, a range of antigens between 99 and 14 kDa were presented. Antigens in the range from 99 to 37 kDa were observed in samples from *P. brasiliensis* species. Regarding *P. restrepiensis*, it is possible to observe that the profile of the antigenic proteins varies between 105 and 21 kDa (Figure 1). As a result, it was possible to verify that the secreted protein extracts from *Paracoccidioides* species present antigenic molecules that induce the production of different antibodies.

Immunoprecipitation of *Paracoccidioides* Exoantigens

After analyzing the differences in the antigenic profiles between *Paracoccidioides* species, we used immunoprecipitation approach to purify the antibody-reactive exoantigens (Supplementary Figure S1) by using the secretome samples from *P. lutzii*,




FIGURE 1 | Analysis of the antigenic profile of proteins secreted by *Paracoccidioides* species through immunoblotting. Secretome samples of *Paracoccidioides* species probed against polyclonal antibodies obtained from animals immunized with the secretomes (Ab). Secretome samples of *Paracoccidioides* species probed against serum from mice immunized only with Freund's adjuvant (C-). kDa, kilodaltons.

P. americana, *P. brasiliensis*, and *P. restrepiensis* species. The immunoprecipitated samples were separated by 12% 1-D SDS-PAGE and demonstrate the presence of immunoreactive proteins, which proves the antigen-antibody binding (Figure 2). These profile differences are observed according to the molecular weight of the bands of proteins that are visualized in the electrophoresis (Figure 2). However, unlike what was seen regarding the immunized serum, the control sample showed no differences in protein profiles. The absence of immunoreactive proteins in the control sample demonstrates that there was no anti-secretome antibody production (Figure 2).

Mass Spectrometry Identification of *Paracoccidioides* Species Exoantigens

To identify the immunoprecipitated exoantigens, proteomic strategies were used. Initially, the samples obtained from the *Paracoccidioides* species were digested with trypsin and submitted to nanoUPLC-MS^E, resulting in the identification of 15 exoantigens for *P. lutzii* (Supplementary Table S1), 14 for *P. americana* (Supplementary Table S2), 17 for *P. brasiliensis* (Supplementary Table S3), and 33 for isolate *P. restrepiensis* (Supplementary Table S4). The immunoprecipitated exoantigens identified in both the immunized and the control serum were removed from the analysis.





FIGURE 2 | Electrophoretic profile of affinity chromatography. One-dimensional (1-D-SDS-PAGE) gel of proteins eluted from Sepharose beads-IgG. Immunoprecipitation using serum from immunized mice (At + Ab) and negative control (C-). Top row: Isolated representatives of *Paracoccidioides* species (*P. lutzii*, *P. americana*, *P. restrepiensis*, and *P. brasiliensis*). The arrows indicate the presence of exoantigens after purification when compared to the negative control. At, antigen; Ab, antibody; IgG HC, immunoglobulin G heavy chain; IgG LC, immunoglobulin G light chain. Electrophoresis stained with Coomassie blue.




The exoantigens identified in *Paracoccidioides* species were functionally classified in biological processes based on UniProt database. In *P. lutzii*, the most abundant functional class was metabolism (27%), followed by protein fate (20%), energy (6%), transcription (7%), protein synthesis (7%), cell cycle and DNA processing (6%), and hypothetical proteins (27%) (Figure 3A). The most abundant functional classes of *P. americana* after the functional classification were protein fate (43%), metabolism (22%), energy (14%), protein with binding function or cofactor requirement (7%), and hypothetical proteins (14%) (Figure 3B). For *P. brasiliensis*, metabolism (59%) was the most represented functional class, followed by protein fate (17%), energy (12%), and protein synthesis (12%) (Figure 3C). The functional classes of *P. restrepiensis* exoantigens were represented by metabolism (37%), protein fate (18%), protein synthesis (18%), cellular transport (3%), cell cycle and DNA processing (6%), cell rescue, defense, and virulence (3%), energy (6%), and hypothetical proteins (9%) (Figure 3D).

To identify signal peptides related to secretion pathways to the extracellular environment by alternative or classic routes, we used SecretomeP and SignalP software, respectively (Supplementary

Tables S1–S4). The data obtained by the SecretomeP software revealed that 53.33% (*P. lutzii*), 28.57% (*P. americana*), 23.52% (*P. brasiliensis*), and 39.39% (*P. restrepiensis*) of identified exoantigens were predicted for secretion by non-classical pathways (Figures 4A–D). Using the SignalP software, it was revealed that 6.66, 14.20, 11.76, and 03.03% exoantigens of *P. lutzii*, *P. americana*, *P. brasiliensis*, and *P. restrepiensis*, respectively, showed signal peptides for secretion by the classical route (Figures 4A–D).

When comparing the exoantigens identified in *Paracoccidioides* species by software OrthoVenn, seven proteins from *P. lutzii* were exclusives in comparison with other fungus species described in our study. Four, seven, and 17 exoantigens of *P. americana*, *P. brasiliensis*, and *P. restrepiensis* were exclusive to these isolates, respectively (Figure 5 and Supplementary Table S5). The proteins 2-methylcitrate synthase mitochondrial, malate dehydrogenase NAD-dependent, and HSP 70-like protein were identified in all species during the immunoproteomics analyses (Supplementary Table S6). Other proteins, such as endo-1,3(4)-beta-glucanase, dihydrolipoyl dehydrogenase, and glutamate carboxypeptidase, were identified in common



between *P. americana*, *P. brasiliensis*, and *P. restrepiensis* species (Supplementary Table S6).


Non-homologous Exoantigens of *Paracoccidioides* Species

A comparative analysis was carried out to verify which molecules were identified as exclusive and conserved among the studied *Paracoccidioides* species. BLASTp-NCBI algorithm was used, and the genome of *P. venezuelensis* was inserted to analyze its exoantigens homology within the *Paracoccidioides* complex. The exoantigens hypothetical protein (PAAG_05807) and glutamate-1-semialdehyde 2,1-aminomutase (PAAG_06925) were exclusive to *P. lutzii* when their homology was compared to the other species of *Paracoccidioides* (Table 1). In addition, the exoantigen hypothetical protein (PAAG_12630) of *P. lutzii* showed low homology to *P. brasiliensis* (72%) and *P. restrepiensis* (76%), but the molecule was not homologous to *P. americana* and *P. venezuelensis*. All other exoantigens analyzed have strong homology between the species of *Paracoccidioides*, demonstrating the presence of these conserved proteins among *Paracoccidioides* species (Supplementary Table S7). In addition, the homology of the *Paracoccidioides* spp. exoantigens was analyzed against proteins of other species and pathogens annotated in the NCBI genome, such as *Homo sapiens* (taxid:9606), *Cryptococcus neoformans* (taxid:5207), *Histoplasma capsulatum* (taxid:5037), *S. schenckii* (taxid:29908),

Coccidioides immitis (taxid:5501), *C. albicans* (taxid:5476), *A. fumigatus* (taxid:746128), *Mycobacterium tuberculosis* (taxid:1773), *Escherichia coli* (taxid:562), *Leishmania braziliensis* (taxid:5660), and *Streptococcus pneumoniae* (taxid:1313). During the analyses of *P. lutzii* exoantigens, the hypothetical protein (PAAG_12701) and hypothetical protein (PAAG_05807) were not homologous to any of the analyzed species, making these molecules excellent biomarkers to be used for the epidemiological monitoring and diagnosis of PCM caused by *P. lutzii*. The exoantigen hypothetical protein (PAAG_12630) presented homology of 92% to *H. capsulatum*, and glutamate-1-semialdehyde 2,1-aminomutase (PAAG_06925) presented low homology to *H. capsulatum* (62%), *C. immitis* (39%), and *C. neoformans* (39%) (Figure 6A). In *P. americana*, the exoantigen serine protease (PABG_00534) showed homology to *H. capsulatum* (74%) and *C. immitis* (51%). In addition, endo-1,3(4)-beta-glucanase (PABG_12341), although homologous to *H. capsulatum* (66%), *C. immitis* (64%), *A. fumigatus* (54%), *S. schenckii* (42%), and *C. albicans* (41%), presented low homology to the antigens of these species (Figure 6B). In *P. brasiliensis*, the exoantigen ribosome biogenesis protein BMS1 (PADG_00459) was homologous to *H. capsulatum* (66%), *C. immitis* (63%), and *A. fumigatus* (65%). In addition, PTH1 family peptidyl-tRNA hydrolase (PADG_05841) was present in several other pathogens, but exhibited low homology (Figure 6C). For the *P. restrepiensis* species, the exoantigen single-strand binding protein family (A0A1E2Y9T2) presented low homology to *H. capsulatum* (73%), *C. immitis* (33%), *A. fumigatus* (45%), *S. schenckii* (60%), and *E. coli* (39%). Another exoantigen in *P. restrepiensis* that showed low homology was oxidoreductase (A0A1D2JMM5). This molecule was homologous to *H. capsulatum* (61%), *C. immitis* (46%), *A. fumigatus* (45%), *S. schenckii* (28%), *C. neoformans* (35%), and *C. albicans* (24%) (Figure 6D). All comparisons of homology between *Paracoccidioides* spp. and other species of pathogens analyzed during the study are listed in Supplementary Table S8.

B-Cell Epitope Prediction of Exoantigens of *Paracoccidioides* Complex

Epitopes are specific regions of an antigen to which an antibody binds. These peptide regions are presented by MHC molecules to T lymphocytes (Abbas et al., 2008). Bioinformatics tools were used during analyses to predict and characterize which B-cell epitopes are present in the exoantigens identified by our immunoproteomics analysis. BCPREDs and ABCpred software were used together to verify if the predicted epitopes would be identified by both tools, leading to confirmation and strengthening of the experimentally obtained data. Initially, the proteins were submitted to BCPREDS software, where the epitopes with the highest score were selected (Supplementary Table S9). Predicted epitopes were also counted for each identified exoantigen, totaling 93 epitopes present in 15 proteins of *P. lutzii* and 158 epitopes present in 14 proteins of *P. americana*. For *P. brasiliensis*, 143 epitopes were predicted in 17 proteins, and there were 232 epitopes in 33 proteins



of *P. restrepiensis*. All of these predicted epitopes are listed in **Supplementary Table S9**.

Subsequently, the epitopes were analyzed using ABCpred software, where the sequences predicted by this tool were selected and compared to the sequences predicted by BCPREDS software. This analysis was performed to verify if these two software tools share the same results, strengthening our experimental data (**Supplementary Table S9**). Thus, all epitopes predicted by both tools were verified and showed high score values, except for the exoantigens PAAG_12701 of *P. lutzii*, which did not show epitope prediction in any software.

In addition, the amount of predicted epitopes for each identified exoantigen was also evaluated. Interestingly, one of the exoantigens that had the largest number of predicted epitopes was endo-1,3 (4) beta-glucanase with 23 epitopes. This antigen was identified in *P. americana*, *P. brasiliensis*, and *P. restrepiensis* species. Next, aminopeptidase 2 presented 18 epitopes and was identified in *P. americana* and *P. brasiliensis* species. Heat shock proteins such as HSP70 and HSP60 presented 17 and 10 predicted epitopes, respectively, and were identified in all species in this study. The glutamate carboxypeptidase was another

exoantigen that showed large numbers of predicted epitopes. A total of 11 epitopes were predicted in this molecule and identified in *P. brasiliensis* and *P. restrepiensis* species. Also, in *P. brasiliensis* and *P. restrepiensis*, the exoantigen dihydrolipoyl dehydrogenase was analyzed and presented nine predicted epitopes (**Supplementary Table S9**).

Specific B-Cell Epitopes of *Paracoccidioides* Species

After verifying which exoantigens were homologous to antigenic molecules in other organisms, the homology of the B lymphocyte epitopes predicted by BCPREDS and ABCpred software was verified to identify which antigenic sequences were conserved or not in other organisms. For this, all sequences of the epitopes listed in **Supplementary Table S9** were analyzed against homologous epitopes of other important fungi and pathogenic bacteria. Proteins with homology to human molecules were also analyzed to identify and avoid the process of cross-reactivity during future tests. Initially, species such as *C. neoformans*, *H. capsulatum*, *S. schenckii*, *C. immitis*, *C. albicans*, *A. fumigatus*, *M. tuberculosis*, *E. coli*, *S. pneumoniae*, *L. braziliensis*, and *H. sapiens* were selected. All sequences of proteins homologous to the exoantigens identified during the immunoproteomic were obtained from the UniProt database. ClustalX software was used to perform the alignment and all of the sequences were manually checked. After analysis of all identified epitopes, we selected only those that did not present homology with other analyzed organisms. In addition, the epitope topology in the protein structure is important. To check which epitopes were hydrophilic and can be present in the protein surface, we used ProtScale software, as performed by Ejazi et al. (2018). Thus, we selected only the hydrophilic, non-homologous to other related organisms and exclusive epitopes of the *Paracoccidioides* complex (**Table 2**). Therefore, five, 11, 10, and 18 epitopes of *P. lutzii*, *P. americana*, *P. brasiliensis*, and *P. restrepiensis*, respectively, were obtained (**Table 2**). In addition, the homology of all epitopes among *Paracoccidioides* species was evaluated to confirm its presence in all species of this fungus.


3-D Analysis of Exclusive Epitopes of *Paracoccidioides* Species

All exoantigens that showed exclusive epitopes common to all *Paracoccidioides* species (**Table 2**) had their 3-D structure built by using I-TASSER software and were refined by ModRefiner and GalaxyWEB algorithms. The position of each identified

TABLE 1 | Level of homology of the exoantigens of *Paracoccidioides* sp. between the species of the complex.

<i>P. lutzii</i> ^{1,2}	<i>P. americana</i> ³ (%)	<i>P. brasiliensis</i> ⁴ (%)	<i>P. restrepiensis</i> ⁵ (%)	<i>P. venezuelensis</i> ⁶ (%)
PAAG_05807 – Hypothetical protein	0	0	0	0
PAAG_12630 – Hypothetical protein	0	72	76	0
PAAG_06925 – Glutamate-1-semialdehyde 2,1-aminomutase	0	0	0	0

¹Access number of *P. lutzii* exoantigens. ²Exoantigens of *P. lutzii*. ³Exoantigens of *P. americana* used for homology analysis. ⁴Exoantigens of *P. brasiliensis* used for homology analysis. ⁵Exoantigens of *P. restrepiensis* used for homology analysis. ⁶Exoantigens of *P. venezuelensis* used for homology analysis. Values in % correspond to the homology levels of exoantigens.



epitope was evaluated in exoantigen models and whether they were present in the surface of the model or not, which possibly facilitates antigen-antibody binding. The epitopes used for the analysis are displayed in **Table 2**, totaling two, four, three, and five exoantigens for *P. lutzii*, *P. americana*, *P. brasiliensis*, and *P. restrepiensis*, respectively. For *P. lutzii*, the exoantigen nucleic acid-binding protein (PAAG_04814 – 3 epitopes) was evaluated. During the analysis of the 3-D model of this exoantigen, it was possible to verify that one epitope was on the surface (**Figure 7A**). In *P. americana*, molecular modeling of four exoantigens was made: serine protease (PABG_00534, one epitope), aminopeptidase 2 (PABG_00710, one epitope), endonuclease G, mitochondrial (PABG_04696, two epitopes), and endo-1,3(4)-beta-glucanase (PABG_12341, seven epitopes). After analysis, it was verified that all epitopes analyzed in the exoantigens were present externally to the analyzed models (**Figure 7B**). Regarding *P. brasiliensis* predicted models, the exoantigens aminopeptidase 2 (PADG_03149, one epitope), carbonic anhydrase (PADG_07674, two epitopes), and endo-1,3(4)-beta-glucanase (PADG_12370, seven epitopes) had their epitopes analyzed in the 3-D models obtained, and it was also possible to verify the presence of the antigens on the surface (**Figure 7C**). In *P. restrepiensis*, five exoantigens had their epitopes analyzed in generated 3-D models. Thus, the exoantigens oxidoreductase (A0A1D2JMM5, 3

epitopes), endonuclease G mitochondrial (A0A1E2XUM2, one epitope), endo-1,3(4)-beta-glucanase (A0A1E2Y5K2, seven epitopes), ATP synthase subunit D (A0A1E2YCS7, five epitopes), and carbonic anhydrase (A0A1E2YFV5, two epitopes) were present in the exposed regions in the 3-D models (**Figure 7D**).

Exclusive Epitopes of Members of the *Paracoccidioides* Complex

After identification of the exclusive epitopes of *Paracoccidioides*, the level of homology of these epitopes was investigated among species. Thus, the FASTA sequences were obtained of proteins of the *Paracoccidioides* species that were homologous to exoantigens identified during the immunoproteomics assays. For this analysis, homologous molecules of the species *P. lutzii*, *P. americana*, *P. brasiliensis*, *P. restrepiensis*, and *P. venezuelensis* were selected. ClustalX software was used to perform the alignment and the homology of these epitopes in the *Paracoccidioides* complex was manually analyzed. Consequently, non-homologous epitopes to other pathogens were identified that were exclusive to each species of *Paracoccidioides* (**Table 3**). In *P. lutzii*, seven epitopes from two proteins, hypothetical protein (PAAG_05807) and hypothetical protein (PAAG_12630),

TABLE 2 | B-cell epitopes hydrophilic, non-homologous, and exclusive of *Paracoccidioides* species.

<i>Paracoccidioides lutzii</i>	
Position ¹	Access number – Protein name
	PAAG_04814 – Nucleic acid-binding protein
77	VPISRPAPDVRPNETIYIGN
166	SRGSANPTPTRSNEPTRTLF
261	ARDSTMDRLRTAPSGSNETF
	PAAG_08322 – Hypothetical protein
33	HIDAREVINPGPVDSPKFFD
264	GTTNPDPSTIKDGSSSFASPP
<i>Paracoccidioides brasiliensis</i>	
Position ¹	Access number – Protein name
	PADG_03149 – Aminopeptidase 2
162	KIEVASNPSVTVNEDNETAT
	PADG_07674 – Carbonic anhydrase
11	STLRTQCTAITTPTRSSINK
43	NKPLPRFPHPCPTRRTISQM
	PADG_12370 – Endo-1,3(4)-beta-glucanase
39	TQFPPTGVRLQATRTPGVYH
82	EDHIGLPHPEYPHLDIVPSK
109	CYYKGNWCPGLPHAEENRIS
130	WLLDQSTGAPPKFATVTSSK
163	AAPSATSRSPPRETATVHPS
203	SSMSHQDIFEPIDKGIPIPSN
551	IGFGPFDPTNGREATLSNAT
<i>Paracoccidioides americana</i>	
Position ¹	Access number – Protein name
	PABG_00534 – Serine protease
81	AYSPIQPPAEVDDPVLVSWA
	PABG_00710 – Aminopeptidase 2
162	KIEVASNPSVTVNEDNETAT
	PABG_04696 – Endonuclease G, mitochondrial
34	LGTPPPSTRLPPQTTSQPTS
56	IPHPKITTPTDASPVPVG
	PABG_12341 – Endo-1,3(4)-beta-glucanase
39	TQFPPTGVRLQATRTPGVYH
82	EDHIGLPHPEYPHLDIVPSK
109	CYYKGNWCPGLPHAEENRIS
130	WLLDQSTGAPPKFATVTSSK
163	AAPSATSRSPPRETATAHPS
203	SSMSHQDIFEPIDKGIPIPSN
551	IGFGPFDPTNGREATLSNAT
<i>Paracoccidioides restrepiensis</i>	
Position ¹	Access number – Protein name
	A0A1D2JMM5 – Oxidoreductase
17	ACQISPQDTTKPPPTVKNRM
55	DVDEVQPTAEATARTENAD
257	HKEADEHRDQVNVHSPERLC
	A0A1E2XUM2 – Endonuclease G mitochondrial
34	LGTPPPSTRLPPQTTSQPTS
	A0A1E2Y5K2 – Endo-1,3 (4)-beta-glucanase
39	TQFPPTGVRLQATRTPGVYH

(Continued)

TABLE 2 | Continued

82	EDHIGLPHPEYPHLDIVPSK
109	CYYKGNWCPGLPHAEENRIS
130	WLLDQSTGAPPKFATVTSSK
163	AAPSATSRSPPRETATVHPS
203	SSMSHQDIFEPIDKGIPIPSN
551	IGFGPFDPTNGREATLSNAT
	A0A1E2YCS7 – ATP synthase subunit D
23	AADSESEAEAGAEEPSIPGG
44	AVTVHSRNEKKARKAIGKLG
80	VINQPDVYRSPSSNTWIIFG
150	EAVEKKDDEEDDGEEVDESG
185	SRKKAIRALKENDNDIVNSI
	A0A1E2YFV5 – Carbonic anhydrase
11	STLRTQCTAITTPTRSSINK
43	NKPLPRFPHPCPTRRTISQM

¹Position of the epitope in the exoantigen.

TABLE 3 | Exclusive epitopes of each species of *Paracoccidioides* complex.

<i>Paracoccidioides lutzii</i>	
Position ¹	Access number – Protein name
	PAAG_05807 – Hypothetical protein
25	NLWRHFNPDTAVVYNHMRL
70	KNSIRKVGFTPLDNSEPEETH
103	SITPIGQSGHHPNLTTSW
	PAAG_12630 – Hypothetical protein
43	EIGGEMRDTKGARENIRVK
73	GFRVRGGEGEGETVQEGS
94	AAGCGRAKGGKRGQRGRAK
146	PMGGVDGGTPGPFGRKREDE
<i>Paracoccidioides brasiliensis</i>	
Position ¹	Access number – Protein name
	PADG_08054 – Malate dehydrogenase NAD-dependent
7	LGASGGIGQVRGSMYTYNTI
28	KTSSTLWASTPKPNSNKMQP
	PADG_12370 – Endo-1,3(4)-beta-glucanase
637	NNYVWMKSDNVNQPDADTPQ
<i>Paracoccidioides restrepiensis</i>	
Position ¹	Access number – Protein name
	A0A1E2Y490 – ER lumen protein retaining receptor 2
42	PSDVHIDSPPEDEVKCLSF
140	RYEFSEVYKAPPYPPPPCIL

¹Position of the epitope in the exoantigen.

were identified (Table 3). In *P. brasiliensis*, three predicted epitopes were exclusive. Two epitopes were present in malate dehydrogenase NAD-dependent (PADG_08054) and one in endo-1,3(4)-beta-glucanase (PADG_12370) (Table 3). For *P. restrepiensis*, two epitopes were identified and belong to ER lumen protein retaining receptor 2 (A0A1E2Y490) (Table 3). Identification of these epitopes needs to be tested to be used during epidemiological monitoring of the disease, supporting the recognition of which species of *Paracoccidioides* is causing PCM.




FIGURE 7 | Prediction of the three-dimensional (3-D) model of the exoantigens of *Paracoccidioides* species by I-TASSER. The models were generated using the structure of the exoantigens identified during the immunoproteomics that presented non-homologous epitopes to other pathogenic organisms. The 3-D representation of predicted non-homologous epitopes was performed after structural refinement and energy minimization using ModRefiner and GalaxyWEB software. 3-D view of the non-homologous and hydrophilic epitopes predicted in (A) *P. lutzii*, (B) *P. americana*, (C) *P. brasiliensis*, and (D) *P. venezuelensis*.

DISCUSSION

In this work, several exoantigens were identified in the representatives of the *Paracoccidioides* complex: *P. lutzii*, *P. americana*, *P. brasiliensis*, and *P. restrepiensis*. These data provide the first large-scale identification of exoantigens of *Paracoccidioides* species. Some of these exoantigens have already been identified and described as antigens of pathogenic fungi, such as formamidase (Borges et al., 2005, 2010), dihydrolipoyl dehydrogenase (Landgraf et al., 2017), HSP60 (Izacc et al., 2001; Cunha et al., 2002), HSP70 (Bromuro et al., 1998; Bisio et al., 2005), serine protease (Parente et al., 2010; Lacerda Pigosso et al., 2017), and isocitrate lyase (Cruz et al., 2011; Silva et al., 2019). This demonstrates the robustness of method used in our immunoproteomics analyses.

Taking into account that our target is the characterization only of secreted antigens, we were able to produce and purify this group of proteins once no DNA from fungal cell lysis was detected in PCR-based assays. Similar results have already been observed in secretome analyses of *Paracoccidioides* sp. (Weber et al., 2012). Subsequently, we verified the profile of secretome obtained, where the molecular weight profiles were found to vary from 16 to 103 kDa, being compatible with the protein profile secreted by *Paracoccidioides* spp. (Weber et al., 2012; de Oliveira et al., 2018), which secreted a range of complex proteins of several sizes. Additionally, it was observed that *Paracoccidioides* species have differences in the exoantigen profiles, with an immunoreactive profile presenting differences between the studied species, although some exoantigens seem to be shared by all species. These antigenic differences may be related to the singular genetic compositions of each species analyzed, as reported (Pigosso et al., 2013; do Amaral et al., 2019). In *Sporothrix* spp., immunoblotting and immunoproteomics approach also have been demonstrating a dissimilar profile between species of complex, highlighting the importance of antigenic molecules as potential biomarkers of serological diagnosis and candidates for vaccine development (Rodrigues et al., 2015).

Currently, immunoprecipitation is a technique used for the purpose of purifying protein extracts, concentrating low-abundance proteins, analyzing protein-protein interactions, posttranslational modification studies, and can also be employed for antigens precipitation (Lund-Johansen et al., 2000). Employing this technique, we found the presence of immunoreactive bands when compared to control samples. Similar results were observed when immunoprecipitation was used for antigen purification with subsequent construction of new diagnostic tests for African trypanosomiasis (Sullivan et al., 2013). In addition, we found that the immunoreactive proteins depict a distinct pattern among the studied species, although some exoantigens were present in all *Paracoccidioides* species, as demonstrated by our proteomics analyses. It can be inferred that these differences can be related to the genetic background, mechanisms involved in the protein expression, as well as the secretion machinery of each *Paracoccidioides* species (Desjardins et al., 2011; Pigosso et al., 2013; de Oliveira et al., 2018; do Amaral et al., 2019; Peres da Silva et al., 2019). Thus, the use of

this technique for the purification of antigenic molecules has been shown to be highly effective for obtaining exoantigens from *Paracoccidioides* species.

Additionally, we characterized the *Paracoccidioides* identified antigenic proteins in functional classes. The distribution of secreted proteins in the functional classes was consistent with previous analyses of secretome of *Paracoccidioides* spp. (Weber et al., 2012; de Oliveira et al., 2018) and with proteins secreted in vesicles in other pathogenic fungi (Albuquerque et al., 2008; Rodrigues et al., 2008, 2014; Oliveira et al., 2010). Also, we verified that the protein profile obtained, including mass range and isoelectric point (IP), is consistent with other *Paracoccidioides* sp. secretomes (Weber et al., 2012). Most of the identified antigenic proteins depicted prediction of subcellular localization in the cytoplasm and mitochondria, which is in accordance with described secretomes. These proteins should perform non-classical functions at other cellular locations, being described as moonlighting proteins (Karkowska-Kuleta and Kozik, 2014). The secretion of these molecules seems to be related to virulence and/or host-pathogen interaction of *Paracoccidioides* spp. once it has been demonstrated that secretion of immunogenic molecules in other pathogens is implicated in virulence during host-pathogen interaction (Oliveira et al., 2010; Campbell et al., 2015; de Oliveira et al., 2018).

As expected, the identification of similar exoantigens among *Paracoccidioides* species was confirmed by mass spectrometry analysis. When comparing the identified proteins, some molecules, such as malate dehydrogenase NAD-dependent, HSP70-like protein, and 2-methylcitrate synthase, were present in all species. On the other hand, several exoantigens were identified as unique in the analyzed species. Previous analysis has shown constitutively and differentially expressed molecules between *Paracoccidioides* species (Pigosso et al., 2013; do Amaral et al., 2019). However, analyses of exclusive proteins among the isolates could possibly be used in the differential diagnosis of PCM, as well as epidemiology studies.

Homology analyses were used to identify unique exoantigens of *Paracoccidioides* species. These analyses were initially employed among the exoantigens identified by nanoUPLC-MS^E, resulting in the identification of exclusive exoantigens to *P. lutzii*. The identification of these unique exoantigens enables its use in the differential diagnosis of PCM caused by *P. lutzii*. Subsequently, all exoantigen sequences were compared to sequences of other fungal and bacterial pathogens to verify the homology of these molecules to other organisms. These homology analyses were also employed on human sequences (Virginio et al., 2014; Rodrigues et al., 2015), aiming to select non-homologous molecules and to avoid possible cross-reactivity in future diagnostic tests. However, a small percentage of the exoantigens analyzed presented low homology, being represented by *P. lutzii* and *P. americana* species. Because homology analysis is performed throughout the exoantigen sequence, we decided to check the homology singly analyzing only the homology of the predicted epitopes. Thus, we identified several exclusive and conserved epitopes in the *Paracoccidioides* complex, since these molecules are candidates for the specific diagnosis of PCM. Similar analyses have already

been performed in other works, where linear B-cell epitopes were employed for serological detection of several pathogenic organisms, such as *Borrelia miyamotoi* (Tokarz et al., 2019), *Toxoplasma gondii* (Gatkowska et al., 2019), *Treponema pallidum* (Ogden et al., 2019), *Leishmania* spp. (Bremer Hinckel et al., 2019; Vale et al., 2019), and several human viruses (Amrun et al., 2019; Mutsvunguma et al., 2019; Yao et al., 2019). This reinforces the importance of these analyses, demonstrating the applications of linear B-cell epitopes in the diagnosis of human pathogenic organisms.

Similar to findings for other fungi, the cell wall of *Paracoccidioides* spp. is composed of a network of glycoproteins and polysaccharides, which act to protect the cell from environmental stressors (De Groot et al., 2005). Recent studies have demonstrated that both the yeast and mycelial phases of *Paracoccidioides* spp. present different amounts of β -(1,3) glucans (Camacho and Nino-Vega, 2017). However, enzymes, such as endo β -(1,3)-glucanase, act by hydrolyzing the β -glucan chains, making these enzymes extremely important because they play a fundamental role in the morphological processes of the fungal cell (Adams, 2004). Endo-1,3(4)-beta-glucanase was found as an antigen secreted in yeasts of *P. americana*, *P. brasiliensis*, and *P. restrepiensis* during our analyses. This enzyme has also been detected as secreted in yeast proteomes of *H. capsulatum* (Garfoot et al., 2017). In addition, the endo-1,3(4)-beta-glucanase showed low homology when compared to other fungi and pathogenic bacteria. Despite the low homology, we decided to investigate the level of homology between the epitopes of this molecule. During analysis, seven specific B-cell epitopes presented as exclusive for the *Paracoccidioides* complex, being excellent candidates for synthesis of new diagnostic tests for PCM.

Interestingly, the carbonic anhydrase (CA) of *Paracoccidioides* spp. was first reported as a secreted antigen in the present study. This is a metalloenzyme and acts by catalyzing the reversible hydration of CO₂ to generate HCO₃⁻ (Supuran, 2008). Additionally, in *Paracoccidioides* sp., four coding genes for carbonic anhydrase have already been described, leading to the synthesis of CA1, CA2, and CA3 proteins for Class β and CA4 protein for Class α . In addition, this enzyme was positively regulated in *Paracoccidioides* yeast when compared to the mycelium phase of the fungus (Tomazett et al., 2016). In *Stenocarpella macrospora*, CA has already been described in the mitochondrial, cytoplasmic, and extracellular compartments (Elleuche and Poggeler, 2010). CA has been identified as an exoantigen in the current work and has been associated with the extracellular compartment. In addition, CA presented exclusive epitopes, which makes this molecule a potential candidate as a PCM biomarker. Additionally, this exoantigen can be characterized as a moonlighting protein because it has a metabolic function in the cytoplasm and, when exported to the extracellular environment, it is probably related to the virulence of the fungus. In addition, CA has been described and is related to morphogenesis and growth of *Paracoccidioides* sp. and *C. neoformans* (Bahn et al., 2005; Costa et al., 2007) during the infectious processes, suggesting the importance of this molecule during host-pathogen interactions.

Amino peptidase 2 was identified as an exoantigen during analyses. These molecules are characterized as exopeptidases and act as cleaving proteins in their N-terminal region (Gonzales and Robert-Baudouy, 1996). In *Aspergillus oryzae* mycelium secretome, amino peptidase was related to collagen degradation processes (Ding et al., 2014). Furthermore, in a recent study, amino peptidase demonstrated its antigenic potential during the expression of recombinant proteins of *Fasciola hepatica* when probed against antibodies from patients with fascioliasis (Mirzadeh et al., 2018) and against anti-amino peptidase monoclonal antibodies from *Taenia pisiformis* (Zhang et al., 2018). The antigenicity of this molecule was also attested to in eggs and adult worms of *Schistosoma mansoni* following mouse infection assays (Maggioli et al., 2018). In *Aspergillus terreus* mycelium exoantigens, this molecule showed a high antigenicity when probed against monoclonal antibodies, demonstrating its antigenic profile (Nayak et al., 2011). During analyses, these molecules were identified in *P. americana* and *P. brasiliensis*, where one epitope was characterized as exclusive of the *Paracoccidioides* complex. These characteristics indicate that this molecule may be an excellent candidate to be tested as a potential specific biomarker for *Paracoccidioides* species infections.

Therefore, further analyses are underway to investigate the potential of these molecules as biomarkers, which may be used in the design of a rapid diagnostic test for PCM, in patient treatment follow-ups, as well as to investigate the immunogenic potential of these molecules and their possible use in the therapy of this important neglected human systemic mycosis.

CONCLUSION

The differences among exoantigen expression profiles were essential to identify specific biomarkers for each *Paracoccidioides* species. Here, the use of an immunoproteomics approach allowed us the characterization of 15, 14, 17, and 33 exoantigens in *P. lutzii*, *P. americana*, *P. brasiliensis*, and *P. restrepiensis*, respectively. Additionally, bioinformatics analyses made it possible to perform a series of other characterizations, allowing us to verify the biological functions, homology of exoantigens, epitopes prediction, as well as strengthen the data accuracy. Two exoantigens were identified and described in this work as unique in *P. lutzii*. Also, 44 epitopes exclusive to the *Paracoccidioides* complex were mapped using bioinformatics. Regarding the epitopes that could be used for epidemiological monitoring of the disease, a total of 12 antigenic sequences were identified in exoantigens in the four *Paracoccidioides* species studied. Therefore, these findings demonstrate that *Paracoccidioides* species have a range of epitopes exclusive to the complex, as well as specific to each fungal species. Finally, the current serological tests could be improved further with combinations of two and more synthetic peptides, as well as modifications that can optimize these tests. It is important to highlight that the reactivity of these proposed epitopes is under evaluation and will be tested against clinical samples to provide a new tool for diagnosis, patient follow-ups, and/or PCM therapy.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation, to any qualified researcher.

ETHICS STATEMENT

The animal study was reviewed and approved by the Comissão de Ética no Uso de Animais (CEUA-UFG) under the registry number 030/2016.

AUTHOR CONTRIBUTIONS

AM, CB, MO, AB, and CA conceptualized and designed the study. AM, LS, VC-L, and JP worked on data acquisition. AM, CB, AB, SW, JP-R, MI, and CA analyzed and interpreted the data. AM, CB, and JP drafted the manuscript. CA, SW, JP, and CB critically revised the manuscript.

FUNDING

This work at Universidade Federal de Goiás was supported in part by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) – Finance Code 001, Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Instituto Nacional de Ciência e Tecnologia (INCT-IPH), Fundação de Amparo à Pesquisa do Estado de Goiás (FAPEG), and FUNDECT/DECIT-MS/CNPq/SES No. 03/2016-PPSUS-MS. We thank the Universidade Federal de Mato Grosso do Sul (UFMS) for the support of publication fee (EDITAL PROPP_RTR/UFMS 109/2019).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2019.02968/full#supplementary-material>

FIGURE S1 | Affinity chromatography obtained by immunoprecipitation. **(A)** Scheme used for purifying and obtaining the antigens using serum from immunized and control animals. **(B)** Scheme demonstrating the interaction of sepharose bead – G protein bound antibody and antibody antigen interaction.

FIGURE S2 | Confirmation by PCR of the extracts secreted by *Paracoccidioides* species. The sensitivity of the PCRs to the formamidase gene were obtained using the genomic DNA of *P. lutzii*, *P. americana*, *P. brasiliensis*, and *P. restrepiensis* in five dilutions ranging from 50 ng to 1 pg (Top panel). C–: Negative control (without genomic DNA). Sec: Cell-free secretome. bp: Base pairs according to the molecular weight marker. The amplicons obtained by PCR were evaluated by 1% agarose gel electrophoresis and stained by GelRed DNA intercalator.

FIGURE S3 | Integrity analysis of the proteome profile of the secreted isolates of *Paracoccidioides* species by one-dimensional electrophoresis. Twenty micrograms of samples were separated by one-dimensional electrophoresis

(SDS-PAGE) at 12%. *P. lutzii*, *P. americana*, *P. restrepiensis*, and *P. brasiliensis*. kDa, Kilodaltons. Coomassie Blue staining was used to visualize the proteins.

TABLE S1 | Identification of exoantigens recognized by serum from animals immunized with *P. lutzii* secretome.

TABLE S2 | Identification of exoantigens recognized by serum from animals immunized with *P. americana* secretome.

TABLE S3 | Identification of exoantigens recognized by serum from animals immunized with *P. brasiliensis* secretome.

TABLE S4 | Identification of exoantigens recognized by serum from animals immunized with *P. restrepiensis* secretome.

TABLE S5 | Exoantigens of *Paracoccidioides* species identified as exclusive during immunoproteomic analyzes.

TABLE S6 | Exoantigens of *Paracoccidioides* species identified as common during immunoproteomic analyzes.

TABLE S7 | Level of homology of the exoantigens of *Paracoccidioides* spp. between the species of the complex.

TABLE S8 | Heat map of homology levels of *Paracoccidioides* exoantigens against other organisms.

TABLE S9 | *In silico* mapping of B-cell epitopes of all exoantigens identified during the immunoproteome by BCPREDS and ABCpreds.

REFERENCES

- Abbas, A. K., Lichtman, A. H., and Pillai, S. (2008). *Imunologia Celular e Molecular*. Amsterdam: Elsevier.
- Adams, D. J. (2004). Fungal cell wall chitinases and glucanases. *Microbiology* 150(Pt 7), 2029–2035. doi: 10.1099/mic.0.26980-0
- Albuquerque, P. C., Nakayasu, E. S., Rodrigues, M. L., Frases, S., Casadevall, A., Zancope-Oliveira, R. M., et al. (2008). Vesicular transport in *Histoplasma capsulatum*: an effective mechanism for trans-cell wall transfer of proteins and lipids in ascomycetes. *Cell Microbiol.* 10, 1695–1710. doi: 10.1111/j.1462-5822.2008.01160.x
- Amrun, S. N., Yee, W. X., Abu Bakar, F., Lee, B., Kam, Y. W., Lum, F. M., et al. (2019). Novel differential linear B-cell epitopes to identify Zika and dengue virus infections in patients. *Clin. Transl. Immunol.* 8:e1066. doi: 10.1002/cti2.1066
- Bahn, Y. S., Cox, G. M., Perfect, J. R., and Heitman, J. (2005). Carbonic anhydrase and CO₂ sensing during *Cryptococcus neoformans* growth, differentiation, and virulence. *Curr. Biol.* 15, 2013–2020. doi: 10.1016/j.cub.2005.09.047
- Bailao, E. F., Parente, J. A., Pigosso, L. L., de Castro, K. P., Fonseca, F. L., Silva-Bailao, M. G., et al. (2014). Hemoglobin uptake by *Paracoccidioides* spp. is receptor-mediated. *PLoS Negl. Trop. Dis.* 8:e2856. doi: 10.1371/journal.pntd.0002856
- Bar, E., Gladiator, A., Bastidas, S., Roschitzki, B., Acha-Orbea, H., Oxenius, A., et al. (2012). A novel Th cell epitope of *Candida albicans* mediates protection from fungal infection. *J. Immunol.* 188, 5636–5643. doi: 10.4049/jimmunol.1200594
- Bisio, L. C., Silva, S. P., Pereira, I. S., Xavier, M. A., Venancio, E. J., Puccia, R., et al. (2005). A new *Paracoccidioides brasiliensis* 70-kDa heat shock protein reacts with sera from paracoccidioidomycosis patients. *Med. Mycol.* 43, 495–503. doi: 10.1080/13693780400029478
- Borges, C. L., Parente, J. A., Barbosa, M. S., Santa, J. M., Bao, S. N., de Sousa, M. V., et al. (2010). Detection of a homotetrameric structure and protein-protein interactions of *Paracoccidioides brasiliensis* formamidase lead to new functional insights. *FEMS Yeast Res.* 10, 104–113. doi: 10.1111/j.1567-1364.2009.00594.x
- Borges, C. L., Pereira, M., Felipe, M. S., de Faria, F. P., Gomez, F. J., Deepe, G. S. Jr., et al. (2005). The antigenic and catalytically active formamidase of *Paracoccidioides brasiliensis*: protein characterization, cDNA and gene cloning, heterologous expression and functional analysis of the recombinant protein. *Microbes Infect.* 7, 66–77. doi: 10.1016/j.micinf.2004.09.011
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248–254. doi: 10.1016/0003-2697(76)90527-3
- Bremer Hinkel, B. C., Marlais, T., Airs, S., Bhattacharyya, T., Imamura, H., Dujardin, J. C., et al. (2019). Refining wet lab experiments with *in silico* searches: a rational quest for diagnostic peptides in visceral leishmaniasis. *PLoS Negl. Trop. Dis.* 13:e0007353. doi: 10.1371/journal.pntd.0007353
- Bromuro, C., La Valle, R., Sandini, S., Urbani, F., Ausiello, C. M., Morelli, L., et al. (1998). A 70-kilodalton recombinant heat shock protein of *Candida albicans* is highly immunogenic and enhances systemic murine candidiasis. *Infect. Immun.* 66, 2154–2162. doi: 10.1128/iai.66.5.2154-2162.1998
- Brummer, E., Castaneda, E., and Restrepo, A. (1993). Paracoccidioidomycosis: an update. *Clin. Microbiol. Rev.* 6, 89–117. doi: 10.1128/cmr.6.2.89
- Camacho, E., and Nino-Vega, G. A. (2017). *Paracoccidioides* spp.: virulence factors and immune-evasion strategies. *Med. Inflamm.* 2017:5313691. doi: 10.1155/2017/5313691
- Campbell, L. T., Simonin, A. R., Chen, C., Ferdous, J., Padula, M. P., Harry, E., et al. (2015). *Cryptococcus* strains with different pathogenic potentials have diverse protein secretomes. *Eukaryot Cell* 14, 554–563. doi: 10.1128/EC.00052-15
- Costa, A. N., Benard, G., Albuquerque, A. L., Fujita, C. L., Magri, A. S., Salge, J. M., et al. (2013). The lung in paracoccidioidomycosis: new insights into old problems. *Clinics* 68, 441–448. doi: 10.6061/clinics/2013(04)02
- Costa, M., Borges, C. L., Bailao, A. M., Meirelles, G. V., Mendonca, Y. A., Dantas, S. F., et al. (2007). Transcriptome profiling of *Paracoccidioides brasiliensis* yeast-phase cells recovered from infected mice brings new insights into fungal response upon host interaction. *Microbiology* 153(Pt 12), 4194–4207. doi: 10.1099/mic.0.2007/009332-0
- Cruz, A. H., Brock, M., Zambuzzi-Carvalho, P. F., Santos-Silva, L. K., Troian, R. F., Goes, A. M., et al. (2011). Phosphorylation is the major mechanism regulating isocitrate lyase activity in *Paracoccidioides brasiliensis* yeast cells. *FEBS J.* 278, 2318–2332. doi: 10.1111/j.1742-4658.2011.08150.x
- Cunha, D. A., Zancope-Oliveira, R. M., Sueli, M., Felipe, S., Salem-Izacc, S. M., Deepe, G. S. Jr., et al. (2002). Heterologous expression, purification, and immunological reactivity of a recombinant HSP60 from *Paracoccidioides brasiliensis*. *Clin. Diagn. Lab. Immunol.* 9, 374–377. doi: 10.1128/cdli.9.2.374-377.2002
- de Curcio, J. S., Silva, M. G., Silva Bailao, M. G., Bao, S. N., Casaletti, L., Bailao, A. M., et al. (2017). Identification of membrane proteome of *Paracoccidioides lutzii* and its regulation by zinc. *Future Sci.* 3:FSO232. doi: 10.4155/fsoa-2017-0044
- De Groot, P. W., Ram, A. F., and Klis, F. M. (2005). Features and functions of covalently linked proteins in fungal cell walls. *Fungal Genet. Biol.* 42, 657–675. doi: 10.1016/j.fgb.2005.04.002
- de Oliveira, A. R., Oliveira, L. N., Chaves, E. G. A., Weber, S. S., Bailao, A. M., Parente-Rocha, J. A., et al. (2018). Characterization of extracellular proteins in members of the *Paracoccidioides* complex. *Fungal Biol.* 122, 738–751. doi: 10.1016/j.funbio.2018.04.001
- Desjardins, C. A., Champion, M. D., Holder, J. W., Muszewska, A., Goldberg, J., Bailao, A. M., et al. (2011). Comparative genomic analysis of human fungal pathogens causing paracoccidioidomycosis. *PLoS Genet.* 7:e1002345. doi: 10.1371/journal.pgen.1002345
- Ding, G. W., Zhou, N. D., and Tian, Y. P. (2014). Over-expression of a proline specific aminopeptidase from *Aspergillus oryzae* JN-412 and its application in collagen degradation. *Appl. Biochem. Biotechnol.* 173, 1765–1777. doi: 10.1007/s12010-014-0963-6
- do Amaral, C. C., Fernandes, G. F., Rodrigues, A. M., Burger, E., and de Camargo, Z. P. (2019). Proteomic analysis of *Paracoccidioides brasiliensis* complex isolates: correlation of the levels of differentially expressed proteins with *in vivo* virulence. *PLoS One* 14:e0218013. doi: 10.1371/journal.pone.0218013
- Ejazi, S. A., Bhattacharyya, A., Choudhury, S. T., Ghosh, S., Sabur, A., Pandey, K., et al. (2018). Immunoproteomic identification and characterization of *Leishmania* membrane proteins as non-invasive diagnostic candidates for clinical visceral leishmaniasis. *Sci. Rep.* 8:12110. doi: 10.1038/s41598-018-30546-y
- Elleuche, S., and Poggeler, S. (2010). Carbonic anhydrases in fungi. *Microbiology* 156(Pt 1), 23–29. doi: 10.1099/mic.0.032581-0
- Fava-Netto, C. (1955). Estudos quantitativos sobre a fixacao do complemento na blastomicose sulamericana com antigenos polissacaridicos. *Arq. Circ. Clin. Exp.* 18, 197–254.

- Franco, M., Sano, A., Kera, K., Nishimura, K., Takeo, K., and Miyaji, M. (1989). Chlamyospore formation by *Paracoccidioides brasiliensis* mycelial form. *Rev. Inst. Med. Trop. São Paulo* 31, 151–157. doi: 10.1590/s0036-46651989000300004
- Garfoot, A. L., Dearing, K. L., VanSchoiack, A. D., Wysocki, V. H., and Rappleye, C. A. (2017). Eng1 and Exg8 Are the Major beta-Glucanases Secreted by the Fungal Pathogen *Histoplasma capsulatum*. *J. Biol. Chem.* 292, 4801–4810. doi: 10.1074/jbc.M116.762104
- Gatkowska, J., Dzitko, K., Ferra, B. T., Holec-Gasior, L., Kawka, M., and Dziadek, B. (2019). The impact of the antigenic composition of chimeric proteins on their immunoprotective activity against chronic toxoplasmosis in mice. *Vaccines* 7:E154. doi: 10.3390/vaccines7040154
- Gautam, P., Sundaram, C. S., Madan, T., Gade, W. N., Shah, A., Sirdeshmukh, R., et al. (2007). Identification of novel allergens of *Aspergillus fumigatus* using immunoproteomics approach. *Clin. Exp. Allergy* 37, 1239–1249. doi: 10.1111/j.1365-2222.2007.02765.x
- Gonzales, T., and Robert-Baudouy, J. (1996). Bacterial aminopeptidases: properties and functions. *FEMS Microbiol. Rev.* 18, 319–344. doi: 10.1111/j.1574-6976.1996.tb00247.x
- Izacc, S. M., Gomez, F. J., Jesuino, R. S., Fonseca, C. A., Felipe, M. S., Deepe, G. S., et al. (2001). Molecular cloning, characterization and expression of the heat shock protein 60 gene from the human pathogenic fungus *Paracoccidioides brasiliensis*. *Med. Mycol.* 39, 445–455. doi: 10.1080/714031053
- Jobbins, S. E., Hill, C. J., D'Souza-Bassel, J. M., Padula, M. P., Herbert, B. R., and Krockenberger, M. B. (2010). Immunoproteomic approach to elucidating the pathogenesis of cryptococcosis caused by *Cryptococcus gattii*. *J. Proteome Res.* 9, 3832–3841. doi: 10.1021/pr100028t
- Karkowska-Kuleta, J., and Kozik, A. (2014). Moonlighting proteins as virulence factors of pathogenic fungi, parasitic protozoa and multicellular parasites. *Mol. Oral Microbiol.* 29, 270–283. doi: 10.1111/omi.12078
- Lacaz, C. D. S., Porto, E., Martins, J. E. C., Heins-Vaccari, E. M., and Melo, N. T. D. (2002). *Tratado de Micologia Médica Lacaz*. São Paulo: Sarvier.
- Lacerda Pigosso, L., Baeza, L. C., Vieira Tomazett, M., Batista Rodrigues, Faleiro, M., Brianezi Dignani de Moura, V. M., et al. (2017). *Paracoccidioides brasiliensis* presents metabolic reprogramming and secretes a serine proteinase during murine infection. *Virulence* 8, 1417–1434. doi: 10.1080/21505594.2017.1355660
- Laemml, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680–685. doi: 10.1038/227680a0
- Landgraf, T. N., Costa, M. V., Oliveira, A. F., Ribeiro, W. C., Panunto-Castelo, A., and Fernandes, F. F. (2017). Involvement of dihydrolipoyl dehydrogenase in the phagocytosis and killing of *Paracoccidioides brasiliensis* by macrophages. *Front. Microbiol.* 8:1803. doi: 10.3389/fmicb.2017.01803
- Lee, P. Y., Gam, L. H., Yong, V. C., Rosli, R., Ng, K. P., and Chong, P. P. (2014). Immunoproteomic analysis of antibody response to cell wall-associated proteins of *Candida tropicalis*. *J. Appl. Microbiol.* 117, 854–865. doi: 10.1111/jam.12562
- Lund-Johansen, F., Davis, K., Bishop, J., and de Waal Malefyt, R. (2000). Flow cytometric analysis of immunoprecipitates: high-throughput analysis of protein phosphorylation and protein-protein interactions. *Cytometry* 39, 250–259. doi: 10.1002/(sici)1097-0320(20000401)39:4<250::aid-cyto2>3.0.co;2-s
- Maggioli, G., Rinaldi, G., Giaudrone, I., Berasain, P., Tort, J. F., Brindley, P. J., et al. (2018). Expression, purification and characterization of two leucine aminopeptidases of the blood fluke *Schistosoma mansoni*. *Mol. Biochem. Parasitol.* 219, 17–23. doi: 10.1016/j.molbiopara.2017.11.006
- Martinez, R. (2015). Epidemiology of paracoccidioidomycosis. *Rev. Inst. Med. Trop. São Paulo* 57(Suppl. 19), 11–20. doi: 10.1590/S0036-46652015000700004
- Martinez, R. (2017). New trends in paracoccidioidomycosis epidemiology. *J. Fungi* 3:1. doi: 10.3390/jof3010001
- McEwen, J. G., Bedoya, V., Patino, M. M., Salazar, M. E., and Restrepo, A. (1987). Experimental murine paracoccidioidomycosis induced by the inhalation of conidia. *J. Med. Vet. Mycol.* 25, 165–175. doi: 10.1080/02681218780000231
- Mendes, R. P. (1994). The gamut of clinical manifestations. *Paracoccidioidomycosis* 2, 233–258. doi: 10.1201/9781351075329-16
- Mirzadeh, A., Yoosyfy, A., Kazemirad, E., Barati, Z., Golkar, M., Babaie, J., et al. (2018). Evaluation of a set of refolded recombinant antigens for serodiagnosis of human fascioliasis. *PLoS One* 13:e0203490. doi: 10.1371/journal.pone.0203490
- Morejon, K. M., Machado, A. A., and Martinez, R. (2009). Paracoccidioidomycosis in patients infected with and not infected with human immunodeficiency virus: a case-control study. *Am. J. Trop. Med. Hyg.* 80, 359–366. doi: 10.4269/ajtmh.2009.80.359
- Munoz, J. F., Farrer, R. A., Desjardins, C. A., Gallo, J. E., Sykes, S., Sakthikumar, S., et al. (2016). Genome diversity, recombination, and virulence across the major lineages of *Paracoccidioides*. *mSphere* 1:e213-16.
- Murad, A. M., and Rech, E. L. (2012). NanoUPLC-MSE proteomic data assessment of soybean seeds using the Uniprot database. *BMC Biotechnol.* 12:82. doi: 10.1186/1472-6750-12-82
- Mutsunguma, L. Z., Rodriguez, E., Escalante, G. M., Muniraju, M., Williams, J. C., Warden, C., et al. (2019). Identification of multiple potent neutralizing and non-neutralizing antibodies against Epstein-Barr virus gp350 protein with potential for clinical application and as reagents for mapping immunodominant epitopes. *Virology* 536, 1–15. doi: 10.1016/j.virol.2019.07.026
- Nayak, A. P., Green, B. J., Janotka, E., Hettick, J. M., Friend, S., Vesper, S. J., et al. (2011). Monoclonal antibodies to hyphal exoantigens derived from the opportunistic pathogen *Aspergillus terreus*. *Clin. Vaccine Immunol.* 18, 1568–1576. doi: 10.1128/CVI.05163-11
- Ogden, N. E., Kurnik, M., Parolo, C., and Plaxco, K. W. (2019). An electrochemical scaffold sensor for rapid syphilis diagnosis. *Analyst* 144, 5277–5283. doi: 10.1039/c9an00455f
- Oliveira, D. L., Freire-de-Lima, C. G., Nosanchuk, J. D., Casadevall, A., Rodrigues, M. L., and Nimrichter, L. (2010). Extracellular vesicles from *Cryptococcus neoformans* modulate macrophage functions. *Infect. Immunol.* 78, 1601–1609. doi: 10.1128/IAI.01171-09
- Parente, J. A., Salem-Izacc, S. M., Santana, J. M., Pereira, M., Borges, C. L., Bailao, A. M., et al. (2010). A secreted serine protease of *Paracoccidioides brasiliensis* and its interactions with fungal proteins. *BMC Microbiol.* 10:292. doi: 10.1186/1471-2180-10-292
- Peres da Silva, R., Longo, L. G. V., Cunha, J., Sobreira, T. J. P., Rodrigues, M. L., Faoro, H., et al. (2019). Comparison of the RNA content of extracellular vesicles derived from *Paracoccidioides brasiliensis* and *Paracoccidioides lutzii*. *Cells* 8:E765. doi: 10.3390/cells8070765
- Pigosso, L. L., Parente, A. F., Coelho, A. S., Silva, L. P., Borges, C. L., Bailao, A. M., et al. (2013). Comparative proteomics in the genus *Paracoccidioides*. *Fungal Genet. Biol.* 60, 87–100. doi: 10.1016/j.fgb.2013.07.008
- Pitarch, A., Nombela, C., and Gil, C. (2016). Seroproteotyping at the *Candida albicans* protein species level unveils an accurate molecular discriminator for candidemia. *J. Proteomics* 134, 144–162. doi: 10.1016/j.jprot.2015.10.022
- Queiroz-Telles, F. (1994). "Paracoccidioides brasiliensis: ultrastructural findings," in *Paracoccidioidomycosis*, ed. M. Franco, (Boca Raton: CRC Press), 27–48. doi: 10.1201/9781351075329-3
- Restrepo, A., and Jimenez, B. E. (1980). Growth of *Paracoccidioides brasiliensis* yeast phase in a chemically defined culture medium. *J. Clin. Microbiol.* 12, 279–281. doi: 10.1128/jcm.12.2.279-281.1980
- Restrepo, A., McEwen, J. G., and Castaneda, E. (2001). The habitat of *Paracoccidioides brasiliensis*: how far from solving the riddle? *Med. Mycol.* 39, 233–241. doi: 10.1080/714031028
- Rodrigues, A. M., Kubitschek-Barreira, P. H., Fernandes, G. F., de Almeida, S. R., Lopes-Bezerra, L. M., and de Camargo, Z. P. (2015). Immunoproteomic analysis reveals a convergent humoral response signature in the *Sporothrix schenckii* complex. *J. Proteomics* 115, 8–22. doi: 10.1016/j.jprot.2014.11.013
- Rodrigues, M. L., Nakayasu, E. S., Almeida, I. C., and Nimrichter, L. (2014). The impact of proteomics on the understanding of functions and biogenesis of fungal extracellular vesicles. *J. Proteomics* 97, 177–186. doi: 10.1016/j.jprot.2013.04.001
- Rodrigues, M. L., Nakayasu, E. S., Oliveira, D. L., Nimrichter, L., Nosanchuk, J. D., Almeida, I. C., et al. (2008). Extracellular vesicles produced by *Cryptococcus neoformans* contain protein components associated with virulence. *Eukaryot Cell* 7, 58–67.
- Sambrook, J., and Russell, D. (2001). *Molecular Cloning: A Laboratory Manual, the Third Edition*. New York, NY: Cold spring harbor laboratory press.
- Shikanai-Yasuda, M. A., Mendes, R. P., Colombo, A. L., Queiroz-Telles, F., Kono, A. S. G., Paniago, A. M. M., et al. (2017). Brazilian guidelines for the clinical management of paracoccidioidomycosis. *Rev. Soc. Bras. Med. Trop.* 50, 715–740. doi: 10.1590/0037-8682-0230-2017

- Shikanai-Yasuda, M. A., Mendes, R. P., Colombo, A. L., Telles, F. D. Q., Kono, A., Paniago, A. M. M., et al. (2018). II consenso brasileiro em paracoccidioidomicose - 2017. *Epidemiol. Serv. Saúde* 27:e0500001.
- Shikanai-Yasuda, M. A., Telles Filho Fde, Q., Mendes, R. P., Colombo, A. L., and Moretti, M. L. (2006). [Guidelines in paracoccidioidomycosis]. *Rev. Soc. Bras. Med. Trop.* 39, 297–310.
- Silva, K. S. F., Lima, R. M., Baeza, L. C., Lima, P. S., Cordeiro, T. M., Charneau, S., et al. (2019). Interactome of glyceraldehyde-3-phosphate dehydrogenase points to the existence of metabolons in *Paracoccidioides lutzii*. *Front. Microbiol.* 10:1537. doi: 10.3389/fmicb.2019.01537
- Sircar, G., Chakrabarti, H. S., Saha, B., and Gupta-Bhattacharya, S. (2012). Identification of aero-allergens from *Rhizopus oryzae*: an immunoproteomic approach. *J. Proteomics* 77, 455–468. doi: 10.1016/j.jprot.2012.09.023
- Sullivan, L., Wall, S. J., Carrington, M., and Ferguson, M. A. (2013). Proteomic selection of immunodiagnostic antigens for human African trypanosomiasis and generation of a prototype lateral flow immunodiagnostic device. *PLoS Negl. Trop. Dis.* 7:e2087. doi: 10.1371/journal.pntd.0002087
- Supuran, C. T. (2008). Carbonic anhydrases—an overview. *Curr. Pharm. Des.* 14, 603–614. doi: 10.2174/138161208783877884
- Tarcha, E. J., Basrur, V., Hung, C. Y., Gardner, M. J., and Cole, G. T. (2006). Multivalent recombinant protein vaccine against coccidioidomycosis. *Infect. Immunol.* 74, 5802–5813. doi: 10.1128/IAI.00961-06
- Teixeira, M. M., Theodoro, R. C., de Carvalho, M. J., Fernandes, L., Paes, H. C., Hahn, R. C., et al. (2009). Phylogenetic analysis reveals a high level of speciation in the *Paracoccidioides* genus. *Mol. Phylogenet. Evol.* 52, 273–283. doi: 10.1016/j.ympev.2009.04.005
- Teixeira, M. M., Theodoro, R. C., Nino-Vega, G., Bagagli, E., and Felipe, M. S. (2014). *Paracoccidioides* species complex: ecology, phylogeny, sexual reproduction, and virulence. *PLoS Pathog.* 10:e1004397. doi: 10.1371/journal.ppat.1004397
- Theodoro, R. C., Teixeira Mde, M., Felipe, M. S., Paduan Kdos, S., Ribolla, P. M., San-Blas, G., et al. (2012). Genus *paracoccidioides*: species recognition and biogeographic aspects. *PLoS One* 7:e37694. doi: 10.1371/journal.pone.0037694
- Tokarz, R., Tagliaferro, T., Caciula, A., Mishra, N., Thakkar, R., Chauhan, L. V., et al. (2019). Identification of immunoreactive linear epitopes of *Borrelia miyamotoi*. *Ticks Tick Borne Dis.* 11:101314. doi: 10.1016/j.ttbdis.2019.101314
- Tomazett, M. V., Zanoelo, F. F., Bailao, E. F., Bailao, A. M., Borges, C. L., and Soares, C. M. (2016). Molecular and biochemical characterization of carbonic anhydrases of *Paracoccidioides*. *Genet. Mol. Biol.* 39, 416–425. doi: 10.1590/1678-4685-GMB-2015-0213
- Turissini, D. A., Gomez, O. M., Teixeira, M. M., McEwen, J. G., and Matute, D. R. (2017). Species boundaries in the human pathogen *Paracoccidioides*. *Fungal Genet. Biol.* 106, 9–25. doi: 10.1016/j.fgb.2017.05.007
- Vale, D. L., Dias, D. S., Machado, A. S., Ribeiro, P. A. F., Lage, D. P., Costa, L. E., et al. (2019). Diagnostic evaluation of the amastin protein from *Leishmania infantum* in canine and human visceral leishmaniasis and immunogenicity in human cells derived from patients and healthy controls. *Diagn. Microbiol. Infect. Dis.* 95, 134–143. doi: 10.1016/j.diagmicrobio.2019.04.015
- Virginio, E. D., Kubitschek-Barreira, P. H., Batista, M. V., Schirmer, M. R., Abdelhay, E., Shikanai-Yasuda, M. A., et al. (2014). Immunoproteome of *Aspergillus fumigatus* using sera of patients with invasive aspergillosis. *Int. J. Mol. Sci.* 15, 14505–14530. doi: 10.3390/ijms150814505
- Wanke, B., and Londero, A. T. (1994). Epidemiology and paracoccidioidomycosis infection. *Paracoccidioidomycosis* 1, 109–120. doi: 10.1201/9781351075329-7
- Weber, S. S., Parente, A. F., Borges, C. L., Parente, J. A., Bailao, A. M., and de Almeida Soares, C. M. (2012). Analysis of the secretomes of *Paracoccidioides* mycelia and yeast cells. *PLoS One* 7:e52470. doi: 10.1371/journal.pone.0052470
- Yao, L., Chen, Y., Wang, X., Bi, Z., Xiao, Q., Lei, J., et al. (2019). Identification of antigenic epitopes in the haemagglutinin protein of H7 avian influenza virus. *Avian Pathol.* 49, 62–73. doi: 10.1080/03079457.2019.1666971
- Zancope-Oliveira, R. M., Pizzini, C. V., de Medeiros Muniz, M., do Valle, A. C. F., and Almeida-Paes, R. (2014). Diagnostic aspects of paracoccidioidomycosis. *Curr. Trop. Med. Rep.* 1, 111–118. doi: 10.1007/s40475-014-0022-y
- Zhang, S., Cai, X., Luo, X., Wang, S., Guo, A., Hou, J., et al. (2018). Molecular cloning and characterization of leucine aminopeptidase gene from *Taenia pisiformis*. *Exp. Parasitol.* 186, 1–9. doi: 10.1016/j.exppara.2018.01.012

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Moreira, Oliveira, Silva, Inácio, Bailão, Parente-Rocha, Cruz-Leite, Paccez, de Almeida Soares, Weber and Borges. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.