



Methanotrophs: Discoveries, Environmental Relevance, and a Perspective on Current and Future Applications

Simon Guerrero-Cruz^{1,2*}, Annika Vaksmaa³, Marcus A. Horn⁴, Helge Niemann^{3,5,6}, Maite Pijuan^{1,2} and Adrian Ho^{4,7*}

¹ Catalan Institute for Water Research (ICRA), Girona, Spain, ² Universitat de Girona, Girona, Spain, ³ Department of Marine Microbiology and Biogeochemistry, NIOZ Royal Netherlands Institute for Sea Research, 't Horntje, Netherlands, ⁴ Institute of Microbiology, Leibniz Universität Hannover, Hannover, Germany, ⁵ Department of Earth Sciences, Faculty of Geosciences, Utrecht University, Utrecht, Netherlands, ⁶ Centre for Arctic Gas Hydrate, Environment and Climate, Department of Geosciences, UiT the Arctic University of Norway, Tromsø, Norway, ⁷ Division of Applied Life Sciences, Gyeongsang National University, Jinju, South Korea

OPEN ACCESS

Edited by:

Sarahi L. Garcia,
Stockholm University, Sweden

Reviewed by:

Antti Juhani Rissanen,
Tampere University, Finland
Marcela Hernandez,
University of Southampton,
United Kingdom

*Correspondence:

Simon Guerrero-Cruz
guerrero.x5452@gmail.com
Adrian Ho
adrian.ho@ifmb.uni-hannover.de

Specialty section:

This article was submitted to
Microbiotechnology,
a section of the journal
Frontiers in Microbiology

Received: 08 March 2021

Accepted: 12 April 2021

Published: 14 May 2021

Citation:

Guerrero-Cruz S, Vaksmaa A,
Horn MA, Niemann H, Pijuan M and
Ho A (2021) Methanotrophs:
Discoveries, Environmental
Relevance, and a Perspective on
Current and Future Applications.
Front. Microbiol. 12:678057.
doi: 10.3389/fmicb.2021.678057

Methane is the final product of the anaerobic decomposition of organic matter. The conversion of organic matter to methane (methanogenesis) as a mechanism for energy conservation is exclusively attributed to the archaeal domain. Methane is oxidized by methanotrophic microorganisms using oxygen or alternative terminal electron acceptors. Aerobic methanotrophic bacteria belong to the phyla Proteobacteria and Verrucomicrobia, while anaerobic methane oxidation is also mediated by more recently discovered anaerobic methanotrophs with representatives in both the bacteria and the archaea domains. The anaerobic oxidation of methane is coupled to the reduction of nitrate, nitrite, iron, manganese, sulfate, and organic electron acceptors (e.g., humic substances) as terminal electron acceptors. This review highlights the relevance of methanotrophy in natural and anthropogenically influenced ecosystems, emphasizing the environmental conditions, distribution, function, co-existence, interactions, and the availability of electron acceptors that likely play a key role in regulating their function. A systematic overview of key aspects of ecology, physiology, metabolism, and genomics is crucial to understand the contribution of methanotrophs in the mitigation of methane efflux to the atmosphere. We give significance to the processes under microaerophilic and anaerobic conditions for both aerobic and anaerobic methane oxidizers. In the context of anthropogenically influenced ecosystems, we emphasize the current and potential future applications of methanotrophs from two different angles, namely methane mitigation in wastewater treatment through the application of anaerobic methanotrophs, and the biotechnological applications of aerobic methanotrophs in resource recovery from methane waste streams. Finally, we identify knowledge gaps that may lead to opportunities to harness further the biotechnological benefits of methanotrophs in methane mitigation and for the production of valuable bioproducts enabling a bio-based and circular economy.

Keywords: methanotrophy, application, methane, resource recovery, microbial ecology, climate change, anaerobic, circular economy

INTRODUCTION TO METHANE AND METHANE MICROBIOLOGY

Methane

Methane represents the most reduced form of carbon, is an important fuel for the global economy, and a greenhouse gas (GHG) in the atmosphere. It has a higher heat retentive capacity compared to CO₂, estimated at 34 times higher in a 100-year timeframe and 86 times higher in a period of 20 years [Intergovernmental Panel on Climate Change (IPCC), 2018]. Furthermore methane concentration has been continuously increasing to approximately 1857 ppb in 2018, 2.6 times higher than in the preindustrial times (Saunois et al., 2020) making it a critical environmental concern for climate change (Myhre et al., 2013; Etminan et al., 2016; Dean et al., 2018).

Methane originates from abiogenic, thermogenic and biogenic microbial sources. In this work, we focus on biogenic methanogenesis by methanogenic archaea; which liberate methane as the end-product from biological decomposition of organic matter as a mean of energy conservation (McInerney and Bryant, 1981; Conrad, 2009; Drake et al., 2009). Methanogenic archaea are present in diverse environments such as wetlands, peatlands, rice agriculture soil, livestock (enteric fermentation in ruminants), landfills, oceans and termites; representing approximately 70% of all sources of methane emissions to the atmosphere (Conrad, 1996; Kirschke et al., 2013). Among these methane sources, agriculture (e.g., rice paddy fields), livestock farming, waste and wastewater treatment (WWT), and fossil fuels are subject to significant human-driven intensification, linked to an expanding global population that demands for supply and in return produce more waste and increase emissions [Intergovernmental Panel on Climate Change (IPCC), 2015; Wolf et al., 2017].

Most recent estimations from data over the past decade (2008–2017) indicate a total average net methane production of 737 Tg CH₄ year⁻¹ from all sources (ranging from 594 to 881 Tg CH₄ year⁻¹) (Figure 1), whereas total terrestrial and aquatic sinks are estimated at an average of 625 Tg CH₄ year⁻¹ (ranging from 500 to 798 Tg CH₄ year⁻¹) leaving a positive net average balance of 112 Tg CH₄ year⁻¹ escaping into the atmosphere (Saunois et al., 2020). However, the net emissions may be underestimated because of methane emissions from the novel and environmentally damaging practice of shale gas extraction (fracking) (Umeozor et al., 2018; Howarth, 2019). Similarly, estimations of methane emissions from wastewater treatment are scarce and accurate determinations are limited (Nguyen et al., 2019) (in detail in Section “Fate of Methane and Application Potential of Anaerobic Methanotrophs in Wastewater Treatment”). In addition, methane estimations are likely to vary significantly with increasing global temperatures, making the estimations, control and mitigation of methane of crucial importance to control global warming (Collins et al., 2018). From all anthropogenic sources, livestock, waste and WWT, and rice cultivation represent approximately 57% of the total anthropogenic methane emissions indicating the need for better understanding and development of methane mitigation

strategies and the microbial processes that regulate the methane cycle (Conrad, 2009). Interestingly, methane has been postulated as the most cost-effective carbon feedstock for microbial chemical production, surpassing the cost competitiveness of chemical methanol or glucose from plant-derived sources. This opens opportunities for the application of methanotrophs to counteract methane emissions while producing valuable compounds derived from their metabolism (Section “Applications of Aerobic Methanotrophs in a Circular Economy”) (Comer et al., 2017). The following sections provide an overview of both methanogenesis and methanotrophy as microbial processes regulating methane fluxes worldwide.


Methanogenesis

During the degradation of organic matter, commonly referred as anaerobic digestion (AD), a limited number of archaeal groups are known to produce methane in the final stage of AD, namely methanogenesis, using CO₂, hydrogen, acetate, or methylated compounds as substrates coupled to the generation of energy for growth and carbon fixation (Liu, 2010; Sieber et al., 2010; Conrad, 2020). Despite substrate variation, methanogenesis is in all cases carried out through the action of the Methyl coenzyme M reductase (McrA) enzyme (Thauer, 1998).

Until very recently, only seven orders from the Euryarchaeota phylum were recognized as methanogenic archaea: Methanobacteriales, Methanocellales, Methanococcales, Methanomicrobiales, Methanomassiliicoccales, Methanosarcinales, and Methanopyrales (Liu, 2010; Enzmann et al., 2018). Today, it is known that putative methanogenic archaea are spread beyond the Euryarchaeota. Through the application of in-depth genomic sequencing the classification and diversity of methanogenic archaea has rapidly evolved. Currently, methanogenic archaea have representatives in 4 recognized superphyla: Euryarchaeota, TACK (Thaumarchaeota, Aigarchaeota, Crenarchaeota, and Korarchaeota), DPANN (Diapherotrites, Parvarchaeota, Aenigmarchaeota, Nanoarchaeota, and Nanohaloarchaeota), and the Asgard superphylum (Evans et al., 2015; Lloyd, 2015; Vanwonterghem et al., 2016; Castelle and Banfield, 2018; Zhou et al., 2018; Berghuis et al., 2019; Dombrowski et al., 2019; Macleod et al., 2019).

Methanotrophy

Methanotrophic microorganisms oxidize methane to harness energy under oxic and anoxic conditions using a range of diverse electron acceptors. Methanotrophy was initially reported in 1906 as an oxygen-dependent process and for almost a century, aerobic methanotrophy was considered as the only biological pathway to oxidize methane and that all methanotrophs belonged to the Proteobacteria phylum (Figure 2; Whittenbury et al., 1970; Hanson and Hanson, 1996). However, discoveries in the last two decades have broadened the view of methanotrophy with the identification of microorganisms outside the proteobacteria phylum and even in the archaea domain, capable of oxidizing methane anaerobically using alternative electron acceptors such as sulfate, nitrite, nitrate, iron and manganese (Figure 2). In



the next sections aerobic and anaerobic methanotrophy and metabolic pathways are described in chronological order.

Aerobic Methanotrophs

Aerobic methane oxidation is catalyzed by particulate and soluble methane monooxygenases (pMMO and sMMO, respectively). These enzymes oxidize methane to methanol, and subsequently, methanol dehydrogenases (MDHs) further oxidize methanol to formaldehyde (Keltjens et al., 2014). After formaldehyde, two more oxidation steps are involved and the intermediates are used for carbon assimilation, for a detailed description on methanotrophy see Dedysh and Knief (2018). Aerobic methanotrophs belong to the Gammaproteobacteria (Type I, with families *Methylococcaceae* and *Methylothermaceae*), Alphaproteobacteria (Type II, with families *Methylocystaceae* and *Beijerinckiaceae*), and the Verrucomicrobia phyla (family *Methylacidiphilaceae*) (Figure 2, based on van Teeseling et al., 2014; Dedysh and Knief, 2018). Verrucomicrobial methanotrophs are likely to play a critical role in methane oxidation in extreme environments where they were initially

discovered (i.e., geothermal and volcanic environments) (Dunfield et al., 2007; Pol et al., 2007; Islam et al., 2008), whereas proteobacterial methanotrophs are also found and active in environments with extreme conditions besides volcanic soils (i.e., acidic wetlands, thermal springs, thermal lakes, and peat soils) (Kolb and Horn, 2012; Islam et al., 2020, 2021; Kaupper et al., 2020; Hogendoorn et al., 2021). While most aerobic methanotrophs possess the pMMO, *Methylocella* and *Methyloferula* (Alphaproteobacteria, Beijerinckiaceae) harbor only the sMMO (Dedysh and Dunfield, 2016). Proteobacterial methanotrophs can be distinguished based on their taxonomy, carbon metabolism, morphology, and ecological life strategies among other features (Hanson and Hanson, 1996; Ho et al., 2013a; Knief, 2015; Dedysh and Knief, 2018; Hakobyan and Liesack, 2020). All pMMO-containing methanotrophs use copper as the catalytic metal in the first step of methane oxidation to methanol; in methanotrophs harboring both the pMMO and sMMO, copper suppresses the expression of the sMMO (Knapp et al., 2007; Trotsenko and Murrell, 2008) which require iron as they belong to a diverse enzymatic




FIGURE 2 | 120 years of methanotrophy depicted in a clock-based timeline, each hour on the clock represents one decade. The most relevant findings are depicted highlighting all recognized methanotrophic groups categorized according to their phylogenetic classification. Seven major milestones in methanotrophy are included, representing the most noteworthy microbial discoveries. (1) In brown, the discovery of aerobic canonical methanotrophy in 1906. On the right, all the known genera according to phylogenetic classification are listed (based on Dedysh and Knief, 2018). (2) In yellow, the discovery of sulfate-dependent anaerobic oxidation of methane (S-dAOM). This process is performed by three defined groups of AN-aerobic ME-thanotrophic archaea (ANME) within the Euryarchaeota phylum. (3) In red, aerobic methanotrophy within the Verrucomicrobia phylum. (4) In blue, nitrate-dependent anaerobic oxidation of methane (N-dAOM) by *Methylomirabilis* bacteria. (5–7) In green, nitrate-, iron-, and manganese-dependent anaerobic oxidation (N-dAOM and Metal-dAOM) of methane by diverse cultured species of the *Methanoperedenaceae* family.

family of di-iron carboxylate enzymes (sDIMO) (Crombie and Murrell, 2014; Nichol et al., 2019). Strikingly, Verrucomicrobial methanotrophs and specific proteobacterial methanotrophs (e.g., *Methylocella*) rely on lanthanide metals for the second step of methanol oxidation as the catalytic driving force in MDHs from the XoxF family (Pol et al., 2014; Farhan Ul Haque et al., 2019; Smith and Wrighton, 2019). For a detailed review on the diversity of sDIMO methanotrophs, see Farhan Ul Haque et al. (2020).

Anaerobic Methanotrophs

The Anaerobic Oxidation of Methane (AOM) was first observed in marine sediments where methane consumption was linked to the presence of sulfate (Knittel and Boetius, 2009). However, the biology of this process was never fully characterized likely due to the lack of advanced cultivation and genomic technologies; furthermore, archaea were not yet discovered at the time thus a potential consortium of bacteria was hypothesized (Reeburgh, 1976; Reeburgh and Heggie, 1977; Zehnder and Brock, 1980). Two decades later, with the reclassification of living organisms


(Woese and Fox, 1977; Fox et al., 1977), archaea were identified as the most likely cause for methane consumption based on carbon isotope studies (Hinrichs et al., 1999). Shortly after, the microbiology behind this process was described as sulfate-dependent anaerobic methane oxidation process (S-dAOM) by a microbial consortium consisting of sulfate-reducing bacteria (SRB) and methanotrophic archaea (Boetius et al., 2000). The identity of the Anaerobic Methanotrophic archaea (ANME) was confirmed soon after, currently classified as ANME 1, 2, 3 (Figure 2; Hinrichs et al., 1999; Boetius et al., 2000; Orphan et al., 2001; Niemann et al., 2006). The mechanism behind the oxidation of methane was identified as a reverse reaction of the canonical final step in the methanogenesis pathway (Krüger et al., 2003; Hallam et al., 2004). Interestingly, the reversibility of the last step of methanogenesis can also be encountered in canonical methanogens where iron minerals are known to stimulate methanotrophy in methanogens (Bar-Or et al., 2017). Furthermore, model methanogens such as *Methanosarcina barkeri* are known to oxidize methane through

extracellular electron transfer (EET), referred as electrogenic anaerobic methane oxidation (Yu et al., 2021).

With the identification of archaea as the responsible microorganisms for the S-dAOM process in marine ecosystems, it became evident that methanotrophy is possible with alternative electron acceptors besides oxygen. In freshwater ecosystems, nitrogen oxyanions such as nitrite and nitrate are more abundant than sulfate, and thermodynamically suitable to support other microbial processes using methane as the energy source, a process commonly referred as nitrogen-dependent anaerobic methane oxidation (N-dAOM) (Reeburgh and Heggie, 1977; In 't Zandt et al., 2018a). After initial indications of methane oxidation occurring under anoxic conditions coupled to denitrification in WWT (Sollo et al., 1976), a microbial consortium of bacteria from the novel NC10 phylum (Holmes et al., 2001) and archaea from the former ANME-2d group performing this process (currently *Methanoperedenaceae*), was identified from anoxic sediments in agricultural soil (Raghoebarsing et al., 2006). The bacterium from the NC10 phylum, "*Candidatus Methyloirabilis oxyfera*" (Figure 2) was identified as the microorganism capable of oxidizing methane at the expense of nitrite as electron acceptor (nitrite-dAOM) (Ettwig et al., 2008). "*Ca. M. oxyfera*" harbored all the genes necessary to perform conventional aerobic methane oxidation via pMMO and MDH enzymes (Ettwig et al., 2010). The presence of these genes was inconsistent for a bacterium of anaerobic nature and highly susceptible to oxygen, it was then postulated that oxygen was formed from the dismutation of nitric oxide produced from nitrite reduction, this oxygen produced intracellularly, would then be used to oxidize methane via an intra-oxygenic pathway (Figure 3) (Wu et al., 2011; Ettwig et al., 2012). To date, the accurate characterization of

the putative nitric oxide dismutase enzyme remains elusive and limited to the detection of these enzymes in environmental samples and candidate genes in enrichment cultures (Wu et al., 2015; Zhu et al., 2017, 2020; Versantvoort et al., 2018). Furthermore, the relevance of lanthanides as an essential metal for the oxidation of methanol, extends into the NC10 phylum where lanthanide-dependent MDH enzymes are seemingly more important and widespread than previously thought (Wu et al., 2015). To date, all bacteria capable of performing nitrite-dAOM with a potential intra-oxygenic pathway belong to the *Methyloirabilis* genus of the NC10 phylum (Versantvoort et al., 2018). *Methyloirabilis* bacteria are widespread in diverse environments, including freshwater and marine ecosystems (He et al., 2015; Padilla et al., 2016; Vaksmaa et al., 2016; Graf et al., 2018; Thamdrup et al., 2019), and in engineered ecosystems such as wastewater treatment plants (WWTPs, Section "Fate of Methane and Application Potential of Anaerobic Methanotrophs in Wastewater Treatment") (Ettwig et al., 2009).

The archaeal partner was identified as a member of the new *Methanoperedenaceae* family (formerly ANME-2d) from the Euryarchaeota phylum and named "*Candidatus Methanoperedens nitroreducens*," capable of coupling the reduction of nitrate to ammonia with nitrite as intermediate, to the oxidation of methane (nitrate-dAOM) via the reverse last step from methanogenesis (Haroon et al., 2013). The intermediate production of nitrite during the reduction of nitrate to ammonia, has demonstrated to be the common link that enables both "*Ca. M. nitroreducens*" and *Methyloirabilis* bacteria to coexist in laboratory enrichments and their simultaneous detection from environmental samples (Figure 3; Vaksmaa et al., 2017; Gambelli et al., 2018). Furthermore, "*Ca. M. nitroreducens*" has been



demonstrated to be capable of performing methane oxidation using iron and manganese (ferrihydrite and birnessite) in the absence of nitrate (Ettwig et al., 2016) while other novel members of the *Methanoperedenaceae* family “*Ca. M. ferrireducens*,” “*Ca. M. manganireducens*,” and “*Ca. M. manganicus*” (Figure 2), do so exclusively with iron and manganese lacking the necessary genetic traits to use nitrate as terminal electron acceptor (Cai et al., 2018b; Leu et al., 2020a). In all members of the *Methanoperedenaceae* family, a complex and abundant network of cytochromes for direct electron transfer is likely to enable methane oxidation (Leu et al., 2020b). After an overview on methanogenesis and methanotrophy, the next section focuses on the environmental prevalence and relevance of the methanotrophs.

ENVIRONMENTAL PREVALENCE AND RELEVANCE OF METHANOTROPHS UNDER MICROAEROPHILIC AND ANAEROBIC CONDITIONS

The ubiquity of (an)aerobic methanotrophs is attributable to their metabolic versatility, thriving in broad temperature (4–95°C) and pH (1–8) ranges (Trotsenko and Khmelenina, 2002; Pol et al., 2007). Particularly, aerobic methanotrophs have been the focus of previous reviews covering environmental detection, prevalence, and relevance in widespread habitats, as well as potential applications (Semrau et al., 2010, 2018; Ho et al., 2013a; Knief, 2015; Strong et al., 2015; Kwon et al., 2019; Kalyuzhnaya et al., 2020). However, more recent evidence indicates that aerobic methanotrophs can thrive under oxygen-limitation, and even under anoxic conditions challenging the conventional view of aerobic methanotrophy. Here, we focused on this aspect to provide a novel perspective of aerobic methanotrophy in environments which may not usually support their physiological requirements. Furthermore, we discuss recent aspects of microbial ecology and novel physiological aspects of anaerobic methanotrophs.

Aerobic Methanotrophs Under Micro-Oxic Conditions

The oxidation of methane to methanol by the MMO-containing methanotrophs requires molecular oxygen serving as the electron acceptor (Ross and Rosenzweig, 2017). Therefore, aerobic methanotrophs are typically active at oxic-anoxic interfaces where methane and oxygen gradients overlap (e.g., soil-overlying water interface in rice paddies and peatlands; Reim et al., 2012; Kaupper et al., 2021). However, the methane-oxygen counter gradient is dynamic, and oxygen could be rapidly depleted by oxygen-respiring organisms. To facilitate methane oxidation under micro-oxic conditions, some aerobic gammaproteobacterial methanotrophs (e.g., *Methylobacter*) may possess high-affinity cytochromes (e.g., cytochrome *bd* ubiquinol-oxidoreductase), enabling methane oxidation at low oxygen levels in the nM range (Skennerton et al., 2015;

Smith et al., 2018; Smith and Wrighton, 2019). The pMMO-containing aerobic methanotrophs also possess hemerythrin, a non-heme oxygen-binding protein acting as an oxygen scavenging transporter to the pMMO which appears to be a widespread feature among pMMO-containing aerobic methanotrophs (Kao et al., 2008; Rahalkar and Bahulikar, 2018). Hemerythrins, initially discovered in *Methylococcus capsulatus* Bath (gammaproteobacterial methanotroph; Kao et al., 2004), have since been detected in other aerobic methanotrophs (e.g., *Methylomicrobium alcaliphilum*; Kalyuzhnaya et al., 2013). The correlation between the expression of pMMO and hemerythrin (Chen et al., 2012) and the increase in hemerythrin expression under oxygen-limited conditions has been reported (Kalyuzhnaya et al., 2013; Yu et al., 2020). Strikingly, some gammaproteobacterial methanotrophs (e.g., *Methyloterricola*, *Methylomagnum*, *Methylomonas*, *Methylobacter*, and *Methylomicrobium*) encode relatively higher hemerythrins and hemerythrin-containing domains compared to than alphaproteobacterial methanotrophs (Rahalkar and Bahulikar, 2018). Numerous environmental studies employing stable isotope probing (SIP) or mRNA-based analyses, showed that gammaproteobacterial methanotrophs are predominant and thrive at oxic-anoxic interfaces with rapid oxygen fluctuations (e.g., lake sediments, mangrove sediments, rice paddies, and peatlands; He et al., 2012; Reim et al., 2012; Shiau et al., 2018; Kaupper et al., 2020, 2021). For instance, vertical stratification along an oxygen gradient in a stratified lake has been observed, supported by metagenomic analysis revealing the genetic potential for aerobic methane oxidation under micro-oxic conditions, with the gammaproteobacterial methanotrophs (e.g., *Methylobacter*) present at the oxic-anoxic interface, and alphaproteobacterial methanotrophs present and stratified at the oxic water layer (Blees et al., 2014; Rissanen et al., 2018, 2020).

Altogether, it seems aerobic methanotrophs are equipped to function under micro-oxic conditions albeit with altered metabolic capabilities. Moreover, under micro-oxic conditions, the gammaproteobacterial methanotroph *Methylomonas denitrificans* can couple methane oxidation to nitrate reduction as an energy conservation strategy, and releasing nitrous oxide (Kits et al., 2015). *Methylomonas* sp. has been detected in anaerobic cultures of nitrate-*daom* microorganisms, where the expression of the *pmoA* gene increased upon exposure to 5% oxygen_{v/v}, highlighting the survival and latency of aerobic methanotrophs under anoxic methane/nitrate-rich conditions (Guerrero-Cruz et al., 2018). Besides nitrate, methane oxidation coupled to iron (in the form of ferrihydrite) and manganese reduction through direct electron transfer *via* cytochromes, or *via* an artificial organic electron acceptor (e.g., anthraquinone-2,6-disulfonate) have been implicated for both gammaproteobacterial (*Methylococcus*, *Methylomonas*) and alphaproteobacterial (*Methylosinus*) methanotrophs under hypoxia (Oswald et al., 2016; Tanaka et al., 2018; Zheng et al., 2020). Proteobacterial methanotrophs (e.g., *Methylomicrobium alcaliphilum* and *Methylocystis*-affiliated) excrete methane-derived organic compounds (e.g., acetate, lactate, succinate) during oxygen-limited growth (Costa et al., 2000; Kalyuzhnaya et al., 2013). The released organic compounds likely fueled

and fostered a complex network of interacting microorganisms comprising both the methanotrophs and non-methanotrophs (He et al., 2020). A complex regulatory mechanism involving an interplay of carbon metabolism (methane and methanol oxidation), denitrification, quorum sensing, and production of secondary metabolite (tundrenone), mediated by the signaling molecules nitric oxide and oxygen has been implicated in the response of *Methylobacter* to hypoxia (Yu et al., 2020). The metabolic versatility exhibited by these aerobic methanotrophs may become advantageous traits in dynamic environments, with rapidly fluctuating oxygen availability. In addition, alphaproteobacterial methanotrophs (*Methylosinus* species) have been co-cultivated alongside strictly anaerobic methanogens under oxygen limitation (In 't Zandt et al., 2018b). Therefore, it is noteworthy that aerobic methanotrophs likely showed species- or even strain-specific metabolic versatility, if parallels are drawn from their N-metabolic capabilities (Hoefman et al., 2014), to cope under oxygen limitation.

Interestingly, *Methylobacter* has been detected to form an active methane-oxidizing community in prolonged dark-anoxic incubations of a sub-Arctic lake sediment, ruling out the possibility that photosynthetic organisms helped sustain aerobic methane oxidation (Martinez-Cruz et al., 2017). Nevertheless, aerobic methanotrophs can thrive in anoxic environments by closely associating with photosynthetic (micro) algae serving as oxygen source (van der Ha et al., 2011; Milucka et al., 2015). Rissanen et al. (2018), demonstrated that light stimulated methane oxidation by gammaproteobacterial methanotrophs, including *Methylobacter*, in the anoxic water layers of an oxygen-stratified lake. More recently, light-induced stimulation of aerobic methane oxidation was detected in a field-based study of a boreal lake (van Grinsven et al., 2021). Similarly, aerobic methanotrophs may also rely on Sphagnum for molecular oxygen, enabling their proliferation in anoxic niches in peatland ecosystems (Raghoebarsing et al., 2005; Ho and Bodelier, 2015); or independently relying on incomplete denitrification (van Grinsven et al., 2020). The interaction of aerobic methanotrophs with their biotic environment, appears to be relevant in modulating community functioning, and may help confer resilience during disturbances prompting the use of alternative strategies and the formation of synergistic/antagonistic interactions (Ho et al., 2014, 2016, 2020; Veraart et al., 2018; Chang et al., 2020; He et al., 2020; Kaupper et al., 2021). Here, such interactions expand the habitat range of the aerobic methanotrophs to encompass unexpected environments which usually do not meet the physiological requirements for aerobic methane oxidation.

Anaerobic Methanotrophs, Physiology Dictates Prevalence Environmental Studies

Anaerobic methanotrophic archaea (ANME) differ vastly in regard to taxonomy (75–92% 16S rRNA similarity) (Knittel et al., 2005; Knittel and Boetius, 2009), cell shape and aggregation mode (Orphan et al., 2002), and even lipid composition (Chevalier et al., 2014). ANME archaea performing S-dAOM

are responsible for 80% of methane oxidation in marine ecosystems where sulfate concentrations are abundant (Hinrichs and Boetius, 2002; Boetius and Wenzhöfer, 2013). Recently, the versatility of ANME archaea has expanded by their use of humic substances, natural components of organic matter with redox-active functional groups such as quinone moieties, that could enable Humic-dAOM. Several studies have used the humic acid analog anthraquinone-2,6-disulfonic acid (AQDS) to demonstrate Humic-dAOM (Coates et al., 1998; Cervantes et al., 2000). Humic-dAOM has been observed in ANME2a and ANME2c in the presence of humic acids and AQDS as electron acceptors in the presence of chelated ferric iron (Scheller et al., 2016). Furthermore, humic substances were shown to stimulate methane oxidation linked to N₂O reducing microbes in coastal sediments in microbial communities composed by *Acinetobacter* and archaea from the Rice Cluster I and an uncultured member of the *Methanomicrobiaceae* (Valenzuela et al., 2020).

Regarding N-dAOM microorganisms, environmental surveys yielded simultaneous detection of both *Methylomirabilis* bacteria and “*Ca. M. nitroreducens*” in a widely diverse range of environments such as, sandy riverbeds (Shen et al., 2019), lake sediments (Lomakina et al., 2020), stratified lakes (Deutzmann et al., 2011, 2014; Graf et al., 2018), various wetlands including peatlands, paddy fields swamps, canals and groundwater (Raghoebarsing et al., 2006; Zhu et al., 2012, 2015; Vaksmaa et al., 2016), estuary and coastal sediments (Chen et al., 2014) among others. For a broad summary on environmental detection see Ding and Zeng (2021). Intriguingly, the first detection of *Methanoperedens* archaea came from the Gulf of Mexico (Mills et al., 2003, 2005); however, N-dAOM in the marine environment was long underestimated due to their low abundance. Recently “*Ca. M. nitroreducens*” and “*Ca. M. oxyfera*,” were identified in cold seeps and in gas hydrate-bearing sediments of the deep sea which suggests N-dAOM is an overlooked sink for methane in marine environments (Jing et al., 2020). However the availability of nitrate and nitrite in the marine environment is much lower than sulfate, and is dependent on the fluxes of nitrogen deposition or degradation of organic matter present in the sediments.

Iron (Fe) as the most abundant element on earth by mass, is widely spread in marine (Beal et al., 2009; Wankel et al., 2012; Egger et al., 2015) and freshwater environments (Sivan et al., 2011; Nordi et al., 2013) indicating the potential environmental relevance of *Methanoperedens* archaea capable of Fe-dAOM (Aromokeye et al., 2020). In this sense, microbial ¹³C carbon assimilation into *Methanoperedens*-like archaea under iron-rich, sulfate-depleted sediment incubations, has been reported (Weber et al., 2017). In addition, sulfate-rich environments could support S-dAOM mediated by *Methanoperedens* archaea, with the possible involvement of iron⁺³ or manganese⁺⁴ according to geochemical evidence and the detection of these archaea in sulfate-rich freshwater lake Cadagno (Schubert et al., 2011). A follow-up study proposed S-dAOM to be carried out by *M. nitroreducens* in a consortium with *Desulfobulbaceae* (Su et al., 2020). The debate whether metal oxides alone can be used as electron acceptors or serve a stimulatory role to support S-dAOM, persists since no genome of the *Methanoperedens* archaea

known to date, contains dissimilatory sulfate reductases (Leu et al., 2020b).

Laboratory Enrichments

Despite the widespread prevalence of AOM microbes in various environments, since their discovery, highly enriched laboratory cultures are very scarce and no pure culture has been reported, limiting physiological experiments. Through optimizations in bioreactor enrichments and cultivation media design, a limited number of cultures with enrichments from 20–83% of *Methanoperedens* archaea and/or *Methylomirabilis* bacteria, are available (Haroon et al., 2013; Guerrero-Cruz et al., 2018; Lu et al., 2018).

Mechanistic and enzymatic machinery to utilize diverse electron acceptors has been reported in several highly enriched laboratory cultures. The duality of “*Ca. M. nitroreducens*” to perform Nitrate-, Fe-, and Mn-dAOM in short term batch cultivations has been shown (Ettwig et al., 2016). However, metal mineral forms cannot be transported over cellular membranes, indicating the need for an extracellular electron transfer (EET) mechanism when no syntrophic partner is involved. “*Ca. M. nitroreducens*” genome encodes for a high number of multi-heme c-type cytochromes (Haroon et al., 2013; Arshad et al., 2015), which could facilitate the electron transfer, but to date, no direct conclusive intermediate interspecies electron transfer has been shown for “*Ca. M. nitroreducens*.” Other *Methanoperedens* archaea utilize iron and manganese in the absence of nitrate reducing genomic traits, where the involvement of cytochromes to mediate electron transfer was proposed based on genomic and transcriptomic experiments (Cai et al., 2018b; Leu et al., 2020a). In transcriptomes of “*Ca. M. manganicus*” and “*Ca. M. manganireducens*,” 23 of the 33, and 9 of the 19 multiheme c-type cytochromes were highly expressed, respectively. These studies propose Menaquinone as electron carrier based on the identification of menaquinone biosynthesis pathway in all the *Methanoperedens* strains. Comparative genomics of 16 genomes of *Methanoperedens* strains revealed that about 88% of genomes contained more than one of the menaquinone cytochrome oxidoreductase complexes and about 63% have at least four (Leu et al., 2020b).

An alternative mechanism for electron transfer is based on conductive nanowire structures (Leu et al., 2020a), similar to “*Ca. M. nitroreducens*” Mnv1 strain (Guerrero-Cruz et al., 2018), where upregulation of archaellum related genes was shown under oxidative stress. “*Ca. M. manganicus*” and “*Ca. M. manganireducens*” genomes encode for genes of the major subunit flagellin (*flaB*), part of the archaellum. Two of the four *flaB* genes were highly expressed during Mn-dAOM experiments in “*Ca. M. manganicus*,” suggesting the involvement of these conductive appendices in electron transfer. In addition to archaellum-mediated AOM, Humic-dAOM has been reported in an enrichment culture of both N-dAOM microorganisms where AQDS and anthraquinone-2-sulfonic acid (AQS) acted as electron acceptors for methane oxidation (Bai et al., 2019). These examples demonstrate with specific evidence the broad versatility of *Methanoperedens* archaea; however, other studies report AOM with alternative electron acceptors with inconclusive

results. Hexavalent chromium (Lu et al., 2016), selenate (Luo et al., 2018; Shi et al., 2020), and bromate (Luo et al., 2017), have been suggested to mediate AOM in cultures containing N-dAOM microorganisms; however, the enrichment and reported activity of side populations of *Geobacter* or canonical aerobic methanotrophs, make the involvement of N-dAOM microorganisms, debatable.

Laboratory bioreactor systems have been fundamental in the discovery and isolation of important novel N- and metal-dAOM microorganisms, yet these systems alter the natural conditions thus opening the possibilities for mutations and induced evolution. Are these changes, natural overlooked features or are these features introduced *in vitro* thus altering natural occurring features? Mn-dAOM representatives, “*Ca. M. manganicus*” and “*Ca. M. manganireducens*” strains, were enriched (up to 50% enrichment) from parent pre-enriched inoculum of “*Ca. M. ferrireducens*” (25% enrichment) and a minimal proportion of “*Ca. M. manganicus*” (Leu et al., 2020a). Similarly, the nitrate-dAOM archaeon “*Ca. M. nitroreducens*,” has several strains described over the last 8 years since its original isolation and is the only member of the Methanoperedenaceae family encoding for *narGHI* genes necessary for nitrate reduction while having the capability to perform iron- and manganese-dAOM under nitrate-depletion (Haroon et al., 2013; Ettwig et al., 2016).

Similarly, “*Ca. M. oxyfera*” enrichments have been subject to adaptation triggered by laboratory conditions since its original isolation, as a result making correlations to natural environments challenging. Early reports established that “*Ca. M. oxyfera*” harbored the canonical aerobic methane oxidation pathways (Ettwig et al., 2010) including methanol dehydrogenase (MDH). MDH in “*Ca. M. oxyfera*” was reported as a heterodimer formed by Calcium-dependent *MxaI*-encoded small subunit and a *XoxF1*-encoded large subunit, furthermore, did not possess the complete PQQ synthesis genes, necessary for the coordination of methanol in the active site of MDH (Wu et al., 2015). XoxF-type enzymes are known to incorporate lanthanide metals in their active sites in order to catalyze the oxidation of methanol (Keltjens et al., 2014). Recent studies have shown that other species of *Methylomirabilis* bacteria have evolved under laboratory conditions from an *Methylomirabilis*-rich inoculum (Ettwig et al., 2009), cultivated with elevated amounts of Cerium. These conditions resulted in the enrichment of “*Ca. Methylomirabilis lanthanidiphila*,” the second highly enriched cultivated member of NC10 phylum, encoding only one XoxF-type MDH highlighting the potential involvement of lanthanides such as cerium in methanol oxidation (Versantvoort et al., 2018). In deep freshwater lakes, other species of *Methylomirabilis* bacteria have been identified harboring lanthanide-dependent MDH although yet uncultured (Graf et al., 2018). These recent findings highlight the MDH gene redundancy in the original “*Ca. M. oxyfera*,” if the role of Calcium-dependent MDH is a metabolic trait in the natural environment or triggered by enrichment conditions where lanthanides were not originally supplied, remains an open question.

Overall, quantification of the AOM rates linked to various electron acceptors is often based on laboratory batch incubations where a single process is investigated. At times, soils are

preincubated to allow depletion of naturally occurring electron acceptors; this however, may affect the community and functioning of the original inoculum.

FATE OF METHANE AND APPLICATION POTENTIAL OF ANAEROBIC METHANOTROPHS IN WASTEWATER TREATMENT

The fate of methane relies on physical conditions such as gradient pressure, depth, and the presence and activity of the diverse array of methanotrophic microorganisms functioning as a biofilter. If the right conditions are met with availability of suitable electron acceptors, methanotrophs will consume methane before it escapes into the atmosphere. A perspective on the fate of methane focusing on the anaerobic methanotrophs and their role in human-influenced ecosystems is given, using wastewater treatment systems as a model.

Carbon Removal in Wastewater Treatment

In wastewater (WW), pollutants are transported from households and/or industrial discharge into wastewater treatment plants (WWTPs). Since modern WWT developed in the late 1800s and formally implemented in the early 1900s, organic carbon removal is performed through microbial aerobic oxidation commonly known as activated sludge (AS) (Alleman and Prakasam, 1983). The AS process takes place in oxidation tanks requiring massive amounts of energy in the form of aeration through mechanical or pneumatic infrastructure. Organic matter (including sludge from AS), is also degraded through anaerobic digestion (AD), which is a sequential set of microbial conversions under reducing conditions in the absence of oxygen namely: hydrolysis, acidification, fermentation, and ultimately methanogenesis (Section “Methanogenesis”). The development of AD was halted for decades due to the lack of fundamental understanding of the microbiology behind methanogenesis at the time (Buswell and Hatfield, 1936; McCarty, 1982). In the 1980’s, researchers achieved the decoupling of the solid retention time (SRT) from the hydraulic retention time (HRT) with the implementation of granular growth. Through the implementation of fluidized bed reactors with granular biomass configuration, the Up-flow anaerobic sludge blanket reactor (UASB) became one of the golden standards in AD technology for full-scale applications (Lettinga et al., 1980; Lettinga and Hulshoff Pol, 1991; Lettinga, 1995). Today, the methane produced from AD in full-scale sewage treatment, is collected as a methane-rich gas mixture termed biogas with remaining environmental concerns (Section “Wastewater Treatment as a Source of Methane Emissions”). Biogas represents a sustainable use of waste for the production of electricity from methane and is one of the first circular economy examples of energy recovery from waste (Bachmann, 2015).


The complexity of the microbial communities in AD, the large volumes of wastewater, and the variations in organic matter load;

pose particular disadvantages prompting to bioaugmentation protocols for biogas. Bioaugmentation is needed to remediate the variation in methane content, and the presence of toxic bioproducts such as hydrogen sulfide (Nzila, 2017; Okoro and Sun, 2019). Carbon removal either through AS or AD is a complex combination of microbial processes under highly concentrated nutrient load and accelerated enrichment conditions, however, the complexity of microbial populations remains vastly unexplored (Ho et al., 2013c; Wu et al., 2019). Only recently, global attempts to characterize WWT microbiology through sequencing technology, were initiated (Karst et al., 2018) aiming at the construction of novel taxonomy databases applied to WWT called MiDAS4 (Nierychlo et al., 2020) for the further understanding of microbial ecology in WWT including anaerobic digestion (AD). With these advancements, we anticipate that microbial engineering of AD will be unraveled and can provide custom-made solutions for waste degradation. This, combined with biotechnological production of tailored products from the wide array of resources contained in WW including volatile fatty acids (VFAs) and methane as a building blocks however, this possibility at the moment remains a black box in WWT engineering. Additionally, biogas represents a low-value and renewable feedstock for a sustainable and circular economy through the application of methanotrophs capable of producing valuable compounds and biopolymers from methane (Section “Applications of Aerobic Methanotrophs in a Circular Economy”) (Wang et al., 2020).

Nitrogen Removal in Wastewater Treatment

Nitrogen as a form of pollution, is mainly represented by ammonia and has traditionally been removed through the canonical nitrification-denitrification process (N/dN). In N/dN, ammonia is oxidized to nitrate and subsequently reduced dinitrogen gas (N₂) covering a large portion of the entire nitrogen cycle in one treatment process (Figure 4A). As in the activated sludge process, nitrification requires large amounts of energy in mixing and sufficient aeration to oxidize all nitrogen-compounds to nitrate. Furthermore, denitrification may require organic carbon (typically methanol) under poor carbon/nitrogen ratios as electron donors for the reduction steps from nitrate to N₂ (Figure 4A). Moreover, under specific conditions nitrous oxide (N₂O), a potent GHG, is produced and released to the atmosphere and current estimates report that 5% of the total N₂O annual emissions come from WWT [Intergovernmental Panel on Climate Change (IPCC), 2014] (a full review on N/dN is given by Thakur and Medhi, 2019). The additional requirements for organic carbon, the large-scale tanks for N/dN processes, and the potential for GHG emissions, render this process unsustainable.

Analogously to the development of the N-dAOM, ammonia oxidation can also occur using nitrite as electron acceptor, a process known as nitrite-dependent anaerobic ammonia oxidation (anammox) performed by anammox bacteria from the phylum Planctomycetes, discovered and isolated from sewage sludge (Mulder et al., 1995; Van De Graaf et al., 1996; Strous et al., 1999a). Soon after their discovery, a series of fundamental



physiology research studies enabled the anammox process to find technological applications to remove ammonia from wastewater. In full-scale applications, half of the ammonia is oxidized to nitrite by aerobic ammonia oxidizers (partial nitrification), which is then used by anammox bacteria to oxidize the remaining ammonia to N_2 , a combination known as partial nitrification/anammox (PN/A) (**Figure 4B**). The PN/A process offers significant advantages over N/dN and biologically, the anammox process represents a shortcut to the nitrogen cycle that has revolutionized ammonia removal worldwide (van der Star et al., 2007; Agrawal et al., 2018). For a full description of the historical developments of anammox, see Kuenen (2020).

Wastewater Treatment as a Source of Methane Emissions


The need for innovative methane removal technologies originates because despite the significant advancements in AD and the possibility of recovering biogas for energy production from waste, some challenges remain. Waste and wastewater treatment technologies remain with open loops that call for novel microbiology-based technologies for GHG mitigation (**Figure 5**). Biological WWTPs, have received recent attention as a source of GHG (Weissenbacher et al., 2010; Daelman et al., 2012, 2013). In anaerobic sludge liquor, methane can be present at levels up to 26 grams per cubic meter (Chen et al., 2015) whereas other calculations from anaerobic systems predict up to 45% methane dissolved in effluents at maximum methane production capacity (Liu Z.H. et al., 2014). Effluents from

WWTP, in particular from AD, carry a fraction of the methane dissolved which subsequently escapes to the atmosphere without control measures (Guisasola et al., 2008; Kampman et al., 2014; Schaum et al., 2016; Nguyen et al., 2019; Ribera-Guardia et al., 2019). Furthermore, in low-temperature waters, up to 60 per cent of methane can remain dissolved in the water and subsequently released into the atmosphere (Noyola et al., 2006; Souza et al., 2011).

With the evidence of methane emissions from WWT, current trends include attempts to estimate the contribution through the implementation of methodologies aiming to accurately determine the footprint of WWT (Guisasola et al., 2009; Rodriguez-Garcia et al., 2012). Current estimations indicate that WWT represents 2.8% of the total global GHG emissions, however the uncertainties in estimations range between 3% to 9% of total anthropogenic emissions (Stern and Kaufmann, 2014; Saunio et al., 2020). In the next section, an overview of the potential application of anaerobic methane and ammonia oxidation is given, emphasizing the research gaps needed to explore the possibilities of implementing a combined process where nitrogen (ammonia) removal and fugitive carbon emissions (methane) can be addressed simultaneously.

Potential Applications of Anaerobic Methanotrophs in Wastewater Treatment

Wastewater treatment represents the ideal niche with the highest loads of nitrogen and carbon input suitable for N-dAOM methanotrophs and anammox to thrive. Anammox bacteria



incorporated in PN/A systems are a clear example of a technology that developed from a novel microbial process characterized in laboratory-scale enrichments, and through physiology studies, reached full-scale realization (Van De Graaf et al., 1997; Strous et al., 1999b; Kuenen, 2020). Some of the microbial aspects relevant for the application of an anaerobic process in wastewater systems are growth, inhibition, and physiology; this knowledge is achieved through the combination of fundamental research on microbial physiology, engineering, genomics and even modeling (Strous et al., 1997; Carvajal-Arroyo et al., 2013; Lotti et al., 2014). Recently, the possibility to remove ammonia from WW coupled to electricity production was demonstrated for anammox bacteria (Shaw et al., 2020), capable of direct extracellular electron transfer (EET) similar to methanogens (Section “Anaerobic Methanotrophs”). EET anammox technology could potentially replace oxygen-limited systems such as PN/A circumventing the requirement for aeration, however this process is at early developmental stage in laboratory systems. EET in *Methanoperedens* archaea has been proposed as a decoupling mechanism from *Methylomirabilis* bacteria (Ding et al., 2017), however the reported enrichment is accompanied by the enrichment of canonical methanotrophs and geobacter under short term incubations. Furthermore, the implementation of EET-dAOM would require high quality methane supply based on the kinetic limitations of *Methanoperedens* archaea, e.g., in hollow membranes (Zhang et al., 2020) (Section “Methane Supply and Reactor Configuration”). For this purpose EET-dAOM is not further discussed.

Nitrogen-dependent anaerobic methane oxidation in WWT was considered a suitable application for simultaneous methane and nitrogen removal almost four decades ago (Davies, 1973; Sollo et al., 1976). Today, after the discovery of the microorganisms responsible for this process in diverse environments including WWT, and their enrichment in laboratory-scale reactors (Ettwig et al., 2009; Hu et al., 2009; Luesken et al., 2011b; Haroon et al., 2013), N-dAOM as a potential application has re-emerged (Wang et al., 2017). The N-dAOM process has significant potential benefits if the microorganisms responsible for this process can be integrated into oxygen-limited systems for ammonia removal (Figure 6). This concept is proposed based on laboratory scale bioreactor enrichments where both N-dAOM and anammox microorganisms co-exists with nitrite as a common link (Figure 7). Bioreactor systems supplied with nitrate, methane and ammonia; have resulted in a coculture of N-dAOM archaea, N-dAOM bacteria and anammox in diverse proportions (Shi et al., 2013; Stultiens et al., 2019). However, when supplied with nitrite, methane and ammonia only a combination of both bacterial partners (N-dAOM bacteria and anammox) was achieved (Luesken et al., 2011a). Based on these reports, models have been developed indicating that both anaerobic processes can coincide and could reach a practical application and performance in WWT (Winkler et al., 2015).

In short, the simultaneous removal of methane and ammonia in oxygen-limited systems relies in the concerted interaction of 4 microbial processes (Figure 6) driven by key microbial groups (Figure 7):

- (i) Aerobic ammonia-oxidizing bacteria (AOB) oxidize half of the ammonia available to nitrite (partial nitrification, PN), requiring half the aeration energy compared to N/dN (Figure 6B).
- (ii) Anammox bacteria, oxidize the remaining ammonia using the nitrite produced by AOB. Anammox however, produces nitrate as a side reaction during carbon fixation (Figures 4B, 6B).
- (iii) Methane is oxidized by nitrite-dAOM bacteria using nitrite most likely under competition with anammox bacteria. Both producing nitrogen gas as final nitrogen-product (Figure 6C).
- (iv) Nitrate-dAOM archaea, will compete with *Methylomirabilis* bacteria to oxidize methane using nitrate (Figure 6C). Nitrate can originate from anammox side reactions or undesirable complete nitrification of ammonia to nitrate or from other nitrate-rich effluents (Figure 6A).

Dissolved methane from diverse effluents, more importantly anaerobic digestion, and biogas mixtures that otherwise would not be used for electricity production; represent an ideal energy source for the combined process (Bandara et al., 2011; Wang et al., 2017), reaching up to 45% in some cases (Liu Z.H. et al., 2014). To successfully combine both processes, physiology and microbial ecology aspects are of vital importance. These aspects include competition for nitrite as an intermediate substrate, doubling times in a mixed culture, affinity and




FIGURE 6 | A schematic representation of the combination of partial-nitritation/anammox (PN/A) and N-dAOM processes in WWT. **(A)** Canonical denitrification where space, energy, and organic carbon, are needed for complete nitrogen removal in WWT. Denitrification produces sludge and GHG emissions as by-products under unfavorable conditions. **(B)** Partial-nitritation/anammox process, as a short cut to complete denitrification. AOB (in yellow) oxidize ammonia to nitrite, anammox bacteria (in red) oxidize ammonia anaerobically using nitrite as electron acceptor. **(C)** Incorporation of N-dAOM processes to PN/A, where both achieve the simultaneous removal of ammonia and methane (represented in green and blue for nitrate- and nitrite-dAOM, respectively). Broken arrows indicate competing processes that are deemed undesirable or processes that are circumvented by the implementation of the PN/A and N-dAOM processes. Adapted from Guerrero-Cruz (2018).




FIGURE 7 | Interactions between anaerobic methane oxidizers performing N-dAOM with anaerobic ammonia oxidation (anammox) incorporated into partial-nitritation systems (PN/A). In existing PN/A systems for ammonia removal, aerobic ammonia oxidizers (AOB, in yellow) oxidize ammonia to nitrite, anammox bacteria (in red) use nitrite to oxidize ammonia to dinitrogen gas, while producing residual nitrate from carbon fixation. “*Ca. Methanoperedens nitroreducens*” (in brown with red outline) converts nitrate to nitrite as common intermediate while oxidizing methane to carbon dioxide and can produce ammonia. Nitrite either produced by AOB or *M. nitroreducens*, is used by *Methylomirabilis* bacteria (in green) for the oxidation of methane to carbon dioxide. Aerobic methane oxidizers (in grey) compete for oxygen with AOB, and can oxidize methane, in competition with anaerobic methane oxidizers.

competition for methane by aerobic and anaerobic methane oxidizers, competition for oxygen between aerobic ammonia and methane oxidizing microorganisms (Figure 7) under oxygen-limited conditions, and more importantly the inhibition of anaerobic microorganisms upon the exposure to oxygen. Most of these microbial metabolic aspects are known for anammox

bacteria (Strous et al., 1999b) but remain to be determined for N-dAOM microorganisms mostly due to the lack of highly enriched cultures. For a summary on the laboratory enrichments and environmental detection of N-dAOM microorganisms since 2006, culture conditions, inoculum sources, and environmental detection see Ding and Zeng (2021). In the following sections, an

overview on the recent discoveries relevant to application is given highlighting the research gaps that need further investigation to point N-dAOM research in the right direction towards becoming a full-scale application.

Nitrite and Nitrate as a Central Link for Anaerobic Ammonia- and Methane-Oxidizers

Nitrite is the common terminal acceptor for Anammox and *Methyloirabilis* bacteria to oxidize ammonia and methane respectively (Figure 7; Strous et al., 1999a; Ettwig et al., 2010), and is the intermediate byproduct during nitrate reduction coupled to methane oxidation by *M. nitroreducens* (Figures 3, 7; Gambelli et al., 2018). This has been corroborated in several laboratory enrichments where nitrite was the defining factor between the co-existence of both N-dAOM microorganisms with anammox bacteria (Ding et al., 2014; Stultiens et al., 2019). The competition for nitrite is dependent on the affinity, whereas inhibition is subject to the formation of nitrite from ammonia oxidation (Figure 7) and is the main engineering parameter that can influence the out-competition of *Methyloirabilis* bacteria as anammox bacteria have a higher affinity for nitrite (Table 1; Winkler et al., 2015). Inhibition would originate from a surplus of nitrite formation under an excess in oxygen supply. *Methyloirabilis* bacteria have a tolerance of approximately 500 μM of nitrite (Guerrero-Cruz, 2018) whereas anammox bacteria have a tolerance ranging in the mM range (>10-fold higher) (Lotti et al., 2012; Table 1).

Growth of N-dAOM Microorganisms

Growth rates, doubling times and yield (the amount of biomass obtained from the energy source) of N-dAOM microorganisms are lower compared to canonical aerobic heterotrophic microorganisms. N-dAOM microorganisms have doubling times estimated from weeks to months based on the quantification of 16S *rRNA* gene over time from environmental studies and long-term incubations (Ettwig et al., 2009; Vaksmaa et al., 2017). Recently, from a limited number of highly enriched cultures of N-dAOM microorganisms, data on growth and substrate kinetics has become available. *Methyloirabilis* bacteria

can double every 5 days under maximum SRT conditions (10 days) and exhibit an estimated yield at steady state of $0.077 \pm 0.027 \text{ C}_{\text{mole}} \text{ CH}_4 \text{ mole}^{-1}$, as determined by indirectly comparing the expected biomass decay versus the total biomass content while monitoring side populations through fluorescent microscopy (Guerrero-Cruz et al., 2019). For “*Ca. M. nitroreducens*” indications of a doubling time between 14–21 days have been obtained under continuous bioreactor incubation under an SRT of 20 days (Guerrero-Cruz, 2018). For comparison, aerobic methanotrophs generally exhibit a doubling time of ~ 30 min to hours (Ogiso et al., 2012). Anammox bacteria have reported doubling times as low as 3 days and a biomass yield of $0.066 \pm 0.01 \text{ C-mole/ammonium mole}$ (Strous et al., 1998; van der Star et al., 2007). Table 1 gives a summary of relevant parameters in literature for anammox bacteria and N-dAOM microorganisms.

Another feature crucial to microbial growth is the capability to form granular biomass structures. Granular growth is essential to ensure the decoupling of SRT from HRT, preventing biomass washout. In addition, granular growth enables a multi-layer growth in oxygen limited systems resulting in the compartmentalization where, anaerobic microorganisms are protected at the core of the granular unit and aerobic bacteria secure the consumption of oxygen in outer layers (Winkler et al., 2015; Speth et al., 2016). Granular growth held a game changing role in the implementation of novel technologies relying on multiple processes occurring simultaneously such as in AD, NEREDA[®], and PN/anammox (Lettinga et al., 1980; Lackner et al., 2014; Pronk et al., 2015; Chen et al., 2020; Trego et al., 2020).

Methanoperedens archaea and *Methyloirabilis* bacteria have the natural capacity to develop granular growth under dynamic conditions in SBR reactors (Luesken et al., 2011a; Guerrero-Cruz et al., 2019). Under continuous stirring in a membrane bioreactor, *Methanoperedens* archaea achieved higher enrichment levels in a granular ultrastructure from a sediment inoculum (Gambelli et al., 2018; Guerrero-Cruz, 2018). With the relevance of granular growth and the potential to form multi-layered microbial processes under oxic conditions, the next section describes the current knowledge of the oxygen effect on these microorganisms.

TABLE 1 | Summary of relevant physiological parameters of both anaerobic ammonia and methane-oxidizing microorganisms.

| | Anammox bacteria | <i>Methyloirabilis</i> bacteria | <i>Methanoperedens</i> archaea |
|-----------------------------------|---|---|---|
| Doubling time (days) | 10 – 12 ^b 11 ^a 3 ^f | <u>25</u> ^e 14 ^c 5 ⁱ | 14–21 d at SRT of 10 d ^{-1-h} (40% archaea enrichment) |
| $\mu_{\text{Max}}(\text{d}^{-1})$ | 0.33 ^f | <u>0.0495</u> ^g 0.14 ⁱ | ND |
| Yield (C-mol/energy source) | 0.066 \pm 0.01 ^a | 0.077 \pm 0.027 ⁱ | ND |
| Methane affinity | NA | <u>92 \pm 5 μM</u> ^e | > 1000 μM ^d (S-dAOM) |
| Methane V_{max} | NA | <u>97 μM</u> ^e 2.6 \pm 0.7 μM ⁱ | ND |
| Ammonia affinity | <5 μM ^a | NA | NA |
| Nitrite affinity | 0.2–5 μM ^b | <u>910 \pm 90 μM</u> ^e 7 μM ⁱ | ND |
| Nitrate affinity | NA | NA | 150 \pm 29 μM ⁱ |
| Nitrite inhibition | 400 gNO ₂ -N m ^{3b} | <u>40 g NO₂-N m^{3e}</u> Estimated at 500 μM ⁱ | Estimated at 500 μM ⁱ |

Values from both real enrichments and mathematical models (underlined) are shown, NA = not applicable, ND = not described. All data has been collected from ^aStrous et al. (1998); ^bvan der Star et al. (2007); ^cEttwig et al. (2009); ^dMeulepas et al. (2009); ^eHe et al. (2013); ^fLotti et al. (2015); ^gWinkler et al. (2015); ^hGuerrero-Cruz (2018); ⁱGuerrero-Cruz et al. (2019); ^jLu et al. (2019).

Oxygen and N-dAOM Microorganisms

Methanoperedens archaea perform the reverse action of methanogenesis and the McrA complex is highly sensitive to oxygen (Thauer, 1998). As such, *Methanoperedens* has been enriched under strict anoxic conditions. Recently, it was demonstrated that an enriched culture of *Methanoperedens* archaea (83% of the total community) expressed the genetic capacity to counteract the adverse effects of oxygen-stress upon 5% v/v oxygen exposure (Guerrero-Cruz et al., 2018). Although further research is needed to characterize how the recovery after oxygen exposure would occur, especially in natural environments. The recent findings corroborate that under oxygen-limited conditions *Methanoperedens* archaea could adapt to partial aeration most specifically in multi-layered granular growth.

Methylomirabilis bacteria are inhibited by oxygen exposure as low as 2% (Luesken et al., 2012) and have demonstrated the capacity to recover methanotrophic and nitrite reduction capacity after exposure to <1.1% oxygen (Kampman et al., 2018). The successful implementation of N-dAOM is subject to oxygen exposure in engineered ecosystems under high hydraulic dynamics conditions, however, the combination with oxygen-limited systems where aerobic ammonia oxidizers consume oxygen for the production of nitrite could be a suitable coupling under granular microbial growth.

Metals and N-dAOM

Metals are essential for microbial activity and growth, aerobic methanotrophs and anaerobic *Methylomirabilis* bacteria are not an exception, requiring copper for the catalytic activity of the pMMO enzyme (Bollinger, 2010; Ho et al., 2013b; Semrau et al., 2018). Recent studies provide direct evidence of the effect of copper on methane oxidation by *Methylomirabilis* bacteria, in laboratory enrichments a concentration of 5–6 micromolar of copper salt in the trace elements, triggered an increase in methane oxidation rates (Hatamoto et al., 2018). Furthermore, higher concentrations of copper and other metals seem to enable higher enrichment ratio of these difficult to cultivate bacteria (Guerrero-Cruz et al., 2019). Besides copper, lanthanide metals are also relevant for the intra-aerobic *Methylomirabilis* methanotrophs such as the novel enriched species “*Ca. Methylomirabilis lanthanidiphila*” and its uncultured environmental analog “*Ca. Methylomirabilis limnetica*” encoding for only a lanthanide-dependent MDH in a way that is not redundant as the first species reported in 2010 (Graf et al., 2018; Versantvoort et al., 2018). This change in MDH enzymes was discussed previously in this review (Section “Anaerobic Methanotrophs, Physiology Dictates Prevalence”), but the concentration of cerium in the trace elements during cultivation is a noteworthy factor. Moreover, iron is perhaps one of the most essential metals in life enabling formation of complexes and a wide spectrum of redox reactions that are crucial to enzymatic processes (Liu J. et al., 2014). Iron is present in cytochrome proteins, essential and highly abundant in different *Methanoperedens* archaea (Arshad et al., 2015; Leu et al., 2020b). Furthermore, increasing iron concentrations has resulted in higher laboratory scale enrichments. Initial attempts to enrich *Methanoperedens* archaea included iron concentrations

below 5 micromolar (Vaksmas et al., 2017), and concentrations of 80 micromolar have resulted in a two-fold increase in enrichment under laboratory scale conditions (Guerrero-Cruz et al., 2018; Lu et al., 2018). Evidently, metals (e.g., copper, cerium, and iron) are crucial for the activity and enrichment of anaerobic N-dAOM microorganisms. To date limited studies have directly compared the effect of trace metal concentrations in enriching these microbes, potentially due to low concentrations in the nature and lack of bioavailable forms. The future applications of these microorganisms must thus account for the role of metals in mix cultures and under varying conditions anticipated in wastewater treatment.

Methane Supply and Reactor Configuration

Methane has a relatively low water solubility (0.03464 Bunsen coefficient at 20°C or 1.237 mM at standard conditions, Wiesenburg and Guinasso, 1979; Haynes, 2011). The poor solubility of methane in effluents from wastewater treatment is a major contributor to methane emissions (Section “Wastewater Treatment as a Source of Methane Emissions”), and this translates to laboratory conditions where the supply of methane is the main limitation for the development of enrichment cultures. Diverse forms of bioreactor design have been implemented to ensure sufficient supply of methane, such as high methane flow, pressurized reactors, membrane-fed reactors or batch incubations. **Table 2** provides a summary of relevant configurations in literature. It is important to note that due to the slow growth of N-dAOM microorganisms, most enrichments are achieved after months or even several years. Furthermore, these enrichments typically are achieved in bioreactors, with volumes not exceeding 10 L.

Laboratory scale membrane bioreactors (MBR) have been proposed as an ideal setup for the enrichment of N-dAOM microorganisms, usually accompanied by anammox bacteria (**Table 2**). MBRs ensure methane supply and biomass retention, however, relying on the external supply of artificial high-purity methane mixtures (>95%), which are costly. Pressurized systems have been used to enrich S-dAOM methanotrophic archaea and have demonstrated to enable higher growth and activity rates by ensuring the availability and saturation of methane above normal conditions of methane (Nauhaus et al., 2002; Timmers et al., 2015). The combination of pressurized reactors and membrane configurations has indeed shown an increase in the activity of N-dAOM microorganisms (Cai et al., 2018a), and these studies highlight the application potential, however the external supply of high-purity methane remains the main obstacle for full-scale application in WWT. When high-quality external supply of methane is required, the potential use of intrinsically-produced methane from AD is overlooked and deviates from the prospective application of N-dAOM microorganisms to mitigate GHG emissions from effluents of AD in WWT (Wang et al., 2017), where commonly a dual treatment configuration is proposed with partial PN/A and N-dAOM as a second step or in combination with anammox (Stultiens et al., 2019). Moreover, membrane-based reactors provide a suitable surface for biofilm growth (Cai

TABLE 2 | Summary of diverse laboratory-scale bioreactor studies combining N-dAOM microorganisms under different bioreactor conditions.

| Reactor type | Microbial community | Performance | Limitation | Reference |
|---|--|---|---|--|
| Membrane biofilm reactor (MBfR). Methane supply at 2 atm Oxygen 100 to 700 mL d ⁻¹ (50 d period). | Derived from Haroon et al. (2013). | 1.5 Kg N m ³ d ⁻¹ (98% removal efficiency) Methane removal was not the focus. | External methane supply (95%). | Liu et al., 2019b |
| Membrane Granular Sludge Reactor Methane supply at 0.1 atm. | N-dAOM archaea (32%) and bacteria (9%), with anammox (27%) approximately. | 16.53 kg N m ⁻³ d ⁻¹ Methane removal was not the focus. | External methane supply (95%). | Fan et al., 2019 |
| Membrane biofilm reactor (MBfR), increase methane pressure 1.44 atm. | N-dAOM archaea and bacteria, with anammox, proportions similar to Haroon et al., 2013. | ≈ 2-fold N-dAOM archaea activity increase (26.1 mM N d ⁻¹). ≈ 5-fold N-dAOM bacteria activity increase (41.4 mM N d ⁻¹). Methane removal was not the focus. | External methane supply (90%). | Cai et al., 2018a |
| Membrane bioreactor (MBR) 19–25 mg methane L ⁻¹ 35.7 mg NH ₄ ⁺ L ⁻¹ . | Not accurately characterized. | 60% nitrogen removal in anaerobic conditions 95% methane removal. | Methane from the effluent of an UASB reactor. | Sánchez et al., 2016 |
| Membrane biofilm reactor (MBfR). Methane at a flow from 0.1 to 0.5 atm. | N-dAOM archaea (74.3%) and bacteria (11.8%), with anammox (5.6%). | 1.3 mM nitrate d ⁻¹ and 2.1 mM d ⁻¹ ammonia. Methane removal was not the focus. | External methane supply (95%). | Fu et al., 2017 |
| Membrane biofilm reactor (MBfR). Methane supply at 2.46 atm. | N-dAOM archaea (50%) and bacteria (20%), with anammox (20%). | 48.8 mM N d ⁻¹ overall. (1.2 Kg N m ³ d ⁻¹) Methane removal was not the focus. | External methane supply (95%). | Xie et al., 2017 |
| Membrane biofilm reactor (MBfR). | N-dAOM archaea (50%) and bacteria (20%), with anammox (20%). | 48.8 mM N d ⁻¹ overall. Methane removal was not the focus. | External methane supply (90%). | Cai et al., 2015, derived from Shi et al. (2013) |
| Sequencing batch reactor, batch methane pressure 0.4 Mpa. | N-dAOM archaea (29%) and bacteria (12%), with anammox (21%). | 4.84 mM d ⁻¹ Nitrate Methane removal was not the focus. | External methane supply (95%). | Ding et al., 2014 |
| Membrane biofilm reactor (MBfR). | N-dAOM archaea (20–30%) and bacteria (20–30%), with anammox (20–30%) | 13.6 mM N d ⁻¹ Methane removal was not the focus. | External methane supply (90%). | Shi et al., 2013 |

et al., 2018a) as opposed to granular-based growth (Guerrero-Cruz et al., 2018). Thus, the stratification proposed as crucial factor for the implementation of anaerobic processes under micro-aerophilic conditions is likely to not occur (Winkler et al., 2015). Furthermore, studies where oxygen exposure at a constant level of 0.01 mg L⁻¹ have shown no negative effects on N-dAOM/partial nitrification/anammox systems (Liu et al., 2019b), but the small scale does not anticipate the effects under high volume of effluent input in real applications.

In this section, relevant aspects affecting the performance and growth of N-dAOM microorganisms were described. Currently the advancement of N-dAOM-based technology is faced with bottlenecks in the following aspects:

- (i) Most work relies on methane-rich artificial gas mixtures (>90 percent). There are no studies with real wastewater, except for one study where in a 50 L pilot plant, dissolved methane was removed (95% removal, 78% methane-containing biogas from a UASB reactor at a rate of 31 L per day) coupled to denitrification (1.43 mM total nitrogen removal). However the role of N-dAOM microorganisms in a dual post-treatment containing oxic and anoxic compartments was not fully

characterized next to the use of synthetic media (Silva-Teira et al., 2017). The bioaugmentation of biogas, and removal of toxic sulfur compounds could improve the use of waste biogas to enable the application of N-dAOM microorganisms when not suitable for electricity production.

- (ii) Studies to optimize fundamental aspects of growth and enrichment conditions are lacking. Currently, evidence on the role of iron, copper, and lanthanide is available (Hatamoto et al., 2018; Lu et al., 2018; Guerrero-Cruz et al., 2019); however, further studies on the requirements of these micronutrients, including up-scaling and modeling aspects in large-scale applications, are still lacking. The availability of these metals could improve the activity, growth, competitiveness, and enable the application of N-dAOM microorganisms in oxygen-limited WWT systems.
- (iii) Methane supply and the gas-liquid transfer are crucial for the successful implementation of N-dAOM microorganisms where *Methylomirabilis* bacteria compete with aerobic methane oxidizers in regard to their methane affinities (Guerrero-Cruz et al., 2019), *Methanoperedens* archaea would need the assistance

of pressurized systems to increase methane saturation above 1.3 mM and reach an equilibrium for the McrA complex to perform methane oxidation at sustainable rates (Cai et al., 2018a).

- (iv) Bioreactor configuration is a limiting factor, where membrane-based bioreactors rely on biofilm formation rich in N-dAOM microorganisms which lacks spatial niche differentiation (Cai et al., 2018a; Liu et al., 2019b). This prevents the formation of compartments that allow for both aerobic and anaerobic processes to co-exist as in the case for granular growth (Winkler et al., 2015; Gambelli et al., 2018; Guerrero-Cruz et al., 2018). Furthermore, biofilm formation is a limiting phenomenon in large scale-membrane systems whereas granular-based bioreactors have successfully been implemented in large-scale applications worldwide (Lettinga et al., 1980; Lackner et al., 2014). A combination of granular-based N-dAOM microorganisms with adequate methane-transfer adaptation through membrane systems or pressurized reactors, could offer significant advantages to apply the N-dAOM process.

After highlighting the diverse factors affecting the performance of N-dAOM microorganisms and knowledge gaps in the research, it is evident that reactor configuration and more importantly the synergistic combination with other processes are crucial research questions. Exploring these aspects can reveal the true value whether N-dAOM microorganisms could be applicable and biotechnologically beneficial. Anaerobic microbes and processes in anoxic compartments are undoubtedly applicable under granular configurations where both aerobic and anaerobic processes occur simultaneously in WWT (Pronk et al., 2015; Winkler et al., 2015). Thus research on complex microbial ecology and interactions is needed for the realization of more integrated WWT technologies. Moreover, diverse operational challenges such as aeration resulting in gas stripping, play a crucial role in the release of methane into the atmosphere. To circumvent these aspects, membrane-based and/or pressurized systems have been discussed as a means of enabling methane-rich conditions for the implementation of N-dAOM microorganisms. Currently, there is no “one fits all” solution that fully integrates all the aspects that challenge the application of N-dAOM from both points of view: microbial ecology and the engineering limitations. Nevertheless, the collection of high quality biogas for membrane-based configurations or biofilm-based oxic/anoxic systems; remain as potential solutions (Liu et al., 2019a). Indisputably, biomass stratification and the controlled supply and retention of methane in the system, are crucial requirements for the coexistence of both aerobic and anaerobic processes (Liu et al., 2019b). Moreover, efficient biogas collection and bioaugmentation are important steps to secure high-quality biogas feed for integrated systems that aim to remove methane and nitrogen contaminants simultaneously where biogas is otherwise not fitted for heat and electricity production. The next section describes alternative uses for methane-rich mixtures.


APPLICATIONS OF AEROBIC METHANOTROPHS IN A CIRCULAR ECONOMY

Currently natural gas is the primary source of methane for industrial and chemical production of methanol and other industrial compounds. However natural gas reservoirs are finite and is thus a non-renewable fuel source. Furthermore, the industrial production of methanol and olefines is not environmentally friendly, requiring large amounts of energy, metal catalysts, and high temperature and pressure conditions (Da Silva, 2016). Methanotrophic microorganisms are natural catalyzers that can perform many biochemical conversions at ambient temperatures and without the need of high energy input compared to industrial processes, making their potential applications relevant sustainable alternatives (Wang et al., 2020). Biogas from AD, is a low-value feedstock for new technologies that harness the potential of methanotrophic bacteria to produce valuable compounds such as methanol, formaldehyde, or short chain fatty acids, and biopolymers (Comer et al., 2017).

The fate and use of methane can follow several paths, and depends on infrastructure, policy, and the adoption of a waste-to-value culture (Figure 8). The production of valuable compounds from methane by methanotrophs has three differentiable targets: (i) Methanotrophs produce methanol as a valuable intermediate compound; (ii) Methanotrophs produce biopolymers as a carbon storage molecule; and (iii) Methanotrophs utilize methane as the building block for cell constituents (lipids or proteins). The application of aerobic methanotrophs for bioremediation and as bio-catalysis for a range of compounds can be found in Strong et al., 2015. Moreover, capitalizing on novel mechanistic insights of the aerobic methanotrophs under oxygen-limiting conditions, the methanotrophs can potentially be applied to produce volatile organic compounds (e.g., acetate, lactate; Kalyuzhnaya et al., 2013), and in methane-driven bioelectrochemical systems (Zheng et al., 2020). Accordingly, aerobic methanotrophs may also drive denitrification in WWTP, either in synergistic interaction with other microorganisms being a source of electron donor (e.g., by excreting acetate under oxygen limitation; Costa et al., 2000) or as a coping strategy under hypoxia (Kits et al., 2015; Zhu et al., 2016). Hence, emerging applications of aerobic methanotrophs may rely on their metabolic capability under oxygen limitation, but this aspect of methanotroph biotechnology needs further exploration. Here, we focus on the role of aerobic methanotrophs under predominantly oxic conditions in the production of value in the form of biopolymers and two examples of valuable chemical intermediates with current proof of principle and with proven potential at industrial scale.

Methane as Energy Source and Chemical Building Block Methanol

Methanol for industrial use is mainly produced from natural gas and to a lesser extent from CO₂ exhaust waste streams through unsustainable thermochemical processes, at a rate of 110 million metric tons per year. Methanol is one of the



most important building blocks for nearly 50000 different chemicals. Approximately one third of it is used for formaldehyde production, followed by synthesis of olefins for the plastic industry (Da Silva, 2016). Biogenic methane from anaerobic digestion could be a source for methanol production through the application of methanotrophs when methane-rich biogas for electricity production is rendered non cost-effective in the near future.

Aerobic methanotrophs oxidize methane to methanol, and its recovery requires the inhibition of the further methanol oxidation step. Ammonium chloride, phosphate, cyclopropanol, CO₂, NaCl, and EDTA are known MMO inhibitors that could potentially aid in the recovery of methanol from methanotrophic cultures (Wang et al., 2020). For a summary of culturing and recovery strategies see Bjorck et al. (2018). *Methylosinus trichosporium* OB3b, is the most extensively studied methanotroph for methanol production using Na-formate and NaCl to inhibit MDH activity reaching 7.7 mM of methanol in 36 h (Sang et al., 2004) or cyclopropanol with methanol levels of 152 mmol g⁻¹ DW cells with a methanol conversion efficiency of 61% (Takeguchi et al., 1997). Regarding methane conversion efficiency in *M. trichosporium* OB3b, studies have

reported 64% (Duan et al., 2011) and 80% yields but the production of methanol at higher efficiency remains a challenge (Han et al., 2013). A species from the genus *Methylocaldum*, was reported to produce methanol from methane-biogas upon 500 ppm of hydrogen sulfide exposure in batch cell suspensions, with a methane conversion efficiency of up to 34% and methanol yields of up to 343 mg L⁻¹ over 105 h incubations (Zhang et al., 2016). Although the application of methanotrophs for the production of methanol seems promising based on laboratory scale evidence, industrial applications are unlikely at this stage due to the need of external electron donors during MDH inhibition, e.g., formate (Sheets et al., 2016), negatively affecting the feasibility of large-scale applications and the continuous performance of methanotrophs.

Proteobacterial methanotrophs have been widely studied, however their mesophilic nature and implementation in non-sterile conditions are still a bottleneck. Verrucomicrobial methanotrophs, are known to thrive at low pH and extremely high temperatures, making them a promising alternative; however, the feasibility to produce methanol from Verrucomicrobia has not been investigated in depth. Attempts to produce methanol through the inhibition of

MDH's requires external supply of electron donors and growth conditions under multiple inhibition strategies, rendering the application of these extremophiles unclear at the low pH these bacteria are known to inhabit (Hogendoorn et al., 2020). Nevertheless, the exploration of Verrucomicrobial strains for methanol production from biogas, could be advantageous when using biogas mixtures that contain hydrogen sulfide as canonical proteobacterial methanotrophs are known to be intolerant to this toxic byproduct of AD biogas (Börjesson, 2001). Another feature of interest in using Verrucomicrobial methanotrophs is the possibility to reduce culture contamination by having operating temperatures as high as 55 degrees and/or lower pH values (Chen and Jiang, 2018). Research in efficient methanol recovery without a negative impact on microbial growth that could compromise culture yields, is crucial to explore biological methanol production, however, remains to be the bottleneck due to metabolic constraints for cellular growth and energy conservation.

Ectoine

Ectoine and hydroxyectoine are molecules produced in response to osmotic stress in numerous bacteria, their chemical properties protect cells during osmotic stress and assist stabilization and preservation of cell integrity without altering cell function. These properties result in a wide spectrum of applications in the cosmetic industry, medicine, dermatology, nutrition, and pharmaceuticals as well as in preservation products for a broad range of cell biology applications (Pastor et al., 2010; Wang et al., 2020). Ectoines are valued at one thousand United States dollar per kilogram, and the current global need is 15 million Kg a year, representing the most profitable market at 15 billion United States dollars a year (Strong et al., 2016). Among the methanotrophs, *Methylomicrobium alcaliphilum* shows the highest reported accumulation of ectoine, at 8.3% of ectoine per gram of biomass (Cantera et al., 2017). Heterotrophic bacteria, however, showed 3 orders of magnitude higher ectoine production rates than methanotrophs (Salar-García et al., 2017). Some of the factors that affect the efficiency of methanotrophs to accumulate ectoine is the susceptibility to mechanical stress in culture enrichments (Cantera et al., 2016). Also, the presence of specific metals is crucial for enzymatic performance. To this end, *Methylomonas* has shown increased activity in the presence of copper, enhancing ectoine production (Cantera et al., 2016). With the significant financial market, the use of methanotrophs seems like a suitable application using methane waste streams while enabling the creation of value from waste.

Methane as Source for Microbial Biopolymers

Biodiesel From Methane-Derived Lipids

Biodiesel production has gained attention as a more environmentally friendly alternative to diesel, currently produced from diverse crops in marginal lands, with particular environmental debate over the use of land in competition with food agriculture [Intergovernmental Panel on Climate Change (IPCC), 2012]. Conversely, biodiesel can be produced from

membrane lipids of methanotrophic bacteria. Methanotrophs, in comparison to other heterotrophs, have an increased lipid content due to the stacked internal membranes, rendering them attractive for biotechnological applications after a culture fed with methane has reached optimal growth yields to secure the maximum ratio of biomass from mol of methane.

Methylomicrobium buryatense 5GB1, a Type I methanotroph, is used as the model microorganism for the production of membrane lipids. Typical to Type I methanotrophs, this microorganism assimilates carbon through the RUMP pathway and fatty acids are produced from Acetyl-CoA (Wang et al., 2020). Up to 95% of the total fatty acids from membrane-bound phospholipids were recovered in *Methylomicrobium buryatense* corresponding to a total fatty acid content of 10% of dry cell weight achieved in batch reactors (Dong et al., 2017). Furthermore, a glycogen knockout strain AP18, has shown to have a higher lipid yield due to the suppression of glycogen production leading to a better feedstock for lipid conversion to biodiesel (Fei et al., 2018). Fatty acids are then extracted using an alkaline hydrolysis at 150°C followed by a hexane extraction where recoveries can reach up to 95% of the methanotrophic culture's total lipids. Lastly, a chemical upgrading of the fatty acids takes place using palladium catalysts for the hydrodeoxygenation process that transforms the lipids into long chain hydrocarbons rich in pentadecane (up to 88% efficiency) (Furimsky, 2000). Chemical upgrading of methanotrophic lipids is essential to remove reactive oxygen (carbon saturation) from the lipid molecule, conveying less reactivity and viscosity, and increasing the calorific capacity of the final product.

Bioplastics

Polyhydroxy alkanooates (PHAs) have become an important biopolymer in the past years as a replacement to chemically produced plastics (Amaro et al., 2019). PHA is a collective term encompassing approximately 100 types of monomeric compounds, all of them being linear polyesters with different degrees of polymerization reaching up to 30 thousand units. Under the right conditions, several types of bacteria can accumulate up to 90% of PHA making bacteria-derived PHA a cost-effective strategy to produce bioplastics (Madison and Huisman, 1999).

Several microbes such as *Haloferax mediterranei* and *Alcaligenes latus* produce high PHA levels, reaching more than 50% of the cells' dry weight. Genetically modified *Escherichia coli* expressing PHA biosynthesis genes (*phbA*, *phbB*, and *phbC*), accumulate PHA from lactose (Park et al., 2001). However, the feedstocks used in the production of PHAs from engineered microbes are expensive and rely on glucose and other carbohydrates, posing additional concerns regarding the sustainable supply chain, and land use for crop production which competes with crops destined for human consumption. Methane is a naturally occurring carbon source for microbial production of PHAs, and some methanotrophic bacteria are efficient at accumulating PHA without the need for genetically engineered metabolism. The possibility to use a cheap carbon source through the application of bacteria to produce bioplastics while mitigating methane emission, broadens the

use of methanotrophs in a circular and bio-based economy. Bioplastics produced from poly-3-hydroxy butyrate-co-3-hydroxyvalerate [P(3HB-3HV)] can be degraded within weeks to a few years depending on the environmental conditions. These timescales are significantly lower than for non-biodegradable plastics such as polypropylene (Mergaert et al., 1994). Diverse products have been manufactured using PHA, for a summary with numerous patents with inventions from P(3HB-3HV) see Madison and Huisman (1999).

The accumulation of PHAs in methanotrophs is a survival response under nutrient starvation which re-routes methane into PHA production instead of converting it to carbon dioxide under normal conditions (Karthikeyan et al., 2015). In nature, the accumulation of PHA occurs under fluctuating nutrient (nitrogen, carbon) availability. In proteobacterial methanotrophs, only Type II are known to possess the genetic capability to express enzymes that produce PHA and contain pathways to direct the TCA cycle into PHA production (Pieja et al., 2011a). The genes *phaA*, *phaB*, and *phaC* encode for the enzymes that perform the condensation of two molecules of Acetyl-CoA, their reduction into (R)-3-hydroxybutyl-CoA, and finally the polymerization into the final PHA molecule, respectively (Peoples and Sinskey, 1989).

Overall, the growth conditions that lead to the accumulation of PHAs are: (i) absence of ammonia or nitrate as nitrogen source, (ii) copper deprivation and (iii) a regime of fed-batch to induce storage of carbon in the cell. *Methylocystis* sp. Rockwell, *Methylocystis* sp. WRR1, *Methylosinus trichosporium* OB3b, *Methylomicrobium album* BG8, and *Methylomonas denitrificans* FJG1 have been subjected to studies aiming to determine the preferred nitrogen (ammonia or nitrate) and carbon (methane or methanol) source (Tays et al., 2018). In general ammonia and methane are preferred as nitrogen and carbon source, respectively; however, the in-depth characterization of each strain is crucial before the application for industrial use where maximum yields, minimum inhibition and maximum efficiency of product are needed. Using a combination of continuous culture, followed by nitrogen limitation under fed-batch has proven to be successful in the enhancement of PHB accumulation in *Methylocystis* sp. GB25 DSM 7674, reaching 51% of cell mass as PHB (Wendlandt et al., 2005). Another key regulator are metals, in particular lanthanides which modulate the activity of the MDH, and play a crucial role in the carbon flow within the cell which have implications on the routes to carbon storage in

the form of PHA (Akberdin et al., 2018). Inherent to the vast diversity of methanotrophs, nutrient conditions and in particular micronutrients, require finetuning according to the metabolic needs of particular strains. **Table 3** provides a summary of those requirements in individual culture cases.

The use of augmented biogas from wastewater is already applied at full-scale (Mango Materials®). However, more research on lower quality biogas use is needed to exploit emerging opportunities with biogas as a substrate while mitigating GHG emissions from waste management.

Single Cell Protein (SCP) From Methanotrophs

Another methanotroph-derived biopolymer includes proteins from single cells (SCP). SCP can be purified from all methanotrophs as all carbon fixation pathways yield structural proteins that form new cells during microbial growth, and its use dates back to the early 1960s as the first bio-based product from methane (Bewersdorff and Dostálek, 1971).

All proteins are formed by polymers of amino acids and depending on their quaternary structure and function, harbor cofactors such as vitamins and metals, making them a suitable feed source for diverse animals regardless of their origin (Skrede et al., 1998, 2003; Øverland et al., 2010). Proteins of microbial origin have better yields compared to other protein sources and offers comparable nutritional quality when used as feed according to FAO recommendations (Anupama and Ravindra, 2000; Ritala et al., 2017). The expanding population growth and the need to feed nearly 10 billion people by 2050 (FAO, 2009), demand for sustainable protein sources. These are crucial in replacing the massive exploitation of marine and freshwater resources where the increase in aquaculture is a recent example of a sustainable practice that still requires significant amounts of protein as feed for produce (Ritala et al., 2017).

Ideal SCP methanotrophic strains should have a fast growth rate with high protein production capacity and easy to cultivate. Furthermore, the methanotrophs should be able to tolerate a wide range of pH and temperatures. The use of methane from natural gas for the production of SCP, is already approved for animal feed in agriculture (Wang et al., 2020). Currently SCP is produced from methane-rich natural gas and commercially applied at an industrial scale (Strong et al., 2015). Other sources deemed ideal for the production of SCP include a variety of solid and liquid waste streams (Wang et al., 2020), whereas biogas from AD remains overlooked.

TABLE 3 | Summary of selected laboratory enrichments and the conditions involved for the production of PHA in methanotrophic bacteria.

| Organism | Cultivation condition | Effect | Reference |
|--|---|--|------------------------|
| <i>Methylocystis</i> sp. GB25 DSM 7674 | Enriched culture (86%) Potassium deficiency. | 45% PHB yield per g of methane. | Helm et al., 2008 |
| <i>Methylocystis</i> sp. GB25 DSM 7674 | Continuous and fed-batch alternation, nitrogen limitation. | 55% PHB yield per g of methane. | Wendlandt et al., 2005 |
| <i>Methylocystis parvus</i> OBPP | Reducing substrates methane and formate | Enable PHB utilization as carbon source. | Pieja et al., 2011b |
| <i>Methylocystis</i> sp. WRR1 | Flasks cultures with Valerate or n-propanol additions. Copper deprivation conditions. | Decrease in melting temperature and crystallinity of PHB-co-HV 67% PHB per g of methane. | Cal et al., 2016 |

For more reports on PHB efficiencies, see Pieja et al. (2017).

A challenge for the successful application of methanotrophs is the poor solubility of methane in the aqueous phase (Bennett et al., 2018). Different reactor configurations that enhance the mass transfer have been tried successfully at a laboratory scale. Petersen et al. (2017) used a forced flow U-loop reactor to study the effects of gas transfer and reported higher efficiencies when compared to canonical stirred tank reactors (STR) or even tubular reactors (Petersen et al., 2017). This application is available at a commercial scale using natural gas as a carbon source for the methanotrophs by UniBio A/S and Calysta Inc., for the production of UniProtein® and FeedKind® respectively. The methanotrophs used are commonly composed of a mix culture with unknown ratios. *Methylococcus capsulatus* or *Methylomonas* sp. as the most common strains described in the literature. For more examples of SCP from bacteria at an industrial scale, see Jones et al. (2020). Current trends are moving from methane-rich sources such as natural gas, to more circular economy approaches using waste streams such as wastewater. The concept of using co-cultures i.e., *Methylococcus capsulatus* with algae *C. sorokiniana*, has been proposed and research under conditions with supplementation of artificial waste biogas (60% methane) for SCP production showed promising results (Rasouli et al., 2018). Implementation of SCP production at a pilot-scale under real wastewater conditions has not been proven successful to date.

After the description of the relevance of anaerobic methanotrophs to counteract GHG emissions, and the role of aerobic methanotrophs in a bio-based and circular economy for a sustainable future, the value offered by methanotrophs seems immeasurable.

CONCLUDING REMARKS

After nearly 120 years of discoveries, diverse (an)aerobic methanotrophs have been identified in widespread environments as key drivers of Earth's carbon fluxes, playing a crucial role as the sole biological methane sink. Methanotrophs determine the fate of methane in both natural and human-impacted ecosystems, where the mitigation of greenhouse gas emissions is recognized as one of the most important environmental development goals. Importantly, methanotrophs have the potential to transform methane into valuable products.

REFERENCES

- Agrawal, S., Seuntjens, D., Cocker, P., De Lackner, S., and Vlaeminck, S. E. (2018). Success of mainstream partial nitrification/anammox demands integration of engineering, microbiome and modeling insights. *Curr. Opin. Biotechnol.* 50, 214–221. doi: 10.1016/j.copbio.2018.01.013
- Akberdin, I. R., Collins, D. A., Hamilton, R., Oshchepkov, D. Y., Shukla, A. K., Nicora, C. D., et al. (2018). Rare earth elements alter redox balance in methylomicrobium alcaliphilum 20ZR. *Front. Microbiol.* 9:2735. doi: 10.3389/fmicb.2018.02735
- Alleman, J. E., and Prakasam, T. B. S. (1983). Reflections on seven decades of activated sludge history. *J. Water Pollut. Cont. Federat.* 55, 436–443.
- Amaro, T. M. M., Rosa, D., Comi, G., and Iacumin, L. (2019). Prospects for the use of whey for polyhydroxyalkanoate (PHA) production. *Front. Microbiol.* 10:992. doi: 10.3389/fmicb.2019.00992
- Anupama, and Ravindra, P. (2000). Value-added food: single cell protein. *Biotechnol. Adv.* 18, 459–479. doi: 10.1016/S0734-9750(00)00045-8
- Aromokeye, D. A., Kulkarni, A. C., Elvert, M., Wegener, G., Henkel, S., Coffinet, S., et al. (2020). Rates and microbial players of iron-driven anaerobic oxidation of methane in methanic marine sediments. *Front. Microbiol.* 10:3041. doi: 10.3389/fmicb.2019.03041
- Arshad, A., Speth, D. R., De Graaf, R. M., Op den Camp, H. J. M., Jetten, M. S. M., and Welte, C. U. (2015). A metagenomics-based metabolic model of nitrate-dependent anaerobic oxidation of methane by Methanoperedens-like archaea. *Front. Microbiol.* 18:1423. doi: 10.3389/fmicb.2015.01423
- Bachmann, N. (2015). *Sustainable Biogas Production in Municipal Wastewater Treatment Plants*. Paris: IEA Bioenergy.
- Bai, Y. N., Wang, X. N., Wu, J., Lu, Y. Z., Fu, L., Zhang, F., et al. (2019). Humic substances as electron acceptors for anaerobic oxidation of methane

Application of methanotrophs for both climate change mitigation and resource recovery in a circular economy face parallel challenges, namely limitations inherent to using methane as a substrate, and the need to revolutionize current chemical-based industrial practices to bio-based solutions. Past research on methanotroph ecology, physiology, and genomics yielded exciting discoveries. Future research direction on novel engineering solutions incorporating methane waste streams from human-influenced ecosystems (WWT), could lead to a new era of methanotrophic biorefineries, move waste management and production systems close to zero net emissions, and reconcile human intervention in the carbon cycle.

AUTHOR CONTRIBUTIONS

SGC, AV, and AH wrote the manuscript. MP, MAH, and HN contributed to the critical review and streamlining of the manuscript, SGC and AH coordinated the manuscript conceptualization. All authors contributed to the final version of the manuscript.

FUNDING

SGC has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie (Grant no. 892322, project MICROWATER). MP received support from the CERCA program for Generalitat de Catalunya and from the Economy and Knowledge Department of the Catalan Government through a Consolidated Research Group (ICRA-TECH - 2017 SGR 1318). AH received funding from the Deutsche Forschungsgemeinschaft (Grant no. HO6234/1-1). HN and AV received funding from the European Research Council (ERC-CoG Grant no. 772923, project VORTEX).

ACKNOWLEDGMENTS

We thank Anniek de Jong and Stefanie Frisch for helpful discussions on methanotrophy and methanogenesis, respectively.

- driven by ANME-2d. *Water Res.* 164, 114935. doi: 10.1016/j.watres.2019.114935
- Bandara, W. M. K. R. T. W., Satoh, H., Sasakawa, M., Nakahara, Y., Takahashi, M., and Okabe, S. (2011). Removal of residual dissolved methane gas in an upflow anaerobic sludge blanket reactor treating low-strength wastewater at low temperature with degassing membrane. *Water Res.* 45, 3533–3540. doi: 10.1016/j.watres.2011.04.030
- Bar-Or, I., Elvert, M., Eckert, W., Kushmaro, A., Vigderovich, H., Zhu, Q., et al. (2017). Iron-Coupled anaerobic oxidation of methane performed by a mixed bacterial-archaeal community based on poorly reactive minerals. *Environ. Sci. Technol.* 51, 12293–12301. doi: 10.1021/acs.est.7b03126
- Beal, E. J., House, C. H., and Orphan, V. J. (2009). Manganese- and iron-dependent marine methane oxidation. *Science* 325, 184–187. doi: 10.1126/science.1169984
- Bennett, R. K., Steinberg, L. M., Chen, W., and Papoutsakis, E. T. (2018). Engineering the bioconversion of methane and methanol to fuels and chemicals in native and synthetic methylophs. *Curr. Opin. Biotechnol.* 50, 81–93. doi: 10.1016/j.copbio.2017.11.010
- Berghuis, B. A., Yu, F. B., Schulz, F., Blainey, P. C., Woyke, T., and Quake, S. R. (2019). Hydrogenotrophic methanogenesis in archaeal phylum Verstraetearchaeota reveals the shared ancestry of all methanogens. *Proc. Natl. Acad. Sci.* 116, 5037–5044. doi: 10.1073/pnas.1815631116
- Bewersdorff, M., and Dostálek, M. (1971). The use of methane for production of bacterial protein. *Biotechnol. Bioeng.* 13, 49–62. doi: 10.1002/bit.260130104
- Björck, C., Dobson, P., and Pandhal, J. (2018). Biotechnological conversion of methane to methanol: evaluation of progress and potential. *AIMS Bioeng.* 5, 1–38. doi: 10.3934/bioeng.2018.1.1
- Blees, J., Niemann, H., Wenk, C. B., Zopf, J., Schubert, C. J., Kirf, M. K., et al. (2014). Micro-aerobic bacterial methane oxidation in the chemocline and anoxic water column of deep south-Alpine Lake Lugano (Switzerland). *Limnol. Oceanogr.* 59, 311–324. doi: 10.4319/lo.2014.59.2.0311
- Boetius, A., and Wenzhöfer, F. (2013). Seafloor oxygen consumption fuelled by methane from cold seeps. *Nat. Geosci.* 6, 725–734. doi: 10.1038/ngeo1926
- Boetius, A., Ravensschlag, K., Schubert, C. J., Rickert, D., Widdel, F., Gieseke, A., et al. (2000). A marine microbial consortium apparently mediating anaerobic oxidation methane. *Nature* 407, 623–626. doi: 10.1038/35036572
- Bollinger, J. M. (2010). Biochemistry: getting the metal right. *Nature* 465, 40–41. doi: 10.1038/465040a
- Börjesson, G. (2001). Inhibition of methane oxidation by volatile sulfur compounds (CH₃SH and CS₂) in landfill cover soils. *Waste Manag. Res.* 19, 314–319. doi: 10.1177/0734242X0101900408
- Buswell, A. M., and Hatfield, W. D. (1936). *Anaerobic Fermentation. Division of the State Water Survey (Illinois), Bulletin*. Urbana, IL: Department of Registration and Education, 32.
- Cai, C., Hu, S., Chen, X., Ni, B. J., Pu, J., and Yuan, Z. (2018a). Effect of methane partial pressure on the performance of a membrane biofilm reactor coupling methane-dependent denitrification and anammox. *Sci. Total Environ.* 639, 278–285. doi: 10.1016/j.scitotenv.2018.05.164
- Cai, C., Hu, S., Guo, J., Shi, Y., Xie, G. J., and Yuan, Z. (2015). Nitrate reduction by denitrifying anaerobic methane oxidizing microorganisms can reach a practically useful rate. *Water Res.* 87, 211–217. doi: 10.1016/j.watres.2015.09.026
- Cai, C., Leu, A. O., Xie, G. J., Guo, J., Feng, Y., Zhao, J. X., et al. (2018b). A methanotrophic archaeon couples anaerobic oxidation of methane to Fe(III) reduction. *ISME J.* 12, 1929–1939. doi: 10.1038/s41396-018-0109-x
- Cal, A. J., Sikkema, W. D., Ponce, M. I., Franqui-Villanueva, D., Riiff, T. J., Orts, W. J., et al. (2016). Methanotrophic production of polyhydroxybutyrate-co-hydroxyvalerate with high hydroxyvalerate content. *Int. J. Biol. Macromol.* 87, 302–307. doi: 10.1016/j.ijbiomac.2016.02.056
- Cantera, S., Lebrero, R., García-Encina, P. A., and Muñoz, R. (2016). Evaluation of the influence of methane and copper concentration and methane mass transport on the community structure and biodegradation kinetics of methanotrophic cultures. *J. Environ. Manag.* 171, 11–20. doi: 10.1016/j.jenvman.2016.02.002
- Cantera, S., Lebrero, R., Rodríguez, E., García-Encina, P. A., and Muñoz, R. (2017). Continuous abatement of methane coupled with ectoine production by *Methylobacterium alcaliphilum* 20Z in stirred tank reactors: a step further towards greenhouse gas biorefineries. *J. Cleaner Product.* 152, 134–141. doi: 10.1016/j.jclepro.2017.03.123
- Caranto, J. D., Vilvert, A. C., and Lancaster, K. M. (2016). *Nitrosomonas europaea* cytochrome P640 is a direct link between nitrification and nitrous oxide emission. *Proc. Natl. Acad. Sci. U.S.A.* 113, 14704–14709. doi: 10.1073/pnas.1611051113
- Carvajal-Arroyo, J. M., Sun, W., Sierra-Alvarez, R., and Field, J. A. (2013). Inhibition of anaerobic ammonium oxidizing (anammox) enrichment cultures by substrates, metabolites and common wastewater constituents. *Chemosphere* 91, 22–27. doi: 10.1016/j.chemosphere.2012.11.025
- Castelle, C. J., and Banfield, J. F. (2018). Major new microbial groups expand diversity and alter our understanding of the tree of life. *Cell* 172, 1181–1197. doi: 10.1016/j.cell.2018.02.016
- Cervantes, F. J., Van Der Velde, S., Lettinga, G., and Field, J. A. (2000). Competition between methanogenesis and quinone respiration for ecologically important substrates in anaerobic consortia. *FEMS Microbiol. Ecol.* 34, 161–171. doi: 10.1016/S0168-6496(00)00091-X
- Chang, J., Kim, D. D., Semrau, J. D., Lee, J., Heo, H., Gu, W., et al. (2020). Enhancement of nitrous oxide emissions in soil microbial consortia via copper competition between proteobacterial methanotrophs and denitrifiers. *Appl. Environ. Microbiol.* 87, e2301–e2320. doi: 10.1128/aem.02301-20
- Chen, C. Q., and Jiang, X. R. (2018). Next generation industrial biotechnology based on extremophilic bacteria. *Curr. Opin. Biotechnol.* 50, 94–100. doi: 10.1016/j.copbio.2017.11.016
- Chen, H., Liu, T., Li, J., Mao, L., Ye, J., Han, X., et al. (2020). Larger anammox granules not only harbor higher species diversity but also support more functional diversity. *Environ. Sci. Technol.* 54, 14664–14673. doi: 10.1021/acs.est.0c02609
- Chen, J., Zhou, Z. C., and Gu, J. D. (2014). Occurrence and diversity of nitrite-dependent anaerobic methane oxidation bacteria in the sediments of the South China Sea revealed by amplification of both 16S rRNA and pmoA genes. *Appl. Microbiol. Biotechnol.* 98, 5685–5696. doi: 10.1007/s00253-014-5733-4
- Chen, K. H. C., Wu, H. H., Ke, S. F., Rao, Y. T., Tu, C. M., Chen, Y. P., et al. (2012). Bacteriohemerythrin bolsters the activity of the particulate methane monooxygenase (pMMO) in *Methylococcus capsulatus* (Bath). *J. Inorg. Biochem.* 111, 10–17. doi: 10.1016/j.jinorgbio.2012.02.019
- Chen, X., Guo, J., Xie, G. J., Liu, Y., Yuan, Z., and Ni, B. J. (2015). A new approach to simultaneous ammonium and dissolved methane removal from anaerobic digestion liquor: a model-based investigation of feasibility. *Water Res.* 85, 295–303. doi: 10.1016/j.watres.2015.08.046
- Chevalier, N., Bouloubassi, I., Stadnitskaia, A., Taphanel, M. H., and Sinnighe Damsté, J. S. (2014). Lipid biomarkers for anaerobic oxidation of methane and sulphate reduction in cold seep sediments of Nyegga pockmarks (Norwegian margin): discrepancies in contents and carbon isotope signatures. *Geo Mar. Lett.* 34, 269–280. doi: 10.1007/s00367-014-0363-5
- Coates, J. D., Ellis, D. J., Blunt-Harris, E. L., Gaw, C. V., Roden, E. E., and Lovley, D. R. (1998). Recovery of humic-reducing bacteria from a diversity of environments. *Appl. Environ. Microbiol.* 64, 1504–1509. doi: 10.1128/aem.64.4.1504-1509.1998
- Collins, W. J., Webber, C. P., Cox, P. M., Huntingford, C., Lowe, J., Sitch, S., et al. (2018). Increased importance of methane reduction for a 1.5 degree target. *Environ. Res. Lett.* 13:54003. doi: 10.1088/1748-9326/aab89c
- Comer, A. D., Long, M. R., Reed, J. L., and Pflieger, B. F. (2017). Flux balance analysis indicates that methane is the lowest cost feedstock for microbial cell factories. *Metab. Eng. Commun.* 5, 26–33. doi: 10.1016/j.meteno.2017.07.002
- Conrad, R. (1996). Soil microorganisms as controllers of atmospheric trace gases (H₂, CO, CH₄, OCS, N₂O, and NO). *Microbiol. Rev.* 60, 609–640. doi: 10.1128/mbr.60.4.609-640.1996
- Conrad, R. (2009). The global methane cycle: recent advances in understanding the microbial processes involved. *Environ. Microbiol. Rep.* 1, 285–292. doi: 10.1111/j.1758-2229.2009.00038.x
- Conrad, R. (2020). Importance of hydrogenotrophic, acetoclastic and methylophic methanogenesis for methane production in terrestrial, aquatic and other anoxic environments: a mini review. *Pedosphere* 30, 25–39. doi: 10.1016/S1002-0160(18)60052-9
- Costa, C., Dijkema, C., Friedrich, M., García-Encina, P., Fernández-Polanco, F., and Stams, A. J. M. (2000). Denitrification with methane as electron donor in oxygen-limited bioreactors. *Appl. Microbiol. Biotechnol.* 53, 754–762. doi: 10.1007/s002530000337

- Crombie, A. T., and Murrell, J. C. (2014). Trace-gas metabolic versatility of the facultative methanotroph *Methylocella silvestris*. *Nature* 510, 148–151. doi: 10.1038/nature13192
- Da Silva, M. J. (2016). Synthesis of methanol from methane: challenges and advances on the multi-step (syngas) and one-step routes (DMTM). *Fuel Process. Technol.* 145:42. doi: 10.1016/j.fuproc.2016.01.023
- Daelman, M. R. J., Van Voorthuizen, E. M., Van Dongen, L. G. J. M., Volcke, E. I. P., and Van Loosdrecht, M. C. M. (2013). Methane and nitrous oxide emissions from municipal wastewater treatment - Results from a long-term study. *Water Sci. Technol.* 67, 2350–2355. doi: 10.2166/wst.2013.109
- Daelman, M. R. J., van Voorthuizen, E. M., van Dongen, U. G. J. M., Volcke, E. I. P., and van Loosdrecht, M. C. M. (2012). Methane emission during municipal wastewater treatment. *Water Res.* 46, 3657–3670. doi: 10.1016/j.watres.2012.04.024
- Davies, T. R. (1973). Isolation of bacteria capable of utilizing methane as a hydrogen donor in the process of denitrification. *Water Res.* 7, 575–579. doi: 10.1016/0043-1354(73)90056-0
- Dean, J. F., Middelburg, J. J., Röckmann, T., Aerts, R., Blauw, L. G., Egger, M., et al. (2018). Methane feedbacks to the global climate system in a warmer world. *Rev. Geophys.* 56, 207–250. doi: 10.1002/2017RG000559
- Dedysh, S. N., and Dunfield, P. F. (2016). "Methylocella," in *Bergey's Manual of Systematics of Archaea and Bacteria*, eds Trujillo, M. E., Dedysh, S., DeVos, P., Hedlund, B., Kämpfer, P., Rainey, F. A., and Whitman, W. B. (Hoboken, NJ: John Wiley & Sons, Inc.). doi: 10.1002/9781118960608.gbm00797.pub2
- Dedysh, S. N., and Knief, C. (2018). *Diversity and Phylogeny of Described Aerobic Methanotrophs. Methane Biocatalysis: Paving the Way to Sustainability*. Manhattan, NY: Springer International Publishing. doi: 10.1007/978-3-319-74866-5_2
- Deutzmann, J. S., Stief, P., Brandes, J., and Schink, B. (2014). Anaerobic methane oxidation coupled to denitrification is the dominant methane sink in a deep lake. *Proc. Natl. Acad. Sci. U.S.A.* 111, 18273–18278. doi: 10.1073/pnas.1411617111
- Deutzmann, J. S., Wörner, S., and Schink, B. (2011). Activity and diversity of methanotrophic acteria at methane seeps in eastern lake Constance sediments. *Appl. Environ. Microbiol.* 77, 2573–2581. doi: 10.1128/AEM.02776-10
- Ding, J., and Zeng, R. J. (2021). Fundamentals and potential environmental significance of denitrifying anaerobic methane oxidizing archaea. *Sci. Total Environ.* 757:143928. doi: 10.1016/j.scitotenv.2020.143928
- Ding, J., Lu, Y.-Z., Fu, L., Ding, Z.-W., Mu, Y., Cheng, S. H., et al. (2017). Decoupling of DAMO archaea from DAMO bacteria in a methane-driven microbial fuel cell. *Water Res.* 110, 112–119. doi: 10.1016/j.watres.2016.12.006
- Ding, Z. W., Ding, J., Fu, L., Zhang, F., and Zeng, R. J. (2014). Simultaneous enrichment of denitrifying methanotrophs and anammox bacteria. *Appl. Microbiol. Biotechnol.* 98, 10211–10221. doi: 10.1007/s00253-014-5936-8
- Dombrowski, N., Lee, J. H., Williams, T. A., Offre, P., and Spang, A. (2019). Genomic diversity, lifestyles and evolutionary origins of DPANN archaea. *FEMS Microbiol. Lett.* 366:Fnz008. doi: 10.1093/femsle/fnz008
- Dong, T., Fei, Q., Genelot, M., Smith, H., Laurens, L. M. L., Watson, M. J., et al. (2017). A novel integrated biorefinery process for diesel fuel blendstock production using lipids from the methanotroph, *Methylomicrobium buryatense*. *Energy Convers. Manag.* 140, 62–70. doi: 10.1016/j.enconman.2017.02.075
- Drake, H. L., Horn, M. A., and Wust, P. K. (2009). Intermediary ecosystem metabolism as a main driver of methanogenesis in acidic wetland soil. *Environ. Microbiol. Rep.* 1, 307–318. doi: 10.1111/j.1758-2229.2009.00050.x
- Duan, C., Luo, M., and Xing, X. (2011). High-rate conversion of methane to methanol by *Methylosinus trichosporium* OB3b. *Biores. Technol.* 102, 7349–7353. doi: 10.1016/j.biortech.2011.04.096
- Dunfield, P. F., Yuryev, A., Senin, P., Smirnova, A. V., Stott, M. B., Hou, S., et al. (2007). Methane oxidation by an extremely acidophilic bacterium of the phylum Verrucomicrobia. *Nature* 540, 879–882. doi: 10.1038/nature06411
- Egger, M., Rasigraf, O., Sapart, C. J., Jilbert, T., Jetten, M. S. M., Röckmann, T., et al. (2015). Iron-mediated anaerobic oxidation of methane in brackish coastal sediments. *Environ. Sci. Technol.* 49, 277–283. doi: 10.1021/es503663z
- Enzmann, F., Mayer, F., Rother, M., and Holtmann, D. (2018). Methanogens: biochemical background and biotechnological applications. *AMB Expr.* 8:531. doi: 10.1186/s13568-017-0531-x
- Etmann, M., Myhre, G., Highwood, E. J., and Shine, K. P. (2016). Radiative forcing of carbon dioxide, methane, and nitrous oxide: a significant revision of the methane radiative forcing. *Geophys. Res. Lett.* 43, 614–623. doi: 10.1002/2016GL071930
- Ettwig, K. F., Butler, M. K., Le Paslier, D., Pelletier, E., Mangenot, S., Kuypers, M. M., et al. (2010). Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* 464, 543–548. doi: 10.1038/nature08883
- Ettwig, K. F., Shima, S., Van De Pas-Schoonen, K. T., Kahnt, J., Medema, M. H., Op Den Camp, H. J. M., et al. (2008). Denitrifying bacteria anaerobically oxidize methane in the absence of Archaea. *Environ. Microbiol.* 10, 3164–3173. doi: 10.1111/j.1462-2920.2008.01724.x
- Ettwig, K. F., Speth, D. R., Reimann, J., Wu, M. L., Jetten, M. S. M., and Keltjens, J. T. (2012). Bacterial oxygen production in the dark. *Front. Microbiol.* 3:273. doi: 10.3389/fmicb.2012.00273
- Ettwig, K. F., Van Alen, T., Van De Pas-Schoonen, K. T., Jetten, M. S. M., and Strous, M. (2009). Enrichment and molecular detection of denitrifying methanotrophic bacteria of the NC10 phylum. *Appl. Environ. Microbiol.* 75, 3656–3662. doi: 10.1128/AEM.00067-09
- Ettwig, K. F., Zhu, B., Speth, D., Keltjens, J. T., Jetten, M. S. M., and Kartal, B. (2016). Archaea catalyze iron-dependent anaerobic oxidation of methane. *Proc. Natl. Acad. Sci. U.S.A.* 113, 12792–12796. doi: 10.1073/pnas.1609534113
- Evans, P. N., Parks, D. H., Chadwick, G. L., Robbins, S. J., Orphan, V. J., Golding, S. D., et al. (2015). Methane metabolism in the archaeal phylum Bathyarchaeota revealed by genome-centric metagenomics. *Science* 350, 434–438. doi: 10.1126/science.aac7745
- Fan, S. Q., Xie, G. J., Lu, Y., Liu, B. F., Xing, D. F., Han, H. J., et al. (2019). Granular Sludge coupling nitrate/nitrite dependent anaerobic methane oxidation with anammox: from proof-of-concept to high rate nitrogen removal. *Environ. Sci. Technol.* 54, 297–305. doi: 10.1021/acs.est.9b02528
- FAO (2009). *How to Feed the World in 2050. Proceedings of the Fsn forum Discussion no. 43*. Rome: FAO.
- Farhan Ul Haque, M., Crombie, A. T., and Murrell, J. C. (2019). Novel facultative *Methylocella* strains are active methane consumers at terrestrial natural gas seeps. *Microbiome* 7:134. doi: 10.1186/s40168-019-0741-3
- Farhan Ul Haque, M., Xu, H.-J., Murrell, J. C., and Crombie, A. (2020). Facultative methanotrophs – diversity, genetics, molecular ecology and biotechnological potential: a mini-review. *Microbiology* 166, 894–908. doi: 10.1099/mic.0.000977
- Fei, Q., Puri, A. W., Smith, H., Dowe, N., and Pienkos, P. T. (2018). Enhanced biological fixation of methane for microbial lipid production by recombinant *Methylomicrobium buryatense*. *Biotechnol. Biofuels* 11:129. doi: 10.1186/s13068-018-1128-6
- Fox, G. E., Magrum, L. J., and Balch, W. E. (1977). Classification of methanogenic bacteria by 16S ribosomal RNA characterization. *Proc. Natl. Acad. Sci. U.S.A.* 74, 4537–4541. doi: 10.1073/pnas.74.10.4537
- Fu, L., Ding, J., Lu, Y. Z., Ding, Z. W., Bai, Y. N., and Zeng, R. J. (2017). Hollow fiber membrane bioreactor affects microbial community and morphology of the DAMO and Anammox co-culture system. *Bioresour. Technol.* 232, 247–253. doi: 10.1016/j.biortech.2017.02.048
- Furimsky, E. (2000). Catalytic hydrodeoxygenation. *Appl. Catal. Gen.* 199, 147–190. doi: 10.1016/S0926-860X(99)00555-4
- Gambelli, L., Guerrero-Cruz, S., Mesman, R. J., Cremers, G., Jetten, M. S. M., Op den Camp, H. J. M., et al. (2018). Community composition and ultrastructure of a nitrate-dependent anaerobic methaneoxidizing enrichment culture. *Appl. Environ. Microbiol.* 84:2186. doi: 10.1128/AEM.02186-17
- Graf, J. S., Mayr, M. J., Marchant, H. K., Tienken, D., Hach, P. F., Brand, A., et al. (2018). Bloom of a denitrifying methanotroph, 'Candidatus Methyloirabilis limnetica', in a deep stratified lake. *Environ. Microbiol.* 20, 2598–2614. doi: 10.1111/1462-2920.14285
- Guerrero-Cruz, S. (2018). *Ecophysiology of Nitrate and Nitrite-Dependent Methane Oxidation*. Nijmegen: Radboud University.
- Guerrero-Cruz, S., Cremers, G., van Alen, T. A., Op den Camp, H. J. M., Jetten, M. S. M., Rasigraf, O., et al. (2018). Response of the anaerobic methanotroph "Candidatus Methanoperedens nitroreducens" to oxygen stress. *Appl. Environ. Microbiol.* 84, e1832–e1818. doi: 10.1128/AEM.01832-18
- Guerrero-Cruz, S., Stultjens, K., van Kessel, M. A. H. J., Versantvoort, W., Jetten, M. S. M., den Camp, H. J. M. O., et al. (2019). Key physiology of a nitrite-dependent methane-oxidizing enrichment culture. *Appl. Environ. Microbiol.* 85, e124–e119. doi: 10.1128/AEM.00124-19

- Guisasola, A., de Haas, D., Keller, J., and Yuan, Z. (2008). Methane formation in sewer systems. *Water Res.* 42, 1421–1430. doi: 10.1016/j.watres.2007.10.014
- Guisasola, A., Sharma, K. R., Keller, J., and Yuan, Z. (2009). Development of a model for assessing methane formation in rising main sewers. *Water Res.* 43, 2874–2884. doi: 10.1016/j.watres.2009.03.040
- Hakobyan, A., and Liesack, W. (2020). Unexpected metabolic versatility among type II methanotrophs in the Alphaproteobacteria. *Biol. Chem.* 401, 1469–1477. doi: 10.1515/hsz-2020-0200
- Hallam, S. J., Putnam, N., Preston, C. M., Detter, J. C., Rokhsar, D., Richardson, P. H., et al. (2004). Reverse methanogenesis: testing the hypothesis with environmental genomics. *Science* 305, 1457–1462. doi: 10.1126/science.1100025
- Han, J. S., Ahn, C. M., Mahanty, B., and Kim, C. G. (2013). Partial oxidative conversion of methane to methanol through selective inhibition of methanol dehydrogenase in methanotrophic consortium from landfill cover soil. *Appl. Biochem. Biotechnol.* 171, 1487–1499. doi: 10.1007/s12010-013-0410-0
- Hanson, R. S., and Hanson, T. E. (1996). Methanotrophic bacteria. *Microbiol. Rev.* 60, 439–471. doi: 10.1128/mmmbr.60.2.439-471.1996
- Haroon, M. F., Hu, S., Shi, Y., Imelfort, M., Keller, J., Hugenholtz, P., et al. (2013). Anaerobic oxidation of methane coupled to nitrate reduction in a novel archaeal lineage. *Nature* 500, 567–570. doi: 10.1038/nature12375
- Hatamoto, M., Nemoto, S., and Yamaguchi, T. (2018). Effects of copper and PQQ on the denitrification activities of microorganisms facilitating nitrite- and nitrate-dependent DAMO reaction. *Int. J. Environ. Res.* 12, 749–753. doi: 10.1007/s41742-018-0118-7
- Haynes, W. M. (2011). *CRC Handbook of Chemistry and Physics*, 92 Edn. Hoboken, NJ: CRC Press. doi: 10.1201/b17379
- He, R., Su, Y., Leewis, M. C., Chu, Y. X., Wang, J., Ma, R. C., et al. (2020). Low O₂ level enhances CH₄-derived carbon flow into microbial communities in landfill cover soils. *Environ. Pollut.* 258:113676. doi: 10.1016/j.envpol.2019.113676
- He, R., Wooller, M. J., Pohlman, J. W., Quensen, J., Tiedje, J. M., and Leigh, M. B. (2012). Diversity of active aerobic methanotrophs along depth profiles of arctic and subarctic lake water column and sediments. *ISME J.* 6, 1937–1948. doi: 10.1038/ismej.2012.34
- He, Z., Cai, C., Geng, S., Lou, L., Xu, X., Zheng, P., et al. (2013). Modeling a nitrite-dependent anaerobic methane oxidation process: parameters identification and model evaluation. *Bioresour. Technol.* 147, 315–320. doi: 10.1016/j.biortech.2013.08.001
- He, Z., Geng, S., Cai, C., Liu, S., Liu, Y., and Hu, B. (2015). Anaerobic oxidation of methane coupled to nitrite reduction by halophilic marine NC10 bacteria. *Appl. Environ. Microbiol.* 81, 5538–5545. doi: 10.1128/AEM.00984-15
- Helm, J., Wendlandt, K. D., Jechorek, M., and Stottmeister, U. (2008). Potassium deficiency results in accumulation of ultra-high molecular weight poly- β -hydroxybutyrate in a methane-utilizing mixed culture. *J. Appl. Microbiol.* 105, 1054–1061. doi: 10.1111/j.1365-2672.2008.03831.x
- Hinrichs, K. U., Hayes, J. M., Sylva, S. P., Brewert, P. G., and DeLong, E. F. (1999). Methane-consuming archaeobacteria in marine sediments. *Nature* 398, 802–805. doi: 10.1038/19751
- Hinrichs, K.-U., and Boetius, A. (2002). *The Anaerobic Oxidation of Methane: New Insights in Microbial Ecology and Biogeochemistry. Ocean Margin Systems*. Berlin: Springer. doi: 10.1007/978-3-662-05127-6_28
- Ho, A., and Bodelier, P. L. E. (2015). Diazotrophic methanotrophs in peatlands: the missing link? *Plant Soil* 389, 419–423. doi: 10.1007/s11104-015-2393-9
- Ho, A., Angel, R., Veraart, A. J., Daebeler, A., Jia, Z., Kim, S. Y., et al. (2016). Biotic interactions in microbial communities as modulators of biogeochemical processes: methanotrophy as a model system. *Front. Microbiol.* 7:1285. doi: 10.3389/fmicb.2016.01285
- Ho, A., De Roy, K., Thas, O., De Neve, J., Hoefman, S., Vandamme, P., et al. (2014). The more, the merrier: heterotroph richness stimulates methanotrophic activity. *ISME J.* 8, 1945–1948. doi: 10.1038/ismej.2014.74
- Ho, A., Kerckhof, F. M., Luke, C., Reim, A., Krause, S., Boon, N., et al. (2013a). Conceptualizing functional traits and ecological characteristics of methane-oxidizing bacteria as life strategies. *Environ. Microbiol. Rep.* 5, 335–345. doi: 10.1111/j.1758-2229.2012.00370.x
- Ho, A., Lüke, C., Reim, A., and Frenzel, P. (2013b). Selective stimulation in a natural community of methane oxidizing bacteria: effects of copper on pmoA transcription and activity. *Soil Biol. Biochem.* 65, 211–216. doi: 10.1016/j.soilbio.2013.05.027
- Ho, A., Mendes, L. W., Lee, H. J., Kaupper, T., Mo, Y., Poehlein, A., et al. (2020). Response of a methane-driven interaction network to stressor intensification. *FEMS Microbiol. Ecol.* 96:180. doi: 10.1093/femsec/iaa180
- Ho, A., Vlaeminck, S. E., Ettwig, K. F., Schneider, B., Frenzel, P., and Boon, N. (2013c). Revisiting methanotrophic communities in sewage treatment plants. *Appl. Environ. Microbiol.* 79, 2841–2846.
- Hoefman, S., van der Ha, D., Boon, N., Vandamme, P., De Vos, P., and Heylen, K. (2014). Niche differentiation in nitrogen metabolism among methanotrophs within an operational taxonomic unit Microbial biochemistry, physiology and metabolism. *BMC Microbiol.* 14:1471. doi: 10.1186/1471-2180-14-83
- Hogendoorn, C., Picone, N., van Hout, F., Vijverberg, S., Poghosyan, L., and Op den Camp, H. J. M. (2021). Draft genome of a novel methanotrophic *Methylobacter* sp. from the volcanic soils of Pantelleria Island. *Antonie Van Leeuwenhoek* 114, 313–324. doi: 10.1007/s10482-021-01525-7
- Hogendoorn, C., Pol, A., Nuijten, G. H. L., and Op den Camp, H. J. M. (2020). Methanol production by “*Methylacidiphilum fumarolicum*” solv under different growth conditions. *Appl. Environ. Microbiol.* 86, E1188–E1120. doi: 10.1128/AEM.01188-20
- Holmes, A. J., Tujula, N. A., Holley, M., Contos, A., James, J. M., Rogers, P., et al. (2001). Phylogenetic structure of unusual aquatic microbial formations in Nullarbor caves, Australia. *Environ. Microbiol.* 3, 256–264. doi: 10.1046/j.1462-2920.2001.00187.x
- Howarth, R. W. (2019). Ideas and perspectives: Is shale gas a major driver of recent increase in global atmospheric methane? *Biogeosciences* 16, 3033–3046. doi: 10.5194/bg-16-3033-2019
- Hu, S., Zeng, R. J., Burow, L. C., Lant, P., Keller, J., and Yuan, Z. (2009). Enrichment of denitrifying anaerobic methane oxidizing microorganisms. *Environ. Microbiol. Rep.* 5, 377–384. doi: 10.1111/j.1758-2229.2009.00083.x
- In ’t Zandt, M. H., de Jong, A. E. E., Slomp, C. P., and Jetten, M. S. M. (2018a). The hunt for the most-wanted chemolithoautotrophic spookmicrobes. *FEMS Microbiol. Ecol.* 94:fy064. doi: 10.1093/femsec/fiy064
- In ’t Zandt, M. H., van den Bosch, T. J. M., Rijkers, R., van Kessel, M. A. H. J., Jetten, M. S. M., and Welte, C. U. (2018b). Co-cultivation of the strictly anaerobic methanogen *Methanosarcina barkeri* with aerobic methanotrophs in an oxygen-limited membrane bioreactor. *Appl. Microbiol. Biotechnol.* 102, 5685–5694. doi: 10.1007/s00253-018-9038-x
- Intergovernmental Panel on Climate Change (IPCC) (2012). *Transport and its Infrastructure. Climate Change 2007*. Geneva: IPCC Intergovernmental Panel on Climate Change. doi: 10.1017/cbo9780511546013.009
- Intergovernmental Panel on Climate Change (IPCC) (2014). *Mitigation of Climate Change: Climate Change 2014*. Geneva: IPCC Intergovernmental Panel on Climate Change. doi: 10.1017/cbo9781107415416
- Intergovernmental Panel on Climate Change (IPCC) (2015). *Agriculture, Forestry and Other Land Use (AFOLU). Climate Change 2014 Mitigation of Climate Change*. Geneva: IPCC Intergovernmental Panel on Climate Change. doi: 10.1017/cbo9781107415416.017
- Intergovernmental Panel on Climate Change (IPCC) (2018). *Global Warming of 1.5°C*. Geneva: IPCC Intergovernmental Panel on Climate Change.
- Islam, T., Gessesse, A., Garcia-Moyano, A., Murrell, J. C., and Øvreås, L. (2020). A Novel moderately thermophilic Type Ib methanotroph isolated from an alkaline thermal spring in the Ethiopian rift valley. *Microorganisms* 8:250. doi: 10.3390/microorganisms8020250
- Islam, T., Hernández, M., Gessesse, A., Murrell, J. C., and Øvreås, L. (2021). A novel moderately thermophilic facultative methylotroph within the class Alphaproteobacteria. *Microorganisms* 9:903077. doi: 10.3390/microorganisms9030477
- Islam, T., Jensen, S., Reigstad, L. J., Larsen, Ø., and Birkeland, N. K. (2008). Methane oxidation at 55°C and pH 2 by a thermoacidophilic bacterium belonging to the *Verrucomicrobia* phylum. *Proc. Natl. Acad. Sci. U.S.A.* 105, 300–304. doi: 10.1073/pnas.0704162105
- Jing, H., Wang, R., Jiang, Q., Zhang, Y., and Peng, X. (2020). Anaerobic methane oxidation coupled to denitrification is an important potential methane sink in deep-sea cold seeps. *Sci. Total Environ.* 748:142459. doi: 10.1016/j.scitotenv.2020.142459

- Jones, S. W., Karpol, A., Friedman, S., Maru, B. T., and Tracy, B. P. (2020). Recent advances in single cell protein use as a feed ingredient in aquaculture. *Curr. Opin. Biotechnol.* 61, 189–197. doi: 10.1016/j.copbio.2019.12.026
- Kalyuzhnaya, M. G., Kumaresan, D., Heimann, K., Caetano, N. S., Visvanathan, C., and Parthiba Karthikeyan, O. (2020). Editorial: methane: a bioresource for fuel and biomolecules. *Front. Environ. Sci.* 8:9. doi: 10.3389/fenvs.2020.00009
- Kalyuzhnaya, M. G., Yang, S., Rozova, O. N., Smalley, N. E., Clubb, J., Lamb, A., et al. (2013). Highly efficient methane biocatalysis revealed in a methanotrophic bacterium. *Nat. Commun.* 4:2785. doi: 10.1038/ncomms3785
- Kampman, C., Piai, L., Temmink, H., Hendrickx, T. L. G., Zeeman, G., and Buisman, C. J. N. (2018). Effect of low concentrations of dissolved oxygen on the activity of denitrifying methanotrophic bacteria. *Water Sci. Technol.* 77, 2589–2597. doi: 10.2166/wst.2018.219
- Kampman, C., Temmink, H., Hendrickx, T. L. G., Zeeman, G., and Buisman, C. J. N. (2014). Enrichment of denitrifying methanotrophic bacteria from municipal wastewater sludge in a membrane bioreactor at 20°C. *J. Hazard. Mater.* 274, 428–435. doi: 10.1016/j.jhazmat.2014.04.031
- Kao, W. C., Chen, Y. R., Yi, E. C., Lee, H., Tian, Q., Wu, K. M., et al. (2004). Quantitative proteomic analysis of metabolic regulation by copper ions in *Methylococcus capsulatus* (Bath). *J. Biol. Chem.* 279, 51554–51560. doi: 10.1074/jbc.M408013200
- Kao, W. C., Wang, V. C. C., Huang, Y. C., Yu, S. S. F., Chang, T. C., and Chan, S. I. (2008). Isolation, purification and characterization of hemerythrin from *Methylococcus capsulatus* (Bath). *J. Inorgan. Biochem.* 102, 1607–1614. doi: 10.1016/j.jinorgbio.2008.02.008
- Karst, S. M., Dueholm, M. S., McIlroy, S. J., Kirkegaard, R. H., Nielsen, P. H., and Albertsen, M. (2018). Retrieval of a million high-quality, full-length microbial 16S and 18S rRNA gene sequences without primer bias. *Nat. Biotechnol.* 36, 190–195. doi: 10.1038/nbt.4045
- Karthikeyan, O. P., Chidambarampadmavathy, K., Cirés, S., and Heimann, K. (2015). Review of sustainable methane mitigation and biopolymer production. *Crit. Rev. Environ. Sci. Technol.* 45, 1579–1610. doi: 10.1080/10643389.2014.966422
- Kaupper, T., Mendes, L. W., Harnisz, M., Krause, S. M. B., Horn, M. A., and Ho, A. (2020). Recovery in methanotrophic activity does not reflect on the methane-driven interaction network after peat mining. *Appl. Environ. Microbiol.* 87:e02355–20. doi: 10.1128/aem.02355–20
- Kaupper, T., Mendes, L. W., Lee, H. J., Mo, Y., Poehlein, A., Jia, Z., et al. (2021). When the going gets tough: Emergence of a complex methane-driven interaction network during recovery from desiccation-rewetting. *Soil Biol. Biochem.* 153:108109. doi: 10.1016/j.soilbio.2020.108109
- Keltjens, J. T., Pol, A., Reimann, J., and Op Den Camp, H. J. M. (2014). PQQ-dependent methanol dehydrogenases: Rare-earth elements make a difference. *Appl. Microbiol. Biotechnol.* 98, 6163–6183. doi: 10.1007/s00253-014-5766-8
- Kirschke, S., Bousquet, P., Ciais, P., Saunoy, M., Canadell, J. G., Dlugokencky, E. J., et al. (2013). Three decades of global methane sources and sinks. *Nat. Geosci.* 6, 813–823. doi: 10.1038/ngeo1955
- Kits, K. D., Klotz, M. G., and Stein, L. Y. (2015). Methane oxidation coupled to nitrate reduction under hypoxia by the Gammaproteobacterium *Methylomonas denitrificans*, sp. nov. type strain FJG1. *Environ. Microbiol.* 17, 3219–3232. doi: 10.1111/1462-2920.12772
- Knapp, C. W., Fowle, D. A., Kulczycki, E., Roberts, J. A., and Graham, D. W. (2007). Methane monooxygenase gene expression mediated by methanobactin in the presence of mineral copper sources. *Proc. Natl. Acad. Sci.* 104, 12040L–12045. doi: 10.1073/pnas.0702879104
- Knief, C. (2015). Diversity and habitat preferences of cultivated and uncultivated aerobic methanotrophic bacteria evaluated based on pmoA as molecular marker. *Front. Microbiol.* 6:1346. doi: 10.3389/fmicb.2015.01346
- Knittel, K., and Boetius, A. (2009). Anaerobic oxidation of methane: progress with an unknown process. *Annu. Rev. Microbiol.* 63, 311–334. doi: 10.1146/annurev.micro.61.080706.093130
- Knittel, K., Lösekann, T., Boetius, A., Kort, R., and Amann, R. (2005). Diversity and distribution of methanotrophic archaea at cold seeps. *Appl. Environ. Microbiol.* 71, 467–479. doi: 10.1128/AEM.71.1.467-479.2005
- Kolb, S., and Horn, M. A. (2012). Microbial CH₄ and N₂O consumption in acidic Wetlands. *Front. Microbiol.* 3:78. doi: 10.3389/fmicb.2012.00078
- Krüger, M., Meyerdiecks, A., Glöckner, F. O., Amann, R., Widdel, F., Kube, M., et al. (2003). A conspicuous nickel protein in microbial mats that oxidize methane anaerobically. *Nature* 426, 878–881. doi: 10.1038/nature02207
- Kuenen, J. G. (2020). Anammox and beyond. *Environ. Microbiol.* 22, 525–536. doi: 10.1111/1462-2920.14904
- Kwon, M., Ho, A., and Yoon, S. (2019). Novel approaches and reasons to isolate methanotrophic bacteria with biotechnological potentials: recent achievements and perspectives. *Appl. Microbiol. Biotechnol.* 103, 1–8. doi: 10.1007/s00253-018-9435-1
- Lackner, S., Gilbert, E. M., Vlaeminck, S. E., Joss, A., Horn, H., and van Loosdrecht, M. C. M. (2014). Full-scale partial nitrification/anammox experiences – An application survey. *Water Res.* 55, 292–303. doi: 10.1016/j.watres.2014.02.032
- Lettinga, G. (1995). Anaerobic digestion and wastewater treatment systems. *Antonie Van Leeuwenhoek* 67, 3–28. doi: 10.1007/BF00872193
- Lettinga, G., and Hulshoff Pol, L. W. (1991). USAB-process design for various types of wastewaters. *Water Sci. Technol.* 24, 87–107. doi: 10.2166/wst.1991.0220
- Lettinga, G., van Velsen, A. F. M., Hobma, S. W., de Zeeuw, W., and Klapwijk, A. (1980). Use of the upflow sludge blanket (USB) reactor concept for biological wastewater treatment, especially for an anaerobic treatment. *Biotechnol. Bioeng.* 22, 699–734. doi: 10.1002/bit.260220402
- Leu, A. O., Cai, C., McIlroy, S. J., Southam, G., Orphan, V. J., Yuan, Z., et al. (2020a). Anaerobic methane oxidation coupled to manganese reduction by members of the *Methanoperedenaceae*. *ISME J.* 14, 1030–1041. doi: 10.1038/s41396-020-0590-x
- Leu, A. O., McIlroy, S. J., Ye, J., Parks, D. H., Orphan, V. J., and Tyson, G. W. (2020b). Lateral gene transfer drives metabolic flexibility in the anaerobic methane-oxidizing archaeal family methanoperedenaceae. *MBio* 11, 1–16. doi: 10.1128/mBio.01325-20
- Liu, J., Chakraborty, S., Hosseinzadeh, P., Yu, Y., Tian, S., Petrik, I., et al. (2014). Metalloproteins containing cytochrome, iron-sulfur, or copper redox centers. *Chem. Rev.* 114, 4366–4469. doi: 10.1021/cr400479b
- Liu, T., Hu, S., and Guo, J. (2019a). Enhancing mainstream nitrogen removal by employing nitrate/nitrite-dependent anaerobic methane oxidation processes. *Crit. Rev. Biotechnol.* 39, 732–745. doi: 10.1080/07388551.2019.1598333
- Liu, T., Hu, S., Yuan, Z., and Guo, J. (2019b). High-level nitrogen removal by simultaneous partial nitrification, anammox and nitrite/nitrate-dependent anaerobic methane oxidation. *Water Res.* 166:115057. doi: 10.1016/j.watres.2019.115057
- Liu, Y. (2010). *Taxonomy of Methanogens. In Handbook of Hydrocarbon and Lipid Microbiology*. Berlin: Springer. doi: 10.1007/978-3-540-77587-4_42
- Liu, Z. H., Yin, H., Dang, Z., and Liu, Y. (2014). Dissolved methane: a hurdle for anaerobic treatment of municipal wastewater. *Environ. Sci. Technol.* 48, 889–890. doi: 10.1021/es405553j
- Lloyd, K. (2015). Beyond known methanogens. *Science* 350:6259. doi: 10.1126/science.aad4066
- Lomakina, A., Pogodaeva, T., Kalmychkov, G., Chernitsyna, S., and Zemskaya, T. (2020). Diversity of NC10 bacteria and ANME-2d archaea in sediments of fault zones at Lake Baikal. *Diversity* 12:10010. doi: 10.3390/d12010010
- Lotti, T., Kleerebezem, R., Abelleira-Pereira, J. M., Abbas, B., and van Loosdrecht, M. C. M. (2015). Faster through training: the anammox case. *Water Res.* 81, 261–268. doi: 10.1016/j.watres.2015.06.001
- Lotti, T., Kleerebezem, R., Lubello, C., and van Loosdrecht, M. C. M. (2014). Physiological and kinetic characterization of a suspended cell anammox culture. *Water Res.* 60, 1–14. doi: 10.1016/j.watres.2014.04.017
- Lotti, T., van der Star, W. R. L., Kleerebezem, R., Lubello, C., and van Loosdrecht, M. C. M. (2012). The effect of nitrite inhibition on the anammox process. *Water Res.* 46, 2559–2569. doi: 10.1016/j.watres.2012.02.011
- Lu, P., Liu, T., Ni, B. J., Guo, J., Yuan, Z., and Hu, S. (2019). Growth kinetics of Candidatus *Methanoperedens nitroreducens* enriched in a laboratory reactor. *Sci. Total Environ.* 659, 442–450. doi: 10.1016/j.scitotenv.2018.12.351
- Lu, Y. Z., Fu, L., Ding, J., Ding, Z. W., Li, N., and Zeng, R. J. (2016). Cr(VI) reduction coupled with anaerobic oxidation of methane in a laboratory reactor. *Water Res.* 102, 445–452. doi: 10.1016/j.watres.2016.06.065
- Lu, Y. Z., Fu, L., Li, N., Ding, J., Bai, Y. N., Samaras, P., et al. (2018). The content of trace element iron is a key factor for competition between anaerobic ammonium oxidation and methane-dependent denitrification processes. *Chemosphere* 198, 370–376. doi: 10.1016/j.chemosphere.2018.01.172

- Luesken, F. A., Sánchez, J., van Alen, T. A., Sanabria, J., Op den Camp, H. J. M., Jetten, M. S. M., et al. (2011a). Simultaneous nitrite-dependent anaerobic methane and ammonium oxidation processes. *Appl. Environ. Microbiol.* 77, 6802–6807. doi: 10.1128/AEM.05539-11
- Luesken, F. A., Van Alen, T. A., Van Der Biezen, E., Frijters, C., Toonen, G., Kampman, C., et al. (2011b). Diversity and enrichment of nitrite-dependent anaerobic methane oxidizing bacteria from wastewater sludge. *Appl. Microbiol. Biotechnol.* 92, 845–854. doi: 10.1007/s00253-011-3361-9
- Luesken, F. A., Wu, M. L., Op den Camp, H. J. M., Keltjens, J. T., Stunnenberg, H., Francoijs, K. J., et al. (2012). Effect of oxygen on the anaerobic methanotroph “*Candidatus Methyloirabilis oxyfera*”: Kinetic and transcriptional analysis. *Environ. Microbiol.* 14, 1024–1034. doi: 10.1111/j.1462-2920.2011.02682.x
- Luo, J. H., Chen, H., Hu, S., Cai, C., Yuan, Z., and Guo, J. (2018). Microbial selenate reduction driven by a denitrifying anaerobic methane oxidation biofilm. *Environ. Sci. Technol.* 52, 4006–4012. doi: 10.1021/acs.est.7b05046
- Luo, J. H., Wu, M., Yuan, Z., and Guo, J. (2017). Biological bromate reduction driven by methane in a membrane biofilm reactor. *Environ. Sci. Technol. Lett.* 4, 562–566. doi: 10.1021/acs.estlett.7b00488
- Macleod, F., Kindler, G. S., Wong, H. L., Chen, R., and Burns, B. P. (2019). Asgard archaea: Diversity, function, and evolutionary implications in a range of microbiomes. *AIMS Microbiol.* 5, 48–61. doi: 10.3934/microbiol.2019.1.48
- Madison, L. L., and Huisman, G. W. (1999). Metabolic engineering of Poly(3-Hydroxyalkanoates): from DNA to Plastic. *Microbiol. Mol. Biol. Rev.* 63, 21–53. doi: 10.1128/mmb.63.1.21-53.1999
- Martinez-Cruz, K., Leewis, M. C., Herriott, I. C., Sepulveda-Jauregui, A., Anthony, K. W., Thalasso, F., et al. (2017). Anaerobic oxidation of methane by aerobic methanotrophs in sub-Arctic lake sediments. *Sci. Total Environ.* 607–608, 23–31. doi: 10.1016/j.scitotenv.2017.06.187
- McCarty, P. L. (1982). “One hundred years of anaerobic treatment. Anaerobic Digestion 1981,” in *Proceedings Symposium, Travemunde*, College Station, TX.
- McInerney, M. J., and Bryant, M. P. (1981). “Basic principles of bioconversions in anaerobic digestion and methanogenesis,” in *Biomass Conversion Processes for Energy and Fuels*, eds S. S. Sofer and O. R. Zaborsky (New York, NY: Plenum), 277–296. doi: 10.1007/978-1-4757-0301-6_15
- Mergaert, J., Anderson, C., Wouters, A., and Swings, J. (1994). Microbial degradation of poly(3-hydroxybutyrate) and poly(3-hydroxybutyrate-co-3-hydroxyvalerate) in compost. *J. Environ. Poly. Degradat.* 2, 177–183. doi: 10.1007/BF02067443
- Meulepas, R. J. W., Jagersma, C. G., Khadem, A. F., Buisman, C. J. N., Stams, A. J. M., and Lens, P. N. L. (2009). Effect of environmental conditions on sulfate reduction with methane as electron donor by an Eckernförde Bay enrichment. *Environ. Sci. Technol.* 43, 6553–6559. doi: 10.1021/es900633c
- Mills, H. J., Hodges, C., Wilson, K., MacDonald, I. R., and Sobczyk, P. A. (2003). Microbial diversity in sediments associated with surface-breaching gas hydrate mounds in the Gulf of Mexico. *FEMS Microbiol. Ecol.* 46, 39–52. doi: 10.1016/S0168-6496(03)00191-0
- Mills, H. J., Martinez, R. J., Story, S., and Sobczyk, P. A. (2005). Characterization of microbial community structure in Gulf of Mexico gas hydrates: comparative analysis of DNA- and RNA-derived clone libraries. *Appl. Environ. Microbiol.* 71, 3235–3247. doi: 10.1128/AEM.71.6.3235-3247.2005
- Milucka, J., Kirf, M., Lu, L., Krupke, A., Lam, P., Littmann, S., et al. (2015). Methane oxidation coupled to oxygenic photosynthesis in anoxic waters. *ISME J.* 9, 1991–2002. doi: 10.1038/ismej.2015.12
- Mulder, A., van de Graaf, A. A., Robertson, L. A., and Kuenen, J. G. (1995). Anaerobic ammonium oxidation discovered in a denitrifying fluidized bed reactor. *FEMS Microbiol. Ecol.* 16, 177–183. doi: 10.1016/0168-6496(94)00081-7
- Myhre, G., Shindell, D., Bréon, F., Collins, W., Fuglestedt, J., Huang, J., et al. (2013). *Anthropogenic and Natural Radiative Forcing. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, 659–740.
- Nauhaus, K., Boetius, A., Krüger, M., and Widdel, F. (2002). In vitro demonstration of anaerobic oxidation of methane coupled to sulphate reduction in sediment from a marine gas hydrate area. *Environ. Microbiol.* 4, 296–305. doi: 10.1046/j.1462-2920.2002.00299.x
- Nguyen, T. K. L., Ngo, H. H., Guo, W., Chang, S. W., Nguyen, D. D., Nghiem, L. D., et al. (2019). Insight into greenhouse gases emissions from the two popular treatment technologies in municipal wastewater treatment processes. *Sci. Total Environ.* 671, 1302–1313. doi: 10.1016/j.scitotenv.2019.03.386
- Nichol, T., Murrell, J. C., and Smith, T. J. (2019). in *Biochemistry and Molecular Biology of Methane Monooxygenase BT – Aerobic Utilization of Hydrocarbons, Oils, and Lipids*, ed. F. Rojo (Manhattan, NY: Springer International Publishing), 99–115. doi: 10.1007/978-3-319-50418-6_5
- Niemann, H., Lösekann, T., De Beer, D., Elvert, M., Nadalig, T., Knittel, K., et al. (2006). Novel microbial communities of the Haakon Mosby mud volcano and their role as a methane sink. *Nature* 443, 854–858. doi: 10.1038/nature05227
- Nierychlo, M., Andersen, K. S., Xu, Y., Green, N., Jiang, C., Albertsen, M., et al. (2020). MiDAS 3: an ecosystem-specific reference database, taxonomy and knowledge platform for activated sludge and anaerobic digesters reveals species-level microbiome composition of activated sludge. *Water Res.* 182:115955. doi: 10.1016/j.watres.2020.115955
- Nordi, K. Á., Thamdrup, B., and Schubert, C. J. (2013). Anaerobic oxidation of methane in an iron-rich Danish freshwater lake sediment. *Limnol. Oceanogr.* 58:546. doi: 10.4319/lo.2013.58.2.0546
- Noyola, A., Morgan-Sagastume, J. M., and López-Hernández, J. E. (2006). Treatment of biogas produced in anaerobic reactors for domestic wastewater: odor control and energy/resource recovery. *Rev. Environ. Sci. Biotechnol.* 5, 93–114. doi: 10.1007/s11157-005-2754-6
- Nzila, A. (2017). Mini review: Update on bioaugmentation in anaerobic processes for biogas production. *Anaerobe* 46, 3–12. doi: 10.1016/j.anaerobe.2016.11.007
- Ogiso, T., Ueno, C., Dianou, D., Van Huy, T., Katayama, A., Kimura, M., et al. (2012). *Methylomonas koyamae* sp. nov., a type I methane-oxidizing bacterium from floodwater of a rice paddy field. *Int. J. Syst. Evol. Microbiol.* 62, 1832–1837. doi: 10.1099/ijs.0.035261-0
- Okoro, O. V., and Sun, Z. (2019). Desulphurisation of biogas: a systematic qualitative and economic-based quantitative review of alternative strategies. *ChemEngineering* 3:76. doi: 10.3390/chemengineering3030076
- Orphan, V. J., House, C. H., Hinrichs, K. U., McKeegan, K. D., and DeLong, E. F. (2001). Methane-consuming archaea revealed by directly coupled isotopic and phylogenetic analysis. *Science* 293, 484–487. doi: 10.1126/science.1061338
- Orphan, V. J., House, C. H., Hinrichs, K. U., McKeegan, K. D., and DeLong, E. F. (2002). Multiple archaeal groups mediate methane oxidation in anoxic cold seep sediments. *Proc. Natl. Acad. Sci. U.S.A.* 99, 7663–7668. doi: 10.1073/pnas.072210299
- Oswald, K., Milucka, J., Brand, A., Hach, P., Littmann, S., Wehrli, B., et al. (2016). Aerobic gammaproteobacterial methanotrophs mitigate methane emissions from oxic and anoxic lake waters. *Limnol. Oceanogr.* 61, S101–S118. doi: 10.1002/lno.10312
- Øverland, M., Tauson, A. H., Shearer, K., and Skrede, A. (2010). Evaluation of methane-utilising bacteria products as feed ingredients for monogastric animals. *Arch. Anim. Nutr.* 64, 171–189. doi: 10.1080/17450391003691534
- Padilla, C. C., Bristow, L. A., Sarode, N., Garcia-Robledo, E., Gómez Ramírez, E., Benson, C. R., et al. (2016). NC10 bacteria in marine oxygen minimum zones. *ISME J.* 10, 2067–2071. doi: 10.1038/ismej.2015.262
- Park, S. J., Ahn, W. S., Green, P. R., and Lee, S. Y. (2001). Biosynthesis of poly(3-hydroxybutyrate-co-3-hydroxyvalerate-co-3-hydroxyhexanoate) by metabolically engineered *Escherichia coli* strains. *Biotechnol. Bioeng.* 74, 81–86. doi: 10.1002/bit.1097
- Pastor, J. M., Salvador, M., Argandoña, M., Bernal, V., Reina-Bueno, M., Csonka, L. N., et al. (2010). Ectoines in cell stress protection: uses and biotechnological production. *Biotechnol. Adv.* 28, 782–801. doi: 10.1016/j.biotechadv.2010.06.005
- Peoples, O. P., and Sinskey, A. J. (1989). Poly-β-hydroxybutyrate biosynthesis in *Alcaligenes eutrophus* H16. Characterization of the genes encoding β-ketothiolase and acetoacetyl-CoA reductase. *J. Biol. Chem.* 264, 15293–15297. doi: 10.1016/S0021-9258(19)84824-X
- Petersen, L. A. H., Villadsen, J., Jørgensen, S. B., and Gernaey, K. V. (2017). Mixing and mass transfer in a pilot scale U-loop bioreactor. *Biotechnol. Bioeng.* 114, 344–354. doi: 10.1002/bit.26084
- Pieja, A. J., Morse, M. C., and Cal, A. J. (2017). Methane to bioproducts: the future of the bioeconomy? *Curr. Opin. Chem. Biol.* 41, 123–131. doi: 10.1016/j.cbpa.2017.10.024

- Pieja, A. J., Rostkowski, K. H., and Criddle, C. S. (2011a). Distribution and Selection of Poly-3-hydroxybutyrate production capacity in methanotrophic *Proteobacteria*. *Microb. Ecol.* 62, 564–573. doi: 10.1007/s00248-011-9873-0
- Pieja, A. J., Sundstrom, E. R., and Criddle, C. S. (2011b). Poly-3-hydroxybutyrate metabolism in the type II methanotroph *Methylocystis parvus* OB8P. *Appl. Environ. Microbiol.* 77, 6012–6019. doi: 10.1128/AEM.00509-11
- Pol, A., Barends, T. R. M., Dietl, A., Khadem, A. F., Eygensteyn, J., Jetten, M. S. M., et al. (2014). Rare earth metals are essential for methanotrophic life in volcanic mudpots. *Environ. Microbiol.* 16, 255–264. doi: 10.1111/1462-2920.12249
- Pol, A., Heijmans, K., Harhangi, H. R., Tedesco, D., Jetten, M. S. M., and Op Den Camp, H. J. M. (2007). Methanotrophy below pH 1 by a new Verrucomicrobia species. *Nature* 450, 874–878. doi: 10.1038/nature06222
- Pronk, M., de Kreuk, M. K., de Bruin, B., Kamminga, P., Kleerebezem, R., and van Loosdrecht, M. C. M. (2015). Full scale performance of the aerobic granular sludge process for sewage treatment. *Water Res.* 84, 207–217. doi: 10.1016/j.watres.2015.07.011
- Raghoebarsing, A. A., Pol, A., Van De Pas-Schoonen, K. T., Smolders, A. J. P., Ettwig, K. F., Rijpstra, W. I. C., et al. (2006). A microbial consortium couples anaerobic methane oxidation to denitrification. *Nature* 440, 918–921. doi: 10.1038/nature04617
- Raghoebarsing, A. A., Smolders, A. J. P., Schmid, M. C., Rijpstra, W. I. C., Wolters-Arts, M., Derksen, J., et al. (2005). Methanotrophic symbionts provide carbon for photosynthesis in peat bogs. *Nature* 436, 1156–1156. doi: 10.1038/nature03802
- Rahalkar, M. C., and Bahulikar, R. A. (2018). Hemerythrins are widespread and conserved for methanotrophic guilds. *Gene Rep.* 11, 250–254. doi: 10.1016/j.genrep.2018.04.008
- Rasouli, Z., Valverde-Pérez, B., D'Este, M., De Francisci, D., and Angelidaki, I. (2018). Nutrient recovery from industrial wastewater as single cell protein by a co-culture of green microalgae and methanotrophs. *Biochem. Eng. J.* 134, 129–135. doi: 10.1016/j.bej.2018.03.010
- Reeburgh, W. S. (1976). Methane consumption in Cariaco Trench waters and sediments. *Earth Planet. Sci. Lett.* 28, 337–344. doi: 10.1016/0012-821X(76)90195-3
- Reeburgh, W. S., and Heggie, D. T. (1977). Microbial methane consumption reactions and their effect on methane distributions in freshwater and marine environments. *Limnol. Oceanogr.* 22, 1–9. doi: 10.4319/lo.1977.22.1.0001
- Reim, A., Lülke, C., Krause, S., Pratscher, J., and Frenzel, P. (2012). One millimetre makes the difference: High-resolution analysis of methane-oxidizing bacteria and their specific activity at the oxic-anoxic interface in a flooded paddy soil. *ISME J.* 6, 2118–2139. doi: 10.1038/ismej.2012.57
- Ribera-Guardia, A., Bosch, L., Corominas, L., and Pijuan, M. (2019). Nitrous oxide and methane emissions from a plug-flow full-scale bioreactor and assessment of its carbon footprint. *J. Clean. Product.* 212, 162–172. doi: 10.1016/j.jclepro.2018.11.286
- Rissanen, A. J., Saarela, T., Jäntti, H., Buck, M., Peura, S., Aalto, S. L., et al. (2020). Vertical stratification patterns of methanotrophs and their genetic controllers in water columns of oxygen-stratified boreal lakes. *FEMS Microbiol. Ecol.* 80, 257–276. doi: 10.1093/femsec/fiaa252
- Rissanen, A. J., Saareheimo, J., Tiirola, M., Peura, S., Aalto, S. L., Karvinen, A., et al. (2018). Gammaproteobacterial methanotrophs dominate methanotrophy in aerobic and anaerobic layers of boreal lake waters. *Aquat. Microb. Ecol.* 97:1874. doi: 10.3354/ame01874
- Ritala, A., Häkkinen, S. T., Toivari, M., and Wiebe, M. G. (2017). Single cell protein-state-of-the-art, industrial landscape and patents 2001-2016. *Front. Microbiol.* 8:2009. doi: 10.3389/fmicb.2017.02009
- Rodríguez-García, G., Hospido, A., Bagley, D. M., Moreira, M. T., and Feijoo, G. (2012). A methodology to estimate greenhouse gases emissions in life cycle inventories of wastewater treatment plants. *Environ. Impact Assess. Rev.* 37, 37–46. doi: 10.1016/j.eiar.2012.06.010
- Ross, M. O., and Rosenzweig, A. C. (2017). A tale of two methane monooxygenases. *J. Biol. Inorgan. Chem.* 22, 307–319. doi: 10.1007/s00775-016-1419-y
- Salar-García, M. J., Bernal, V., Pastor, J. M., Salvador, M., Argandoña, M., Nieto, J. J., et al. (2017). Understanding the interplay of carbon and nitrogen supply for ectoines production and metabolic overflow in high density cultures of *Chromohalobacter salexigens*. *Microb. Cell Fact.* 16:23. doi: 10.1186/s12934-017-0643-7
- Sánchez, A., Rodríguez-Hernández, L., Buntner, D., Esteban-García, A. L., Tejero, I., and Garrido, J. M. (2016). Denitrification coupled with methane oxidation in a membrane bioreactor after methanogenic pre-treatment of wastewater. *J. Chem. Technol. Biotechnol.* 91, 2950–2958. doi: 10.1002/jctb.4913
- Sang, G. L., Jae, H. G., Hee, G. K., Oh, J. I., Young, M. K., and Si, W. K. (2004). Optimization of methanol biosynthesis from methane using *Methylosinus trichosporium* OB3b. *Biotechnol. Lett.* 26, 947–950. doi: 10.1023/B:bile.0000025908.19252.63
- Saunio, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., et al. (2020). The global methane budget 2000-2017. *Earth Syst. Sci. Data* 12, 1561–1623. doi: 10.5194/essd-12-1561-2020
- Schaum, C., Fundneider, T., and Cornel, P. (2016). Analysis of methane emissions from digested sludge. *Water Sci. Technol.* 73, 1599–1607. doi: 10.2166/wst.2015.644
- Scheller, S., Yu, H., Chadwick, G. L., McGlynn, S. E., and Orphan, V. J. (2016). Artificial electron acceptors decouple archaeal methane oxidation from sulfate reduction. *Science* 351, 703–707. doi: 10.1126/science.aad7154
- Schubert, C. J., Vazquez, F., Lösekann-Behrens, T., Knittel, K., Tonolla, M., and Boetius, A. (2011). Evidence for anaerobic oxidation of methane in sediments of a freshwater system (Lago di Cadagno). *FEMS Microbiol. Ecol.* 76, 26–38. doi: 10.1111/j.1574-6941.2010.01036.x
- Semrau, J. D., DiSpirito, A. A., and Yoon, S. (2010). Methanotrophs and copper. *FEMS Microbiol. Rev.* 34, 496–531. doi: 10.1111/j.1574-6976.2010.00212.x
- Semrau, J. D., DiSpirito, A. A., Gu, W., and Yoon, S. (2018). Metals and methanotrophy. *Appl. Environ. Microbiol.* 84, e2289–e2217. doi: 10.1128/AEM.02289-17
- Shaw, D. R., Ali, M., Katuri, K. P., Gralnick, J. A., Reimann, J., Mesman, R., et al. (2020). Extracellular electron transfer-dependent anaerobic oxidation of ammonium by anammox bacteria. *Nat. Commun.* 11:2058. doi: 10.1038/s41467-020-16016-y
- Sheets, J. P., Ge, X., Li, Y. F., Yu, Z., and Li, Y. (2016). Biological conversion of biogas to methane using methanotrophs isolated from solid-state anaerobic digestate. *Bioresour. Technol.* 201, 50–57. doi: 10.1016/j.biortech.2015.11.035
- Shen, L., Dong, O. L., Zhu, Y., and Trimmer, M. (2019). Active pathways of anaerobic methane oxidation across contrasting riverbeds. *ISME J.* 13, 752–766. doi: 10.1038/s41396-018-0302-y
- Shi, L. D., Lv, P. L., Wang, M., Lai, C. Y., and Zhao, H. P. (2020). A mixed consortium of methanotrophic archaea and bacteria boosts methane-dependent selenate reduction. *Sci. Total Environ.* 732:139310. doi: 10.1016/j.scitotenv.2020.139310
- Shi, Y., Hu, S., Lou, J., Lu, P., Keller, J., and Yuan, Z. (2013). Nitrogen removal from wastewater by coupling anammox and methane-dependent denitrification in a membrane biofilm reactor. *Environ. Sci. Technol.* 47, 11577–11583. doi: 10.1021/es402775z
- Shiau, Y. J., Cai, Y., Lin, Y., Te Jia, Z., and Chiu, C. Y. (2018). Community structure of active aerobic methanotrophs in red mangrove (*Kandelia obovata*) soils under different frequency of tides. *Microb. Ecol.* 75, 761–770. doi: 10.1007/s00248-017-1080-1
- Sieber, J. R., McInerney, M. J., Plugge, C. M., Schink, B., and Gunsalus, R. P. (2010). *Methanogenesis: Syntrophic Metabolism. In Handbook of Hydrocarbon and Lipid Microbiology*. Berlin: Springer. doi: 10.1007/978-3-540-77587-4_22
- Silva-Teira, A., Sánchez, A., Buntner, D., Rodríguez-Hernández, L., and Garrido, J. M. (2017). Removal of dissolved methane and nitrogen from anaerobically treated effluents at low temperature by MBR post-treatment. *Chem. Eng. J.* 326, 970–979. doi: 10.1016/j.cej.2017.06.047
- Sivan, O., Adler, M., Pearson, A., Gelman, F., Bar-Or, I., John, S. G., et al. (2011). Geochemical evidence for iron-mediated anaerobic oxidation of methane. *Limnol. Oceanogr.* 56, 1536–1544. doi: 10.4319/lo.2011.56.4.1536
- Skennerton, C. T., Ward, L. M., Michel, A., Metcalfe, K., Valiente, C., Mullin, S., et al. (2015). Genomic reconstruction of an uncultured hydrothermal vent Gammaproteobacterial Methanotroph (Family *Methylothermaceae*) indicates multiple adaptations to oxygen limitation. *Front. Microbiol.* 6:1425. doi: 10.3389/fmicb.2015.01425
- Skrede, A., Berge, G. M., Storebakken, T., Herstad, O., Aarstad, K. G., and Sundstøl, F. (1998). Digestibility of bacterial protein grown on natural gas in mink,

- pigs, chicken and Atlantic salmon. *Anim. Feed Sci. Technol.* 76, 103–116. doi: 10.1016/S0377-8401(98)00208-9
- Skrede, A., Faaland Schøyen, H., Svihus, B., and Storebakken, T. (2003). The effect of bacterial protein grown on natural gas on growth performance and sensory quality of broiler chickens. *Canad. J. Anim. Sci.* 83:4141. doi: 10.4141/A02-047
- Smith, G. J., and Wrighton, K. C. (2019). Metagenomic approaches unearth methanotroph phylogenetic and metabolic diversity. *Curr. Issues Mol. Biol.* 33, 57–84. doi: 10.21775/cimb.033.057
- Smith, G. J., Angle, J. C., Solden, L. M., Borton, M. A., Morin, T. H., Daly, R. A., et al. (2018). Members of the genus *Methylobacter* are inferred to account for the majority of aerobic methane oxidation in oxic soils from a freshwater wetland. *MBio* 9, e815–e818. doi: 10.1128/mBio.00815-18
- Sollo, F. W., Mueller, H. F., and Larson, T. E. (1976). Denitrification of wastewater effluents with methane. *J. Water Pollut. Control Feder.* 48, 1840–1842.
- Souza, C. L., Chernicharo, C. A. L., and Aquino, S. F. (2011). Quantification of dissolved methane in UASB reactors treating domestic wastewater under different operating conditions. *Water Sci. Technol.* 64, 2259–2264. doi: 10.2166/wst.2011.695
- Speth, D. R., In'T Zandt, M. H., Guerrero-Cruz, S., Dutilh, B. E., and Jetten, M. S. M. (2016). Genome-based microbial ecology of anammox granules in a full-scale wastewater treatment system. *Nat. Commun.* 7:11172. doi: 10.1038/ncomms11172
- Stern, D. I., and Kaufmann, R. K. (2014). Anthropogenic and natural causes of climate change. *Clim. Change* 122, 257–269. doi: 10.1007/s10584-013-1007-x
- Strong, P. J., Kalyuzhnaya, M., Silverman, J., and Clarke, W. P. (2016). A methanotroph-based biorefinery: potential scenarios for generating multiple products from a single fermentation. *Bioresour. Technol.* 215, 314–323. doi: 10.1016/j.biortech.2016.04.099
- Strong, P. J., Xie, S., and Clarke, W. P. (2015). Methane as a resource: can the methanotrophs add value? *Environ. Sci. Technol.* 49, 4001–4018. doi: 10.1021/es504242n
- Strous, M., Fuerst, J. A., Kramer, E. H. M., Logemann, S., Muyzer, G., Van De Pas-Schoonen, K. T., et al. (1999a). Missing lithotroph identified as new planctomycete. *Nature* 400, 446–449. doi: 10.1038/22749
- Strous, M., Heijnen, J. J., Kuenen, J. G., and Jetten, M. S. M. (1998). The sequencing batch reactor as a powerful tool for the study of slowly growing anaerobic ammonium-oxidizing microorganisms. *Appl. Microbiol. Biotechnol.* 50, 589–596. doi: 10.1007/s002530051340
- Strous, M., Kuenen, J. G., and Jetten, M. S. M. (1999b). Key physiology of anaerobic ammonium oxidation. *Appl. Environ. Microbiol.* 65, 3248–3250. doi: 10.1128/aem.65.7.3248-3250.1999
- Strous, M., Van Gerven, E., Kuenen, J. G., and Jetten, M. (1997). Effects of aerobic and microaerobic conditions on anaerobic ammonium-oxidizing (anammox) sludge. *Appl. Environ. Microbiol.* 63, 2446–2448. doi: 10.1128/aem.63.6.2446-2448.1997
- Stultiens, K., Cruz, S. G., van Kessel, M. A. H. J., Jetten, M. S. M., Kartal, B., and Op den Camp, H. J. M. (2019). Interactions between anaerobic ammonium- and methane-oxidizing microorganisms in a laboratory-scale sequencing batch reactor. *Appl. Microbiol. Biotechnol.* 103, 6783–6795. doi: 10.1007/s00253-019-09976-9
- Su, G., Zopfi, J., Yao, H., Steinle, L., Niemann, H., and Lehmann, M. F. (2020). Manganese/iron-supported sulfate-dependent anaerobic oxidation of methane by archaea in lake sediments. *Limnol. Oceanogr.* 65, 863–875. doi: 10.1002/lno.11354
- Takeguchi, M., Furuto, T., Sugimori, D., and Okura, I. (1997). Optimization of methanol biosynthesis by *Methylosinus trichosporium* OB3b: an approach to improve methanol accumulation. *Appl. Biochem. Biotechnol.* 68, 143–152. doi: 10.1007/BF02785987
- Tanaka, K., Yokoe, S., Igarashi, K., Takashino, M., Ishikawa, M., Hori, K., et al. (2018). Extracellular electron transfer via outer membrane cytochromes in a methanotrophic bacterium *Methylococcus capsulatus* (Bath). *Front. Microbiol.* 9:2905. doi: 10.3389/fmicb.2018.02905
- Tays, C., Guarnieri, M. T., Sauvageau, D., and Stein, L. Y. (2018). Combined effects of carbon and nitrogen source to optimize growth of proteobacterial methanotrophs. *Front. Microbiol.* 9:2239. doi: 10.3389/fmicb.2018.02239
- Thakur, I. S., and Medhi, K. (2019). Nitrification and denitrification processes for mitigation of nitrous oxide from waste water treatment plants for biovalorization: challenges and opportunities. *Bioresour. Technol.* 282, 502–513. doi: 10.1016/j.biortech.2019.03.069
- Thamdrup, B., Steinsdóttir, H. G. R., Bertagnolli, A. D., Padilla, C. C., Patin, N. V., Garcia-Robledo, E., et al. (2019). Anaerobic methane oxidation is an important sink for methane in the ocean's largest oxygen minimum zone. *Limnol. Oceanogr.* 64, 2569–2585. doi: 10.1002/lno.11235
- Thauer, R. K. (1998). Biochemistry of methanogenesis: a tribute to Marjory Stephenson:1998 marjory stephenson prize lecture. *Microbiology* 144, 2377–2406. doi: 10.1099/00221287-144-9-2377
- Timmers, P. H. A., Gieteling, J., Widjaja-Greefkes, H. C. A., Plugge, C. M., Stams, A. J. M., Lens, P. N. L., et al. (2015). Growth of anaerobic methane-oxidizing archaea and sulfate-reducing bacteria in a high-pressure membrane capsule bioreactor. *Appl. Environ. Microbiol.* 81, 1286–1296. doi: 10.1128/AEM.03255-14
- Trego, A. C., O'Sullivan, S., Quince, C., Mills, S., Ijaz, U. Z., and Collins, G. (2020). Size Shapes the active microbiome of the methanogenic granules, corroborating a biofilm life cycle. *MSystems* 5:e00323. doi: 10.1128/msystems.00323-20
- Trotsenko, Y. A., and Khmelenina, V. N. (2002). Biology of extremophilic and extremotolerant methanotrophs. *Arch. Microbiol.* 177, 123–131. doi: 10.1007/s00203-001-0368-0
- Trotsenko, Y. A., and Murrell, J. C. B. T. (2008). Metabolic aspects of aerobic obligate methanotrophy. *Adv. Appl. Microbiol.* 63, 183–229. doi: 10.1016/S0065-2164(07)00005-6
- Umeozor, E. C., Jordaan, S. M., and Gates, I. D. (2018). On methane emissions from shale gas development. *Energy* 152, 594–600. doi: 10.1016/j.energy.2018.03.151
- Vakmsaa, A., Guerrero-Cruz, S., van Alen, T. A., Cremers, G., Ettwig, K. F., Lüke, C., et al. (2017). Enrichment of anaerobic nitrate-dependent methanotrophic 'Candidatus *Methanoperedens nitroreducens*' archaea from an Italian paddy field soil. *Appl. Microbiol. Biotechnol.* 101, 7075–7084. doi: 10.1007/s00253-017-8416-0
- Vakmsaa, A., Lüke, C., van Alen, T., Valè, G., Lupotto, E., Jetten, M. S. M., et al. (2016). Distribution and activity of the anaerobic methanotrophic community in a nitrogen-fertilized Italian paddy soil. *FEMS Microbiol. Ecol.* 92:181. doi: 10.1093/femsec/fw181
- Valenzuela, E. I., Padilla-Loma, C., Gómez-Hernández, N., López-Lozano, N. E., Casas-Flores, S., and Cervantes, F. J. (2020). Humic substances mediate anaerobic methane oxidation linked to nitrous oxide reduction in wetland sediments. *Front. Microbiol.* 11:587. doi: 10.3389/fmicb.2020.00587
- Van De Graaf, A. A., De Bruijn, P., Robertson, L. A., Jetten, M. S. M., and Kuenen, J. G. (1996). Autotrophic growth of anaerobic ammonium-oxidizing microorganisms in a fluidized bed reactor. *Microbiology* 142:2187. doi: 10.1099/13500872-142-8-2187
- Van De Graaf, A. A., De Bruijn, P., Robertson, L. A., Jetten, M. S. M., and Kuenen, J. G. (1997). Metabolic pathway of anaerobic ammonium oxidation on the basis of 15N studies in a fluidized bed reactor. *Microbiology* 143, 2415–2421. doi: 10.1099/00221287-143-7-2415
- van der Ha, D., Bundervoet, B., Verstraete, W., and Boon, N. (2011). A sustainable, carbon neutral methane oxidation by a partnership of methane oxidizing communities and microalgae. *Water Res.* 45, 2845–2854. doi: 10.1016/j.watres.2011.03.005
- van der Star, W. R. L., Abma, W. R., Blommers, D., Mulder, J. W., Tokutomi, T., Strous, M., et al. (2007). Startup of reactors for anoxic ammonium oxidation: experiences from the first full-scale anammox reactor in Rotterdam. *Water Res.* 41, 4149–4163. doi: 10.1016/j.watres.2007.03.044
- van Grinsven, S., Oswald, K., Wehrli, B., Jegge, C., Zopfi, J., Lehmann, M. F., et al. (2021). Methane oxidation in the waters of a humics-rich boreal lake stimulated by photosynthesis, nitrite, Fe(III) and humics. *Biogeoosci. Discuss* [Preprint]. doi: 10.5194/bg-2021-3
- van Grinsven, S., Sinnighe Damsté, J. S., Abdala Asbun, A., Engelmann, J. C., Harrison, J., and Villanueva, L. (2020). Methane oxidation in anoxic lake water stimulated by nitrate and sulfate addition. *Environ. Microbiol.* 22, 766–782. doi: 10.1111/1462-2920.14886
- van Teeseling, M. C. F., Pol, A., Harhangi, H. R., van der Zwart, S., Jetten, M. S. M., Op den Camp, H. J. M., et al. (2014). Expanding the verrucomicrobial methanotrophic world: description of three novel species

- of *Methylacidimicrobium* gen. nov. *Appl. Environ. Microbiol.* 80, 6782–6791. doi: 10.1128/AEM.01838-14
- Vanwonterghem, L., Evans, P., Parks, D. H., Jensen, P. D., Woodcroft, B. J., Hugenholtz, P., et al. (2016). Methylophilic methanogenesis discovered in the archaeal phylum Verstraearchaeota. *Nat. Microbiol.* 1:16170. doi: 10.1038/nmicrobiol.2016.170
- Veraart, A. J., Garbeva, P., Van Beersum, F., Ho, A., Hordijk, C. A., Meima-Franke, M., et al. (2018). Living apart together - Bacterial volatiles influence methanotrophic growth and activity. *ISME J.* 12, 1163–1166. doi: 10.1038/s41396-018-0055-7
- Versantvoort, W., Guerrero-Cruz, S., Speth, D. R., Frank, J., Gambelli, L., Cremers, G., et al. (2018). Comparative genomics of Candidatus *Methylomirabilis* species and description of “Ca. *Methylomirabilis lanthanidiphila*”. *Front. Microbiol.* 9:1672. doi: 10.3389/fmicb.2018.01672
- Wang, S., An, Z., and Wang, Z.-W. (2020). Bioconversion of methane to chemicals and fuels by methane-oxidizing bacteria. *Adv. Bioener.* 5, 169–247. doi: 10.1016/b5.aibe.2020.04.005
- Wang, Y., Wang, D., Yang, Q., Zeng, G., and Li, X. (2017). Wastewater opportunities for denitrifying anaerobic methane oxidation. *Trends Biotechnol.* 35, 799–802. doi: 10.1016/j.tibtech.2017.02.010
- Wankel, S. D., Adams, M. M., Johnston, D. T., Hansel, C. M., Joye, S. B., and Girguis, P. R. (2012). Anaerobic methane oxidation in metalliferous hydrothermal sediments: influence on carbon flux and decoupling from sulfate reduction. *Environ. Microbiol.* 14, 2726–2740. doi: 10.1111/j.1462-2920.2012.02825.x
- Weber, H. S., Habicht, K. S., and Thamdrup, B. (2017). Anaerobic methanotrophic archaea of the ANME-2d cluster are active in a low-sulfate, iron-rich freshwater sediment. *Front. Microbiol.* 8:619. doi: 10.3389/fmicb.2017.00619
- Weissenbacher, N., Takacs, I., Murthy, S., Fuerhacker, M., and Wett, B. (2010). Gaseous Nitrogen and carbon emissions from a full-scale deammonification plant. *Water Environ. Res.* 82, 169–175. doi: 10.2175/106143009x447867
- Wendlandt, K. D., Geyer, W., Mirschel, G., and Al-Haj Hemidi, F. (2005). Possibilities for controlling a PHB accumulation process using various analytical methods. *J. Biotechnol.* 117, 119–129. doi: 10.1016/j.jbiotec.2005.01.007
- Whittenbury, R., Phillips, K. C., and Wilkinson, J. F. (1970). Enrichment, isolation and some properties of methane-utilizing bacteria. *J. Gen. Microbiol.* 61, 205–218. doi: 10.1099/00221287-61-2-205
- Wiesenburg, D. A., and Guinasso, N. L. (1979). Equilibrium solubilities of methane, carbon monoxide, and hydrogen in water and sea water. *J. Chem. Eng. Data* 24, 356–360. doi: 10.1021/jc60083a006
- Winkler, M. K. H., Ettwig, K. F., Vannacke, T. P. W., Stultjens, K., Bogdan, A., Kartal, B., et al. (2015). Modelling simultaneous anaerobic methane and ammonium removal in a granular sludge reactor. *Water Res.* 15, 323–331. doi: 10.1016/j.watres.2015.01.039
- Woese, C. R., and Fox, G. E. (1977). Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proc. Natl. Acad. Sci. U.S.A.* 74, 5088–5090. doi: 10.1073/pnas.74.11.5088
- Wolf, J., Asrar, G. R., and West, T. O. (2017). Revised methane emissions factors and spatially distributed annual carbon fluxes for global livestock. *Carb. Bal. Manage* 12:13021. doi: 10.1186/s13021-017-0084-y
- Wu, L., Ning, D., Zhang, B., Li, Y., Zhang, P., Shan, X., et al. (2019). Global diversity and biogeography of bacterial communities in wastewater treatment plants. *Nat. Microbiol.* 4, 1193–1195. doi: 10.1038/s41564-019-0426-5
- Wu, M. L., Ettwig, K. F., Jetten, M. S. M., Strous, M., Keltjens, J. T., and Van Niftrik, L. (2011). A new intra-aerobic metabolism in the nitrite-dependent anaerobic methane-oxidizing bacterium Candidatus “*Methylomirabilis oxyfera*”. *Biochem. Soc. Transact.* 39, 243–248. doi: 10.1042/BST0390243
- Wu, M. L., Wessels, H. J. C. T., Pol, A., Op den Camp, H. J. M., Jetten, M. S. M., van Niftrik, L., et al. (2015). XoxF-type methanol dehydrogenase from the anaerobic methanotroph “Candidatus *Methylomirabilis oxyfera*”. *Appl. Environ. Microbiol.* 81, 1442–1451. doi: 10.1128/AEM.03292-14
- Xie, G. J., Cai, C., Hu, S., and Yuan, Z. (2017). Complete nitrogen removal from synthetic anaerobic sludge digestion liquor through integrating anammox and denitrifying anaerobic methane oxidation in a membrane biofilm reactor. *Environ. Sci. Technol.* 51, 819–827. doi: 10.1021/acs.est.6b04500
- Yu, L., He, D., Zhang, E., He, Q., Li, J., Jason Ren, Z., et al. (2021). Electricity from anaerobic methane oxidation by a single methanogenic archaeon *Methanosarcina barkeri*. *Chem. Eng. J.* 405:126691. doi: 10.1016/j.cej.2020.126691
- Yu, Z., Pesesky, M., Zhang, L., Huang, J., Winkler, M., and Chistoserdova, L. (2020). A Complex interplay between nitric oxide, quorum sensing, and the unique secondary metabolite tundrenone constitutes the hypoxia response in methylobacter. *MSystems* 5, E770–E719. doi: 10.1128/mSystems.00770-19
- Zehnder, A. J. B., and Brock, T. D. (1980). Anaerobic methane oxidation: occurrence and ecology. *Appl. Environ. Microbiol.* 39, 194–204. doi: 10.1128/aem.39.1.194-204.1980
- Zhang, W., Ge, X., Li, Y. F., Yu, Z., and Li, Y. (2016). Isolation of a methanotroph from a hydrogen sulfide-rich anaerobic digester for methanol production from biogas. *Process Biochem.* 51, 838–844. doi: 10.1016/j.procbio.2016.04.003
- Zhang, X., Rabiee, H., Frank, J., Cai, C., Stark, T., Virdis, B., et al. (2020). Enhancing methane oxidation in a bioelectrochemical membrane reactor using a soluble electron mediator. *Biotechnol. Biof.* 13:173. doi: 10.1186/s13068-020-01808-7
- Zheng, Y., Wang, H., Liu, Y., Zhu, B., Li, J., Yang, Y., et al. (2020). Methane-dependent mineral reduction by aerobic methanotrophs under hypoxia. *Environ. Sci. Technol. Lett.* 7, 606–612. doi: 10.1021/acs.estlett.0c00436
- Zhou, Z., Pan, J., Wang, F., Gu, J. D., and Li, M. (2018). Bathyarchaeota: globally distributed metabolic generalists in anoxic environments. *FEMS Microbiol. Rev.* 42, 639–655. doi: 10.1093/femsre/fuy023
- Zhu, B., Bradford, L., Huang, S., Szalay, A., Leix, C., Weissbach, M., et al. (2017). Unexpected diversity and high abundance of putative nitric oxide dismutase (n.d.) genes in contaminated aquifers and wastewater treatment systems. *Appl. Environ. Microbiol.* 83:2750. doi: 10.1128/AEM.02750-16
- Zhu, B., van Dijk, G., Fritz, C., Smolders, A. J. P., Pol, A., Jetten, M. S. M., et al. (2012). Anaerobic oxidation of methane in a minerotrophic peatland: enrichment of nitrite-dependent methane-oxidizing bacteria. *Appl. Environ. Microbiol.* 78, 8657–8665. doi: 10.1128/AEM.02102-12
- Zhu, B., Wang, Z., Kanaparthi, D., Kublik, S., Ge, T., Casper, P., et al. (2020). Long-read amplicon sequencing of nitric oxide dismutase (n.d.) genes reveal diverse oxygenic denitrifiers in agricultural soils and lake sediments. *Microb. Ecol.* 80, 243–247. doi: 10.1007/s00248-020-01482-0
- Zhu, J., Wang, Q., Yuan, M., Tan, G.-Y. A., Sun, F., Wang, C., et al. (2016). Microbiology and potential applications of aerobic methane oxidation coupled to denitrification (AME-D) process: a review. *Water Res.* 90, 203–215. doi: 10.1016/j.watres.2015.12.020
- Zhu, G., Zhou, L., Wang, Y., Wang, S., Guo, J., Long, X. E., et al. (2015). Biogeographical distribution of denitrifying anaerobic methane oxidizing bacteria in Chinese wetland ecosystems. *Environ. Microbiol. Rep.* 7, 128–138. doi: 10.1111/1758-2229.12214

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Guerrero-Cruz, Vaksmaa, Horn, Niemann, Pijuan and Ho. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.