



Effect of a *Monascus* sp. Red Yeast Rice Extract on Germination of Bacterial Spores

Marketa Husakova, Michaela Plechata, Barbora Branska and Petra Patakova*

Department of Biotechnology, University of Chemistry and Technology Prague, Prague, Czechia

OPEN ACCESS

Edited by:

Wanping Chen,
Georg-August-University Goettingen,
Germany

Reviewed by:

Yanli Feng,
Hubei Normal University, China
Yanchun Shao,
Huazhong Agricultural University,
China

*Correspondence:

Petra Patakova
petra.patakova@vscht.cz

Specialty section:

This article was submitted to
Food Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 26 March 2021

Accepted: 27 April 2021

Published: 24 May 2021

Citation:

Husakova M, Plechata M,
Branska B and Patakova P (2021)
Effect of a *Monascus* sp. Red Yeast
Rice Extract on Germination
of Bacterial Spores.
Front. Microbiol. 12:686100.
doi: 10.3389/fmicb.2021.686100

The pink-red color of traditional sausages (cured meat) is the result of nitrite addition and the formation of nitrosomyoglobin. However, the pleasant color of processed meat products is a side effect of nitrite addition while the main anticipated goal is to suppress the germination of clostridial spores. The fungus *Monascus* is known as a producer of oligoketide pigments, which are used in Asian countries, especially in China, for coloring foods, including meat products. Although, different biological activities of *Monascus* pigments have been tested and confirmed in many studies, their effect on germination of bacterial spores has never been investigated. This study is focused on testing the activity of red yeast rice (RYR) extract, containing monascin, rubropunctatin, rubropunctamine complexes and monascuspiloin as the main pigments, on germination of *Clostridium* and *Bacillus* spores. It was found that addition of nitrite alone, at the permitted concentration, had no effect on spore germination. However, the combined effects of nitrite with NaCl, tested after addition of pickling salt, was efficient in inhibiting the germination of *C. beijerinckii* spores but had no effect on *B. subtilis* spores. In contrast, total suppression of *C. beijerinckii* spore germination was reached after addition of RYR extract to the medium at a concentration of 2% v/v. For *B. subtilis*, total inhibition of spore germination was observed only after addition of 4% v/v RYR extract to the medium containing 1.3% w/w NaCl.

Keywords: *Monascus*, red yeast rice, bacterial spores germination, nitrite, *Clostridium beijerinckii*, *Bacillus subtilis*

INTRODUCTION

Red yeast rice (RYR) is rice fermented by the fungus *Monascus*, which is prepared for different applications using different *Monascus* species (for a recent review see Zhu et al., 2019). RYR has various synonyms such as hong-qu, beni-koji or ang-kak in languages of Asian countries, where the product is popular. In Europe, RYR is only permitted as a food supplement on the condition that it is prepared using *Monascus purpureus* and the preparation (food supplement) should contain 10 mg of monacolin K, administered daily, in order to guarantee the effect as described in the health claim “Monacolin K from red yeast rice contributes to the maintenance of normal blood cholesterol levels” (Commission Regulation (EU) No 432/2012, 2012). As a dose of 10 mg of monacolin K corresponds to the lowest therapeutically effective dose of statins in prescription drugs, the required amount of monacolin K in RYR food supplements was reconsidered by the EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS) in

2018 (EFSA Panel on Food Additives and Nutrient Sources added to Food [ANS] et al., 2018) but with no clear conclusion. Nevertheless, the main concern associated with the use of RYR is its potential contamination with citrinin, a mycotoxin whose toxicological effects on people have not been fully elucidated (de Oliveira Filho et al., 2017). By the Commission Regulation (EU) 2019/1901, 2019, the maximum tolerated citrinin concentration in RYR was set to 100 µg/kg.

The fungus *Monascus* is especially known for its production of red pigments, which are used in certain Asian countries, such as China, Japan or Philippines, for food coloring. As the color red is associated with many different fruits, vegetables and meat products, *Monascus* pigments are mostly used for coloring cakes or other sweet products, fruit yogurts or other fermented milk products and processed meat. The suitability of *Monascus* pigments for coloring meat products, particularly with regard to color, texture, smell and other sensory parameters of the products, has already been proven in the scientific literature (Leistner et al., 1991; Fabre et al., 1993; Yu et al., 2015; Seong et al., 2017). In addition, inhibitory effects of *Monascus* pigments or *Monascus* extracts on vegetative bacterial cells, e.g., *Staphylococcus aureus*, *Escherichia coli*, *Bacillus subtilis* (Kim et al., 2006; Vendruscolo et al., 2014; Zhao et al., 2016) have been demonstrated. In addition, the safety of *Monascus* pigments for human consumption has been confirmed in several studies (Bianchi, 2005; Yu et al., 2008; Mohan Kumari et al., 2009).

The aim of the study was to test whether an ethanol extract of RYR, having a red color and containing a mixture of *Monascus* pigments but without citrinin and monacolin K, might suppress germination of bacterial spores. In traditional food processing, nitrite salts have been added to meat products in order to achieve total inhibition of germination of *Clostridium botulinum* spores. Nitrite salts are also responsible for the pleasing red color of processed meat products, caused by the formation of nitrosomyoglobin. In the work described for the first time here, *Clostridium beijerinckii* and *Bacillus subtilis* spores were used as models of anaerobic and aerobic bacterial spore formers, respectively.

MATERIALS AND METHODS

RYR Preparation

Monascus sp. DBM 4361, isolated from a non-sterile dried red fermented rice sample, was maintained on Potato-Dextrose agar (VWR Chemicals) slants at 4°C. The strain was deposited at the Department of Biochemistry and Microbiology (DBM), University of Chemistry and Technology Prague.

An amount of rice (Giana, Thailand) (150 g) was washed with hot water, then boiled for 1 min. The rice was evenly divided into three autoclavable plastic bags, which were closed with a metal ring and a cotton plug. The use of plastic bags instead of glass vessels enabled manual separation of rice kernels after sterilization without opening of the bags. The cultivation was inspired by soy koji preparation (Lotong and Suwanarit, 1983). The bags were placed in a beaker sealed with aluminum foil and sterilized at 121°C for 20 min. Sterilization was repeated after 24 h

to eliminate contamination by spore-forming bacteria. Spores from the *Monascus* culture (mixture of ascospores and conidia, because the strain formed both asexual and sexual spores, see **Figure 1**) were transferred to sterile water using a sterile loop. The sterile rice in the bags was inoculated with 5 mL of the spore suspension. Cultivation of the fungus on rice was performed for 10 days at 30°C. The rice was mixed by hand daily.

Extraction of Pigments, pH Estimation

The RYR (**Figure 2**) (5 g) was extracted with 25 mL of 70% ethanol and distilled water in 250 mL Erlenmeyer flasks for 1 h, at 30°C, with shaking (laboratory shaker Infors, 100 rpm). The mixture was then filtered through Whatman 1 filter paper. Pooled ethanol extracts from 3 flasks were concentrated using a rotary vacuum evaporator (Boeck) (max. temperature 55°C), so that all ethanol was evaporated. The remaining water extract contained insoluble pigmented particles, which were collected by filtration and dissolved in 96% ethanol. The resulting ethanol extract, having dry matter concentration 9.4 mg/mL, was used for all microbiological assays and was analyzed by HPLC. The pH of pooled water extracts from 3 flasks was measured and shown to be 4.9.

RYR Extract Analysis

Spectrophotometric Analysis of the RYR Extract

The RYR extract was diluted 200-fold with 96% ethanol to adjust the absorbance to 0.1–1.0 at 330–600 nm. The absorbance of the sample was detected using a spectrophotometer (Varian Cary 50 Bio). The results were represented as an absorbance spectrum or as an absorbance value at a selected wavelength, the dilution factor being taken into consideration. As a blank, 96% ethanol was used.

UHPLC Analysis of Pigments, Citrinin, and Monacolin K

UHPLC (Agilent Technologies 1260 Infinity II) was used to determine *Monascus* pigments, citrinin and monacolin K. The



FIGURE 1 | Mycelium, conidia and cleistothecia of *Monascus* sp. DBM 4361. An example of a conidium is marked with a black arrow, the red arrow shows a cleistothecium with ascospores. The specimen was prepared from the fungus grown on a PDA agar slant for 5 days at 30°C; magnification was 400×.



FIGURE 2 | Red yeast rice fermented with *Monascus* sp. DBM 4361 for 10 days at 30°C.

following conditions were used: Kinetex Polar C18, 100A, 150 × 4.6 mm column; the mobile phase: 0.025% H₃PO₄ in water:acetonitrile at a ratio of 60:40; isocratic elution at a flow rate 1.5 mL/min; injection volume 5 μL. For determination of yellow, orange and red pigments, a photodiode detector set at 390, 470, and 500 nm resp. was used. The presence of monacolin K was detected at 237 nm. For the determination of the mycotoxin citrinin, the fluorescence detector setting was 331 nm for excitation and 500 nm for emission. For analysis, the extract sample was diluted 10-fold with 96% ethanol.

The standards of mycotoxin citrinin (Sigma-Aldrich), yellow pigment monascins (Sigma-Aldrich), and orange pigment rubropunctatin (1717 CheMall Corporation) were used as reference samples. A rubropunctamine laboratory standard was prepared from the rubropunctatin standard by reaction with NH₄OH (Penta). Unknown yellow, orange and red pigments were identified based on their absorption spectra and quantified as equivalents to their respective standards, i.e. monascins, rubropunctatin and rubropunctamine. For identification of individual compounds in the chromatogram, previous results (Patrovsky et al., 2019) were used.

HPLC-MS Analysis

HPLC-HRMS (Accella 600 Thermo Fisher Scientific) was used to determine molecular weights of the *Monascus* pigments. The following conditions were used: Luna Omega Polar 1.6 μm, 50 × 2.1 mm Phenomenex column; mobile phase: 0.1% HCOOH in water:methanol; gradient elution (A:B in ratio from 90:10 to 5:95) at a flow rate of 300 μL/min; injection volume 5 μL; ESI-positive mode; LTQ Orbitrap Velos mass analyzer (Thermo

Scientific). For the analysis, the extract sample was diluted 200-fold with a solution of 90% water 10% methanol.

Spore Germination Assays

Preparation of Spore Suspensions

Clostridium beijerinckii NCIMB 8052 and *Bacillus subtilis* DBM 3006 were stored in the form of spore suspensions in sterile distilled water at 4°C. Spores of *C. beijerinckii* NCIMB 8052 were obtained after 48 h incubation of the culture in 250 mL Erlenmeyer flasks containing 100 mL of TYA medium in an anaerobic chamber (Concept 400, Ruskin Technology, United Kingdom) at 37°C. TYA medium contained in g/L: glucose 40, yeast extract (Merck) 2, tryptone (Sigma-Aldrich) 6, potassium dihydrogenphosphate 0.5, ammonium acetate 3, magnesium sulfate heptahydrate, 0.3, ferrous sulfate heptahydrate 0.01; the pH of the medium was adjusted prior to sterilization in the autoclave (20 min, 121°C, 0.1 MPa) to 6.8. Spores of *B. subtilis* DBM 3006 were obtained after 48 h incubation of the culture in 250 mL Erlenmeyer flasks containing 50 mL of MP broth shaken on a rotary shaker (New Brunswick Scientific) at 300 rpm and 30°C. MP broth contained in g/L: meat extract (Roth) 3, peptone (Roth) 5; the pH of the medium was adjusted prior to sterilization in the autoclave (20 min, 121°C) to 7.0. Salts, HCl and NaOH for medium preparation and pH adjustment were purchased from Penta, Czech Republic. After cultivation, spores of both bacterial cultures were harvested by centrifugation (Hettich MIKRO 220R) for 5 min, 5,000 rpm, at 4°C. Spores were washed with sterile water, centrifuged under the same conditions and re-suspended in 20 mL of sterile water. Finally, the spore suspension was pipetted in 1 mL portions into Eppendorf tubes, which were stored at 4°C. All spore handling was performed under aseptic conditions using sterile tools and materials. Spore concentrations were estimated by flow cytometry (Branska et al., 2018) to be 2 × 10⁸ spores/mL and 8 × 10⁸ spores/mL for *C. beijerinckii* and *B. subtilis*, respectively. Prior to inoculation of medium for germination assays, spore suspensions were heat shocked (80°C for 30 s followed by cooling on ice for 2 min) to accelerate germination.

C. beijerinckii Germination Assay

The cultivation tests were performed in 20 mL test tubes containing 9.8 mL of medium inoculated with 0.2 mL heat shocked spore suspension of *C. beijerinckii*. Unmodified TYA medium was used as a positive control. The cultivation tests were performed in triplicate, in the anaerobic chamber, at 37°C for 48 h. Different combinations of agents were added to the TYA medium to test their effect on spore germination (see **Table 1**).

B. subtilis Germination Assay

The cultivation tests were performed on microcultivation plates using the Bioscreen C device (LabSystem) with intermittent shaking (30 s every 3 min) at 30°C for 24 h. Each well of the plate was filled with 196 μL of medium and 4 μL of *B. subtilis* heat shocked spore suspension. In each well, optical density was measured at 600 nm, every 30 min. Unmodified MP medium was used as a positive control. Each cultivation test was performed in

TABLE 1 | Design of *C. beijerinckii* germination assays in individual test tubes.

Assay component	TYA (μL)	96% ethanol (μL)	RYR extract (μL)	Citrinin (μL)	Inoculum Heat-shocked spores (μL)
Code	Type of experiment				
TYA	Positive control	9,800	–	–	200
TYA	Negative control	10,000	–	–	–
TYA + et	Control	9,600	200	–	200
TYA + RYR	Test	9,600	–	200	200
TYA + cit	Test	9,780	–	–	20
TYA + RYR + cit	Test	9,580	–	200	20
		TYAS1 (μL)	96% ethanol (μL)	RYR extract (μL)	Inoculum Heat-shocked spores (μL)
TYAS1	Test	9,800	–	–	200
TYAS1 + et	Control	9,600	200	–	200
TYAS1 + RYR	Test	9,600	–	200	200
		TYAS2 (μL)			Inoculum Heat-shocked spores (μL)
TYAS2	Test	9,800	–	–	200
		TYAN1 (μL)			Inoculum Heat-shocked spores (μL)
TYAN1	Test	9,800	–	–	200
		TYAN2 (μL)			Inoculum Heat-shocked spores (μL)
TYAN2	Test	9,800	–	–	200
		TYA + NaNO₂ (μL)			Inoculum Heat-shocked spores (μL)
TYA + NaNO ₂	Test	9,800	–	–	200
		TYA + NaNO₃ (μL)			Inoculum Heat-shocked spores (μL)
TYA + NaNO ₃	Test	9,800	–	–	200

TYA, unmodified TYA medium with 20 g/L glucose; TYAS1, TYAS2, TYA medium with the addition of 1.3 and 2.0% w/w NaCl, respectively, TYAN1, TYAN2, TYA medium with the addition of 1.3 and 2.0% w/w Praganda butcher's nitrite pickling mixture (K + S company, Czech Republic); TYA + NaNO₂ and TYA + NaNO₃, TYA media with the addition of 300 mg/L NaNO₂ or NaNO₃; citrinin was added from stock solution, of concentration 100 mg/L in dimethylsulfoxide; et, ethanol; cit, citrinin.

6 wells. Different combinations of agents were added to the MP medium to test their effect on spore germination (see **Table 2**).

RESULTS

C. beijerinckii Germination

Clostridium beijerinckii spores were chosen as a model substituting for *Clostridium botulinum* spores because both species belong to the same Cluster I (*sensu stricto*) of the *Clostridium* genus (Cruz-Morales et al., 2019). Heat-shocked spores of *C. beijerinckii* were inoculated to 2% by volume to the medium in test tubes and were allowed to germinate under anaerobic conditions. The TYA medium and the inoculation ratio were chosen to guarantee reliable spore germination based on previous experience with the strain (Kolek et al., 2016). The compositions of different media were designed (see **Table 1**) to be able to compare any effect of nitrite salts with the potential effect of the RYR extract, not containing citrinin and monacolin K. Nitrites are only allowed to be added to meat products in a mixture with sodium chloride in the form of

NaNO₂ (E249 food additive) or KNO₂ (E250 food additive), in an amount not exceeding 150 mg of nitrite per 1 kg of a standard meat product (only in some national specialities produced in different EU countries can the amount of nitrite be higher, up to 300 mg/kg, for certain salamis and bacons and up to 500 mg/kg for herrings and sprouts); for a survey of rules valid in the EU for the addition of nitrite/nitrate to meat products (see Honikel, 2008).

Nitrite, in the form of NaNO₂, was added to the culture, either independently or as a component of Praganda nitrite pickling salt. In addition, NaNO₃ (E251 food additive) was tested because its addition is permitted and applied in cheeses to a maximum concentration of 150 mg/kg, with the aim of suppressing germination of *Clostridium butyricum* spores. While even the addition of NaNO₂ or NaNO₃ alone at a concentration of 300 mg/kg had no effect on the germination of *C. beijerinckii* spores, addition of nitrite pickling salt to the recommended concentration for various products, i.e., 2% (w/w) of the pickling salt or 1.3% (w/w) as recommended for products with low salt content, resulted in total inhibition of spore germination (see **Table 3**, experiment

TABLE 2 | Design of *B. subtilis* germination assays in individual wells of a microcultivation plate.

Assay component	MP (μL)	96% ethanol (μL)	RYR extract (μL)	Citrinin (μL)	Inoculum Heat-shocked spores (μL)	
Code	Type of experiment					
MP	Positive control	196	–	–	–	4
MP	Negative control	200	–	–	–	–
MP + et1	Control	192	4	–	–	4
MP + et2	Control	188	8	–	–	4
MP + RYR1	Test	192	–	4	–	4
MP + RYR2	Test	188	–	8	–	4
MP + cit	Test	196	–	–	0.8	4
MP + RYR1 + cit	Test	192	–	4	0.8	4
		MPS1 (μL)	96% ethanol (μL)	RYR extract (μL)		Inoculum Heat-shocked spores (μL)
MPS1	Test	196	–	–	–	4
MPS1 + et1	Control	192	4	–	–	4
MPS1 + et2	Control	188	8	–	–	4
MPS1 + RYR1	Test	192	–	4	–	4
MPS1 + RYR2	Test	188	–	8	–	4
		MPN2 (μL)				Inoculum Heat-shocked spores (μL)
MPN2	Test	196	–	–	–	4

MP, unmodified MP medium; MPS1, MP medium with addition of 1.3% w/w NaCl; MPN2, MP medium with addition of 2.0% w/w Praganda butcher's nitrite pickling mixture (K + S company, Czech Republic); citrinin was added from a stock solution, of concentration 100 mg/L in dimethylsulfoxide.

TABLE 3 | Results of *C. beijerinckii* germination assays.

Code	Type of experiment	Experiment description	Result
TYA	Positive control	–	+++
TYA	Negative control	–	–
TYA + et	Control	Control ethanol addition	+++
TYA + RYR	Test	Addition of RYR extract in concentration 2% v/v	–
TYA + cit	Test	Addition of citrinin in concentration 2,000 μg/kg	+++
TYA + RYR + cit	Test	Combined addition of citrinin in concentration 2,000 μg/kg and RYR in concentration 2% v/v	–
TYAS1	Test	Addition of NaCl in concentration 1.3% w/w	++-*
TYAS1 + et	Control	Combined addition of NaCl in concentration 1.3% w/w and ethanol in concentration 2% v/v	–
TYAS1 + RYR	Test	Combined addition of NaCl in concentration 1.3% w/w and RYR in concentration 2% v/v	–
TYAS2	Test	Addition of NaCl in concentration 2% w/w	–
TYAN1	Test	Addition of nitrite pickling salt in concentration 1.3% w/w	+--*
TYAN2	Test	Addition of nitrite pickling salt in concentration 2% w/w	–
TYA + NaNO ₂	Test	Addition of alone NaNO ₂ in concentration 300 mg/L	+++
TYA + NaNO ₃	Test	Addition of alone NaNO ₂ in concentration 300 mg/L	+++

The experimental code correlates with the experimental design given in **Table 1**. All experiments were performed in triplicate, spore outgrowth observed in all 3 test tubes is marked as +++, no growth as –. *In case of dubious results, the cultivation test was repeated but if the result was the same, i.e., ++– or +– this means unreliable suppression of germination. The most significant results are highlighted.

codes TYA + NaNO₂; TYA + NaNO₃; TYAN2 and TYAN1). As *Clostridium beijerinckii* is sensitive to high concentrations of NaCl (Branska et al., 2020), the independent effect of NaCl addition was also tested. While addition of NaCl to 2% w/w suppressed spore germination, 1.3% w/w did not reliably suppress germination in all cases (see **Table 3**, code TYAS2 and TYAS1).

Addition of 2% v/v RYR extract was tested with standard TYA medium and with medium containing 1.3% w/w NaCl; addition of ethanol to the same concentration (2% v/v) was tested as a control (**Table 3**, codes TYA + RYR, TYA + Et, TYAS1 + RYR, TYAS1 + Et). While addition of the RYR extract suppressed spore germination, ethanol did not. The RYR extract did not contain citrinin but, because it is known that




FIGURE 3 | Demonstration of growth in test tubes. Positive control, i.e., outgrowth of *C. beijerinckii* spores in TYA medium is shown on the left while suppression of spore germination after addition of RYR extract to 2% v/v is shown on the right. A test tube with medium was photographed (A) prior to inoculation, (B) after inoculation, and (C) after 48 h cultivation.

citrinin has certain antimicrobial properties, the citrinin effect was tested at a concentration of 2,000 $\mu\text{g}/\text{kg}$. This concentration of citrinin reflects the amount that was EU-permitted in an RYR food supplement until 2019, after which the limit was reconsidered and adjusted to 100 $\mu\text{g}/\text{kg}$. Citrinin was tested alone or in combination with RYR (Table 3, codes TYA + cit, TYA + RYR + cit).

Typical test tube growth characteristics of *C. beijerinckii* exhibiting high turbidity, foam and development of bubbles of fermentation gas (mixture of CO_2 and hydrogen) as well as color of the medium after addition of the RYR extract are shown in Figure 3. Only the ability to grow (indicated as + or –) was tested in the *C. beijerinckii* germination assay. Determination of optical density was not performed in order to not disturb the anaerobic atmosphere. Changed morphology from vegetative cells to spores influences OD values, so comparisons between tests might be misleading. Spores in TYA medium (positive control experiment) started to germinate 16–18 h after inoculation, and in other cases, germination started with different delays.

B. subtilis Germination


To follow the growth of spore formers after germination, *Bacillus subtilis* germination assays were performed, even if these spores do not normally occur in meat products (Figure 4). Nevertheless, *B. subtilis* spores are more resistant to adverse environmental effects in comparison with the *C. beijerinckii* spores, therefore the design of experiments had to be different (Table 2) in order to inhibit germination. Total suppression of spore germination was achieved only after addition of the RYR extract to 4% v/v in medium containing 1.3% w/w NaCl (Figure 4D, MPS1 + RYR2). In other cases, growth was always detected, even if, in some cases, germination was delayed by up to 10 h (Figures 4B,C,

MP + RYR2; MP + RYR1 + Cit). Surprisingly, addition of nitrite pickling salt to the recommended 2% w/w did not inhibit germination of the spores (Figure 4A, MPN2).

RYR Extract Analysis

The absorption spectrum of the RYR extract is shown in Figure 5. Values of absorption found by spectrophotometric analysis at 390, 470, and 500 nm, corresponding to assumed absorption maxima of yellow, orange and red pigments, were 98, 58, and 70, respectively. Monascin (yellow), rubropunctatin (orange) and rubropunctamine (red) were identified in the RYR extract by UHPLC analysis (Figure 6), while their analogs with seven carbon side chains, i.e., ankaflavin (yellow), monascorubrin (orange) and monascorubramine (red) were not detected; neither was citrinin or monacolin K. The detected yellow pigments were quantified as monascin equivalents (1m220 mg/L), orange pigments as rubropunctatin equivalents (336 mg/L) and red pigments as rubropunctamine equivalents (408 mg/L). However, other compounds labeled as yellow I and red I–VI were found and for their putative identification, HPLC-MS analysis and already published m/z data on different *Monascus* metabolites were used (for survey of the *Monascus* pigments data see Chen et al., 2019). While red I–red VI are probably rubropunctamine derivatives that were formed by the reaction of rubropunctatin with available amino group containing compounds, yellow I was identified as monascupiloin (m/z 360.4).

The RYR extract was added to TYA and MP medium to concentrations of 2 and 4% v/v, respectively, and the concentrations of pigments added to the medium are shown in Table 4. The water extract of the RYR had a pH 4.9 but the pH of TYA or MP medium was 6.8 or 7.0. To determine whether the pigment profile was the same as the original RYR extract in




the culture medium, samples of medium containing RYR mixed in a 1:1 volume ratio were analyzed by UHPLC. As expected, rubropunctatin was absent in the samples while monascin and monascuspiloin remained (data not shown).

DISCUSSION

Nitrite and nitrate addition to meat products, together with NaCl, is traditional in European countries and is considered to be of low impact on human health even if cancerogenic nitrosamines can be formed in the acidic environment of the

human stomach after ingestion of nitrite/nitrate containing food (Honikel, 2008; EFSA Panel on Food Additives Nutrient Sources added to Food [ANS] et al., 2017). It is believed that the benefits of stable red color, antioxidant and antimicrobial effects of nitrite/nitrate outweigh potential risks. Nevertheless, within this study it was found that addition of nitrite or nitrate alone, to the permitted concentration, did not suppress spore germination. Similar observations were documented in other studies for the germination of *Clostridium perfringens* (Labbe and Duncan, 1970), *Clostridium botulinum* (Sofos et al., 1979) or cheese associated clostridia including *C. beijerinckii* (Ávila et al., 2014) spores. However, combined effect of nitrite with NaCl tested after addition of the pickling salt both at the standard concentration of pickling salt (2% w/w) and at the level recommended for low salt products (1.3% w/w) was efficient in inhibiting the germination of *C. beijerinckii* spores but had no effect on germination of *B. subtilis* spores. However, total suppression of germination of *C. beijerinckii* spores was also achieved after addition of RYR extract to TYA medium, to a concentration of 2% v/v while total suppression of germination of *B. subtilis* spores was only achieved after addition of 4% v/v RYR extract to MP containing 1.3% w/w NaCl. These results suggest that the RYR extract might substitute for nitrite salts in inhibiting germination of *Clostridium* spores.

Within the study, the ethanol effect on bacterial spore germination was confirmed (Setlow et al., 2002) as well as the effect of NaCl (Nagler et al., 2014) and the synergistic effect of






FIGURE 6 | RYR extract chromatogram from UHPLC analysis. Detected compounds are given in black while expected but undetected compounds are given in gray color. Conditions of UHPLC analysis are given in section “RYR Extract Analysis.”

different agents (Nerandzic et al., 2015). Even if the RYR extract did not contain citrinin, its effect at 2,000 $\mu\text{g/L}$ (the permitted concentration of citrinin in RYR food supplements in the EU until 2019) was tested, but if applied independently, had no effect on spore germination.

Monascus sp. DBM 4361, used for the preparation of RYR, was not classified on the species level but it might be *M. pilosus*

because its characteristics, i.e., no citrinin production and formation of both conidia and ascospores, corresponds with an already described strain, *M. pilosus* MS-1, also isolated from the RYR (Feng et al., 2016). The absence of citrinin production was also found in *M. pilosus* NBRC4520 (Higa et al., 2020). In the RYR extract, there were found three of six iconic *Monascus* pigments; in particular monascin, rubropunctatin and rubropunctamine, together with rubropunctamine complexes with different amino group-containing compounds and monascuspiloin, a yellow pigment with a structure similar to monascin that has already been described as the metabolite of *M. pilosus* M93 (Chen et al., 2012). Interestingly, only monascin and rubropunctatin, i.e., pigments with a shorter five carbon side chain, but not their analogs (ankaflavin and monascorubrin with seven carbon side chains) were found. During the biosynthesis of pigments by *M. ruber* M7 (Chen et al., 2017), MrPigJ and MrPigK subunits of fatty acid synthase were found as the proteins responsible for the integration of β -ketoctanoic or β -ketodecanoic acid moieties into the structure of the pigments. However, the selection of the particular fatty acid moiety was considered to be random or not yet understood. Homologous genes to MrPigJ, MrPigK were described in *M. purpureus* (MpFasA2, MpFasB2) (Balakrishnan et al., 2013) and other *Monascus* strains (Guo et al., 2019) but not in *M. pilosus*. Even in the newest review of azaphilone

TABLE 4 | Concentration of pigments quantified as monascin, rubropunctatin, and rubropunctamine equivalents added to TYA and MP media.

Medium code*	Yellow pigments (concentration equivalent to Monascin in mg/L)	Orange pigments (concentration equivalent to Rubropunctatin in mg/L)	Red pigments (concentration equivalent to Rubropunctamine in mg/L)
TYA + RYR	24.4	6.7	8.2
TYAS1 + RYR			
MP + RYR1	24.4	6.7	8.2
MPS1 + RYR1			
MP + RYR2	48.8	13.4	16.4
MPS1 + RYR2			

*Medium codes are the same as used in Tables 1, 2.

biosynthesis (Pavesi et al., 2021), factors that determine the selection of particular fatty acid moieties are not described.

For the detection of substances in *Monascus* extracts, it is not sufficient to use the absorption spectrum or to determine absorbance values of the extract at the absorption maxima of individual pigments, typically at 390, 470, and 500 nm. It is really necessary to analyze extracts by HPLC or other analytical method (cf. Figures 5, 6). Notwithstanding, the determination of individual compounds in *Monascus* extracts is difficult because it depends *inter alia* on the pH of the extract (Shi et al., 2016).

After the addition of the RYR extract to medium, the original pH of the RYR extract changed from 4.9 to 6.8 or 7.0, resulting in the reaction of rubropunctatin with available amino group containing compounds, such as amino acids. It is possible that this reaction might contribute to the suppression of spore germination in the medium. Amino acids such as L-alanine or glycine are known germinants (factors stimulating germination) of bacterial spores (Setlow, 2014; Bhattacharjee et al., 2016) and if their amount was decreased it might affect germination. In addition, it was reported that *Monascus* red pigment derivatives had pronounced effects on the growth of Gram positive bacteria, including *B. subtilis* (Kim et al., 2006). The assumed cause of inhibition was adsorption of pigment derivatives onto the surface of cells, limiting oxygen uptake, where their MIC values were found to be 4–8 µg/L. In our assay, rubropunctatin and a mixture of red pigment derivatives, to concentrations of 13.4 mg/L and 16.4 mg/L, respectively, were added to MP culture medium used for outgrowth of *B. subtilis* spores (see Table 4) and *B. subtilis* was cultured under aerobic conditions; the above aerobic effect (Kim et al., 2006) might also apply here. Orange and red *Monascus* pigments, at concentrations of 10–20 mg/L, inhibited

growth of Gram negative bacteria (Vendruscolo et al., 2014), which corresponds with our findings. Yellow pigments monascin and monascuspiloin, detected in the RYR, were found to have anticarcinogenic effects (Akihisa et al., 2005; Chen et al., 2012; Chiu et al., 2012) but their antibacterial effect was never tested.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

AUTHOR CONTRIBUTIONS

MH performed cultivation experiments, prepared samples for analyses, participated in analysis, performed data evaluation, and revised the manuscript. MP assisted with cultivation experiments, preparation of samples for analyses, participated in analysis, and revised the manuscript. BB developed the UHPLC method for the pigments analysis and revised the manuscript. PP conceived the study, designed and coordinated it, analyzed the data, and wrote the manuscript. All authors read and approved the final manuscript version.

FUNDING

Financial support from specific University of Chemistry and Technology in Prague research, Project No. MEYS No 8-SVV/2021.

REFERENCES

- Akihisa, T., Tokuda, H., Ukiya, M., Kiyota, A., Yasukawa, K., Sakamoto, N., et al. (2005). Anti-tumor-initiating effects of monascin, an azaphilone pigment from the extract of *Monascus pilosus* fermented rice (red-mold rice). *Chem. Biodivers* 2, 1305–1309. doi: 10.1002/cbdv.200590101
- EFSA Panel on Food Additives and Nutrient Sources added to Food [ANS], Younes, M., Aggett, P., Aguilar, F., Crebelli, R., Dusemund, B. (2018). Scientific opinion on the safety of monacolins in red yeast rice. *EFSA J.* 16:e05368.
- Ávila, M., Gómez-Torres, N., Hernández, M., and Garde, S. (2014). Inhibitory activity of reuterin, nisin, lysozyme and nitrite against vegetative cells and spores of dairy-related *Clostridium* species. *Int. J. Food Microbiol.* 172, 70–75. doi: 10.1016/j.ijfoodmicro.2013.12.002
- Balakrishnan, B., Karki, S., Chiu, S.-H., Kim, H.-J., Suh, J.-W., Nam, B., et al. (2013). Genetic localization and in vivo characterization of a *Monascus* azaphilone pigment biosynthetic gene cluster. *Appl. Microbiol. Biotechnol.* 97, 6337–6345. doi: 10.1007/s00253-013-4745-9
- Bhattacharjee, D., McAllister, K. N., and Sorg, J. A. (2016). Germinants and their receptors in *Clostridia*. *J. Bacteriol.* 198, 2767–2775. doi: 10.1128/JB.00405-16
- Bianchi, A. (2005). Extracts of *Monascus purpureus* beyond statins —Profile of efficacy and safety of the use of extracts of *Monascus purpureus*. *Chin. J. Integr. Med.* 11, 309–313. doi: 10.1007/BF02835797
- Branska, B., Fořtová, L., Dvořáková, M., Liu, H., Patakova, P., Zhang, J., et al. (2020). Chicken feather and wheat straw hydrolysate for direct utilization in biobutanol production. *Renewable Energy* 145, 1941–1948. doi: 10.1016/j.renene.2019.07.094
- Branska, B., Pechacova, Z., Kolek, J., Vasylikivska, M., and Patakova, P. (2018). Flow cytometry analysis of *Clostridium beijerinckii* NRRL B-598 populations exhibiting different phenotypes induced by changes in cultivation conditions. *Biotechnol. Biofuels* 11:99. doi: 10.1186/s13068-018-1096-x
- Chen, R.-J., Hung, C.-M., Chen, Y.-L., Wu, M.-D., Yuan, G.-F., and Wang, Y.-J. (2012). Monascuspiloin induces apoptosis and autophagic cell death in human prostate cancer cells via the Akt and AMPK signaling pathways. *J. Agric. Food Chem.* 60, 7185–7193. doi: 10.1021/jf3016927
- Chen, W., Chen, R., Liu, Q., He, Y., He, K., Ding, X., et al. (2017). Orange, red, yellow: biosynthesis of azaphilone pigments in *Monascus* fungi. *Chem. Sci.* 8, 4917–4925. doi: 10.1039/c7sc00475c
- Chen, W., Feng, Y., Molnár, I., and Chen, F. (2019). Nature and nurture: confluence of pathway determinism with metabolic and chemical serendipity diversifies *Monascus* azaphilone pigments. *Nat. Product Rep.* 36, 561–572. doi: 10.1039/C8NP00060C
- Chiu, H.-W., Fang, W.-H., Chen, Y.-L., Wu, M.-D., Yuan, G.-F., Ho, S.-Y., et al. (2012). Monascuspiloin enhances the radiation sensitivity of human prostate cancer cells by stimulating endoplasmic reticulum stress and inducing autophagy. *PLoS One* 7:e40462. doi: 10.1371/journal.pone.0040462
- Commission Regulation (EU) 2019/1901, (2019). Commission Regulation (EU) 2019/1901 of 7 November 2019 Amending Regulation (EC) No 1881/2006 as Regards Maximum Levels of Citrinin in Food Supplements Based on Rice Fermented with Red Yeast *Monascus Purpureus*. Available online at: <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=OJ:L:2019:293:FULL&from=EN>
- Commission Regulation (EU) No 432/2012, (2012). Commission Regulation (EU) No 432/2012 of 16 May 2012 Establishing a List of Permitted Health Claims Made on Foods, Other Than Those Referring to the Reduction of Disease Risk and to Children's Development and Health. Available

- online at: <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32012R0432&from=CS>
- Cruz-Morales, P., Orellana, C. A., Moutafis, G., Moonen, G., Rincon, G., Nielsen, L. K., et al. (2019). Revisiting the evolution and taxonomy of *Clostridia*, a phylogenomic update. *Genome Biol. Evol.* 11, 2035–2044. doi: 10.1093/gbe/evz096
- de Oliveira Filho, J. W. G., Islam, M. T., Ali, E. S., Uddin, S. J., Santos, J. V. D. O., de Alencar, M. V. O. B., et al. (2017). A comprehensive review on biological properties of citrinin. *Food Chem. Toxicol.* 110, 130–141. doi: 10.1016/j.fct.2017.10.002
- EFSA Panel on Food Additives Nutrient Sources added to Food [ANS], Aguilar, F., Crebelli, R., Di Domenico, A., Dusemund, B., Frutos, M. J. (2017). Re-evaluation of potassium nitrite (E 249) and sodium nitrite (E 250) as food additives. *EFSA J* 15:e04786. doi: 10.2903/j.efsa.2017.4786
- Fabre, C. E., Santerre, A. L., Loret, M. O., Baberian, R., Pareilleux, A., Goma, G., et al. (1993). Production and food applications of the red pigments of *Monascus ruber*. *J. Food Sci.* 58, 1099–1102. doi: 10.1111/j.1365-2621.1993.tb06123.x
- Feng, Y., Chen, W., and Chen, F. (2016). A *Monascus pilosus* MS-1 strain with high-yield monacolin K but no citrinin. *Food Sci. Biotechnol.* 25, 1115–1122. doi: 10.1007/s10068-016-0179-3
- Guo, X., Li, Y., Zhang, R., Yu, J., Ma, X., Chen, M., et al. (2019). Transcriptional regulation contributes more to *Monascus* pigments diversity in different strains than to DNA sequence variation. *World J. Microbiol. Biotechnol.* 35, 1–13. doi: 10.1007/s11274-019-2711-0
- Higa, Y., Kim, Y.-S., Altaf-Ul-Amin, M., Huang, M., Ono, N., and Kanaya, S. (2020). Divergence of metabolites in three phylogenetically close *Monascus* species (*M. pilosus*, *M. ruber*, and *M. purpureus*) based on secondary metabolite biosynthetic gene clusters. *BMC Genomics* 21:679. doi: 10.1186/s12864-020-06864-9
- Honikel, K.-O. (2008). The use and control of nitrate and nitrite for the processing of meat products. *Meat Sci.* 78, 68–76. doi: 10.1016/j.meatsci.2007.05.030
- Kim, C., Jung, H., Kim, Y. O., and Shin, C. S. (2006). Antimicrobial activities of amino acid derivatives of monascus pigments. *FEMS Microbiol. Lett.* 264, 117–124. doi: 10.1111/j.1574-6968.2006.00451.x
- Kolek, J., Branska, B., Drahokoupil, M., Patakova, P., and Melzoch, K. (2016). Evaluation of viability, metabolic activity and spore quantity in clostridial cultures during ABE fermentation. *FEMS Microbiol. Lett.* 363:fnw031. doi: 10.1093/femsle/fnw031
- Labbe, R. G., and Duncan, C. L. (1970). Growth from spores of *Clostridium perfringens* in the presence of sodium nitrite. *Appl. Microbiol.* 19, 353–359.
- Leistner, L., Fink-Gremmels, J., and Dresel, J. (1991). “*Monascus* extract—A possible alternative to nitrite in meats,” in *Proceedings of 37th International Congress of Meat Science and Technology*, (Kulmbach: Federal Centre for Meat Research), 1252–1256.
- Lotong, N., and Suwanarit, P. (1983). Production of soy sauce koji mold spore inoculum in plastic bags. *Appl. Environ. Microbiol.* 46, 1224–1226. doi: 10.1128/AEM.46.5.1224-1226.1983
- Mohan Kumari, H. P., Akhilender Naidu, K., Vishwanatha, S., Narasimhamurthy, K., and Vijayalakshmi, G. (2009). Safety evaluation of *Monascus purpureus* red mould rice in albino rats. *Food Chem. Toxicol.* 47, 1739–1746. doi: 10.1016/j.fct.2009.04.038
- Nagler, K., Setlow, P., Li, Y.-Q., and Moeller, R. (2014). High salinity alters the germination behavior of *Bacillus subtilis* spores with nutrient and nonnutrient germinants. *Appl. Environ. Microbiol.* 80, 1314–1321. doi: 10.1128/AEM.03293-13
- Nerandzic, M. M., Sankar, C. T., Setlow, P., and Donskey, C. J. (2015). A cumulative spore killing approach: synergistic sporocidal activity of dilute peracetic acid and ethanol at low pH against *Clostridium difficile* and *Bacillus subtilis* spores. *Open Forum Infect. Dis.* 3:ofv206. doi: 10.1093/ofid/ofv206
- Patrovsky, M., Sinovska, K., Branska, B., and Patakova, P. (2019). Effect of initial pH, different nitrogen sources, and cultivation time on the production of yellow or orange *Monascus purpureus* pigments and the mycotoxin citrinin. *Food Sci. Nutr.* 7, 3494–3500. doi: 10.1002/fsn3.1197
- Pavesi, C., Flon, V., Mann, S., Leleu, S., Prado, S., and Franck, X. (2021). Biosynthesis of azaphilones: a review. *Nat. Prod. Rep.* doi: 10.1039/D0NP00080A
- Seong, P. N., Ba, H. V., Kim, Y. S., Kang, S. M., Cho, S. H., Kim, J. H., et al. (2017). Effects of additions of *Monascus* and laccase acid on the color and quality properties of nitrite-free emulsion sausage during refrigerated storage. *Korean J. Food Sci. Anim. Resour.* 37, 10–17. doi: 10.5851/kosfa.2017.37.1.10
- Setlow, B., Loshon, C. A., Genest, P. C., Cowan, A. E., Setlow, C., and Setlow, P. (2002). Mechanisms of killing spores of *Bacillus subtilis* by acid, alkali and ethanol. *J. Appl. Microbiol.* 92, 362–375. doi: 10.1046/j.1365-2672.2002.01540.x
- Setlow, P. (2014). Germination of spores of *Bacillus* species: what we know and do not know. *J. Bacteriol.* 196, 1297–1305. doi: 10.1128/JB.01455-13
- Shi, K., Chen, G., Pistolozzi, M., Xia, F., and Wu, Z. (2016). Improved analysis of *Monascus* pigments based on their pH-sensitive UV-Vis absorption and reactivity properties. *Food Addit. Contam. Part A* 33, 1396–1401. doi: 10.1080/19440049.2016.1214289
- Sofos, J., Busta, F., and Allen, C. (1979). Sodium nitrite and sorbic acid effects on *Clostridium botulinum* spore germination and total microbial growth in chicken frankfurter emulsions during temperature abuse. *Appl. Environ. Microbiol.* 37, 1103–1109. doi: 10.1128/AEM.37.6.1103-1109.1979
- Vendruscolo, F., Tosin, I., Giachini, A. J., Schmidell, W., and Ninow, J. L. (2014). Antimicrobial activity of *Monascus* pigments produced in submerged fermentation. *J. Food Process. Preserv.* 38, 1860–1865. doi: 10.1111/jfpp.12157
- Yu, C.-C., Wang, J.-J., Lee, C.-L., Lee, S.-H., and Pan, T.-M. (2008). Safety and mutagenicity evaluation of nanoparticulate red mold rice. *J. Agric. Food Chem.* 56, 11038–11048. doi: 10.1021/jf801335u
- Yu, X., Wu, H., and Zhang, J. (2015). Effect of *Monascus* as a nitrite substitute on color, lipid oxidation, and proteolysis of fermented meat mince. *Food Sci. Biotechnol.* 24, 575–581. doi: 10.1007/s10068-015-0075-2
- Zhao, G.-P., Li, Y.-Q., Yang, J., and Cui, K.-Y. (2016). Antibacterial characteristics of orange pigment extracted from *Monascus* pigments against *Escherichia coli*. *Czech J. Food Sci.* 34, 197–203. doi: 10.17221/430/2015-CJFS
- Zhu, B., Qi, F., Wu, J., Yin, G., Hua, J., Zhang, Q., et al. (2019). Red yeast rice: a systematic review of the traditional uses, chemistry, pharmacology, and quality control of an important chinese folk medicine. *Front. Pharmacol.* 10:1449. doi: 10.3389/fphar.2019.01449

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Husakova, Plechata, Branska and Patakova. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.