



Corrigendum: Asymptomatic Malaria Infection Is Maintained by a Balanced Pro- and Anti-inflammatory Response

OPEN ACCESS

Edited and reviewed by:

Rachel Daniels,
Harvard University, United States

*Correspondence:

Augustina Frimpong
afrimpong@noguchi.ug.edu.gh;
tinafrimp@gmail.com
Kwadwo Asamoah Kusi
akusi@noguchi.ug.edu.gh

Specialty section:

This article was submitted to
Microbial Immunology,
a section of the journal
Frontiers in Microbiology

Received: 26 March 2021

Accepted: 20 April 2021

Published: 19 May 2021

Citation:

Frimpong A, Amponsah J, Adjokatseh AS, Agyemang D, Bentum-Ennin L, Ofori EA, Kyei-Baafour E, Akyea-Mensah K, Adu B, Mensah GI, Amoah LE and Kusi KA (2021) Corrigendum: Asymptomatic Malaria Infection Is Maintained by a Balanced Pro- and Anti-inflammatory Response. *Front. Microbiol.* 12:686435. doi: 10.3389/fmicb.2021.686435

Augustina Frimpong^{1,2,3*}, Jones Amponsah², Abigail Sena Adjokatseh⁴, Dorothy Agyemang⁴, Lutterodt Bentum-Ennin², Ebenezer Addo Ofori^{1,2}, Eric Kyei-Baafour², Kwadwo Akyea-Mensah², Bright Adu², Gloria Ivy Mensah⁵, Linda Eva Amoah^{1,2} and Kwadwo Asamoah Kusi^{1,2,4*}

¹ West African Centre for Cell Biology of Infectious Pathogens (WACCBIP), University of Ghana, Accra, Ghana, ² Department of Immunology, Noguchi Memorial Institute for Medical Research, College of Health Sciences, University of Ghana, Accra, Ghana, ³ African Institute for Mathematical Sciences, Accra, Ghana, ⁴ Department of Biochemistry, Cell and Molecular Biology, College of Basic and Applied Sciences, University of Ghana, Accra, Ghana, ⁵ Department of Bacteriology, Noguchi Memorial Institute for Medical Research, College of Health Sciences, University of Ghana, Accra, Ghana

Keywords: microscopic, *Plasmodium*, anti-inflammatory cytokines, pro-inflammatory cytokines, asymptomatic malaria, submicroscopic

A Corrigendum on

Asymptomatic Malaria Infection Is Maintained by a Balanced Pro- and Anti-inflammatory Response

by Frimpong, A., Amponsah, J., Adjokatseh, A. S., Agyemang, D., Bentum-Ennin, L., Ofori, E. A., et al. (2020) *Front. Microbiol.* 11:559255. doi: 10.3389/fmicb.2020.559255

In the original article, there was a mistake in **Figure 1C** as published. The graph provided was a duplicate of **Figure 1E**. The corrected **Figure 1** and caption appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2021 Frimpong, Amponsah, Adjokatseh, Agyemang, Bentum-Ennin, Ofori, Kyei-Baafour, Akyea-Mensah, Adu, Mensah, Amoah and Kusi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 1 | Profile of pro-inflammatory mediators during microscopic and submicroscopic malaria. Scatter plot graphs are plotted showing the median fluorescence intensities (MFI) of **(A-F)** TNF- α , IL-6, IFN- γ , IL-17A, IL-12p70, and Granzyme B in plasma samples collected from uninfected controls ($n = 18$), patients with microscopic asymptomatic malaria ($n = 38$) and submicroscopic malaria ($n = 22$). Plots show median and interquartile ranges. Significant differences are denoted by * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, ns = not significant.