



# Diversifying Isoprenoid Platforms via Atypical Carbon Substrates and Non-model Microorganisms

David N. Carruthers<sup>1,2</sup> and Taek Soon Lee<sup>1,2\*</sup>

<sup>1</sup>Joint BioEnergy Institute, Emeryville, CA, United States, <sup>2</sup>Biological Systems and Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA, United States

## OPEN ACCESS

### Edited by:

João M. P. Jorge,  
Universidade Nova de Lisboa,  
Portugal

### Reviewed by:

Chonglong Wang,  
Soochow University, China  
Guang Zhao,  
Shandong University, China

### \*Correspondence:

Taek Soon Lee  
tslee@lbl.gov

### Specialty section:

This article was submitted to  
Microbiotechnology,  
a section of the journal  
Frontiers in Microbiology

**Received:** 07 October 2021

**Accepted:** 10 November 2021

**Published:** 02 December 2021

### Citation:

Carruthers DN and Lee TS (2021)  
Diversifying Isoprenoid Platforms via  
Atypical Carbon Substrates and  
Non-model Microorganisms.  
Front. Microbiol. 12:791089.  
doi: 10.3389/fmicb.2021.791089

Isoprenoid compounds are biologically ubiquitous, and their characteristic modularity has afforded products ranging from pharmaceuticals to biofuels. Isoprenoid production has been largely successful in *Escherichia coli* and *Saccharomyces cerevisiae* with metabolic engineering of the mevalonate (MVA) and methylerythritol phosphate (MEP) pathways coupled with the expression of heterologous terpene synthases. Yet conventional microbial chassis pose several major obstacles to successful commercialization including the affordability of sugar substrates at scale, precursor flux limitations, and intermediate feedback-inhibition. Now, recent studies have challenged typical isoprenoid paradigms by expanding the boundaries of terpene biosynthesis and using non-model organisms including those capable of metabolizing atypical C1 substrates. Conversely, investigations of non-model organisms have historically informed optimization in conventional microbes by tuning heterologous gene expression. Here, we review advances in isoprenoid biosynthesis with specific focus on the synergy between model and non-model organisms that may elevate the commercial viability of isoprenoid platforms by addressing the dichotomy between high titer production and inexpensive substrates.

**Keywords:** isoprenoids, metabolic engineering, synthetic biology, non-model organisms, C1 metabolism, terpenes

## INTRODUCTION

Isoprenoids are ubiquitous across all domains of life and span a wide and varied range of natural products. Isoprenoids are characterized by condensation of the five carbon precursor molecules isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP), which are typically generated through either the mevalonate (MVA) or methylerythritol phosphate (MEP) pathways. The ease with which specialized synthases and cytochromes can conjugate or decorate these precursors has led to a uniquely diverse class of chemicals. Estimates of natural isoprenoid compounds in the last several decades have steadily increased from 20,000 (Chappell, 1995) to over 70,000 (Moser and Pichler, 2019). The advent of advanced sequencing, -omics, and bioinformatics technologies coupled with protein structural software and flux balance analyses have facilitated a veritable revolution in synthetic biology and assured the continued elucidation of isoprenoid compounds through bioprospecting and biosynthetic efforts.

Isoprenoids serve a number of critical roles both as primary and secondary metabolites. Primary metabolites are essential to cell survival and propagation. They include carotenoids

that serve as auxiliary molecules for photoprotection and antioxidants (carotene, lycopene, lutein, and zeaxanthin) as well as sterols that help maintain membrane structure. Other isoprenoids function as components of dolichols, quinones, and essential proteins that aid in glycosylation and electron transport (Chappell, 1995). Secondary isoprenoid metabolites impart a non-essential benefit to cells usually by providing some defensive benefit or, in higher plants, hormone signaling. As for isoprenoids, these include pigments, fragrances, essential oils, and defensive chemicals that are most prominent in higher plants. Many secondary metabolites have attracted particular interest due to their applications as pharmaceuticals [e.g., artemisinin (Ro et al., 2006) and pacilitaxel (Biggs et al., 2016)], nutraceuticals, biofuels [e.g., isoprenol (Kang et al., 2019), prenol (Zheng et al., 2013), bisabolene, and limonene (Alonso-Gutierrez et al., 2013)], and cosmetics (Schempp et al., 2018). Hybrid technologies have capitalized on isoprenoid versatility through semi-synthetic approaches to generate elastomers (Della Monica and Kleij, 2020). Collectively, the bioproduction of these chemicals has enabled access to multibillion dollar chemical markets.

Microbial pathway engineering has proven especially successful in *Escherichia coli* and *Saccharomyces cerevisiae*, which have produced many of the aforementioned isoprenoid compounds. *E. coli* and *S. cerevisiae* maintain certain metabolic advantages including a fast growth phenotype, historical breadth of knowledge, ease of transformation and hence heterologous protein expression, substrate specificity, and published successes of bioproduction (Vickers et al., 2017; Ward et al., 2018). These advantages are complemented by specialized synthetic biology strategies that enable tuning of ribosome binding site and promoter strength, codon optimization of heterologous proteins, protein fusions, and the knocking out of competing pathways. In recent years, this has been accomplished by systematic gene downregulation using regulatable CRISPR interference systems (Kim et al., 2016; Tian et al., 2019) that express a modified dCas9 protein for fine-tuning of the overall pathway and optimization of target production. Furthermore, *E. coli* endogenously generates isoprenoids through the MEP pathway while *S. cerevisiae* utilizes its native MVA pathway, together enabling researchers to combine synthetic biological toolkits with the abundance of information of these strains to facilitate high-titer isoprenoid production. As for downstream isoprenoid functionalization, these metabolic chassis are genetically tractable whereas many natural isoprenoid production pathways are prevalent in recalcitrant organisms that make high-titer production infeasible. Only recently have certain non-model organisms been engineered to yield comparable or higher isoprenoid titers than in *E. coli* and *S. cerevisiae*.

Despite the clear successes of isoprenoid production, *E. coli* and *S. cerevisiae* have significant disadvantages that limit successful bioproduction at scale. Precursor limitations, either the availability of IPP and DMAPP for direct synthesis of isoprenoids or the availability of MVA/MEP precursors, have been identified as a major obstacle to advancing isoprenoid synthesis (Zu et al., 2020). Fine-tuning of metabolic pathways within the cell to balance cofactor supply by downregulation

or upregulation of select enzymes has been identified as a major engineering opportunity and, although generally successful, often involves strain-specific and product-guided strategies (Zu et al., 2020). Scaling of successful production is also limited by the necessity of episomal expression systems, which are ill-suited for industrial production due to the necessity of selective markers and their general instability.

A second major challenge in industrial isoprenoid biosynthesis is simultaneously increasing titer, rate, and yield while reducing the environmental and monetary burden intrinsic to industrial production. Precursor limitations are also complicated by isoprenoid production platforms that rely on sugar-based metabolism. Although sugars like glucose and glycerol provide high MVA/MEP flux by generation of G3P/pyruvate or acetyl-CoA, respectively, the high production costs are prohibitive to competition with petroleum-derived analogs. The greatest cost drivers of isoprenoid biosynthesis stem from carbon feed, which accounts for over 90% of production costs, and product yield (Wu and Maravelias, 2018).

A promising solution to exorbitant substrate costs is carbon source switching, especially to carbon waste streams like cellulosic biomass or C1 substrates (e.g., methane, methanol, carbon dioxide, and formate). Recent estimates assert that sugar switching from glucose to pretreated cellulosic biomass could yield a 53% decrease in cost (Wu and Maravelias, 2018) with further gains if organisms can simultaneously consume multiple substrates (e.g., hexose and pentose sugars). Growth and production on atypical carbon sources and native generation of secondary metabolites is prevalent among many microorganisms. Recent advances have capitalized on the diversity of microbial carbon assimilation pathways, especially in the elucidation of synthetic and natural C1 metabolic pathways that enable access to cheap, abundant carbon sources (Aldridge et al., 2021).

Many archaea have also evolved a robust array of resistance strategies to cope with inhibitory chemicals and conditions. These include tolerance mechanisms (efflux pumps, heat tolerance, membrane modifications, and general stress resistance; Dunlop, 2011) that facilitate extremophilic growth in inhospitable environments like anaerobic conditions or deep sea vents. These mechanisms in some cases directly involve secondary metabolite production and even enable enhanced tolerance to secondary metabolite toxicity (Dunlop et al., 2011). Toxicity tolerance is an appealing phenotype for biofuel production systems as well as for survival on substrates that are typically toxic to many microorganisms like pretreated lignocellulosic biomass (Dunlop et al., 2011). To date, bioproduction on pretreated cellulosic biomass has proven challenging due to the associated toxicity of the substrate, especially the prevalence of aromatic compounds. In response, researchers have begun focusing on resilient bioproduction chassis like *Pseudomonas putida* and *Rhodospiridium toruloides* that can readily degrade aromatic compounds (Yaegashi et al., 2017; Johnson et al., 2019).

Conversely, microbes with unique phenotypes tend to have limited metabolic toolkits available. Next generation sequencing technologies have expedited exploration and characterization of novel organisms from unique environments, yet direct

engineering of such organisms for production remains a fundamental challenge. Neither *E. coli* or *S. cerevisiae* naturally accumulate isoprenoids at high titer and bioproduction is often limited to heavily modified strains with inducible episomal expression systems. Even so, the highest production of isoprenoids has been achieved in *E. coli* and *S. cerevisiae* (Moser and Pichler, 2019). As a result, there is a significant disparity between model and non-model isoprenoid production.

Addressing the disparity between the prevalence of nature's clever solutions to environmental challenges and the genetic tractability of those organisms remains a principal obstacle in isoprenoid bioproduction. In many instances, it poses the question of whether to heterologously express pathways in common metabolic chassis or to optimize pathways *in situ*, both of which come with drawbacks. In this review, we highlight recent advances in core understanding of isoprenoid synthesis, namely the elucidation of the archaeal MVA pathways, precursor flux modulation, and how those discoveries have contributed to novel isoprenoid production schemes. We then explore the exchange between lessons learned in the metabolic engineering of *E. coli* and *S. cerevisiae* and of non-model microorganisms with natural predispositions for atypical and economical carbon substrates. Pathways include C1 metabolism in methylotrophic organisms (*Methanosarcina* sp., *Methylorubrum extorquens*) and phototrophic microbes (cyanobacteria, purple non-sulfur bacteria, diatoms, and green algae) capable of fixing CO<sub>2</sub>. We also explore advances in engineering of oleaginous yeast naturally capable of efficient lipid toleration and accumulation (*R. toruloides* and *Y. lipolytica*) and finally soil bacteria with special focus on their propensity for survival on and degradation of aromatic substrates (*B. subtilis* and *P. putida*). Collectively, these advances move isoprenoid biosynthesis toward economic and environmental feasibility.

## ADVANCES IN ISOPRENOID PATHWAY CONSTRUCTION:

All isoprenoids are generated from the common cellular precursors acetyl-CoA or glyceraldehyde 3-phosphate (G3P) and pyruvate *via* either the MVA or MEP pathway, respectively (Figure 1). These pathways share no homology and are evolutionarily distinct. Comparisons of the MVA and MEP pathway efficiencies, cofactors, and energetic requirements have been well documented in previous reviews (Dugar and Stephanopoulos, 2011; Yadav et al., 2012). The recent characterization of archaeal MVA pathways, shunts, and alternative precursors for these pathways have harbored the development of unique and more efficient routes for isoprenoid production (Kang et al., 2017; Hayakawa et al., 2018; Clomburg et al., 2019).

### Mevalonate Pathway

The MVA pathway is native to eukaryotes, some ancient and often predatory gram-positive bacteria (Pasternak et al., 2013), as well as, with some significant deviations, archaea (Boucher


et al., 2004). The canonical MVA pathway commences with a Claisen condensation of two acetyl-CoA thioester molecules followed by five sequential enzymatic steps that ultimately yield IPP. IPP is then converted to DMAPP by the isopentenyl diphosphate isomerase (IDI) for further condensation into isoprenoid compounds.

Over the last decade, the origin of the archaeal MVA pathway – either progenating from horizontal gene transfer or a cenancestor – has been hotly debated. However, recent analysis of monophylogenetic candidate phyla radiation and DPANN (Diapherotrites, Parvarcheota, Aenigmarchaeota, Nanoarchaeota, and Nanohaloarchaeota) have provided conclusive evidence to support an extant ancestral MVA in all domains of life (Castelle and Banfield, 2018). Most notably, the archaeal pathway lacks PMK, PMD, and IDI1. Instead, archaea have an alternative IDI2 similar in function to IDI1 and rely upon the recently discovered isopentenyl phosphate kinase (Dellas et al., 2013) to generate IPP through unique MVA intermediates. Specifically, three distinctive archaeal MVA pathways have been elucidated: the haloarchaea-type MVA, the thermoplasma-type MVA, and the archaeal MVA pathway that is conserved throughout the kingdom (Hayakawa et al., 2018; Thomas et al., 2019; Yoshida et al., 2020) as depicted in Figure 1.

Beyond the perplexities of phylogenetic classification (hereditary, horizontal gene transfer, etc), the elucidation of archaeal MVA pathways and their associated enzymes has proven instrumental in optimizing *S. cerevisiae*/*E. coli* production titers by capitalizing upon enzyme promiscuity or efficiency. Collectively, heterologous expression and fine-tuning of the MVA pathway to minimize flux bottlenecks has included the expression of genes across different domains. Overexpression of HMGS and HMGR from *Staphylococcus aureus* (Tsuruta et al., 2009) as well as a kinase from the archaeal *M. mazei* (Primak et al., 2011), for example, was successfully shown to improve C5 isoprenoid accumulation and laid the groundwork for longer chain isoprenoid production *via* the MVA pathway (George et al., 2015).

### Methylerythritol Phosphate Pathway

The MEP pathway is native to most Gram-negative bacteria and cyanobacteria as well as to algae and higher plants, but in the latter eukaryotes it is compartmentalized in the plastid. Despite being theoretically more efficient than the MVA pathway, the MEP pathway is more tightly regulated and challenging to engineer. Studies have elucidated rate limiting enzymatic steps in the MEP pathway, namely IDI and DXS, for  $\beta$ -carotene production (Yuan et al., 2006). However, overexpression of MEP pathway genes can also have deleterious effects on actual isoprenoid synthesis due to accumulation of intermediates. Our fundamental understanding of MEP pathway regulation is incomplete, encompassing some feedback and feedforward mechanisms between downstream isoprenoids and MEP intermediates (Bitok and Meyers, 2012). Studies in higher plants and algae, which have demonstrated that circadian light/dark cycling have a significant influence on pathway regulation, further complicate our understanding (Vranová et al., 2012).



**FIGURE 1** | A depiction of isoprenoid synthesis through the core 6 enzyme MVA and 7 enzyme MEP pathways. Also depicted are the newly discovered archaeal branches from the MVA pathway. The thermoarchaeal-type branch begins with mevalonic acid whereas the archaeal and haloarchaeal-type branches stem from MVAP. Typically, isoprenoids are synthesized by acetyl-CoA, pyruvate, and G3P, however IPP and DMAPP can also be synthesized from C5 alcohols, D-ribulose or D-ribulose-5-phosphate, and a synthetic route in which HMG-CoA is ultimately converted to prenol. AibAB, 3-methylglutaconyl-coenzyme decarboxylase; AtoB, acetyl-CoA acetyltransferase; BMD, bisphosphomevalonate decarboxylase; cbjALD, 3-methylcrotonyl-CoA reductase; DXR, 1-deoxy-D-xylulose 5-phosphate reductase; DXS, 1-deoxy-D-xylulose 5-phosphate synthase; HMGR, 3-hydroxy-3-methylglutaryl-CoA reductase; HMGS, 3-hydroxy-3-methylglutaryl-CoA synthase; IDI, isopentenyl diphosphate isomerase; IPK, isopentenyl phosphate kinase; IspD, 2-C-methyl-D-erythritol 4-phosphate cytidylyltransferase; IspE, 4-diphosphocytidyl-2-C-methyl-D-erythritol kinase; IspF, 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase; IspG, 4-hydroxy-3-methylbut-2-enyl diphosphate synthase; IspH, 4-hydroxy-3-methylbut-2-enyl diphosphate reductase; LiuC, 3-hydroxy-3-methylglutaryl CoA dehydratase; M3K, Mevalonate 3-kinase; M3P5K, Mevalonate 2-phosphate-kinase; nDXP, 1-deoxyxylulose-5-phosphate synthase; PMD, phosphomevalonate decarboxylase; PMK, phosphomevalonate kinase; tAHMP, anhydromevalonate diphosphate decarboxylase; ThiM, hydroxyethylthiazole kinase; XK, xylulose kinase; YahK, aldehyde reductase.

A recent metabolic control analysis employed -omics studies with recombineering to show that, normalized to DXS flux, IspG is the rate limiting step for isoprene synthesis with other enzymes increasing linearly with DXS concentration (Volke et al., 2019). This is an important finding as isoprene is the simplest hemiterpene and therefore a good reporter for MEP tuning. Yet production did not increase with overexpression of IspG and IspH, suggesting instead that other cofactors may be limiting (Volke et al., 2019). Indeed, careful balancing of IspG and IspH expression has shown enhanced  $\beta$ -carotene and  $\alpha$ -lycopene production (Li et al., 2017), suggesting that pathway tuning should be based on an intricate, product-driven approach (e.g., different tuning for isoprene vs. higher chain length terpenoids) rather than an intuitive, generalizable rule. In general, careful expression balancing has been the most successful strategy to MEP pathway optimization due to the complexity of regulatory mechanisms in *E. coli*, though even careful balancing in other organisms like the cyanobacterium *Synechococcus elongatus* has proven challenging (Englund et al., 2018).

## Synthetic Isoprenoid Production Pathways

Although essential to isoprenoid production, high IPP and DMAPP accumulation is toxic and can result in significant growth inhibition (George et al., 2018). This dilemma has led to a number of clever strategies for synthetic “growth-decoupled” and “bypass” isoprenoid production routes that comprise components of MVA/MEP pathways but avoid IPP/DMAPP accumulation. Many of these strategies have been informed or directly use elements from the recently elucidated archaeal MVA pathways, either through direct codon optimized expression or as templates for engineering promiscuous activity. A mevalonate decarboxylase from *Halobacterium volcanii*, for example, was expected to demonstrate conversion of MVAP to IP and employed as a template to rationally design PMDs for C5 alcohol production. The strategy successfully enhanced isoprenol production by bypassing intracellular IPP accumulation (Vannice et al., 2014; Kang et al., 2016). Further mutagenesis of a *S. cerevisiae* PMD in tandem with an endogenous phosphokinase resulted in an IPP-bypass pathway that yields IP and ultimately the highest isoprenol titer reported at 10.8 g/L (Kang et al., 2017, 2019).

Retro-biosynthetic approaches postulated that archaeal IPKs could enable phosphorylation of the C5 alcohols (isoprenol and prenol) into IPP and DMAPP, respectively. Direct feeding of alcohols for production of isoprenoid precursors could thereby decouple isoprenoid production from central carbon metabolism. Of particular interest were IPKs from *Halobacterium volcanii*, *Methanothermobacter thermautotrophicus*, *Thermoplasma acidophilum*, and *Methanocaldococcus jannaschii* (Chatzivasileiou et al., 2019). In one study, IPKs from the latter three archaea were screened for activity and cloned into an *E. coli* strain harboring a  $\beta$ -carotene production pathway (Liu et al., 2016). Expression of the IPK from *T. acidophilum* and feeding of 2 mM prenol

resulted in a 45% increase in  $\beta$ -carotene production and was further improved by site-specific mutagenesis to 97% (Liu et al., 2016). Growth-decoupled production of lycopene was also demonstrated by overexpressing a codon-optimized *T. acidophilum* IPK paired with an endogenous *E. coli* phosphatase PhoN, with titers nearing 190 mg/L in a mixture of 2.5 mM prenol and isoprenol (Chatzivasileiou et al., 2019; Clomburg et al., 2019; Lund et al., 2019). IPK-mediated production of carotenoid and neurosporene was also improved in *E. coli* by 18-fold and 45-fold, respectively, by decoupling terpene synthesis from central carbon metabolism through production on C5 alcohols (Rico et al., 2019).

Several other strategies utilize two upper MVA pathway genes (*E. coli* AtoB, *Staphylococcus aureus* HMGS) prior to diverging with the expression of the hydratase LiuC and *Myxococcus xanthus* decarboxylase AibAB (Clomburg et al., 2019; Eiben et al., 2020). From there, Eiben et al. demonstrated 80 mg/L isopentanol production through subsequent expression of *M. xanthus* AibC and *Clostridium acetobutylicum* AdhE2 (Eiben et al., 2020). In a more holistic approach, AibAB was followed with expression of the *Clostridium beijerinckii* acyl-CoA reductase (cbjALD), and *E. coli* YahK to generate prenol at the highest titer reported for biological production (Figure 1; Clomburg et al., 2019). Conversion of prenol to DMAP was then accomplished by the *E. coli* hydroxyethylthiazole kinase (ThiM) and finally converted to DMAPP by the *M. thermoautotrophicus* IPK (Clomburg et al., 2019). The unique approach employed by Clomburg et al. succinctly demonstrates how novel enzymes wrought by recent discoveries can be instrumental in designing pathways that circumvent metabolic bottlenecks to yield high tier production platforms.

To conclude, several studies have explored novel precursors to the MEP pathway using ribulose. The initial step in the MEP pathway, condensation of G3P and pyruvate with DXS, results in production of DXP with the loss of a CO<sub>2</sub> molecule or one sixth of total carbon (Kirby et al., 2015). A D-ribulose 5-phosphate shunt by nDXP was initially explored in *E. coli* by a semi-rational approach, which identified *yajO* and *ribB* gene mutants as candidate enzymes and improved carbon efficiency by direct conversion of C5 sugars to C5 MEP intermediates. Expression of the nDXP shunt enabled a 4-fold increase in MEP derived bisabolene production (Kirby et al., 2015). This approach was further demonstrated in *P. putida* by expression of the mutant *ribB* gene, but with low efficiency (Hernandez-Arranz et al., 2019). In an analogous work, promiscuous activity of fructose-6-phosphate aldolase in *E. coli* was used to generate D-ribulose from the glycolaldehyde and hydroxyacetone. Another DXP shunt overexpressed a native xylulose kinase (King et al., 2017). These novel shunts, like the archaeal informed MVA pathways, have the potential to alleviate precursor flux limitations. While the MVA pathway modifications have had clear success, it has yet to be determined whether these shunts can address the regulatory challenges associated with MEP derived isoprenoid production.

## ADVANCES IN ISOPRENOID FUNCTIONALIZATION

The C5 precursors IPP and DMAPP are dephosphorylated, cyclized, and modified to create a structurally diverse group of over 70,000 chemicals through a coordinated enzyme network (Kirby and Keasling, 2009; Moser and Pichler, 2019). The first stage or module of isoprenoid biosynthesis is characterized by the successive addition of the diphosphate precursor *via* head-to-head or tail-to-head condensation. The second module is an operation or series of operations conducted by terpene synthases (TSs) in which the terpenoid skeleton is dephosphorylated and cyclized. The third module involves further decoration by cytochrome P450s (CYPs), acetyltransferases, methyltransferases, dehydrogenases, and in some cases, glycosylations. This overall framework is consistently repeated in nature with some variations (Zhou and Pichersky, 2020). In this section we discuss recent advances in the functionalization of isoprenoids. Broad ranges of chemical production targets have been demonstrated and scaled from biofuels to pharmaceuticals by heterologous expression of prenyl diphosphate synthases, TSs, and CYPs.

### Cytochrome P450s

Heme-thiolate monooxygenases or CYPs are an interesting class of enzymes that functionalize terpenes through oxygenation reactions (hydroxylation, dealkylation, demethylation, decarboxylation, cyclization, C-C bond cleavage, among others) and present an important opportunity for generating highly decorated natural products. To date, over 300,000 CYPs have been discovered, with less than 1% actually characterized (Li et al., 2020; Liu et al., 2020). There is particular interest to produce CYP-derived terpenoids in microbial chassis due to the high barriers of slow growth and costly deconstruction inherent to native plant extraction.

Engineering CYPs has significant implications for novel and unnatural bioproducts (Helfrich et al., 2019; Xiao et al., 2019). The range of oxygenated terpenes is complemented by the sheer expanse of CYP availability in plants (Zhou and Pichersky, 2020). For example, CYPs are critical for the production of bioactive molecules with high pharmacological impacts. Case studies of microbial expression include production of precursor intermediates to artemisinin (CYP71AV1, aaCPR) and taxadiene (CYP725A4, tcCPR), which are natively produced by wormwood (*Artemisia annua*) and the Pacific yew tree (*Taxus brevifolia*) and were heterologously expressed in *E. coli* and *S. cerevisiae*, respectively. Indeed, the diversity and complexity of plant TSs, presented elsewhere (Karunanithi and Zerbe, 2019) offer tremendous potential as candidates for microbial production.

These specific examples demonstrate successful engineering of CYPs for pharmaceutical production, however functional plant CYP expression in microbes has proved challenging. Unfortunately, *E. coli* cannot naturally perform most posttranslational modifications and expression of membrane bound proteins like CYPs generates inclusion bodies or

aggregates of insoluble proteins. The production of oxygenated taxanes in *E. coli* (Biggs et al., 2016), for example, required extensive engineering of the CYP redox-partner cytochrome P450 reductase (CPR) pairings, N-terminal modifications for better solubility, and had significant repercussions on upstream MEP pathway balance. These pairings and modifications are necessary for any heterologous CYP expression in *E. coli* and vary depending on the selected proteins. Remarkably, although yeasts are naturally capable of many posttranslational modifications, express native CPRs, and require less N-terminal modification, meta-analyses have shown that *E. coli* studies tend to have higher yield CYPs than *S. cerevisiae* despite the necessity of many more genetic modifications (Hausjell et al., 2018). Many CYP-reductase pairings have been explored in *E. coli* and *S. cerevisiae* as listed in **Table 1**.

In recent years, microbial expression of CYPs has produced many variable length terpenoids (**Figure 2**). Of special significance are the monoterpene perillyl alcohol, an anti-cancer drug, from limonene (Alonso-Gutierrez et al., 2013) and the sesquiterpenes nootkatone, a pharmaceutical, from valencene (Guo et al., 2018) and zerumbone, an antioxidant, from  $\alpha$ -humulene (Zhang et al., 2018). Of further interest are the pharmacologically relevant diterpenes taxadiene and oxygenated casbenes as well as triterpenoids glycerrhetol from 11-oxo-b-amyrin and betulinic acid from lupeol.

There have been many breakthroughs in CYP-derived terpenes in the last 5 years that may provide guidance in the engineering of other CYP expression systems. Their interest has led to toolkits for enhanced CYP selection for targeted product engineering to streamline oxyfunctionalization of terpenes (Hernandez-Ortega et al., 2018). Challenges remain with regards to CYP promiscuity, enantiomeric purity, and, perhaps most importantly, production titers. Hopefully, such tools will guide hypotheses and penetrate into the library of known but currently inaccessible plant bioactive terpenes for use in high titer therapeutic production.

### Atypical Terpenoid Production

Several studies have investigated novel isoprenoid chain lengths through the expression of unique S-adenosyl-methionine (SAM)-dependent methyltransferases that methylate the fundamental IPP/DMAPP building blocks and thereby break the natural C5 dogma. In a pioneering study, a GPP methyltransferase (2meGPP) from the cyanobacterium *Pseudanabaena limnetica* was expressed in *S. cerevisiae* with its native 2-methylisoborneol synthase and then with seven distinct plant monoterpene synthases (Ignea et al., 2018). Each synthase generated a unique fingerprint of novel C11 compounds. By demonstrating a range of C11 targets, the authors provided a proof of concept for future enzyme optimization strategies toward specific C11 targets. In another work, IPP SAM-dependent methyltransferases enable conversion of IPP to C6 and C7 prenyl diphosphates with a methyltransferase from *Streptomyces monomycini*, which could then generate C11, C16, and C17 terpenes as well

**TABLE 1** | CYP expression, reductase pairing, and production of oxygenated terpenoids in various microbial hosts.

CYP	Reductase pair	Source organism(s)	Expression organism	Precursor	Product	Titer	Ref.
CYP71AV1	aaCPR	<i>Artemisia annua</i>	<i>S. cerevisiae</i>	Amorphadiene	Artemisinic acid	100mg/L	Ro et al., 2006
CYP71AV1	aaCPR	<i>A. annua</i>	<i>E. coli</i>	Amorphadiene	Artemisinic acid	5.8mg/L	Chang et al., 2007
CYP706B1	ctCPR	<i>Candida tropicalis</i>	<i>E. coli</i>	Cadinene	8-hydroxycadinene	105mg/L	
CYP71BA1	atCPR	<i>Zingiber zerumbet</i> , <i>A. thaliana</i>	<i>E. coli</i>	$\alpha$ -humulene	8-hydroxy-a-humulene	2.972 $\mu$ g/L	Yu et al., 2011
CYP76AH1	atCPR1	<i>Salvia miltiorrhiza</i> , <i>A. thaliana</i>	<i>S. cerevisiae</i>	Miltiradiene	Ferruginol	10.5mg/L	Guo et al., 2013
CYP153A6	mCPR	<i>Mycobacterium sp.</i>	<i>E. coli</i>	Limonene	Perillyl alcohol	100mg/L	Alonso-Gutierrez et al., 2013
HPO	atCPR	<i>Hyoscyamus muticus</i> , <i>P. pastoris</i> <i>A. thaliana</i>		(+)-valencene	(+)-nootkanone	208mg/L	Wriessnegger et al., 2014
CYP725A4	tcCPR	<i>Taxus cuspidata</i>	<i>E. coli</i>	Paclitaxel	Oxygenated taxanes	570mg/L	Biggs et al., 2016
CYP726A20	jcCPR1	<i>Jatropha curcas</i>	<i>S. cerevisiae</i>	Casbene	Oxidized casbanes	~1g/L	Wong et al., 2018
CYP716A12	atCPR	<i>Callitropsis nootkatensis</i> , <i>A. thaliana</i>	<i>Y. lipolytica</i>	(+)-valencene	(+)-nootkanone	978.2 $\mu$ g/L	Guo et al., 2018
CYP71BA1	atCPR	<i>Z. zerumbet</i> ; <i>A. thaliana</i>	<i>S. cerevisiae</i>	a-humulene	A-humulene 8-hydroxylase; zerumbone	40mg/L	Zhang et al., 2018
CYP716A12	atCPR	<i>Medicago truncatula</i> , <i>A. thaliana</i>	<i>Y. lipolytica</i>	Lupeol	Betulinic acid	26.53mg/L	Sun et al., 2018
CYP716A47	pgCPR1	<i>Panax ginseng</i> , <i>A. thaliana</i>	<i>S. cerevisiae</i>	Dammarenediol II	Protopanaxidiol	11.02g/L	Wang et al., 2019
CYP716A12	mtCPR	<i>M. truncatula</i>	<i>Phaeodactylum tricornutum</i>	Lupeol	Betulinic acid	0.1mg/L	D'Adamo et al., 2019
BcABA1	bcCPR1	<i>Botrytis cinerea</i>	<i>S. cerevisiae</i>	FPP	Abscisic acid	4.7mg/L	Otto et al., 2019
BcABA2							
CYP72A63	atCPR1, mtCPR2, mtCPR3, guCPR2	<i>A. thaliana</i> , <i>M. truncatula</i> , <i>Glycyrrhiza uralensis</i>	<i>S. cerevisiae</i>	11-oxo-b-amyrin	Glycyrrhetol	31.8mg/L	Sun et al., 2020

Culture conditions, scale, and medium vary significantly. HPO, *hyoscyamus muticus prenaspirodiene oxygenase*; CPR, *cytochrome P450 reductase*.


as zeaxanthin-like C41, C42, and C43 compounds depending on methylation (Drummond et al., 2019). A product-driven approach was also able to elucidate a C-methyltransferase that generated sodoferin, an atypical C16 sesquiterpene (von Reuss et al., 2018). The work demonstrated that sodoferin, which is produced naturally and perhaps exclusively by a *Serratia plymuthica*, could be produced in *E. coli* through heterologous expression of the methyltransferase and TS (von Reuss et al., 2018). A final, completely divergent approach utilizes a thiolase from lepidoptera (butterflies and moths) that naturally produces juvenile hormones in the form of C16 methylated diterpenoids (Eiben et al., 2019). Specifically, the thiolase PhaA condenses a propionyl-CoA with an acetyl-CoA as opposed to the standard AtoB of the MVA pathway, which condenses two acetyl-CoA substrates.

It is probable that the control of specific methylation sites decreases with compound size such that targeting specific triterpenoids would remain an obstacle. In the case of C11 targets, site-directed mutagenesis of the monoterpene synthases did enable higher selectivity (Ignea et al., 2018), which is encouraging for future engineering. These unique approaches have expanded the boundaries of isoprenoid synthesis well beyond the C5 rule, though admittedly practical applications of these novel compounds have yet to be realized.

## Meroterpenoids

Partial isoprenoids or meroterpenoids are a class of compounds containing an isoprenoid chain paired with another structure and may have beneficial bioactive properties. Broadly, meroterpenoids include cytokinins, quinones, steroids, and porphyrins like heme A and chlorophyll a. The optimization of heterologous meroterpenoids poses a unique engineering challenge as the isoprenoid compound must be cogenerated with another structure, then converted to the terpenoid by a specified synthase.

A good case study is the production of cannabinoids in *S. cerevisiae*. Cannabinoids are of commercial interest but, like many natural products, suffer from low *in planta* yields. In a recent publication, the production of olivetolic acid from acetyl-CoA was engineered using a six gene pathway (Luo et al., 2019). Prenylation of olivetolic acid using a *Cannabis sativa* prenyltransferase (csPT4) and further heterologous synthases led to *in vivo* production of cannabigerolic acid,  $\Delta$ 9-tetrahydrocannabinolic acid, cannabidiolic acid,  $\Delta$ 9-tetrahydrocannabivarinic acid, and cannabidivarinic acid (Luo et al., 2019). The work not only presents a novel production scheme in *S. cerevisiae* but demonstrates the ease with which transgenic elements can be translated into production chassis.



Another relevant example is the production of prenylated flavonoids, which are derived from cyclic amino acid biosynthesis and can serve as nutraceuticals and medicines. They, again, are found in low natural abundance in plant species like *Sophora flavescens*, a shrub and *Humulus lupulus*, brewing hops (Yang et al., 2015). Production of naringenin in *S. cerevisiae* coupled with overexpression of a plant flavonoid prenyltransferase enabled production of the pharmaceutically relevant 8-prenylnaringenin (Levisson et al., 2019). Although both strategies were limited to yeast, they underline the flexibility of engineering

meroterpenoid production in microbes to address commercial needs and provide a promising opportunity for accessing low abundance natural products.

Unique meroterpenoids are also generated in high natural abundance in certain microbes. Archaea differ from bacteria primarily in membrane composition. Archaea produce isoprenoid-derived glycerol lipid ethers (namely ester linked sn-glycerol 3-phosphates rather than ether linked sn-glycerol 1-phosphates) that facilitate growth in unique environments. Full reduction of these long length C20/C40 membrane isoprenoid

chains is accomplished by downstream geranylgeranyl reductases (GGR). While they provide an evolutionary advantage for survival in extreme conditions, membrane isoprenoids may also be utilized to generate unsaturated chemicals of interest (Jain et al., 2014). Archaeal lipids, namely archaeol and caldarchaeol, have been identified as potentially valuable for the formation of archaeosomes. Archaeosomes are lipid vesicles composed of archaeal derived lipids and have shown higher physicochemical stability than liposomes, a conventional drug delivery system. As a result, archaeosomes have been singled out as a possible adjuvant and could prove particularly valuable in slow release drug delivery systems (Caforio and Driessen, 2017).

## INDUSTRIAL PRODUCTION FROM C1 CHEMICAL FEEDSTOCKS

C1 substrates are typically generated as industrial and petrochemical byproducts and, in general, C1 substrates are stable, abundant, and inexpensive. Advances in sequestration and hydrogenation of atmospheric CO<sub>2</sub> via heterogeneous catalysts have enabled the conversion of emissions into valuable C<sub>2+</sub> substrates (Ye et al., 2019). Metabolic engineering of organisms capable of C1 growth is an enticing opportunity for achieving cost parity with petrochemical products while simultaneously improving sustainability metrics. C1 metabolism may be subdivided into phototrophic, methylotrophic, or formatotrophic microbes that consume CO<sub>2</sub>, methane/methanol, and formate/formic acid, respectively. Here, we describe recent approaches for converting C1 substrates into isoprenoid precursors with specific attention to works demonstrating isoprenoid production. The generalized pathways for C1 metabolism are illustrated in **Figure 3**.

### Methylotrophic Isoprenoid Production


Methane is an abundant byproduct of many chemical processes including fracking and petroleum drilling (Clomburg et al., 2017). In terms of greenhouse gas emissions, methane is approximately 20 times more potent than CO<sub>2</sub> and excess capacity is flared at refineries, leading to an increase in direct CO<sub>2</sub> emissions and a loss of revenue (Conrado and Gonzalez, 2014). Methane is also generated through anaerobic digestion of waste biomass. To date, many successful chemical platforms utilize methane as a feedstock to generate alcohols, carboxylic acids, as well as other common C<sub>2</sub>/C<sub>3</sub> chemicals (Kuhl et al., 2012). While biological conversion rates tend to be lower, certain methylotrophic organisms have arisen as potential candidates to capitalize upon methane/methanol availability for more complex bioproduction. In this section, we discuss recent advances in isoprenoid biosynthesis in methylotrophic cell factories across domains.

Archaea are well regarded for their ability to thrive in nutrient-limited anaerobic and extreme conditions. As a result, many archaea have evolved highly efficient strategies for C1 assimilation. On one hand this has helped elucidate more

efficient MVA pathways as previously described, but on the other it makes the engineering of tightly regulated archaeal pathways that are geared toward energy conservation thermodynamically challenging. Nonetheless, one study demonstrated production of isoprene from methanol in *Methanosarcina acetivorans* and *Methanosarcina barkeri* under anaerobic conditions and showed a redirection of electron flux from membrane precursors in favor of isoprene accumulation (Aldridge et al., 2021). The diverted isoprene accounted for 4% of total carbon flux.

More substantial success has been achieved in methylotrophic bacteria, many of which thrive in more mesophilic conditions. Methylotrophic bacteria are predominantly divided into two types: Type I assimilates formaldehyde using the RuMP cycle and Type II assimilates formaldehyde via the serine cycle. *Methylorubrum extorquens* AM1 (formerly *Methylobacterium extorquens* AM1), a Type II methylotroph, has been studied for over 60 years such that many genetic tools are available (-omics data, metabolic networks, genome-scale model; Schrader et al., 2009). *M. extorquens* fermentations have produced methanol-derived products ranging from 1-butanol (Hu and Lidstrom, 2014) to polymers (Orita et al., 2014). A series of stepwise optimizations in *M. extorquens* AM1 included heterologous expression of the *M. xanthus* MVA genes, an FPP synthase from *S. cerevisiae*,  $\alpha$ -humulene synthase from *Zingiber zerumbet*, and reduced carotenoid flux. Combinedly, these modifications resulted in the accumulation of 1.65 g/L  $\alpha$ -humulene on methanol in fed-batch cultivation, which stands as the highest titer reported (Sonntag et al., 2015). Other works have explored high titer production of the MVA pathway intermediates, including 2.59 g/L mevalonic acid from methanol on minimal media using a mevalonate biosensor strategy (Liang et al., 2017), which suggests that high titer production of other isoprenoids is also possible. As another attractive feature, *M. extorquens* harbors the ethylmalonyl-CoA pathway (EMCP) that includes a series of anaerobic activated CoA esters useful for pathway remodeling (Schrader et al., 2009; Schada von Borzyskowski et al., 2018). Interpathway metabolite exchange has risen as a major avenue for further C1 metabolism, especially for formatotrophs as discussed later.

Recently, the methylotrophic yeast *Pichia pastoris* (formerly known as *Komagataella phaffii*) has arisen as a promising candidate for isoprenoid production on methanol. *P. pastoris* maintains several unique characteristics including a tightly regulated and highly expressed alcohol oxidase AOX1, which catalyzes oxidation of methanol to formaldehyde. AOX1 is strongly induced by methanol but repressed by glucose and glycerol (Hartner and Glieder, 2006). As a result, fed-batch production schemes have been designed to partition high cell density growth on glycerol/glucose and production on methanol. This strategy has successfully demonstrated the production of 714 mg/L lycopene (Zhang et al., 2020), (+)-ambrein, squalene (Moser et al., 2018), and 208 mg/L (+)-nootkatone (Wriessnegger et al., 2014). Interestingly, the latter study successfully used an approach for CYP production that had failed in *S. cerevisiae*. More involved engineering of *P. pastoris* has demonstrated *de novo* production using heterologously expressed



Calvin-Benson-Bassham (CBB) Cycle enzymes along with native genes in the xylulose monophosphate (XuMP) cycle and deletion of certain fatty acid enzymes, ultimately yielding a mutant exhibiting autotrophic growth on CO<sub>2</sub> (Gassler et al., 2020).

Methylotrophic production has garnered special interest for reducing bioproduction costs either through valorization of commercial waste streams or CO<sub>2</sub> conversion. **Table 2** provides a list of methylotrophic production strains, their respective C1 substrates, and product titers. In most cases, titers are

significantly lower than comparative production in *E. coli* or *S. cerevisiae* and methylotrophic cultures require significant supplementation with a rich medium that may somewhat reduce the benefits of C1 production.

Past attempts at methylotrophic production have historically been hindered by low carbon and energy efficiencies and addressing these issues by leveraging RuMP/Serine cycles with the CBB cycle have been long postulated (Conrado and Gonzalez, 2014). Recently, major breakthroughs in *P. pastoris* demonstrated

**TABLE 2** | Production of isoprenoids by methylotrophic organisms.

Organism	Feedstock	Product	Titer	Ref.
<i>Methylomonas</i> sp. 16a	Methane	astaxanthin	2.4 mg/g CDW	Ye et al., 2007
<i>P. pastoris</i>	Methanol**	(+)-nootkanone	208 mg/L	Wriessnegger et al., 2014
<i>P. pastoris</i>	Methanol**	(+)-valencene	136 mg/L	
<i>M. extorquens</i>	Methanol	a-humulene	1.65 g/L	Sonntag et al., 2015
<i>P. pastoris</i>	Methanol**	(+)-ambrein	105 mg/L	Moser et al., 2018
<i>P. pastoris</i>	Methanol**	squalene	58 mg/L	
<i>P. pastoris</i>	Methanol**	lycopene	714 mg/L	Zhang et al., 2020
<i>M. alcaliphilum</i>	Methane	$\alpha$ -humulene	0.75 mg/g CDW	Nguyen et al., 2020
<i>M. acetivorans</i>	Methanol*	isoprene	0.954 mM/L	Aldridge et al., 2021
<i>M. barkeri</i>	Methanol*	isoprene	36.0 $\mu$ M/L	

\*Complex medium; \*\*Carbon sources switched from glycerol to methanol in fed-batch fermentation.

*de novo* production using heterologously expressed CBB cycle enzymes, overexpressed native genes in the XuMP cycle, and deletions of certain fatty acid enzymes. The mutant was ultimately capable of autotrophic growth on CO<sub>2</sub> (Gassler et al., 2020). Likewise, artificial methanotrophy and formatotrophy have been explored in a complex rewiring of *E. coli* (Bennett et al., 2018; Chen et al., 2020; Kim et al., 2020). These seminal works are excellent examples of how systems biology can be applied to tune precursors, adapt strains, and incorporate well-defined isoprenoid pathways for higher production efficiency at lower substrate costs.

## Phototrophic Isoprenoid Production

Phototrophic growth is characterized by the photosynthetic conversion of CO<sub>2</sub> to sugars *via* complex photoreductive reactions and the Calvin-Benson-Bassham cycle. Photosynthetic organisms naturally produce carotenoids in high concentrations to cope with excess intracellular reactive oxygen species. Specifically, lutein is used in non-photochemical quenching of chlorophyll triplets during photosynthesis (Dall'Osto et al., 2006), zeaxanthin for heat dissipation and photoprotection, while carotene and chlorophyll absorb light. In many cases, gains from the engineering of the genetic architecture of photosynthesis (e.g., light harvesting complexes, RuBisCo) have been limited. Nonetheless, cyanobacteria, which only exhibit the MEP pathway, have become major targets for metabolic engineering due to their genetic plasticity and malleability with respect to isoprenoid precursors, especially through carbon sinks.

Engineered carbon sinks operate on the hypothesis that carbon fixation reactions are faster than downstream carbon utilizing growth reactions such that the accumulation of intracellular carbon metabolites reduces NADPH consumption and ultimately inhibits photosynthesis (Oliver and Atsumi, 2015). Introduction of exogenous genes demonstrated a 1.8-fold increase in carbon yield for the generation of 2,3-butanediol

(Oliver and Atsumi, 2015). Others have shown sucrose, ethylene, and isobutyraldehyde production all while enhancing photosynthetic activity through more optimal use of the electron transfer chain (Ducat et al., 2012; Santos-Merino et al., 2021). This effect was also found to be additive when multiple sinks were introduced, suggesting that “sink engineering” could be conceptually applied to secondary metabolite synthesis through downstream modifications of the MEP pathway (Santos-Merino et al., 2021). Several works have capitalized on this upregulation of photosynthesis by combining this source/sink approach with computationally informed modification of limonene synthase, resulting in a 100-fold production improvement in limonene production (Wang et al., 2016) and potential applications for other isoprenoids.

Cyanobacterial studies have also made improvements through direct modification of native isoprenoid pathway genes in combination with a product-specific terminal synthase. *Synechococcus elongatus* and *Synechocystis* sp. PCC 6803 have been primary targets for production with recent attempts focusing on generation of isoprene, with a hallmark study demonstrating 1.26 g/L production (Gao et al., 2016; Yang et al., 2016) albeit over several weeks. This feat was accomplished through the overexpression of MEP pathway enzymes though, more importantly, by bioprospecting for a more efficient isoprene synthase. Comprehensive analyses of the MEP pathway metabolic bottlenecks in *S. elongatus* have also been studied by a systematic investigation of each enzymatic step in the MEP pathway, specifically using isoprene as a simple reporter for MEP flux (Englund et al., 2018). The work found that the regulatory circuitry of the *S. elongatus* MEP pathway is, like that of many other MEP pathway harboring organisms, complex and that a simple overexpression of select pathway genes does not necessarily equate to higher/lower production. Despite this complexity, products including squalene (Pattanaik et al., 2020), bisabolene (Sebesta and Peebles, 2020; Rodrigues and Lindberg, 2021), and  $\alpha$ -farnesene (Lee et al., 2017) have been produced in *S. elongatus* through some combination of *idi*, *dxs*, *ispA*, and terminal synthase overexpression. It is possible that cyanobacteria could benefit from acetyl-CoA/pyruvate precursor rebalancing. In one study, overexpression of a pyruvate dehydrogenase increased the pool of available acetyl-CoA for isopropanol production (Hirokawa et al., 2020) and could, in theory, be applied to facilitate a heterologously expressed MVA pathway or, in the reverse, to enhance pyruvate accumulation.

Unlike cyanobacteria, eukaryotic algae maintain both the MVA pathway, located in the cytosol, and the MEP pathway, which is sequestered to the chloroplast in proximity to CO<sub>2</sub>-derived metabolites from photosynthesis. Algae have been hailed as candidate bioproduction microbes for many years due to their propagation in many media and thus potential for growth in wastewater streams like agricultural runoff rich in phosphorus and nitrogen. In general, however, eukaryotic algae are notably more challenging to engineer due to their comparably smaller metabolic toolkits and robust regulatory mechanisms on metabolic flux. Studies have established their propensity for some algal isoprenoid production in low titers including patchoulol (Lauersen et al., 2016), bisabolene (Wichmann et al.,

2018), and mixed diterpenoids (Lauersen et al., 2018) in the modal alga *Chlamydomonas reinhardtii* with CO<sub>2</sub> as the sole carbon source. Another alga, *Dunaliella salina*, has been singled out due to its resilience to highly saline environments that serve as a natural antibiotic against contaminants like protozoa, bacteria, dinoflagellates, and other algae. *D. salina* also naturally accumulates β-carotene under abiotic stress and remains one of the few commercially exploited green algae (Borowitzka, 2013; Fachet et al., 2020) along with *Haematococcus pluvialis* for astaxanthin production. Lastly, *Botryococcus braunii*, a colonial green alga, is rich in isoprenoid derived lipids that consist of 35% dry cell weight (DCW) biomass. The isoprenoids generated are characterized by race and consist of either Botryococcones (C30–C37), methylated squalenes (C31–C34), or odd-number n-alkadienes or trienes (C23–C33; Metzger and Largeau, 2005). Despite their unique composition, broad attempts to culture and optimize isoprenoid production have been limited in part due to slow growth comparative to other green algae (Morales-Sánchez et al., 2017). Somewhat remarkably, both commercial successes stem from unmodified organisms that simply generate isoprenoids under abiotic stress conditions.

Diatoms are a unique subset of algae with a characteristic cell-wall composed of silica. Certain diatoms are capable of generating highly branched isoprenoids (HBIs) like trienes, tetraenes, and pentaenes intrinsic to some diatoms with potential for pharmaceutical or biofuel usage (Athanasakoglou et al., 2019), possibly generated by promiscuous activity of diatom specific farnesyl pyrophosphate synthases (Ferriols et al., 2015). A specific diatom, *Haslea ostrearia* maintains a plastidal MEP cycle with a cytosolic MVA pathway and has demonstrated significant crosstalk between these localized elements, suggesting complex regulatory mechanisms perhaps in response to external stimuli and pose potential opportunities to tune both pathways for downstream C5 precursors depending on target terpenoids.

A final distinctive group of phototrophic organisms are purple non-sulfur bacteria, which are identified by a unique color that stems from a combination of pigmented carotenoids. In particular, *Rhodobacter sphaeroides* is a well-established isoprenoid producer, with industrial production of sesquiterpenes valencene and nootkanone demonstrated by BASF (Beekwilder et al., 2014; Schempp et al., 2018). Like many bacteria, *R. sphaeroides* accumulates polyhydroxybutyrate (PHB), a biopolymer with industrial bioplastic applications in of itself, under nitrogen limited conditions. Elimination of the PHB biosynthetic pathway (*phaC1*, *phaC2*) and expression of the heterologous MVA pathway contributed to increased flux through the isoprenoid pathway under nitrogen limited conditions (Orsi et al., 2020b).

As a whole, photosynthetic organisms remain tantalizingly elusive for high titer heterologous isoprenoid production despite advances in “sink engineering” and successes in the production of certain short chain biofuels.

## Formatrophic Production Pathways

Formate remains an enticing C1 substrate due to the relative ease with which it may be generated. Proposed strategies include

the hydration of syngas, the hydrogenation of CO<sub>2</sub>, and electrochemical reduction of CO<sub>2</sub> using, preferably, renewable generated electricity (Yishai et al., 2016). Bioproduction on formate remains challenging, though recent works have attempted to address this challenge by mapping natural pathways within the context of microbial metabolism (Bar-Even, 2016). The intrinsic nature of formate as an intermediate and availability of natural formate assimilation pathways like the serine, reductive acetyl-CoA, RuMP, XuMP, and reductive glycine pathways have led to the proposal of many synthetic pathways that could theoretically outperform their natural counterparts (Bar-Even, 2016). This hypothesis was encouraged by a previous study that determined formate, not formaldehyde, was the major branch point in *M. extorquens* methylotrophy (Crowther et al., 2008). In particular, this suggested that direct feeding of formate could be energetically beneficial due to the affiliated reduction of NAD<sup>+</sup> in aldehyde dehydrogenase thereby further supporting formatrophic pathways (Crowther et al., 2008).

Acting on this hypothesis, *M. extorquens* genes encoding formate-THF ligase, methenyl-THF cyclohydrolase, and methylene-THF dehydrogenase were heterologously expressed in *E. coli* to enable growth on formate through the serine cycle. In combination with downstream modifications, the strain was capable of 90 mg/L ethanol production on sugar-free formate minimal medium by adaptive laboratory evolution (ALE; Kim et al., 2019). In a subsequent study, expression of the reductive glycine pathway (rGlyP) in *E. coli* enabled growth on methanol and formate (Kim et al., 2020). Despite clear demonstration of formate-based growth here and *M. extorquens* isoprenoid production on methanol above, few formate derived isoprenoid compounds have been shown. A single exception was a study of the archaea *Methanococcus maripaludis*, which is capable of growth on H<sub>2</sub>, CO<sub>2</sub>, formate, and acetate as substrates under strict anaerobic conditions. Heterologous expression of a geraniol synthase enabled production of 4.0 mg/g and 2.8 mg/g geraniol on H<sub>2</sub>/CO<sub>2</sub> and formate feeds, respectively (Lyu et al., 2016). Although meager, this represents a baseline for further isoprenoid production and, with the addition of the groundbreaking production of formatrophic *E. coli* works, likely represents the first of many formate-based production strains.

So far, we have described a number of routes for isoprenoid production on C1 substrates, including several instances in which whole pathways have been translated between organisms. Life cycle assessment (LCA) and techno-economic analysis (TEA) will both be critical in quantifying the relative process level sustainability and monetary impacts, validating whether modified microbes are competitive with conventional production on glucose or from petroleum, and prioritizing future optimization opportunities based on projected impact gains. Growth and production on C1 substrates are inherently more sustainable than on pure sugar substrates, however the sustainability of the entire process from cradle-to-gate will be contingent on nontrivial improvement of production titer, rate, and yield. While LCAs and TEAs are common in CO<sub>2</sub>-derived biofuel production, they remain uncommon for all other C1 substrates. Indeed, the first LCA/TEA of a methane-derived bioproduct was only recently published (Fei et al., 2020). Nonetheless this

initial study provides a baseline for future valorization of other C1-derived chemicals and, hopefully, represents a first effort to quantify the economic and sustainability advantages of C1 substrates.

## ISOPRENOID PRODUCTION ON LIGNOCELLULOSIC CARBON SOURCES

Certain microbes are capable of valorizing more complex waste streams due to unique evolutionary predispositions. Here, we describe two strains of oleaginous yeasts, *Yarrowia lipolytica* and *Rhodospiridium toruloides*, capable of high titer isoprenoid production from woody biomass and waste cooking oil (WCO). Lastly, we describe two prototypical isoprenoid production platforms: *P. putida*, which is a prime candidate for conversion of pretreated lignocellulosic biomass, and *B. subtilis*, a candidate bacteria renown for high titer protein production.

### *Yarrowia lipolytica*

The oleaginous yeast *Yarrowia lipolytica* can naturally assimilate many atypical carbon sources including glycerol, organic acids, succinate, citrate, and even WCO. Likewise, *Y. lipolytica* is of keen interest due to its natural accumulation of  $\beta$ -carotene, farnesene, and linalool. Multi-copy pathway integration has proven especially successful in targeted isoprenoid overproduction (Xie et al., 2015). A recent work applied a random chromosomal integration approach of multiple MVA pathway operons, cofactor modulation, and culture condition tuning produced 25.55 g/L  $\alpha$ -farnesene on YPD complex medium over 20 days in fed-batch production with significant byproduct formation (Liu et al., 2019). A similar strategy led to 6.5 g/L production of  $\beta$ -carotene by chromosomal integration of multiple copies of CarB, CarRP, and GGPPS in fed-batch production with over 40 g/L lipid byproduct (Larroude et al., 2018). Other reports of note include high squalene production at titers of 531.6 mg/L (Gao et al., 2017) and 402.4 mg/L (Arnesen et al., 2020). Building upon previous limonene demonstrations with neryl diphosphate synthase (tNPPS1) from *Agastache rugosa* (Korean mint) and limonene synthase from *Solanum lycopersicum* (tomato; Cao et al., 2016), *Y. lipolytica* ultimately yielded 165.3 mg/L limonene on glycerol/citrate (Cheng et al., 2019). More comprehensive descriptions of *Y. lipolytica* regulatory changes for production have also been published (Arnesen et al., 2020).

*Y. lipolytica* is also capable of converting fatty acids into C2 substrates through the beta-oxidation pathway and has high native lipid tolerance. Recent works have demonstrated high lipid production of modified *Y. lipolytica* on pretreated lignocellulosic biomass (0.11 g lipids/g sugars), even approaching efficiencies observed on glucose (Yook et al., 2020). In fact, *Y. lipolytica* has shown up to 90% DCW lipid accumulation (Park et al., 2018), which demonstrates an encouraging propensity for lipid tolerance. This tolerance has been harnessed by works that have grown *Y. lipolytica* strains on WCO. Growth on WCO increased lipolytic activity (Domínguez et al., 2010) and, in one study, a *Y. lipolytica* strain expressed D-limonene synthase

(*Citrus limon*) and L-limonene synthase (*Mentha spicata*) to yield 2.4 mg/L of each enantiomer on WCO (Pang et al., 2019). Although this strain has produced only 11 mg/L of each enantiomer on complex medium, this stands as an excellent proof of concept for future ALE and optimization studies on WCO.

Xylose catabolism and overcoming catabolite repression are major boundaries to bioproduction on lignocellulosic biomass (Sun et al., 2021). One study showed that carbon source switching enabled production of 20.6 mg/L and 15.1 mg/L limonene in *Y. lipolytica* from xylose and a 50% lignocellulosic biomass 50% YP rich medium broth, respectively (Yao et al., 2020). This feat was accomplished by overexpression of a native xylulose synthase with heterologous expression of xylitol dehydrogenase and xylulose reductase from *Scheffersomyces stipitis* (Yao et al., 2020). Together, these modifications provided increased G3P production and, ultimately, increased flux through the MVA pathway.

### *Rhodospiridium toruloides*

*R. toruloides* has attracted attention due to natural high titer lipid and carotenoid accumulation, namely torularhodin, torulene,  $\gamma$ -carotene, and  $\beta$ -carotene, as a convenient carbon storage mechanism under nitrogen-limited conditions (Park et al., 2018). Originally isolated from wood pulp, *R. toruloides* can also metabolize many components of lignocellulosic biomass and has shown simultaneous uptake not only of pentose and hexose sugars, but of p-coumaric acid and aromatic motifs analogous to lignin, which suggest that it could be adapted for direct consumption of lignin (Yaegashi et al., 2017). These traits are further complemented by its ability to thrive on various pretreatment conditions. For example, growth has been demonstrated on ionic liquid (choline  $\alpha$ -ketoglutarate) and alkaline pretreated cellulosic biomass, with the latter accumulating 680 mg/L  $\alpha$ -bisabolene in fed-batch reactor conditions (Yaegashi et al., 2017). Further optimization of the  $\alpha$ -bisabolene synthase cassette yielded 4-fold increased titer on lignocellulosic biomass, reaching a final titer of 2.2 g/L on corn stover hydrolysate (Kirby et al., 2021). 1,8-cineole was also accumulated to a titer of 1.4 g/L on the same substrate, both of which represent titers that, even without significant core metabolic rewiring or downregulation, outstrip comparative *E. coli* and *S. cerevisiae* production. Importantly, pilot scaling of *R. toruloides* to a 1,000 L bioreactor for lipid production has been successfully shown (Soccol et al., 2017). Collectively, these traits establish *R. toruloides* as a potential microbial host for lignin valorization. The translation of successful pilot scale *R. toruloides* lipid production platforms to strains with tuned lipid reflex pathways could elevate the yeast to an industrially competitive isoprenoid production platform.

### *Pseudomonas putida*

As noted with *Y. lipolytica*, tolerance to and simultaneous uptake of multiple carbon substrates is a key phenotype for successful bioproduction on lignocellulosic biomass. The soil bacterium *Pseudomonas putida* maintains significant advantages

over common production chassis due to its natural biodegradation pathways and oxidative stress tolerance, which has contributed to its broad proliferation in many environmental niches. Several studies have explored substrate tolerance through toxicity adaptive laboratory evolution (TALE) of *P. putida* (Mohamed et al., 2020; Lim et al., 2021). A recent work integrated three different xylose pathways (Dahms, Isomerase, and Weimberg) on plasmids to enable growth on xylose, a prominent component of degraded hemicellulose (Bator et al., 2019). The combination of pathway expression and ALE resulted in improved tolerance and hence improved growth rate (Bator et al., 2019).

Other production schemes have exploited the natural aromatic tolerance of *P. putida* for growth on substrates like toluene, m-xylene, and p-xylene (Nikel and de Lorenzo, 2018). Comparatively, *P. putida* maintains better *de novo* tolerance toward products that are typically toxic to other organisms (Mi et al., 2014). For example, the saprophytic uptake of organic nutrients and high tolerance to oxidative stress is ideal for biofuel production candidates (Kim and Park, 2014). These traits coupled with overexpression of efflux pumps have shown increased tolerance to short chain C4 and C5 alcohols, which could prove especially valuable for production of isoprenoid biofuels (Basler et al., 2018).

In a hallmark bioproduction study, 2.21 g/L of mevalonate were generated by *P. putida* in M9 minimal medium supplemented with 7.5 g/L 2,3-butanediol by overexpression of the upper MVA pathway enzymes, namely the native AtoB and the MvaE/MvaS from *Enterococcus faecalis* (Yang et al., 2020). Mevalonate production on 2,3-butanediol proved 6.61- and 8.44-fold higher than production on glucose and glycerol, respectively, though with manageable growth inhibition (Yang et al., 2020). Overall, *P. putida* isoprenoid production has historically been limited to zeaxanthin and geranic acid such that only recently have studies begun addressing MEP/heterologous MVA precursor limitations. One such study exhibited metabolic rerouting of central carbon metabolism from the EMP to ED cycles for better precursor management, namely efficient pyruvate production (Sánchez-Pascuala et al., 2019). This strategy led to a 2-fold increase in carotenoid yield on glucose with plasmid expression of a lycopene synthesis pathway but without any modification to the endogenous MEP pathway (Sánchez-Pascuala et al., 2019).

It is clear that *P. putida* has high innate tolerance to toxic substrates and, as in nature, can adapt to adverse growth conditions. The next, critical stages of realizing *P. putida* as a chemical production platform will be combining advances in ALE, precursor availability, and pathway tuning to enhance terpene synthesis on atypical carbon substrates.

## **Bacillus subtilis**

*Bacillus subtilis* is one of the best characterized gram-positive bacteria to date and has been an attractive bioproduction candidate due to high titer protein production and high secreting properties. Largely, the industrial focus has been on the production of biologics and enzymes (Pham et al., 2019). *B. subtilis* maintains a faster growth rate than *S. cerevisiae*, a

robust metabolism on diverse carbon substrates, and has also shown natural isoprene production at titers comparable to *E. coli* (Zhang et al., 2015). Unlike other chassis organisms like *P. putida*, *B. subtilis* is generally recognized as safe (GRAS), a designation that reduces regulatory boundaries to commercialization. Collectively, these factors suggest that *B. subtilis* could be an excellent candidate for isoprenoid production. Unfortunately, production studies remain relatively limited in part due to a poorly defined metabolic toolkit, which has historically been hampered by a limited subset of selection/counterselection markers that have made genetic manipulation challenging.

Mirroring in *E. coli* from the early 2000s, recent production studies demonstrated that incorporation of amorphadiene synthase (ADS) with overexpression of DXS and IDI led to the accumulation of 20 mg/L amorphadiene (Zhou et al., 2013). This titer has dramatically improved to 116 mg/L using a CRISPR-cas9 system without culture medium optimization (Song et al., 2021) and then to 416 mg/L (Pramastya et al., 2021) with pyruvate supplementation. Another recent study overexpressed the entire MEP pathway excluding IDI, a taxadiene synthase, and a heterologous GGPPS in *B. subtilis*, leading to an accumulation of 17.8 mg/L taxadiene (Abdallah et al., 2019). Expression of a squalene synthase from *Bacillus megaterium* also enabled 7.5 mg/L production of squalene, which can serve as a precursor to other triterpenoids (Song et al., 2020). Although far from competitive with *E. coli* and *S. cerevisiae*, these initial demonstrations have provided a basis of isoprenoid production in *B. subtilis*. The publication by Song et al. is of particular interest due to their application of CRISPR-cas9 to circumvent boundaries that have historically limited the establishment of *B. subtilis* as an isoprenoid production workhorse. In theory, this approach could be easily translated to the production of other isoprenoid targets.

## **PERSPECTIVES AND CONCLUSION**

The rapid expansion of -omics studies, deep sequencing, and pathway engineering have facilitated bioprospecting of more efficient enzymes, robust combinatorial approaches for tailored isoprenoid production, and the design of altogether novel production pathways. Such tools have also facilitated the exploration of plant derived CYPs and terminal synthases whose subsequent expression has expanded the microbial isoprenoid repertoire to more pharmacologically relevant as well as entirely synthetic terpenoids. In this review, we focused on improvements to isoprenoid precursor biosynthesis and translation of enzymes or pathways between organisms, which could assist in overcoming current major barriers to commercial viability (Zu et al., 2020). Specifically, we highlighted how atypical carbon sources and non-model organisms harbor metabolic advantages that could be harnessed to reduce substrate costs and the associated emissions of bioproduction. Co-substrate utilization by certain organisms as in the case of *R. toruloides* and *P. putida* has the potential to unlock lignocellulosic biomass and many

methylotrophs could tap into inexpensive and highly abundant substrates.

We have also described works that capitalized upon the modularity of isoprenoid advances through heterologous expression of entire pathways. Systems engineering strategies are of particular interest for C1 metabolism. The translation of successful whole systems engineering strategies from *E. coli* and *S. cerevisiae* to non-model organisms will prove useful in further optimization. For example, the entire MVA pathway had been expressed in *E. coli* many years ago (Martin et al., 2003) and a decade later, the entire MEP pathway has been expressed in *S. cerevisiae* conversely (Kirby et al., 2016). Both strains have also been extensively mapped through metabolic flux analysis (MFA) which has proven pivotal in metabolic engineering (Orth et al., 2010). The translation of systems engineering strategies like MFA and genome-scale modeling to other organisms will undoubtedly help to inform and improve isoprenoid production in non-model organisms. An MFA of *R. sphaeroides*, for example, showed a mutualistic coupling between its MEP and MVA pathways (Orsi et al., 2020a). Remarkably the true extent of the MEP pathway – MVA pathway relationship could not be resolved as gene knockouts tended to have unpredictable effects on C13 product partitioning but suggested complex regulatory interactions. Nonetheless further work could shed light on how such combined MVA/MEP pathway systems could prove beneficial (Orsi et al., 2020a). Similarly, a metabolic flux reconstruction of *Dunaliella salina* established baseline carbon metabolism during carotenogenesis (Fachet et al., 2020), a critical step in elucidating metabolic bottlenecks. TALE has also proven a powerful strategy for increasing resistance to toxicity of high titer products especially with alcohols. TALE has now been applied to *P. putida* and enhanced toxicity tolerance against the lignocellulosic aromatics such as p-coumaric acid and ferulic acid (Lim et al., 2020; Mohamed et al., 2020). The application of machine learning approaches has enabled extrapolation and gap filling in genome-scale models for rationally designed engineering strategies of non-canonical organisms, as demonstrated to great effect in *Y. lipolytica* (Czajka et al., 2021). And, finally, C1 assimilation pathways have been thoroughly explored, synthetic and natural routes hypothesized (Bar-Even, 2016), then optimal pathways have been heterologously expressed in conventional production chassis (Kim et al., 2020). Having shown adapted growth on C1 substrates there is now a tremendous opportunity to further develop strains for isoprenoid production especially given the comparative sustainability and cost reduction of such substrates with respect to production on refined sugars.

Consortial approaches are also valuable by improving total system productivity. Microbial consortia have proven successful for short chain alcohol production from lignocellulosic biomass (Minty et al., 2013) and have recently been explored in the cross-feeding of methane-derived organic acids produced by *Methylococcus capsulatus* to *E. coli* for the generation of mevalonate at 60 mg/L (Lee et al., 2021). Building upon CYP optimization, an *E. coli* and *S. cerevisiae* consortium produced 33 mg/L oxygenated taxanes in a

consortia where *E. coli* consumes xylose and produces acetate and the precursor taxadiene for consumption and further conjugation in *S. cerevisiae*, respectively (Zhou et al., 2015). Another group produced 0.32 g mevalonate/g ethanol in *P. putida* batch experiments (Yang et al., 2019) that, paired with the aforementioned successes in ALE, could provide another promising cross-feeding consortial bioproduction strategy. Finally, isoprenoid production has been expanded to 2,3-butanediol (Yang et al., 2020), which could facilitate consortial bioproduction by subdividing pathways between members.

Exploration of the microbial tree of life has continued to yield an abundant natural diversity of protein homologues, pathway shunts, and mechanisms with which targeted production of isoprenoids has been demonstrably improved. The principal challenge of isoprenoid bioproduction in the next decade will be bridging the knowledge gap between conventional high titer bioproduction on pure sugar substrates and non-model comparatively low titer production on affordable substrates.

## AUTHOR CONTRIBUTIONS

DC and TSL contributed to the writing of this review and conceived the topic. DC prepared the initial draft, reviewed, and edited the draft. TSL directed the writing process and wrote and reviewed the draft. All authors contributed to the article and approved the submitted version.

## FUNDING

This work was supported by the DOE Joint BioEnergy Institute (<http://www.jbei.org>), supported by the US Department of Energy, Office of Science, Office of Biological and Environmental Research, through contract DE-AC0205CH11231 with Lawrence Berkeley National Laboratory.

## ACKNOWLEDGMENTS

The United States Government retains and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes. The views and opinions of the authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights.

## REFERENCES

- Abdallah, I. I., Pramastya, H., van Merkerk, R., and Sukrasno, and Quax, W. J., (2019). Metabolic engineering of *Bacillus subtilis* toward taxadiene biosynthesis as the first committed step for taxol production. *Front. Microbiol.* 10:218. doi: 10.3389/fmicb.2019.00218
- Aldridge, J., Carr, S., Weber, K. A., and Buan, N. R. (2021). Anaerobic production of isoprene by engineered methanosarcina species archaea. *Appl. Environ. Microbiol.* 87, e02417–e02420. doi: 10.1128/AEM.02417-20
- Alonso-Gutierrez, J., Chan, R., Batth, T. S., Adams, P. D., Keasling, J. D., Petzold, C. J., et al. (2013). Metabolic engineering of *Escherichia coli* for limonene and perillyl alcohol production. *Metab. Eng.* 19, 33–41. doi: 10.1016/j.ymben.2013.05.004
- Arnesen, J. A., Kildegaard, K. R., Cernuda Pastor, M., Jayachandran, S., Kristensen, M., and Borodina, I. (2020). *Yarrowia lipolytica* strains engineered for the production of terpenoids. *Front. Bioeng. Biotechnol.* 8:945. doi: 10.3389/fbioe.2020.00945
- Athanasakoglou, A., Grypioti, E., Michailidou, S., Ignea, C., Makris, A. M., Kalantidis, K., et al. (2019). Isoprenoid biosynthesis in the diatom *Haslea ostrearia*. *New Phytol.* 222, 230–243. doi: 10.1111/nph.15586
- Bar-Even, A. (2016). Formate assimilation: the metabolic architecture of natural and synthetic pathways. *Biochemistry* 55, 3851–3863. doi: 10.1021/acs.biochem.6b00495
- Basler, G., Thompson, M., Tullman-Ercek, D., and Keasling, J. (2018). A *Pseudomonas putida* efflux pump acts on short-chain alcohols. *Biotechnol. Biofuels* 11:136. doi: 10.1186/s13068-018-1133-9
- Bator, I., Wittgens, A., Rosenau, F., Tiso, T., and Blank, L. M. (2019). Comparison of three xylose pathways in *Pseudomonas putida* KT2440 for the synthesis of valuable products. *Front. Bioeng. Biotechnol.* 7:480. doi: 10.3389/fbioe.2019.00480
- Beekwilder, J., van Houwelingen, A., Cankar, K., van Dijk, A. D. J., de Jong, R. M., Stoopen, G., et al. (2014). Valencene synthase from the heartwood of Nootka cypress (*Callitropsis nootkatensis*) for biotechnological production of valencene. *Plant Biotechnol. J.* 12, 174–182. doi: 10.1111/pbi.12124
- Bennett, R. K., Gonzalez, J. E., Whitaker, W. B., Antoniewicz, M. R., and Papoutsakis, E. T. (2018). Expression of heterologous non-oxidative pentose phosphate pathway from *Bacillus methanolicus* and phosphoglucose isomerase deletion improves methanol assimilation and metabolite production by a synthetic *Escherichia coli* methylotroph. *Metab. Eng.* 45, 75–85. doi: 10.1016/j.ymben.2017.11.016
- Biggs, B. W., Lim, C. G., Sagliani, K., Shankar, S., Stephanopoulos, G., De Mey, M., et al. (2016). Overcoming heterologous protein interdependency to optimize P450-mediated Taxol precursor synthesis in *Escherichia coli*. *Proc. Natl. Acad. Sci. U. S. A.* 113, 3209–3214. doi: 10.1073/pnas.1515826113
- Bitok, J. K., and Meyers, C. F. (2012). 2C-methyl-d-erythritol 4-phosphate enhances and sustains cyclodiphosphate synthase IspF activity. *ACS Chem. Biol.* 7, 1702–1710. doi: 10.1021/cb300243w
- Borowitzka, M. A. (2013). High-value products from microalgae—their development and commercialisation. *J. Appl. Phycol.* 25, 743–756. doi: 10.1007/s10811-013-9983-9
- Boucher, Y., Kamekura, M., and Doolittle, W. F. (2004). Origins and evolution of isoprenoid lipid biosynthesis in archaea. *Mol. Microbiol.* 52, 515–527. doi: 10.1111/j.1365-2958.2004.03992.x
- Caforio, A., and Driessen, A. J. M. (2017). Archaeal phospholipids: structural properties and biosynthesis. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* 1862, 1325–1339. doi: 10.1016/j.bbalip.2016.12.006
- Cao, X., Lv, Y.-B., Chen, J., Imanaka, T., Wei, L.-J., and Hua, Q. (2016). Metabolic engineering of oleaginous yeast *Yarrowia lipolytica* for limonene overproduction. *Biotechnol. Biofuels* 9:214. doi: 10.1186/s13068-016-0626-7
- Castelle, C. J., and Banfield, J. F. (2018). Major new microbial groups expand diversity and alter our understanding of the tree of life. *Cell* 172, 1181–1197. doi: 10.1016/j.cell.2018.02.016
- Chang, M. C. Y., Eachus, R. A., Trieu, W., Ro, D.-K., and Keasling, J. D. (2007). Engineering *Escherichia coli* for production of functionalized terpenoids using plant P450s. *Nat. Chem. Biol.* 3, 274–277. doi: 10.1038/nchembio875
- Chappell, J. (1995). The biochemistry and molecular biology of isoprenoid metabolism. *Plant Physiol.* 107, 1–6. doi: 10.1104/pp.107.1.1
- Chatzivasileiou, A. O., Ward, V., Edgar, S. M., and Stephanopoulos, G. (2019). Two-step pathway for isoprenoid synthesis. *Proc. Natl. Acad. Sci. U. S. A.* 116, 506–511. doi: 10.1073/pnas.1812935116
- Chen, F. Y.-H., Jung, H.-W., Tsuei, C.-Y., and Liao, J. C. (2020). Converting *Escherichia coli* to a synthetic methylotroph growing solely on methanol. *Cell* 182, 933–946.e14. doi: 10.1016/j.cell.2020.07.010
- Cheng, B.-Q., Wei, L.-J., Lv, Y.-B., Chen, J., and Hua, Q. (2019). Elevating limonene production in oleaginous yeast *Yarrowia lipolytica* via genetic engineering of limonene biosynthesis pathway and optimization of medium composition. *Biotechnol. Bioprocess Eng.* 24, 500–506. doi: 10.1007/s12257-018-0497-9
- Clomburg, J. M., Crumbley, A. M., and Gonzalez, R. (2017). Industrial biomufacturing: the future of chemical production. *Science* 355:aag0804. doi: 10.1126/science.aag0804
- Clomburg, J. M., Qian, S., Tan, Z., Cheong, S., and Gonzalez, R. (2019). The isoprenoid alcohol pathway, a synthetic route for isoprenoid biosynthesis. *Proc. Natl. Acad. Sci. U. S. A.* 116, 12810–12815. doi: 10.1073/pnas.1821004116
- Conrado, R. J., and Gonzalez, R. (2014). Chemistry: Envisioning the bioconversion of methane to liquid fuels. *Science* 343, 621–623. doi: 10.1126/science.1246929
- Crowther, G. J., Kosály, G., and Lidstrom, M. E. (2008). Formate as the main branch point for methylotrophic metabolism in *Methylobacterium extorquens* AM1. *J. Bacteriol.* 190, 5057–5062. doi: 10.1128/JB.00228-08
- Czajka, J. J., Oyetunde, T., and Tang, Y. J. (2021). Integrated knowledge mining, genome-scale modeling, and machine learning for predicting *Yarrowia lipolytica* bioproduction. *Metab. Eng.* 67, 227–236. doi: 10.1016/j.ymben.2021.07.003
- D'Adamo, S., Schiano di Visconte, G., Lowe, G., Szaub-Newton, J., Beacham, T., Landels, A., et al. (2019). Engineering the unicellular alga *Phaeodactylum tricornutum* for high-value plant triterpenoid production. *Plant Biotechnol. J.* 17, 75–87. doi: 10.1111/pbi.12948
- Dall'Osto, L., Lico, C., Alric, J., Giuliano, G., Havaux, M., and Bassi, R. (2006). Lutein is needed for efficient chlorophyll triplet quenching in the major LHCII antenna complex of higher plants and effective photoprotection in vivo under strong light. *BMC Plant Biol.* 6:32. doi: 10.1186/1471-2229-6-32
- Della Monica, F., and Kleij, A. W. (2020). From terpenes to sustainable and functional polymers. *Polym. Chem.* 11, 5109–5127. doi: 10.1039/D0PY00817F
- Dellas, N., Thomas, S. T., Manning, G., and Noel, J. P. (2013). Discovery of a metabolic alternative to the classical mevalonate pathway. *elife* 2:e00672. doi: 10.7554/eLife.00672
- Domínguez, A., Deive, F. J., Angeles Sanromán, M., and Longo, M. A. (2010). Biodegradation and utilization of waste cooking oil by *Yarrowia lipolytica* CECT 1240. *Eur. J. Lipid Sci. Technol.* 112, 1200–1208. doi: 10.1002/ejlt.201000049
- Drummond, L., Kschowak, M. J., Breitenbach, J., Wolff, H., Shi, Y.-M., Schrader, J., et al. (2019). Expanding the isoprenoid building block repertoire with an IPP methyltransferase from *Streptomyces monomycinii*. *ACS Synth. Biol.* 8, 1303–1313. doi: 10.1021/acssynbio.8b00525
- Ducat, D. C., Avelar-Rivas, J. A., Way, J. C., and Silver, P. A. (2012). Rerouting carbon flux to enhance photosynthetic productivity. *Appl. Environ. Microbiol.* 78, 2660–2668. doi: 10.1128/AEM.07901-11
- Dugar, D., and Stephanopoulos, G. (2011). Relative potential of biosynthetic pathways for biofuels and bio-based products. *Nat. Biotechnol.* 29, 1074–1078. doi: 10.1038/nbt.2055
- Dunlop, M. J. (2011). Engineering microbes for tolerance to next-generation biofuels. *Biotechnol. Biofuels* 4:32. doi: 10.1186/1754-6834-4-32
- Dunlop, M. J., Dossani, Z. Y., Szmids, H. L., Chu, H. C., Lee, T. S., Keasling, J. D., et al. (2011). Engineering microbial biofuel tolerance and export using efflux pumps. *Mol. Syst. Biol.* 7:487. doi: 10.1038/msb.2011.21
- Eiben, C. B., de Rond, T., Bloszies, C., Gin, J., Chiniquy, J., Baidoo, E. E. K., et al. (2019). Mevalonate pathway promiscuity enables noncanonical terpene production. *ACS Synth. Biol.* 8, 2238–2247. doi: 10.1021/acssynbio.9b00230
- Eiben, C. B., Tian, T., Thompson, M. G., Mendez-Perez, D., Kaplan, N., Goyal, G., et al. (2020). Adenosine triphosphate and carbon efficient route to second generation biofuel isopentanol. *ACS Synth. Biol.* 9, 468–474. doi: 10.1021/acssynbio.9b00402
- Englund, E., Shabestary, K., Hudson, E. P., and Lindberg, P. (2018). Systematic overexpression study to find target enzymes enhancing production of terpenes in *Synechocystis* PCC 6803, using isoprene as a model compound. *Metab. Eng.* 49, 164–177. doi: 10.1016/j.ymben.2018.07.004

- Fachet, M., Witte, C., Flassig, R. J., Rihko-Struckmann, L. K., McKie-Krisberg, Z., Polle, J. E. W., et al. (2020). Reconstruction and analysis of a carbon-core metabolic network for *Dunaliella salina*. *BMC Bioinform.* 21:1. doi: 10.1186/s12859-019-3325-0
- Fei, Q., Liang, B., Tao, L., Tan, E. C. D., Gonzalez, R., Henard, C., et al. (2020). Biological valorization of natural gas for the production of lactic acid: techno-economic analysis and life cycle assessment. *Biochem. Eng. J.* 158:107500. doi: 10.1016/j.bej.2020.107500
- Ferriols, V. M. E. N., Yaginuma, R., Adachi, M., Takada, K., Matsunaga, S., and Okada, S. (2015). Cloning and characterization of farnesyl pyrophosphate synthase from the highly branched isoprenoid producing diatom *Rhizosolenia setigera*. *Sci. Rep.* 5:10246. doi: 10.1038/srep10246
- Gao, X., Gao, F., Liu, D., Zhang, H., Nie, X., and Yang, C. (2016). Engineering the methylerythritol phosphate pathway in cyanobacteria for photosynthetic isoprene production from CO<sub>2</sub>. *Energy Environ. Sci.* 9, 1400–1411. doi: 10.1039/C5EE03102H
- Gao, S., Tong, Y., Zhu, L., Ge, M., Zhang, Y., Chen, D., et al. (2017). Iterative integration of multiple-copy pathway genes in *Yarrowia lipolytica* for heterologous β-carotene production. *Metab. Eng.* 41, 192–201. doi: 10.1016/j.ymben.2017.04.004
- Gassler, T., Sauer, M., Gasser, B., Egermeier, M., Troyer, C., Causon, T., et al. (2020). The industrial yeast *Pichia pastoris* is converted from a heterotroph into an autotroph capable of growth on CO<sub>2</sub>. *Nat. Biotechnol.* 38, 210–216. doi: 10.1038/s41587-019-0363-0
- George, K. W., Thompson, M. G., Kang, A., Baidoo, E., Wang, G., Chan, L. J. G., et al. (2015). Metabolic engineering for the high-yield production of isoprenoid-based C<sub>5</sub> alcohols in *E. coli*. *Sci. Rep.* 5:11128. doi: 10.1038/srep11128
- George, K. W., Thompson, M. G., Kim, J., Baidoo, E. E. K., Wang, G., Benites, V. T., et al. (2018). Integrated analysis of isopentenyl pyrophosphate (IPP) toxicity in isoprenoid-producing *Escherichia coli*. *Metab. Eng.* 47, 60–72. doi: 10.1016/j.ymben.2018.03.004
- Guo, X., Sun, J., Li, D., and Lu, W. (2018). Heterologous biosynthesis of (+)-nootkatone in unconventional yeast ' *Yarrowia lipolytica*. *Biochem. Eng. J.* 137, 125–131. doi: 10.1016/j.bej.2018.05.023
- Guo, J., Zhou, Y. J., Hillwig, M. L., Shen, Y., Yang, L., Wang, Y., et al. (2013). CYP76AH1 catalyzes turnover of miltiradiene in tanshinones biosynthesis and enables heterologous production of ferruginol in yeasts. *Proc. Natl. Acad. Sci. U. S. A.* 110, 12108–12113. doi: 10.1073/pnas.1218061110
- Hartner, F. S., and Glieder, A. (2006). Regulation of methanol utilisation pathway genes in yeasts. *Microb. Cell Factories* 5:39. doi: 10.1186/1475-2859-5-39
- Hausjell, J., Halbwirth, H., and Spadiut, O. (2018). Recombinant production of eukaryotic cytochrome P450s in microbial cell factories. *Biosci. Rep.* 38:BSR20171290. doi: 10.1042/BSR20171290
- Hayakawa, H., Motoyama, K., Sobue, F., Ito, T., Kawaide, H., Yoshimura, T., et al. (2018). Modified mevalonate pathway of the archaeon *Aeropyrum pernix* proceeds via trans-anhydromevalonate 5-phosphate. *Proc. Natl. Acad. Sci. U. S. A.* 115, 10034–10039. doi: 10.1073/pnas.1809154115
- Helfrich, E. J. N., Lin, G.-M., Voigt, C. A., and Clardy, J. (2019). Bacterial terpene biosynthesis: challenges and opportunities for pathway engineering. *Beilstein J. Org. Chem.* 15, 2889–2906. doi: 10.3762/bjoc.15.283
- Hernandez-Arranz, S., Perez-Gil, J., Marshall-Sabey, D., and Rodriguez-Concepcion, M. (2019). Engineering *Pseudomonas putida* for isoprenoid production by manipulating endogenous and shunt pathways supplying precursors. *Microb. Cell Factories* 18:152. doi: 10.1186/s12934-019-1204-z
- Hernandez-Ortega, A., Vinaixa, M., Zebec, Z., Takano, E., and Scrutton, N. S. (2018). A toolbox for diverse oxyfunctionalisation of monoterpenes. *Sci. Rep.* 8:14396. doi: 10.1038/s41598-018-32816-1
- Hirokawa, Y., Kubo, T., Soma, Y., Saruta, F., and Hanai, T. (2020). Enhancement of acetyl-CoA flux for photosynthetic chemical production by pyruvate dehydrogenase complex overexpression in *Synechococcus elongatus* PCC 7942. *Metab. Eng.* 57, 23–30. doi: 10.1016/j.ymben.2019.07.012
- Hu, B., and Lidstrom, M. E. (2014). Metabolic engineering of *Methylobacterium extorquens* AM1 for 1-butanol production. *Biotechnol. Biofuels* 7:156. doi: 10.1186/s13068-014-0156-0
- Ignea, C., Pontini, M., Motawia, M. S., Maffei, M. E., Makris, A. M., and Kampranis, S. C. (2018). Synthesis of 11-carbon terpenoids in yeast using protein and metabolic engineering. *Nat. Chem. Biol.* 14, 1090–1098. doi: 10.1038/s41589-018-0166-5
- Jain, S., Caforio, A., and Driessen, A. J. M. (2014). Biosynthesis of archaeal membrane ether lipids. *Front. Microbiol.* 5:641. doi: 10.3389/fmicb.2014.00641
- Johnson, C. W., Salvachúa, D., Rorrer, N. A., Black, B. A., Vardon, D. R., St. John, P. C., et al. (2019). Innovative chemicals and materials from bacterial aromatic catabolic pathways. *Joule* 19, 1523–1537. doi: 10.1016/j.joule.2019.05.011
- Kang, A., George, K. W., Wang, G., Baidoo, E., Keasling, J. D., and Lee, T. S. (2016). Isopentenyl diphosphate (IPP)-bypass mevalonate pathways for isoprenol production. *Metab. Eng.* 34, 25–35. doi: 10.1016/j.ymben.2015.12.002
- Kang, A., Meadows, C. W., Canu, N., Keasling, J. D., and Lee, T. S. (2017). High-throughput enzyme screening platform for the IPP-bypass mevalonate pathway for isoprenol production. *Metab. Eng.* 41, 125–134. doi: 10.1016/j.ymben.2017.03.010
- Kang, A., Mendez-Perez, D., Goh, E.-B., Baidoo, E. E. K., Benites, V. T., Beller, H. R., et al. (2019). Optimization of the IPP-bypass mevalonate pathway and fed-batch fermentation for the production of isoprenol in *Escherichia coli*. *Metab. Eng.* 56, 85–96. doi: 10.1016/j.ymben.2019.09.003
- Karunanithi, P. S., and Zerbe, P. (2019). Terpene synthases as metabolic gatekeepers in the evolution of plant terpenoid chemical diversity. *Front. Plant Sci.* 10:1166. doi: 10.3389/fpls.2019.01166
- Kim, S. K., Han, G. H., Seong, W., Kim, H., Kim, S.-W., Lee, D.-H., et al. (2016). CRISPR interference-guided balancing of a biosynthetic mevalonate pathway increases terpenoid production. *Metab. Eng.* 38, 228–240. doi: 10.1016/j.ymben.2016.08.006
- Kim, S., Lindner, S. N., Aslan, S., Yishai, O., Wenk, S., Schann, K., et al. (2020). Growth of *E. coli* on formate and methanol via the reductive glycine pathway. *Nat. Chem. Biol.* 16, 538–545. doi: 10.1038/s41589-020-0473-5
- Kim, J., and Park, W. (2014). Oxidative stress response in *Pseudomonas putida*. *Appl. Microbiol. Biotechnol.* 98, 6933–6946. doi: 10.1007/s00253-014-5883-4
- Kim, S.-J., Yoon, J., Im, D.-K., Kim, Y. H., and Oh, M.-K. (2019). Adaptively evolved *Escherichia coli* for improved ability of formate utilization as a carbon source in sugar-free conditions. *Biotechnol. Biofuels* 12:207. doi: 10.1186/s13068-019-1547-z
- King, J. R., Woolston, B. M., and Stephanopoulos, G. (2017). Designing a new entry point into isoprenoid metabolism by exploiting fructose-6-phosphate aldolase side reactivity of *Escherichia coli*. *ACS Synth. Biol.* 6, 1416–1426. doi: 10.1021/acssynbio.7b00072
- Kirby, J., Dietzel, K. L., Wichmann, G., Chan, R., Antipov, E., Moss, N., et al. (2016). Engineering a functional 1-deoxy-D-xylulose 5-phosphate (DXP) pathway in *Saccharomyces cerevisiae*. *Metab. Eng.* 38, 494–503. doi: 10.1016/j.ymben.2016.10.017
- Kirby, J., Geiselman, G. M., Yaegashi, J., Kim, J., Zhuang, X., Tran-Gyamfi, M. B., et al. (2021). Further engineering of *R. toruloides* for the production of terpenes from lignocellulosic biomass. *Biotechnol. Biofuels* 14:101. doi: 10.1186/s13068-021-01950-w
- Kirby, J., and Keasling, J. D. (2009). Biosynthesis of plant isoprenoids: perspectives for microbial engineering. *Annu. Rev. Plant Biol.* 60, 335–355. doi: 10.1146/annurev.arplant.043008.091955
- Kirby, J., Nishimoto, M., Chow, R. W. N., Baidoo, E. E. K., Wang, G., Martin, J., et al. (2015). Enhancing Terpene yield from sugars via novel routes to 1-deoxy-D-xylulose 5-phosphate. *Appl. Environ. Microbiol.* 81, 130–138. doi: 10.1128/AEM.02920-14
- Kuhl, K. P., Cave, E. R., Abram, D. N., and Jaramillo, T. F. (2012). New insights into the electrochemical reduction of carbon dioxide on metallic copper surfaces. *Energy Environ. Sci.* 5:7050. doi: 10.1039/c2ee21234j
- Larroude, M., Celinska, E., Back, A., Thomas, S., Nicaud, J.-M., and Ledesma-Amaro, R. (2018). A synthetic biology approach to transform *Yarrowia lipolytica* into a competitive biotechnological producer of β-carotene. *Biotechnol. Bioeng.* 115, 464–472. doi: 10.1002/bit.26473
- Laursen, K. J., Baier, T., Wichmann, J., Wördenweber, R., Mussnug, J. H., Hübner, W., et al. (2016). Efficient phototrophic production of a high-value sesquiterpenoid from the eukaryotic microalga *Chlamydomonas reinhardtii*. *Metab. Eng.* 38, 331–343. doi: 10.1016/j.ymben.2016.07.013
- Laursen, K. J., Wichmann, J., Baier, T., Kampranis, S. C., Pateraki, I., Møller, B. L., et al. (2018). Phototrophic production of heterologous diterpenoids and a hydroxy-functionalized derivative from *Chlamydomonas reinhardtii*. *Metab. Eng.* 49, 116–127. doi: 10.1016/j.ymben.2018.07.005

- Lee, H., Baek, J. I., Lee, J.-Y., Jeong, J., Kim, H., Lee, D.-H., et al. (2021). Syntrophic co-culture of a methanotroph and heterotroph for the efficient conversion of methane to mevalonate. *Metab. Eng.* 67, 285–292. doi: 10.1016/j.ymben.2021.07.008
- Lee, H. J., Lee, J., Lee, S.-M., Um, Y., Kim, Y., Sim, S. J., et al. (2017). Direct conversion of CO<sub>2</sub> to  $\alpha$ -farnesene using metabolically engineered *Synechococcus elongatus* PCC 7942. *J. Agric. Food Chem.* 65, 10424–10428. doi: 10.1021/acs.jafc.7b03625
- Levisson, M., Araya-Cloutier, C., de Bruijn, W. J. C., van der Heide, M., Salvador López, J. M., Daran, J.-M., et al. (2019). Toward developing a yeast cell factory for the production of prenylated flavonoids. *J. Agric. Food Chem.* 67, 13478–13486. doi: 10.1021/acs.jafc.9b01367
- Li, Q., Fan, F., Gao, X., Yang, C., Bi, C., Tang, J., et al. (2017). Balanced activation of IspG and IspH to eliminate MEP intermediate accumulation and improve isoprenoids production in *Escherichia coli*. *Metab. Eng.* 44, 13–21. doi: 10.1016/j.ymben.2017.08.005
- Li, Z., Jiang, Y., Guengerich, F. P., Ma, L., Li, S., and Zhang, W. (2020). Engineering cytochrome P450 enzyme systems for biomedical and biotechnological applications. *J. Biol. Chem.* 295, 833–849. doi: 10.1016/S0021-9258(17)49939-X
- Liang, W.-F., Cui, L.-Y., Cui, J.-Y., Yu, K.-W., Yang, S., Wang, T.-M., et al. (2017). Biosensor-assisted transcriptional regulator engineering for *Methylobacterium extorquens* AM1 to improve mevalonate synthesis by increasing the acetyl-CoA supply. *Metab. Eng.* 39, 159–168. doi: 10.1016/j.ymben.2016.11.010
- Lim, H. G., Eng, T., Banerjee, D., Alarcon, G., Lau, A. K., Park, M.-R., et al. (2021). Generation of *Pseudomonas putida* KT2440 strains with efficient utilization of xylose and galactose via adaptive laboratory evolution. *ACS Sustain. Chem. Eng.* 9, 11512–11523. doi: 10.1021/acsschemeng.1c03765
- Lim, H. G., Fong, B., Alarcon, G., Magurudeniya, H. D., Eng, T., Szubin, R., et al. (2020). Generation of ionic liquid tolerant *Pseudomonas putida* KT2440 strains via adaptive laboratory evolution. *Green Chem.* 22, 5677–5690. doi: 10.1039/D0GC01663B
- Liu, Y., Jiang, X., Cui, Z., Wang, Z., Qi, Q., and Hou, J. (2019). Engineering the oleaginous yeast *Yarrowia lipolytica* for production of  $\alpha$ -farnesene. *Biotechnol. Biofuels* 12:296. doi: 10.1186/s13068-019-1636-z
- Liu, Y., Yan, Z., Lu, X., Xiao, D., and Jiang, H. (2016). Improving the catalytic activity of isopentenyl phosphate kinase through protein coevolution analysis. *Sci. Rep.* 6:24117. doi: 10.1038/srep24117
- Liu, X., Zhu, X., Wang, H., Liu, T., Cheng, J., and Jiang, H. (2020). Discovery and modification of cytochrome P450 for plant natural products biosynthesis. *Synth. Syst. Biotechnol.* 5, 187–199. doi: 10.1016/j.synbio.2020.06.008
- Lund, S., Hall, R., and Williams, G. J. (2019). An artificial pathway for isoprenoid biosynthesis decoupled from native hemiterpene metabolism. *ACS Synth. Biol.* 8, 232–238. doi: 10.1021/acssynbio.8b00383
- Luo, X., Reiter, M. A., d Espaux, L., Wong, J., Denby, C. M., Lechner, A., et al. (2019). Complete biosynthesis of cannabinoids and their unnatural analogues in yeast. *Nature* 567, 123–126. doi: 10.1038/s41586-019-0978-9
- Lyu, Z., Jain, R., Smith, P., Fetchko, T., Yan, Y., and Whitman, W. B. (2016). Engineering the autotroph *Methanococcus marisplacidis* for geraniol production. *ACS Synth. Biol.* 5, 577–581. doi: 10.1021/acssynbio.5b00267
- Martin, V. J. J., Pitera, D. J., Withers, S. T., Newman, J. D., and Keasling, J. D. (2003). Engineering a mevalonate pathway in *Escherichia coli* for production of terpenoids. *Nat. Biotechnol.* 21, 796–802. doi: 10.1038/nbt833
- Metzger, P., and Largeau, C. (2005). *Botryococcus braunii*: a rich source for hydrocarbons and related ether lipids. *Appl. Microbiol. Biotechnol.* 66, 486–496. doi: 10.1007/s00253-004-1779-z
- Mi, J., Becher, D., Lubuta, P., Dany, S., Tusch, K., Schewe, H., et al. (2014). De novo production of the monoterpene geranic acid by metabolically engineered *Pseudomonas putida*. *Microb. Cell Factories* 13:170. doi: 10.1186/s12934-014-0170-8
- Minty, J. J., Singer, M. E., Scholz, S. A., Bae, C.-H., Ahn, J.-H., Foster, C. E., et al. (2013). Design and characterization of synthetic fungal-bacterial consortia for direct production of isobutanol from cellulosic biomass. *Proc. Natl. Acad. Sci. U. S. A.* 110, 14592–14597. doi: 10.1073/pnas.1218447110
- Mohamed, E. T., Werner, A. Z., Salvachúa, D., Singer, C. A., Szostkiewicz, K., Rafael Jiménez-Díaz, M., et al. (2020). Adaptive laboratory evolution of *Pseudomonas putida* KT2440 improves p-coumaric and ferulic acid catabolism and tolerance. *Metab. Eng. Commun.* 11:e00143. doi: 10.1016/j.mec.2020.e00143
- Morales-Sánchez, D., Martínez-Rodríguez, O. A., and Martínez, A. (2017). Heterotrophic cultivation of microalgae: production of metabolites of commercial interest. *J. Chem. Technol. Biotechnol.* 92, 925–936. doi: 10.1002/jctb.5115
- Moser, S., and Pichler, H. (2019). Identifying and engineering the ideal microbial terpenoid production host. *Appl. Microbiol. Biotechnol.* 103, 5501–5516. doi: 10.1007/s00253-019-09892-y
- Moser, S., Strohmeier, G. A., Leitner, E., Plocek, T. J., Vanhessche, K., and Pichler, H. (2018). Whole-cell (+)-ambrein production in the yeast *Pichia pastoris*. *Metab. Eng. Commun.* 7:e00077. doi: 10.1016/j.mec.2018.e00077
- Nguyen, A. D., Kim, D., and Lee, E. Y. (2020). Unlocking the biosynthesis of sesquiterpenoids from methane via the methylerythritol phosphate pathway in methanotrophic bacteria, using  $\alpha$ -humulene as a model compound. *Metab. Eng.* 61, 69–78. doi: 10.1016/j.ymben.2020.04.011
- Nikel, P. I., and de Lorenzo, V. (2018). *Pseudomonas putida* as a functional chassis for industrial biocatalysis: From native biochemistry to trans-metabolism. *Metab. Eng.* 50, 142–155. doi: 10.1016/j.ymben.2018.05.005
- Oliver, J. W. K., and Atsumi, S. (2015). A carbon sink pathway increases carbon productivity in cyanobacteria. *Metab. Eng.* 29, 106–112. doi: 10.1016/j.ymben.2015.03.006
- Orita, I., Nishikawa, K., Nakamura, S., and Fukui, T. (2014). Biosynthesis of polyhydroxyalkanoate copolymers from methanol by *Methylobacterium extorquens* AM1 and the engineered strains under cobalt-deficient conditions. *Appl. Microbiol. Biotechnol.* 98, 3715–3725. doi: 10.1007/s00253-013-5490-9
- Orsi, E., Beekwilder, J., Peek, S., Eggink, G., Kengen, S. W. M., and Weusthuis, R. A. (2020a). Metabolic flux ratio analysis by parallel 13C labeling of isoprenoid biosynthesis in *Rhodobacter sphaeroides*. *Metab. Eng.* 57, 228–238. doi: 10.1016/j.ymben.2019.12.004
- Orsi, E., Mougiakos, I., Post, W., Beekwilder, J., Dompé, M., Eggink, G., et al. (2020b). Growth-uncoupled isoprenoid synthesis in *Rhodobacter sphaeroides*. *Biotechnol. Biofuels* 13:123. doi: 10.1186/s13068-020-01765-1
- Orth, J. D., Thiele, I., and Palsson, B. Ø. (2010). What is flux balance analysis? *Nat. Biotechnol.* 28, 245–248. doi: 10.1038/nbt.1614
- Otto, M., Teixeira, P. G., Vizcaino, M. I., David, F., and Siewers, V. (2019). Integration of a multi-step heterologous pathway in *Saccharomyces cerevisiae* for the production of abscisic acid. *Microb. Cell Factories* 18:205. doi: 10.1186/s12934-019-1257-z
- Pang, Y., Zhao, Y., Li, S., Zhao, Y., Li, J., Hu, Z., et al. (2019). Engineering the oleaginous yeast *Yarrowia lipolytica* to produce limonene from waste cooking oil. *Biotechnol. Biofuels* 12:241. doi: 10.1186/s13068-019-1580-y
- Park, Y.-K., Nicaud, J.-M., and Ledesma-Amaro, R. (2018). The engineering potential of *Rhodospiridium toruloides* as a workhorse for biotechnological applications. *Trends Biotechnol.* 36, 304–317. doi: 10.1016/j.tibtech.2017.10.013
- Pasternak, Z., Pietrokovski, S., Rotem, O., Gophna, U., Lurie-Weinberger, M. N., and Jurkevitch, E. (2013). By their genes ye shall know them: genomic signatures of predatory bacteria. *ISME J.* 7, 756–769. doi: 10.1038/ismej.2012.149
- Pattanaik, B., Englund, E., Nolte, N., and Lindberg, P. (2020). Introduction of a green algal squalene synthase enhances squalene accumulation in a strain of *Synechocystis* sp. PCC 6803. *Metab. Eng. Commun.* 10:e00125. doi: 10.1016/j.mec.2020.e00125
- Pham, J. V., Yilma, M. A., Feliz, A., Majid, M. T., Maffetone, N., Walker, J. R., et al. (2019). A review of the microbial production of bioactive natural products and biologics. *Front. Microbiol.* 10:1404. doi: 10.3389/fmicb.2019.01404
- Pramastya, H., Xue, D., Abdallah, I. I., Setroikromo, R., and Quax, W. J. (2021). High level production of amorphadiene using *Bacillus subtilis* as an optimized terpenoid cell factory. *New Biotechnol.* 60, 159–167. doi: 10.1016/j.nbt.2020.10.007
- Primak, Y. A., Du, M., Miller, M. C., Wells, D. H., Nielsen, A. T., Weyler, W., et al. (2011). Characterization of a feedback-resistant mevalonate kinase from the archaeon *Methanosarcina mazei*. *Appl. Environ. Microbiol.* 77, 7772–7778. doi: 10.1128/AEM.05761-11
- Rico, J., Duquesne, K., Petit, J.-L., Mariage, A., Darii, E., Peruch, F., et al. (2019). Exploring natural biodiversity to expand access to microbial terpene synthesis. *Microb. Cell Factories* 18:23. doi: 10.1186/s12934-019-1074-4
- Ro, D.-K., Paradise, E. M., Ouellet, M., Fisher, K. J., Newman, K. L., Ndungu, J. M., et al. (2006). Production of the antimalarial drug precursor artemisinic acid in engineered yeast. *Nature* 440, 940–943. doi: 10.1038/nature04640

- Rodrigues, J. S., and Lindberg, P. (2021). Metabolic engineering of *Synechocystis* sp. PCC 6803 for improved bisabolene production. *Metab. Eng. Commun.* 12:e00159. doi: 10.1016/j.mec.2020.e00159
- Sánchez-Pascuala, A., Fernández-Cabezón, L., de Lorenzo, V., and Nikel, P. I. (2019). Functional implementation of a linear glycolysis for sugar catabolism in *Pseudomonas putida*. *Metab. Eng.* 54, 200–211. doi: 10.1016/j.mben.2019.04.005
- Santos-Merino, M., Torrado, A., Davis, G. A., Röttig, A., Bibby, T. S., Kramer, D. M., et al. (2021). Improved photosynthetic capacity and photosystem I oxidation via heterologous metabolism engineering in cyanobacteria. *Proc. Natl. Acad. Sci. U. S. A.* 118. doi: 10.1073/pnas.2021523118
- Schada von Borzyskowski, L., Sonntag, F., Pöschel, L., Vorholt, J. A., Schrader, J., Erb, T. J., et al. (2018). Replacing the Ethylmalonyl-CoA pathway with the glyoxylate shunt provides metabolic flexibility in the central carbon metabolism of *Methylobacterium extorquens* AM1. *ACS Synth. Biol.* 7, 86–97. doi: 10.1021/acssynbio.7b00229
- Schempp, F. M., Drummond, L., Buchhaupt, M., and Schrader, J. (2018). Microbial cell factories for the production of terpenoid flavor and fragrance compounds. *J. Agric. Food Chem.* 66, 2247–2258. doi: 10.1021/acs.jafc.7b00473
- Schrader, J., Schilling, M., Holtmann, D., Sell, D., Filho, M. V., Marx, A., et al. (2009). Methanol-based industrial biotechnology: current status and future perspectives of methylotrophic bacteria. *Trends Biotechnol.* 27, 107–115. doi: 10.1016/j.tibtech.2008.10.009
- Sebesta, J., and Peebles, C. A. (2020). Improving heterologous protein expression in *Synechocystis* sp. PCC 6803 for alpha-bisabolene production. *Metab. Eng. Commun.* 10:e00117. doi: 10.1016/j.mec.2019.e00117
- Soccol, C. R., Dalmas Neto, C. J., Soccol, V. T., Sydney, E. B., da Costa, E. S. F., Medeiros, A. B. P., et al. (2017). Pilot scale biodiesel production from microbial oil of *Rhodospiridium toruloides* DEBB 5533 using sugarcane juice: performance in diesel engine and preliminary economic study. *Bioresour. Technol.* 223, 259–268. doi: 10.1016/j.biortech.2016.10.055
- Song, Y., Guan, Z., van Merkerk, R., Pramastya, H., Abdallah, I. I., Setroikromo, R., et al. (2020). Production of squalene in *Bacillus subtilis* by squalene synthase screening and metabolic engineering. *J. Agric. Food Chem.* 68, 4447–4455. doi: 10.1021/acs.jafc.0c00375
- Song, Y., He, S., Abdallah, I. I., Jopkiewicz, A., Setroikromo, R., van Merkerk, R., et al. (2021). Engineering of multiple modules to improve amorphaadiene production in *Bacillus subtilis* using CRISPR-Cas9. *J. Agric. Food Chem.* 69, 4785–4794. doi: 10.1021/acs.jafc.1c00498
- Sonntag, F., Kroner, C., Lubuta, P., Peyraud, R., Horst, A., Buchhaupt, M., et al. (2015). Engineering *Methylobacterium extorquens* for de novo synthesis of the sesquiterpenoid  $\alpha$ -humulene from methanol. *Metab. Eng.* 32, 82–94. doi: 10.1016/j.mben.2015.09.004
- Sun, W., Xue, H., Liu, H., Lv, B., Yu, Y., Wang, Y., et al. (2020). Controlling chemo- and regioselectivity of a plant P450 in yeast cell toward rare licorice triterpenoid biosynthesis. *ACS Catal.* 10, 4253–4260. doi: 10.1021/acscatal.0c00128
- Sun, T., Yu, Y., Wang, K., Ledesma-Amaro, R., and Ji, X.-J. (2021). Engineering *Yarrowia lipolytica* to produce fuels and chemicals from xylose: a review. *Bioresour. Technol.* 337:125484. doi: 10.1016/j.biortech.2021.125484
- Sun, J., Zhang, C., Nan, W., Li, D., Ke, D., and Lu, W. (2018). Glycerol improves heterologous biosynthesis of betulinic acid in engineered *Yarrowia lipolytica*. *Chem. Eng. Sci.* 196, 82–90. doi: 10.1016/j.ces.2018.10.052
- Thomas, S. T., Louie, G. V., Lubin, J. W., Lundblad, V., and Noel, J. P. (2019). Substrate specificity and engineering of mevalonate 5-phosphate decarboxylase. *ACS Chem. Biol.* 14, 1767–1779. doi: 10.1021/acscchembio.9b00322
- Tian, T., Kang, J. W., Kang, A., and Lee, T. S. (2019). Redirecting metabolic flux via combinatorial multiplex CRISPRi-mediated repression for isopentenol production in *Escherichia coli*. *ACS Synth. Biol.* 8, 391–402. doi: 10.1021/acssynbio.8b00429
- Tsuruta, H., Paddon, C. J., Eng, D., Lenihan, J. R., Horning, T., Anthony, L. C., et al. (2009). High-level production of amorpha-4,11-diene, a precursor of the antimalarial agent artemisinin, in *Escherichia coli*. *PLoS One* 4:e4489. doi: 10.1371/journal.pone.0004489
- Vannice, J. C., Skaff, D. A., Keightley, A., Addo, J. K., Wyckoff, G. J., and Mizioroko, H. M. (2014). Identification in *Haloferax volcanii* of phosphomevalonate decarboxylase and isopentenyl phosphate kinase as catalysts of the terminal enzyme reactions in an archaeal alternate mevalonate pathway. *J. Bacteriol.* 196, 1055–1063. doi: 10.1128/JB.01230-13
- Vickers, C. E., Williams, T. C., Peng, B., and Cherry, J. (2017). Recent advances in synthetic biology for engineering isoprenoid production in yeast. *Curr. Opin. Chem. Biol.* 40, 47–56. doi: 10.1016/j.cbpa.2017.05.017
- Volke, D. C., Rohwer, J., Fischer, R., and Jennewein, S. (2019). Investigation of the methylerythritol 4-phosphate pathway for microbial terpenoid production through metabolic control analysis. *Microb. Cell Factories* 18:192. doi: 10.1186/s12934-019-1235-5
- von Reuss, S., Domik, D., Lemfack, M. C., Magnus, N., Kai, M., Weise, T., et al. (2018). Sodorifen biosynthesis in the rhizobacterium *Serratia plymuthica* involves methylation and cyclization of MEP-derived farnesyl pyrophosphate by a SAM-dependent C-methyltransferase. *J. Am. Chem. Soc.* 140, 11855–11862. doi: 10.1021/jacs.8b08510
- Vranová, E., Coman, D., and Grussem, W. (2012). Structure and dynamics of the isoprenoid pathway network. *Mol. Plant* 5, 318–333. doi: 10.1093/mp/sss015
- Wang, X., Liu, W., Xin, C., Zheng, Y., Cheng, Y., Sun, S., et al. (2016). Enhanced limonene production in cyanobacteria reveals photosynthesis limitations. *Proc. Natl. Acad. Sci. U. S. A.* 113, 14225–14230. doi: 10.1073/pnas.1613340113
- Wang, P., Wei, W., Ye, W., Li, X., Zhao, W., Yang, C., et al. (2019). Synthesizing ginsenoside Rh2 in *Saccharomyces cerevisiae* cell factory at high-efficiency. *Cell Discov.* 5:5. doi: 10.1038/s41421-018-0075-5
- Ward, V. C. A., Chatzivasileiou, A. O., and Stephanopoulos, G. (2018). Metabolic engineering of *Escherichia coli* for the production of isoprenoids. *FEMS Microbiol. Lett.* 365. doi: 10.1093/femsle/fny079
- Wichmann, J., Baier, T., Wentnagel, E., Lauersen, K. J., and Kruse, O. (2018). Tailored carbon partitioning for phototrophic production of (E)- $\alpha$ -bisabolene from the green microalga *Chlamydomonas reinhardtii*. *Metab. Eng.* 45, 211–222. doi: 10.1016/j.mben.2017.12.010
- Wong, J., de Rond, T., d Espaux, L., van der Horst, C., Dev, I., Rios-Solis, L., et al. (2018). High-titer production of lathyrane diterpenoids from sugar by engineered *Saccharomyces cerevisiae*. *Metab. Eng.* 45, 142–148. doi: 10.1016/j.mben.2017.12.007
- Wriessnegger, T., Augustin, P., Engleder, M., Leitner, E., Müller, M., Kaluzna, I., et al. (2014). Production of the sesquiterpenoid (+)-nootkatone by metabolic engineering of *Pichia pastoris*. *Metab. Eng.* 24, 18–29. doi: 10.1016/j.mben.2014.04.001
- Wu, W., and Maravelias, C. T. (2018). Synthesis and techno-economic assessment of microbial-based processes for terpenes production. *Biotechnol. Biofuels* 11:294. doi: 10.1186/s13068-018-1285-7
- Xiao, H., Zhang, Y., and Wang, M. (2019). Discovery and engineering of cytochrome p450s for terpenoid biosynthesis. *Trends Biotechnol.* 37, 618–631. doi: 10.1016/j.tibtech.2018.11.008
- Xie, D., Jackson, E. N., and Zhu, Q. (2015). Sustainable source of omega-3 eicosapentaenoic acid from metabolically engineered *Yarrowia lipolytica*: from fundamental research to commercial production. *Appl. Microbiol. Biotechnol.* 99, 1599–1610. doi: 10.1007/s00253-014-6318-y
- Yadav, V. G., De Mey, M., Lim, C. G., Ajikumar, P. K., and Stephanopoulos, G. (2012). The future of metabolic engineering and synthetic biology: towards a systematic practice. *Metab. Eng.* 14, 233–241. doi: 10.1016/j.mben.2012.02.001
- Yaegashi, J., Kirby, J., Ito, M., Sun, J., Dutta, T., Mirsiaghi, M., et al. (2017). *Rhodospiridium toruloides*: a new platform organism for conversion of lignocellulose into terpene biofuels and bioproducts. *Biotechnol. Biofuels* 10:241. doi: 10.1186/s13068-017-0927-5
- Yang, C., Gao, X., Jiang, Y., Sun, B., Gao, F., and Yang, S. (2016). Synergy between methylerythritol phosphate pathway and mevalonate pathway for isoprene production in *Escherichia coli*. *Metab. Eng.* 37, 79–91. doi: 10.1016/j.mben.2016.05.003
- Yang, J., Im, Y., Kim, T. H., Lee, M. J., Cho, S., Na, J.-G., et al. (2020). Engineering *Pseudomonas putida* KT2440 to convert 2,3-butanediol to mevalonate. *Enzym. Microb. Technol.* 132:109437. doi: 10.1016/j.enzmictec.2019.109437
- Yang, X., Jiang, Y., Yang, J., He, J., Sun, J., Chen, F., et al. (2015). Prenylated flavonoids, promising nutraceuticals with impressive biological activities. *Trends Food Sci. Technol.* 44, 93–104. doi: 10.1016/j.tifs.2015.03.007
- Yang, J., Son, J. H., Kim, H., Cho, S., Na, J.-G., Yeon, Y. J., et al. (2019). Mevalonate production from ethanol by direct conversion through acetyl-CoA using recombinant *Pseudomonas putida*, a novel biocatalyst for terpenoid production. *Microb. Cell Factories* 18:168. doi: 10.1186/s12934-019-1213-y

- Yao, F., Liu, S.-C., Wang, D.-N., Liu, Z.-J., Hua, Q., and Wei, L.-J. (2020). Engineering oleaginous yeast *Yarrowia lipolytica* for enhanced limonene production from xylose and lignocellulosic hydrolysate. *FEMS Yeast Res.* 20:foaa046. doi: 10.1093/femsyr/foaa046
- Ye, R.-P., Ding, J., Gong, W., Argyle, M. D., Zhong, Q., Wang, Y., et al. (2019). CO<sub>2</sub> hydrogenation to high-value products via heterogeneous catalysis. *Nat. Commun.* 10:5698. doi: 10.1038/s41467-019-13638-9
- Ye, R. W., Yao, H., Stead, K., Wang, T., Tao, L., Cheng, Q., et al. (2007). Construction of the astaxanthin biosynthetic pathway in a methanotrophic bacterium *Methylomonas* sp. strain 16a. *J. Ind. Microbiol. Biotechnol.* 34, 289–299. doi: 10.1007/s10295-006-0197-x
- Yishai, O., Lindner, S. N., Gonzalez de la Cruz, J., Tenenboim, H., and Bar-Even, A. (2016). The formate bio-economy. *Curr. Opin. Chem. Biol.* 35, 1–9. doi: 10.1016/j.cbpa.2016.07.005
- Yook, S. D., Kim, J., Gong, G., Ko, J. K., Um, Y., Han, S. O., et al. (2020). High-yield lipid production from lignocellulosic biomass using engineered xylose-utilizing *Yarrowia lipolytica*. *Glob. Change Biol. Bioenergy.* 12, 670–679. doi: 10.1111/gcbb.12699
- Yoshida, R., Yoshimura, T., and Hemmi, H. (2020). Reconstruction of the “Archaeal” Mevalonate pathway from the Methanogenic Archaeon *Methanosarcina mazei* in *Escherichia coli* cells. *Appl. Environ. Microbiol.* 86, e02889–e02819. doi: 10.1128/AEM.02889-19
- Yu, F., Okamoto, S., Harada, H., Yamasaki, K., Misawa, N., and Utsumi, R. (2011). Zingiber zerumbet CYP71BA1 catalyzes the conversion of  $\alpha$ -humulene to 8-hydroxy- $\alpha$ -humulene in zerumbone biosynthesis. *Cell. Mol. Life Sci.* 68, 1033–1040. doi: 10.1007/s00018-010-0506-4
- Yuan, L. Z., Rouvière, P. E., Larossa, R. A., and Suh, W. (2006). Chromosomal promoter replacement of the isoprenoid pathway for enhancing carotenoid production in *E. coli*. *Metab. Eng.* 8, 79–90. doi: 10.1016/j.ymben.2005.08.005
- Zhang, X., Guan, H., Dai, Z., Guo, J., Shen, Y., Cui, G., et al. (2015). Functional analysis of the isopentenyl diphosphate isomerase of *Salvia miltiorrhiza* via color complementation and RNA interference. *Molecules* 20, 20206–20218. doi: 10.3390/molecules201119689
- Zhang, C., Liu, J., Zhao, F., Lu, C., Zhao, G.-R., and Lu, W. (2018). Production of sesquiterpenoid zerumbone from metabolic engineered *Saccharomyces cerevisiae*. *Metab. Eng.* 49, 28–35. doi: 10.1016/j.ymben.2018.07.010
- Zhang, X., Wang, D., Duan, Y., Zheng, X., Lin, Y., and Liang, S. (2020). Production of lycopene by metabolically engineered *Pichia pastoris*. *Biosci. Biotechnol. Biochem.* 84, 463–470. doi: 10.1080/09168451.2019.1693250
- Zheng, Y., Liu, Q., Li, L., Qin, W., Yang, J., Zhang, H., et al. (2013). Metabolic engineering of *Escherichia coli* for high-specificity production of isoprenol and prenol as next generation of biofuels. *Biotechnol. Biofuels* 6:57. doi: 10.1186/1754-6834-6-57
- Zhou, F., and Pichersky, E. (2020). More is better: the diversity of terpene metabolism in plants. *Curr. Opin. Plant Biol.* 55, 1–10. doi: 10.1016/j.pbi.2020.01.005
- Zhou, K., Qiao, K., Edgar, S., and Stephanopoulos, G. (2015). Distributing a metabolic pathway among a microbial consortium enhances production of natural products. *Nat. Biotechnol.* 33, 377–383. doi: 10.1038/nbt.3095
- Zhou, K., Zou, R., Zhang, C., Stephanopoulos, G., and Too, H.-P. (2013). Optimization of amorphaadiene synthesis in *Bacillus subtilis* via transcriptional, translational, and media modulation. *Biotechnol. Bioeng.* 110, 2556–2561. doi: 10.1002/bit.24900
- Zu, Y., Prather, K. L., and Stephanopoulos, G. (2020). Metabolic engineering strategies to overcome precursor limitations in isoprenoid biosynthesis. *Curr. Opin. Biotechnol.* 66, 171–178. doi: 10.1016/j.copbio.2020.07.005

**Conflict of Interest:** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Publisher’s Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Carruthers and Lee. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.