



OPEN ACCESS

EDITED BY

Huancai Lin,
Sun Yat-sen University, China

REVIEWED BY

Jintae Lee,
Yeungnam University,
South Korea
Jules-Roger Kuate,
University of Dschang, Cameroon

*CORRESPONDENCE

Hans P. Steenackers
hans.steenackers@kuleuven.be

SPECIALTY SECTION

This article was submitted to
Antimicrobials, Resistance and
Chemotherapy, a section of the journal
Frontiers in Microbiology

RECEIVED 05 July 2022

ACCEPTED 18 November 2022

PUBLISHED 06 January 2023

CITATION

Villanueva X, Zhen L, Ares JN, Vackier T,
Lange H, Crestini C and
Steenackers HP (2023) Effect of chemical
modifications of tannins on their
antimicrobial and antibiofilm effect against
Gram-negative and Gram-positive bacteria.
Front. Microbiol. 13:987164.
doi: 10.3389/fmicb.2022.987164

COPYRIGHT

© 2023 Villanueva, Zhen, Ares, Vackier,
Lange, Crestini and Steenackers. This is an
open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that
the original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

Effect of chemical modifications of tannins on their antimicrobial and antibiofilm effect against Gram-negative and Gram-positive bacteria

Xabier Villanueva¹, Lili Zhen^{2,3}, José Nunez Ares⁴, Thijs Vackier¹, Heiko Lange^{3,5}, Claudia Crestini^{3,6} and Hans P. Steenackers^{4*}

¹Faculty of Bioscience Engineering, Centre of Microbial and Plant Genetics (CMPG), KU Leuven, Heverlee, Belgium, ²Department of Chemical Science and Technologies, University of Rome 'Tor Vergata', Rome, Italy, ³CSGI – Center for Colloid and Surface Science, Sesto Fiorentino, Italy, ⁴Division of Mechatronics, Biostatistics and Sensors (MeBioS), Department of Biosystems (BIOSYST), KU Leuven, Heverlee, Belgium, ⁵Department of Earth and Environmental Sciences, University of Milano-Bicocca, Milan, Italy, ⁶Department of Molecular Science and Nanosystems, Ca' Foscari University of Venice, Venice, Italy

Background: Tannins have demonstrated antibacterial and antibiofilm activity, but there are still unknown aspects on how the chemical properties of tannins affect their biological properties. We are interested in understanding how to modulate the antibiofilm activity of tannins and in delineating the relationship between chemical determinants and antibiofilm activity.

Materials and methods: The effect of five different naturally acquired tannins and their chemical derivatives on biofilm formation and planktonic growth of *Salmonella* Typhimurium, *Pseudomonas aeruginosa*, *Escherichia coli* and *Staphylococcus aureus* was determined in the Calgary biofilm device.

Results: Most of the unmodified tannins exhibited specific antibiofilm activity against the assayed bacteria. The chemical modifications were found to alter the antibiofilm activity level and spectrum of the tannins. A positive charge introduced by derivatization with higher amounts of ammonium groups shifted the anti-biofilm spectrum toward Gram-negative bacteria, and derivatization with lower amounts of ammonium groups and acidifying derivatization shifted the spectrum toward Gram-positive bacteria. Furthermore, the quantity of phenolic OH-groups per molecule was found to have a weak impact on the anti-biofilm activity of the tannins.

Conclusion: We were able to modulate the antibiofilm activity of several tannins by specific chemical modifications, providing a first approach for fine tuning of their activity and antibacterial spectrum.

KEYWORDS

tannins, antibiofilm activity, structure–activity relationships, *Salmonella* Typhimurium, *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*

Introduction

Plant-derived tannins have been used from ancient times in leather industry because of their ability of making leather last over time, and the name “tannin” comes from the use of these chemicals for “tanning” leather (Falcão and Araújo, 2011; Pizzi, 2019). Accordingly, it was hypothesized that the resistance of leather to microbial decomposition could be explained by this use of tannin-rich compounds in leather curation and several studies have pointed to multiple additional pharmacological properties of tannins, including anti-inflammatory and anti-cancer effects, which are contributing to a renewed interest in these products as a source of new bio-based pharmaceuticals (Widsten et al., 2014; Vu et al., 2017; Pizzi, 2019; Farha et al., 2020). Tannins have been shown to inhibit bacterial growth of different Gram-positive and Gram-negative bacteria (Taguri et al., 2004; Ekambaram et al., 2016; Slobodníková et al., 2016; Chandra et al., 2017; Vu et al., 2017), and are shown to be able to disperse biofilms (Trentin et al., 2013). On some occasions this antibiofilm activity is specific and independent from the ability to inhibit bacterial growth (Ulrey et al., 2014). Common examples of tannins with antibacterial activity are tannic acid (Chung et al., 1998), ellagic acid (De et al., 2018) and epigallocatechin gallate (Akiyama, 2001).

From a molecular point of view, tannins can be divided in two main groups: condensed and hydrolysable tannins. Hydrolysable tannins are esters of gallic acid with a core sugar, often glucose or quinic acid. Tannic acid is the most prominent representative of the family of hydrolysable tannins comprising a glucose center (Koleckar et al., 2008; Ekambaram et al., 2016). Condensed tannins are oligomeric and polymeric proanthocyanidins, consisting of flavan-3-ol units, linked by carbon-carbon bonds not susceptible to hydrolytic cleavage (Koleckar et al., 2008; Versari et al., 2013). The scaffold of the subclass of tannins called complex tannins is very similar to those found in condensed and hydrolysable tannins, where a flavan-3-ol unit is linked to gallic acid in a monomeric or polymeric system (Falcão and Araújo, 2011). However, there is no reference in literature that this kind of differentiation has any effect on the level and kind of antimicrobial activity of the tannins.

To date, there is no clear understanding on the factors that explain the antimicrobial activity of tannins. It has been suggested that the observable activity could be explained by the presence of free phenolic hydroxyl groups which can affect, for example, enzymatic activity *via* covalent or non-covalent linking (Scalbert, 1991). In this respect, it has been seen that phenolic compounds can have antimicrobial effects against *Pseudomonas aeruginosa* and *Staphylococcus aureus* (Jagani et al., 2009; Lahiri et al., 2019). This ultimately means that the typical phenolic character of the tannins could play an important role for the antimicrobial activity (Mori et al., 1987; Baba and Malik, 2015). Other mechanisms of action for the antimicrobial activity of tannins have also been described, in particular for tannic acid, like disruption of peptidoglycan formation (Dong et al., 2018), iron chelation (Chung et al., 1998), membrane disruption (Kim et al., 2010), efflux pump inhibition (Tintino et al., 2016) and fatty acid synthesis (Wu et al., 2010).

It has also been shown in previous literature that tannins have the ability to reduce biofilm formation (Jagani et al., 2009; Klug et al., 2017; Dettweiler et al., 2019; Lahiri et al., 2019). Biofilms are conglomerates of bacteria, usually at an interphase (solid-air, solid-liquid, liquid-air), that are surrounded by a protective mesh of extracellular polymeric substances (EPS). This enhances the ability of the bacteria to survive dehydration, disinfectants and antibiotics (Rabin et al., 2015; Roy et al., 2018). The mechanisms of protection include reduced penetration of antimicrobial compounds, reduced bacterial metabolism, induction of efflux pumps and more frequent horizontal gene transfer (Rabin et al., 2015).

Regarding the antibiofilm activity of tannins, it has been described that some of them have a biofilm-specific mechanism of action, such as inhibition of quorum sensing (QS) in *P. aeruginosa* by the tannin-rich fractions of *Terminalia catappa* (Taganna et al., 2011) and *T. chebulata* (Sarabhai et al., 2013), and induction of transglucosylase activity in *S. aureus* by tannic acid (Payne et al., 2013). This type of biofilm-specific behavior is actually desired, because the lack of direct growth inhibition decreases the selective pressure toward resistance phenotypes (Allen et al., 2010; Imperi et al., 2019; Dieltjens et al., 2020).

Because of this reduced potential of resistance development, in the current study we evaluated selected chemical variants of tannins for their ability to inhibit biofilm formation without inhibiting planktonic growth. A diverse range of five commercially obtained tannins, three condensed tannins and two hydrolysable tannins, were included. The tannins were selected because they have been structurally thoroughly characterized in prior works and they are representative of the known molecular diversity of tannins. Also, we investigated how different chemical modifications change the activity level and spectrum of the tannins. Understanding the effect of structural features on activity allows to enhance or finetune activity. The functional groups chosen for derivatization were selected for known capacity (i) to interfere with membranes, i.e., ammonium groups; and (ii) to alter modes of interactions with surfaces, i.e., carboxylic acids and poly(ethylene glycol). The objective was to study the consequences of such chemical derivatizations at the molecular level and at the level of intermolecular interaction possibilities and obtain hints regarding matches and mismatches between tannin-type and tannin functionalization with respect to tuning anti-biofilm activities. The scarcity of structure-activity relationship (SAR) research in the field of tannins and bioactive phenolic compounds from plant sources as antimicrobial compounds highlights the value of this work (Fang et al., 2016; Bouarab-Chibane et al., 2019).

Materials and methods

Tannin assays and modifications

Assayed tannins

Five commercially available tannin extracts, comprising three condensed and two hydrolysable tannins were used. The three

condensed tannins were Omnivin 20R (monomeric (epi) catechin, **Vv-20**), Omnivin WG (procyanidins (62%)/proflisetidins (34%), **Vv**) and Mimosa ATO ME (prorobinetidins (33%)/proflisetidins (67%), **Am**), and the two hydrolysable tannins were Tanal 01 (tannic acid, **Ta-01**) and Tanal 04 (galloylquinic acid, **Ta-04**). The chemical structures of these tannins were elucidated in detail as reported elsewhere (Zhen et al., 2021a). Figure 1A gives an overview of the structural features.

Modification of tannins

The five selected tannins were chemically modified by derivatizing them *via* their phenolic functionalities, as reported in detail elsewhere (Zhen et al., 2021b), with different levels of specific functional motifs: (i) hydroxy-*N,N,N*-trimethylpropanyl-3-aminium chloride (C_3NMe_3Cl -eq), (ii) hydroxypropyl-1-carboxylic acid (C_3COOH -eq), and (iii) oligomeric ethylene glycol polyether (PEG₅₀₀-eq), whereby “eq.” in the compounds listed in Figure 1B indicates the equivalents of the functional motif that were used for the chemical modification (Zhen et al., 2021b). These chemical functionalizations gave several properties to the tannins: (i) C_3NMe_3Cl -eq added positive charges to the tannin molecule, (ii) C_3COOH -eq, as a weak acid, potentially added negative charges to the molecule, and (iii) PEG₅₀₀-eq polymerized the tannin molecules. During the various functionalizations, control tannins were re-isolated from blank reactions. Tannins labeled as “Blank-W” are tannins isolated from blank reactions performed in water and tannins labeled as “Blank-D” are tannins isolated from blank reaction performed in dimethylformamide.

Because of the low solubility of the tannins in aqueous media, the dry compounds were first suspended in dimethyl sulfoxide (DMSO) at a stock concentration of 60 g/L and from there diluted to the desired concentration in the following experiments. A 1% v/v concentration of DMSO was never exceeded in order to prevent potential effects of DMSO on bacterial growth or biofilm formation.

All the tests were done in aerobic conditions in growth media with a pH of ~7.4 and a salinity range of 0.025–0.5% w/v.

Study of antibiofilm activities of tannins

The antibiofilm activities of the tannins were determined using 96-well plates with a cover lid with pegs for growing the bacterial biofilms in presence of several concentration of the assayed tannins. These assays, divided in a 1st exploratory screening (Section “Antibiofilm screening”) and a 2nd validation experiment (Section “Validation anti-biofilm screening: Experimental design and statistical analysis”), are explained in detail in the following section.

Generation of biofilms of each microorganism

Biofilms of *Salmonella enterica* var. Typhimurium ATCC14028, *Pseudomonas aeruginosa* PA14, *Escherichia coli* TG1 and *Staphylococcus aureus* SH1000 were grown in the Calgary biofilm

device *via* a protocol that was previously described (Janssens et al., 2008; Steenackers et al., 2011). The bacteria were grown overnight (ON) in LB broth at 37°C. These ON cultures were then diluted 1/100 in diluted (1/20) Tryptic Soy Broth (TSB, Thermo Fisher Scientific) for Gram-negative bacteria and in undiluted TSB for *S. aureus*. 100 µl of growth medium or a solution of the tannins in growth medium was added to the wells of a 96-well Calgary device (Nunc nos. 269789 and 445497, Thermo Fisher Scientific). The diluted ON cultures were then added to the wells to obtain a starting inoculum of 10⁶ cfu/ml in a final volume of 200 µl of growth medium per well. Also, both for the exploratory screening test and validation of anti-biofilm activity, one row of the plate was filled with inoculum with growth media without tannin and another row was filled with media without bacteria. To account for potential effects of the compounds themselves, the same tannin-concentration was added to separate control 96-well Calgary biofilm plates without the inoculation of the bacteria. Afterwards, all plates were incubated in a wet chamber for the appropriate time and temperature: 48 h at 37°C for *S. aureus*, 24 h at 25°C for the Gram-negatives.

Biofilm formation and planktonic growth were determined by crystal violet staining of the pegs and OD₆₀₀ measurements of the base plate of the device, respectively. Specifically, after incubation the covers of the plates (which contain the pegs) were removed and washed once with PBS. The pegs were then stained with 200 µl per well of 0.1% v/v of crystal violet (CV, VWR International) for 30 min. After staining, the excess of CV was washed once with 200 µl per well of distilled water and let dry for 30 min. Finally, the CV was recovered in a new 96-well plate with 200 µl per well of 30% v/v glacial acetic and the optical density at 570 nm (OD₅₇₀) measured in a plate reader. The optical density at 600 nm (OD₆₀₀) of the bacteria in the base plates was measured to determine the growth of the planktonic cells.

Antibiofilm screening

In a preliminary experiment, the tannins were tested in two-fold dilution series ranging from 600 to 9.38 mg/L in 3 independent replicates. The obtained raw OD₅₇₀ (for the biofilm formation) and OD₆₀₀ (for the planktonic growth) were corrected by the OD of the tannins incubated in absence of bacterial inoculum and then normalized using the average OD of the bacterial inoculum incubated in absence of tannin, thus being converted to percentage of biofilm formation (OD₅₇₀) and percentage of bacterial growth (OD₆₀₀).

The BIC₅₀ and IC₅₀ (the compound concentration required to inhibit the biofilm formation and the bacterial growth by 50%, respectively) were calculated by applying a log [tannin] vs. percentage of biofilm formation or percentage of planktonic growth non-linear regression (four-parameters) using the statistical package GraphPad 8.0.

Validation anti-biofilm screening: Experimental design and statistical analysis

Based on the information from the preliminary experiment, a definitive antibiofilm experiment in the Calgary biofilm device




FIGURE 1

(A) Chemical structure of commercially available condensed and hydrolysable tannins used in this study (40) and (B) their chemically derivatized structures as described elsewhere (41). Exemplary structural aspects are shown; synthetic route leads to generation of both primary and secondary aliphatic alcohols within the total C_3 linker moiety connecting the functional to the tannin (41). (i) C_3NMe_3Cl -eq – hydroxy-*N,N,N*-trimethylpropanyl-3-amminium chloride; (ii) C_3CO_2H -eq – hydroxypropyl-1-carboxylic acid; and (iii) PEG_{500} -eq – oligomeric ethylene glycol polyether (PEG_{500} -eq).

was set up with eight independent repeats. We first defined the factors under study, i.e., the parameters that potentially have an influence on the formation of biofilm. The considered parameters are: (i) original unmodified tannin: **Vv-20**, **Vv**, **Am**, **Ta-01**, **Ta-04**; (ii) concentration of tannin (mg/L): 9.38, 79.69, 150; (iii) chemical

modifier: C_3CO_2H , PEG, C_3NMe_3Cl , Blank-D (blank reaction with dimethylformamide), Blank-W (blank reaction with water), unmodified; and (iv) concentration of applied chemical modifier: low and high (low: 0.05 and 0.1 eq of chemical substitution; high: 0.25 and 0.5 eq of chemical substitution). In order to minimize

effects of plate-to-plate variation, we applied an optimal randomized experimental design with the statistical package JMP 15.0 to reduce experimental noise and confounding factors. Different to the preliminary experiment, only three concentrations were assayed, which were chosen to best capture the antibiofilm activity of the tannins: 9.38, 79.69, and 150 mg/L. The remaining part of the protocol of this experiment is identical to that of the preliminary experiment described above.

To determine the effect of the chemical derivatizations on the antibiofilm and antibacterial effect of the unmodified tannins on each of the assayed bacteria, an ANOVA test with Tukey post-hoc test comparing the unmodified tannins with their respective derivatized tannins was done based on the obtained biofilm formation and planktonic growth levels.

Relationship between the phenolic hydroxyl content and antibiofilm effect

To determine the effect of phenolic hydroxyl (OH) content on the antimicrobial and antibiofilm effect of the different tannins, a simple linear regression between the phenolic OH content of the tannins and the level of biofilm formation (or planktonic growth) was performed. To better calculate this correlation, we used the different tannin assayed concentrations and the obtained mmol of phenolic OH per gram of material to calculate the mmol of phenolic OH present in the system for each tannin at each assayed concentration, and we correlated this value with the respective percentage of biofilm inhibition and planktonic growth inhibition. The phenolic OH content of the tannins was determined *via* ³¹P NMR as described elsewhere (Zhen et al., 2021a).

Results and discussion

Exploratory screening to determine the concentration test range

A diverse range of five commercially obtained tannins, and five types of derivatives, were selected and screened for antibiofilm activity. The condensed tannins were previously identified as mixtures of (epi) catechins and fisetinidols (Zhen et al., 2021a) and consisted of (i) the essentially monomeric **Vv-20**, (ii) the low oligomeric **Vv** and (iii) the higher oligomeric **Am**. The hydrolysable tannins consisted of two large tannins: (i) the “typical” tannic acid **Ta-01** and (ii) the galloquinic acid derivative **Ta-04**. All tannins were previously functionalized with a positive charge introducing ammonium salt hydroxy-*N,N,N*-trimethylpropanyl-3-aminium chloride (C₃NMe3Cl-0.1 and C₃NMe3Cl-0.5), an acidifying hydroxypropyl-1-carboxylic acid (C₃COOH-0.1 and C₃COOH-0.1-0.5) motive and a polymerizing oligomeric ethylene glycol polyether (PEG₅₀₀-0.05; Figure 1). The preventive activity of both the natural and derivatized tannins against the biofilm formation and planktonic growth of the Gram-negative species

S. Typhimurium, *P. aeruginosa*, *E. coli* and the Gram-positive species *S. aureus* was evaluated by means of the Calgary biofilm device. In a first set of exploratory experiments, two-fold serial dilutions (from 600 till 4.96 mg/L) of the tannins were evaluated in order to obtain a first glance on activity spectrum and active concentration range. Activities against all four bacterial species were observed, with BIC₅₀ values ranging from 4.69 to 545.8 mg/L and IC₅₀ values ranging from 37.5 to 459.2 mg/L (Supplementary Table S1). As such this experiment allowed to determine the test concentrations for future validation experiments: 9.38 mg/L was the lowest assayed concentration in the preliminary screening and the most active tannins exhibited BIC₅₀ equal to or below that value; 150 mg/L was the concentration at which almost all tannins with antibiofilm effect were active; and 79.69 mg/L is the average of those two concentrations. Such validation experiments were required because the exploratory experiments only had three independent repeats and this did not provide sufficient statistical power to distinguish the levels of activity of the different tannins and delineate the relationship between the chemistry and the antibiofilm level of the tannins. Furthermore, there was a statistically significant plate-to-plate variation between the controls of each plate (see Supplementary Figures S1, S2).

Extensive randomized validation screening at limited number of concentrations

To allow a multivariate analysis considering tannin scaffold, derivatization and concentration as well as bacterial target species, the previous antibiofilm and antimicrobial experiments were repeated in one experiment with eight repeats per condition, but only focusing on the three tannin concentrations that could capture best the antibiofilm effect of the tannins: 9.38, 79.69, and 150 mg/L. In order to minimize previously observed effects of plate-to-plate variation, these experiments were designed in a randomized way, i.e., all the tannins were distributed through all the plates in a random fashion. This allowed to decrease the random error and the possibility of confounding factors, a necessary step for doing a complex statistical analysis that allows to link the different chemical characteristics. In what follows we will first focus on the unmodified tannins, after which we will elaborate on the effect of chemical derivatization.

To determine the percentual inhibition of biofilm formation and planktonic growth of the assayed tannins against the selected bacteria, the bacterial growth and biofilm development in absence of tannin was taken as reference. As can be seen in Supplementary Figure S3, the assayed bacteria exhibited differences in their biofilm formation and bacterial growth, *Staphylococcus aureus* being the bacterium that reached the highest planktonic population density and formed most biofilm. On the contrary, the assayed Gram-negative bacteria showed less biofilm and less planktonic growth in comparison with *S. aureus*.

Effect of unmodified tannin scaffold on the antibiofilm activity level and spectrum

In [Figure 2](#) it can be seen that the commercially available natural tannins showed different antibiofilm activities against the four assayed bacterial species. All unmodified tannins can be considered to have “broad spectrum activity,” since all of them exhibited statistically significant antibiofilm activity against Gram-positive and Gram-negative bacteria at least at the concentration of 150 mg/L, according to an ANOVA test with Tukey post-hoc analysis. Also, most of the assayed tannins exhibited a concentration-dependent activity, depending on the assayed tannin and the tested bacteria.

However, there were clear differences in the degree of antibiofilm activity of each tannin. Starting from the condensed tannins, the monomeric **Vv-20** exhibited significant antibiofilm activity against all the assayed bacteria at all the assayed concentrations in a clear dose-dependent way, a dose of 79.69 mg/L being sufficient for inhibiting biofilm formation more than 50% against the four assayed bacteria. Also, **Vv-20** was the most effective tannin against *S. Typhimurium*, with more than 75% of biofilm inhibition at 79.69 and 150 mg/L, and against *E. coli*, with more than 80% biofilm inhibition at 79.69 mg/L. Regarding the low oligomeric **Vv**, the antibiofilm activity was preferential against *P. aeruginosa* and *S. aureus*, for which it exhibited potent dose-dependent antibiofilm activity. This compound was able to inhibit more than 50% of biofilm formation by *S. aureus* both at 79.69 and 150 mg/L, and is the unmodified tannin with highest effect against *P. aeruginosa*, displaying antibiofilm activity of more than 90% at 79.69 mg/L and more than 95% at 150 mg/L. On the contrary, **Vv** was less effective against *S. Typhimurium* and *E. coli* and was able to inhibit <30% of biofilm formation of both bacterial species, regardless of the assayed concentration. Contrary to the previous two unmodified condensed tannins, the high oligomeric **Am** had preferential antibiofilm activity against *S. Typhimurium* and *S. aureus*, but also had significant antibiofilm activity against *P. aeruginosa*. Moreover, **Am** is the unmodified tannin with the highest antibiofilm activity against *S. aureus*, with more than 85% of antibiofilm activity at 79.69 and 150 mg/L, also inhibiting more than 50% of biofilm formation of *S. Typhimurium* both at 79.69 and 150 mg/L, and inhibiting *P. aeruginosa* biofilm formation in more than 30% at 79.69 mg/L and more than 50% at 150 mg/L. On the contrary, **Am** had unnoticeable inhibitory activity against *E. coli* biofilms at any of the assayed concentrations.

With respect to the hydrolysable tannins, tannic acid, **Ta-01** exhibited preferential activity against *P. aeruginosa* and *S. Typhimurium*. The biofilm inhibitory activity of **Ta-01** ranged from 40% at 9.38 mg/L to 60% at 150 mg/L against *S. Typhimurium*, and from 15% at 9.38 mg/L to more than 90% at 150 mg/L against *P. aeruginosa*. On the contrary, the galloquinic acid derivative, **Ta-04** exhibited preferential activity against *S. aureus*. While the antibiofilm activity of **Ta-04** against *S. Typhimurium* and *E. coli*

was dose dependent (reaching a maximum of 50% biofilm inhibition against both bacteria at a concentration of 150 mg/L), the antibiofilm activity of **Ta-04** against *S. aureus* was not dose dependent, with more than 70% biofilm inhibition at the three assayed concentrations.


Importantly, the unmodified tannins were in general not found to have antibacterial activity against the planktonic bacteria, except for the low oligomeric condensed **Vv** against *S. Typhimurium* and the hydrolysable tannic acid **Ta-01** against *E. coli* at the highest concentration of compound ([Figure 2](#)). In previous reports, it was found that hydrolysable tannins similar to galloylquinic acid, hence similar to **Ta-04**, exhibit broad spectrum antibiofilm activity ([Widsten et al., 2014](#); [Ekambaram et al., 2016](#); [Puljula et al., 2020](#)). Those same reports, however, also suggest that hydrolysable tannins have antibacterial effects against planktonic Gram positive and Gram bacteria, which was not observed in our experiment. On the other hand, tannic acid ([Lee et al., 2013](#)) and 1,2,3,4,6-penta-O-galloyl- β -D-glucopyranose ([Lin et al., 2011](#)), both hydrolysable tannins, have been reported to inhibit biofilm formation of *S. aureus* without inhibiting planktonic growth, which is consistent with the results observed for tannic acid **Ta-01**. In our assay, also galloylquinic acid **Ta-04** showed such activity. This selective activity against biofilms offers opportunities for potential applications, such as the titanium-tannin composite coating for implants developed by [Shukla and Bhatena \(2015\)](#), that allows sustained release of the tannin.

Effect of chemical substitutions on the antibiofilm and antibacterial activity of the tannins

The aim of the modifications was to partially functionalize *via* ether linkages the phenolic OH-groups of each tannin molecule and to add tannin-alien functionalities at various levels to test the possibility to modulate the native activity of tannins toward biofilms and planktonic bacteria.

[Figure 3](#) shows the effect of derivatization on the antibiofilm activity. The differences in antibiofilm activity between the different modifications were assessed *via* the Tukey test for multiple comparisons, by comparing if there were differences in the maximum activity (i.e., the antibiofilm activity at the highest assayed concentration). If there were no differences in the maximum activity, the activity at lower concentrations was also evaluated, thus allowing to determine if a derivatization was able to obtain the same effect as the unmodified tannin, but at a lower concentration.

As a general conclusion, it could be established that most of the chemical derivatizations, but especially positive charge introducing **C₃NMe₃Cl-0.1** reduce the antibiofilm activity of the tannins, while some of them can shift the activity spectrum toward preferential activity against Gram-positive or Gram-negative bacteria. More importantly, it must be noted that, in general, the chemical derivatizations increased the activity against



planktonic bacteria and decreased the activity against bacterial biofilms of the unmodified tannins, thus decreasing the biofilm specificity of the assayed tannins (see [Supplementary Figures S4–S7](#) for a direct comparison between growth inhibitory effect and biofilm inhibitory capability of the assayed tannins against each bacterial species). This effect will be discussed in more detail in Section “Biofilm selectivity of the antibiofilm effect of the assayed tannins.”

Two derivatizations generally shifted the antibiofilm spectrum toward the Gram-negative group of bacteria: polymerizing PEG₅₀₀-0.05 and positive charge-introducing C₃NMe₃Cl-0.5. Contrarily, derivatization with acidifying C₃COOH-0.1 and C₃COOH-0.5 in general decreased the antibiofilm activity against Gram-negative bacteria, while it retained activity against *S. aureus* for larger tannins *Am*, *Ta-01* and *Ta-04*, and increased activity against *S. aureus* for monomeric and low oligomeric *Vv-20* and


Vv. The other derivatizations in general decreased the activity against *S. aureus*. However, this effect of derivatizations on the spectrum and potency was highly dependent on the specific tannin submitted to the derivatization.

In more detail, it can be seen that the blank reaction (both in water -Blank-W- and in dimethylformamide -Blank-D-) already modified the antibiofilm effect of the assayed tannins, a phenomenon that could be attributed to removal of impurities that affect the antibiofilm effect of the tannins. It can be seen that Blank-W conditions increased the maximum antibiofilm effect of Vv against *Salmonella* Typhimurium, but decreased the maximum antibiofilm effect of Vv-20 and Vv against *P. aeruginosa* and decreases the antibiofilm effect of Vv-20 against *E. coli* at 79.69 mg/L without affecting its maximum antibiofilm effect. It can also be seen that Blank-D conditions only affected the

antibiofilm activity of Ta-04, by reducing its antibiofilm effect against Gram-negative bacteria.

Regarding actual substitutions, it can be seen that the effect of a derivatization with positive charge introducing ammonium groups, i.e., C₃NMe₃Cl, was different depending on the equivalents of chemical substitution.

On the one hand, C₃NMe₃Cl-0.5, the high equivalent derivatization, decreased the maximum antibiofilm effect of condensed tannins against *S. aureus* but did not affect the anti-staphylococcal effect of hydrolysable tannins. However, C₃NMe₃Cl-0.5 derivatization had a tannin-dependent effect against Gram-negative bacteria. It did not affect the antibiofilm effect of monomeric Vv-20 against any Gram-negative bacteria but increased the maximum antibiofilm effect of low oligomeric Vv against *Salmonella* Typhimurium while decreasing the



maximum antibiofilm effect of **Vv** against *P. aeruginosa*. It increased the effect of high oligomeric **Am** against all the Gram-negative bacteria at 79.69 mg/L without significantly changing the maximum effect compared to the unmodified tannin. It increased the maximum effect of tannic acid **Ta-01** against *E. coli* and *Salmonella* Typhimurium but decreased the effect against *P. aeruginosa* at 79.69 mg/L and it significantly decreased the maximum antibiofilm effect of galloquinic acid derivative **Ta-04** against *Salmonella* Typhimurium. On the other hand, $C_3NMe_3Cl-0.1$, the low equivalent derivatization, reduced the maximum effect of all tannins (with the exception of the galloquinic acid derivative **Ta-04**, which was not affected) against Gram-negative bacteria without affecting the antibiofilm effect against *S. aureus*.

Contrary to C_3NMe_3Cl , there were no big differences between different levels of derivatizations with acidifying motif CH_3COOH , i.e., $CH_3COOH-0.1$ and $CH_3COOH-0.5$, in terms of their impact on the activities compared to the unmodified tannins. For *S. aureus*, neither derivatization level affected the antibiofilm effect exerted by larger tannins **Am**, **Ta-01**, and **Ta-04**, comparable to the $C_3NMe_3Cl-0.1$ derivatization, while $CH_3COOH-0.1$ increased the maximum effect of low oligomeric **Vv**. $CH_3COOH-0.5$ had a similar effect on essentially monomeric **Vv-20**. For the Gram-negative bacteria, both derivatizations equally decreased the maximum antibiofilm effect of condensed tannins **Vv-20** and **Am**, as well as for hydrolysable tannin **Ta-04** against *S. Typhimurium*, but only $CH_3COOH-0.1$ significantly decreased the maximum antibiofilm effect against *S. Typhimurium* of **Vv**. While both derivatization levels drastically decreased the maximum antibiofilm effect of condensed tannins against *P. aeruginosa*, only $CH_3COOH-0.1$ decreased the maximum antibiofilm effect of tannic acid **Ta-01**. Neither derivatization level significantly affected the antibiofilm effect of hydrolysable **Ta-04**. This effect was similar regarding *E. coli* because neither derivatization level affected the maximum antibiofilm effect of hydrolysable tannins, while $CH_3COOH-0.1$ decreased the maximum antibiofilm effect of all condensed tannins. $CH_3COOH-0.5$ only significantly decreased the maximum antibiofilm effect of **Vv-20** against *E. coli*.

Regarding derivatization with polymerizing PEG_{500} , it can be seen that $PEG_{500}-0.05$ in general did not affect the antibiofilm effect of the assayed tannins. The only exceptions are the increase in the maximum antibiofilm effect of low oligomeric **Vv** against *Salmonella* Typhimurium and the decrease in the antibiofilm effect of monomeric **Vv-20** against *S. aureus* at 79.69 mg/L without changing the maximum antibiofilm effect. It has to be taken in account that, due to technical issues, it was not possible to analyze the effect of polymerizing $PEG_{500}-0.05$ derivatization on **Ta-04**. Polymerizing $PEG_{500}-0.05$ derivatization proved difficult to analyze in terms of loading for **Ta-04** (Zhen et al., 2021b).

No literature data are yet available that would describe the effects of chemical modifications of tannins on their antibiofilm,

or more generally antibacterial activity. With the aim of elucidating a hypothesis about the reason behind the effect of the derivatizations on our assayed tannins in comparison to the parent tannins and the blanks, we decided to look into other non-tannin organic compounds.

One of these examples targets the effect of the high equivalent derivatization $C_3NMe_3Cl-0.5$ in shifting the antibiofilm spectrum toward Gram-negative bacteria, which may be associated with addition of positive charges in the form of ammonium groups to the tannin molecules. This allows for comparison of our results with data obtained by Dalcin et al. (2017), who discovered that nanoencapsulation of dihydromyricetin within the polycationic polymer Eudragit RS 100® not only increased its antibiofilm activity against *P. aeruginosa*, but that the polymer itself had antibiofilm activity. This finding goes in accordance with a previous publication of Campanac et al. (2002) which states that cationic quaternary ammonium compounds were more effective against *P. aeruginosa* than *S. aureus* biofilm. Also, Gao et al. (2019) showed a decreased biofilm formation in *E. coli* and *S. aureus* using positively charged nanoaggregates based on zwitterionic pillar-[5] arene, requiring a 10 times lower concentration of nanoaggregate to decrease biofilm formation in *E. coli* compared to *S. aureus*. These results suggest that equipping the tannins with positive charge-introducing ammonium groups may give preferential action against Gram-negative bacteria. However, tannins should be derivatized with enough positive charge-introducing ammonium groups to obtain this shift toward inhibition of Gram-negative bacteria since the low equivalent derivatization $C_3NMe_3Cl-0.1$ appeared to lower the activity against Gram-negative bacteria.

Regarding the effect of acidifying CH_3COOH derivatizations, there is a precedent of the effect of several substitutions on the antibiofilm effect of anthraquinones against methicillin-resistant *S. aureus* (MRSA; Song et al., 2021). This study shows that a carboxyl group at position 2 of the anthraquinone molecule increases both the antibiofilm and the antimicrobial activity, which is partially in agreement with our data that shows that CH_3COOH derivatizations increase the antibiofilm activity of low oligomeric **Vv-20** and essentially monomeric **Vv** without affecting the bacterial growth. Relatedly, Warraich et al. (2020) found that the acidic D-amino acids D-aspartic acid and D-glutamic acid were effective in dispersing and inhibiting biofilm formation in *S. aureus*, and they attributed this effect to the negative charges introduced by carboxyl groups of the molecules under growth conditions.

Finally, regarding polymerizing PEG derivatizations, there are several studies about the potential of PEG cross-linked hydrogels for wound healing because of their antimicrobial, pro-angiogenesis and pro-epithelization capabilities (Chen et al., 2019), but there is no indication regarding the effect of an introduction of a PEG-motif on the antibiofilm capability of an organic compound.

Biofilm selectivity of the antibiofilm effect of the assayed tannins

In the last years, there has been increasing research on non-lethal antimicrobial targets against several bacterial species, from virulence factors to biofilm formation, including inhibition of regulatory mechanisms such as quorum sensing and production of public goods (Defoirdt et al., 2013; Defoirdt, 2016; Maura et al., 2016; Totsika, 2016). The rationale behind this research is the assumption that if bacterial viability is not affected, the selective pressure will be lower and thus the risk for emergence of antimicrobial resistance will be lower too (Kalia et al., 2014; Dieltjens et al., 2020; Hemmati et al., 2020). However, there is still discussion about the effectiveness of this “resistant-proof” approach (García-Contreras et al., 2016).

The heatmap of Figure 4 shows the anti-planktonic activity of the tannins at the assayed concentrations. We defined that the antibiofilm activity of a particular tannin was biofilm specific in case the tannin did not exhibit significant anti-planktonic effect at that concentration (Dos Santos Goncalves et al., 2014; Vijayakumar and Thirunanasambandham, 2021). Most of the tannins with antibiofilm activity did not exhibit anti-planktonic effects, indicating that they are biofilm specific. This is particularly true for unmodified tannins, of which only **Vv** and **Ta-01** exhibited anti-planktonic activity against *Salmonella* Typhimurium and *E. coli*, respectively. Also, unmodified tannins re-isolated from blank reactions, i.e., Blank-W or Blank-D, with the exception of Blank-D against *Salmonella* Typhimurium, did not reduce the planktonic growth of the assayed bacteria.

Regarding the effect of derivatizations on the biofilm specificity of the tannins, this was both derivatization and species dependent. Derivatization with positive charge introducing $C_3NMe_3Cl-0.1$ not only generally decreased the antibiofilm activity of the assayed tannins, but also increased the anti-planktonic activity specifically against Gram-negative bacteria. Antibiofilm specificity of tannins derivatized with $C_3NMe_3Cl-0.5$ was highly dependent on both the assayed bacteria and the derivatized tannin: the antibiofilm effect was non-specific when the derivatization was applied to **Vv-20**, but it was biofilm-specific for **Vv**, **Am** and **Ta-01**. Also, tannins derivatized with $C_3NMe_3Cl-0.5$ most strongly affected the growth of *P. aeruginosa* except for **Am**- $C_3NMe_3Cl-0.5$, which did not affect the growth of *P. aeruginosa* at any concentration. Only a derivatization of **Ta-04**, and here especially with $C_3NMe_3Cl-0.5$, led to a dose-dependent anti-planktonic effect, thus exhibiting non-specific antibiofilm activity at higher concentrations. With respect to a functionalization with the acidifying element CH_3COOH at various concentrations, it can be stated that these in this study did not significantly change the anti-planktonic behavior of the derivatized **Vv**, **Am** and **Ta-01** against the assayed bacteria, but increased the antibacterial effect of **Vv-20** and **Ta-04** against planktonic bacteria. However, one notable exception are the tannins that

were modified with crosslinking $PEG_{500}-0.05$, whose antibiofilm activity against Gram negative bacteria was highly correlated with the ability to inhibit the planktonic growth (Figure 4).


As a general summary, tannin derivatization did not affect biofilm specificity against *S. aureus* but affected the biofilm specificity against Gram-negative bacteria in a tannin-specific and derivatization specific manner. More importantly, there was no correlation in the assayed tannins between the degree of inhibition of planktonic growth and the degree of antibiofilm effect, a situation that goes in accordance with some previous reports (Janecki and Kolodziej, 2010; Payne et al., 2013; Hricoviniová et al., 2021).

Relation between antibiofilm effect and phenolic hydroxyl group content of the tannins

One of the potential consequences of the chemical derivatization of the tannins are changes in the content of free phenolic hydroxyl groups present in the chemical structure of the tannins, since functionalization occurs at these hydroxyl groups. This is potentially important, because in previous literature it has been described that the biological activity of polyphenols could be mediated by their phenolic hydroxyl groups (Ultee et al., 2002; Ito et al., 2005; Ben Arfa et al., 2006; Veldhuizen et al., 2006; Alves et al., 2013; d'Ávila Farias et al., 2014; Salar et al., 2017). Particularly regarding hydrolysable tannins, Taguri linked the degree of antibacterial activity to the presence of galloyl groups (Taguri et al., 2004). Because of this, we aimed to determine if there was a significant impact of the derivatizations on the phenolic hydroxyl (OH) content of the tannins, and if the phenolic OH content affected the antibiofilm.

In a first approach, we studied the impact of derivatizations on the phenolic OH content of the tannins. As can be seen in Table 1, derivatizations with $C_3NMe_3Cl-0.1$ decreased the phenolic OH content of the unmodified tannin. This is important, since derivatization with $C_3NMe_3Cl-0.1$ significantly decreased the antibiofilm effect of tannins, mostly against Gram negative bacteria. In a similar trend, derivatizations with CH_3COOH tended to decrease the phenolic content of condensed tannins: derivatization with CH_3COOH decreased the antibiofilm effect against Gram negative bacteria of condensed tannins, but, interestingly, not of still larger hydrolysable tannins.

In a second approach, we hence studied potential correlations between the phenolic OH content and the antibiofilm activity and we observed a weak correlation between the phenolic OH content and the antibiofilm activity against Gram negative bacteria and, to a lesser extent, against *S. aureus* (see Supplementary Figure S8). These data, combined with the previously mentioned effect of the different derivatizations on the phenolic OH content of the tannins,



suggest that the phenolic OH content of the tannins is more important for the antibiofilm effect against Gram negative bacteria than against *S. aureus*. However, we did not observe a correlation between the antibacterial activity against planktonic bacteria and the phenolic OH content of the assayed tannins (see [Supplementary Figure S9](#)), which is in accordance with the study [Kim et al. \(2020\)](#), who did not find a significant correlation between the total phenolic content of several plant extracts from Chinese traditional medicine and the antimicrobial activity against *S. aureus*. However, this contradicts a previous study from [Vattem et al. \(2004\)](#), who found a linear correlation between the phenolic content of

cranberry pomace and the antimicrobial activity against *Listeria monocytogenes*, *Vibrio parahaemolyticus* and *E. coli*.

It is important to note that the previous studies are focused on the antibacterial effect against planktonic bacteria, since there are no previous studies pointing to the effect of the hydroxyl or phenolic content on the capability of tannins to inhibit biofilms. However, research on other classes of polyphenolic compounds showed that the quantity and the position of the hydroxyl groups affects the capabilities of polyphenolic compounds to disperse biofilm, which could explain the results obtained in this research with the assayed tannins. In more detail, the research of Cho et al. on the capability of flavonoids to disperse *S. aureus* biofilms

TABLE 1 Phenolic OH content (in mmol phenolic OH/g materials) of unmodified and derivatized tannins.

Tannin	Unmodified	Blank-W	Blank-D	C ₃ NMe ₃ Cl-0.1	C ₃ NMe ₃ Cl-0.5	CH ₃ COOH-0.1	CH ₃ COOH-0.5	PEG ₅₀₀ -0.05
Vv-20	10.37	7.85	8.79	4.35	N.d. ^a	6.19	6.15	7.03
Vv	5.38	6.83	11.7	3.26	5.21	1.51	5.09	7.11
Am	8.35	7.64	10.27	0.45	N.d. ^a	5.76	3.76	8.03
Ta-01	12.56	■ ^b	10.99	1.55	7.22	10.05	11.61	8.87
Ta-04	10.49	■ ^b	6.89	6	N.d. ^a	9.89	9	6.77

The values were obtained using ³¹P-NMR. ^aN.d.: Not possible to calculate phenolic OH content due to solubility issues.

^b■ Blank-W reaction was not performed with Ta-01 and Ta-04 (hydrolysable tannins).

showed that there is a direct correlation between the quantity of hydroxyl substitutions of the flavonoid molecules and their antibiofilm capability (Cho et al., 2015), and the research of Kim et al. with phorbaketals against *S. aureus* biofilms indicated that the position and quantity of hydroxyls substitutions in the molecule affects their antibiofilm capacity (Kim et al., 2021), which is also consistent with the results of Arshia et al. (2017).

Conclusion

Our work provides a clear understanding on which chemical modifications can be made to enhance the activity level or change the activity spectrum of natural tannins, and which chemical modifications are inconvenient for increasing their antibiofilm activity. This is one of the few studies that uses a systematic and statistical analysis to correlate specific chemical characteristics of modified tannins with their antibiofilm activity (Ge et al., 2003; Peeters et al., 2016; Kemege et al., 2017; Bouarab-Chibane et al., 2019; De Paiva et al., 2019; Jia et al., 2019).

From our work, it can be concluded that tannins not only have good activity against biofilms of different bacterial species, but that this antibiofilm activity is in most cases also biofilm specific. More important, we could identify that modifying the tannins with C₃NMe₃Cl-0.5 and PEG₅₀₀-0.05 can increase the antibiofilm activity against Gram-negative bacteria, although this often coincides with a decrease in activity against Gram-positive bacteria. Modifying the tannins with C₃NMe₃Cl-0.1, CH₃COOH-0.1 and CH₃COOH-0.5 generally decreases the effect against Gram-negatives, without affecting the activity against *S. aureus*. We can thus modulate the spectrum and the antibiofilm potency of tannins by the applied chemical modifications.

We could identify a weak correlation between the antibiofilm effect and the content of phenolic hydroxyl groups for Gram-negative bacteria and, to a lesser extent, for *S. aureus*. However, exploring the mode of action against other bacteria is a necessary and interesting avenue to explore further based on the initial insights generated in this work, pointing to a more complex interplay between functionalization, type of tannin in the sense of exposed galloyl units, and tannin size. A continued exploration of the possible mechanisms of actions of these compounds and the possible modifications that can be made to enhance their effect is

necessary to better optimize the antibiofilm potential of the tannins.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary material, further inquiries can be directed to the corresponding author.

Author contributions

XV performed the experimental biological work, data analysis, and article writing. LZ performed the chemical manipulations and provided the chemical analysis of the assayed tannins. JA developed the experimental design and contributed to the statistical analysis. TV contributed to the article writing and data analysis. HL was part of the coordination of the research project, and contributed to the results analysis and the writing of the text. CC contributed to the research project funding and coordination. HS contributed to project funding, project coordination, article writing, and data analysis. All authors contributed to the article and approved the submitted version.

Funding

This work was supported by H2020-MSCA-ITN-2016-BIOCLEAR project (grant agreement no. 722871). HL would like to additionally thank the MIUR for the Grant “Dipartimento di Eccellenza 2018-2022” to the Department of Earth and Environmental Sciences of the University of Milano-Bicocca. CC acknowledges the Ca’Foscari FPI 2019 funding.

Acknowledgments

We would like to thank A.S. Ajinomoto OmniChem N.V. in Belgium for generously providing samples of commercialized tannins. We especially like to thank David De Coster for his technical support during the experimental work.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2022.987164/full#supplementary-material>

References

- Akiyama, H. (2001). Antibacterial action of several tannins against *Staphylococcus aureus*. *J. Antimicrob. Chemother.* 48, 487–491. doi: 10.1093/jac/48.4.487
- Allen, H. K., Donato, J., Wang, H. H., Cloud-Hansen, K. A., Davies, J., and Handelsman, J. (2010). Call of the wild: antibiotic resistance genes in natural environments. *Nat. Rev. Microbiol.* 8, 251–259. doi: 10.1038/nrmicro2312
- Alves, M. J., Ferreira, I. C. F. R., Froufe, H. J. C., Abreu, R. M. V., Martins, A., and Pintado, M. (2013). Antimicrobial activity of phenolic compounds identified in wild mushrooms, SAR analysis and docking studies. *J. Appl. Microbiol.* 115, 346–357. doi: 10.1111/jam.12196
- Arshia, K. A. K., Khan, K. M., Ahmed, A., Taha, M., and Perveen, S. (2017). Antibiofilm potential of synthetic 2-amino-5-chlorobenzophenone Schiff bases and its confirmation through fluorescence microscopy. *Microb. Pathog.* 110, 497–506. doi: 10.1016/j.micpath.2017.07.040
- Baba, S. A., and Malik, S. A. (2015). Determination of total phenolic and flavonoid content, antimicrobial and antioxidant activity of a root extract of *Arisaema jacquemontii* Blume. *J. Taibah Univ. Sci.* 9, 449–454. doi: 10.1016/j.jtusci.2014.11.001
- Ben Arfa, A., Combes, S., Preziosi-Belloy, L., Gontard, N., and Chalier, P. (2006). Antimicrobial activity of carvacrol related to its chemical structure. *Lett. Appl. Microbiol.* 43, 149–154. doi: 10.1111/j.1472-765X.2006.01938.x
- Bouarab-Chibane, L., Forquet, V., Lantéri, P., Clément, Y., Léonard-Akkari, L., Oulahal, N., et al. (2019). Antibacterial properties of polyphenols: characterization and QSAR (quantitative structure–activity relationship) models. *Front. Microbiol.* 10:829. doi: 10.3389/fmicb.2019.00829
- Campanac, C., Pineau, L., Payard, A., Baziard-Mouyset, G., and Roques, C. (2002). Interactions between biocide cationic agents and bacterial biofilms. *Antimicrob. Agents Chemother.* 46, 1469–1474. doi: 10.1128/AAC.46.5.1469-1474.2002
- Chandra, H., Bishnoi, P., Yadav, A., Patni, B., Mishra, A., and Nautiyal, A. (2017). Antimicrobial resistance and the alternative resources with special emphasis on plant-based antimicrobials—a review. *Plan. Theory* 6:16. doi: 10.3390/plants6020016
- Chen, H., Cheng, R., Zhao, X., Zhang, Y., Tam, A., Yan, Y., et al. (2019). An injectable self-healing coordinative hydrogel with antibacterial and angiogenic properties for diabetic skin wound repair. *NPG Asia Mater* 11, 3. doi: 10.1038/s41427-018-0103-9
- Cho, H. S., Lee, J.-H., Cho, M. H., and Lee, J. (2015). Red wines and flavonoids diminish *Staphylococcus aureus* virulence with anti-biofilm and anti-hemolytic activities. *Biofouling* 31, 1–11. doi: 10.1080/08927014.2014.991319
- Chung, K.-T., Lu, Z., and Chou, M. (1998). Mechanism of inhibition of tannic acid and related compounds on the growth of intestinal bacteria. *Food Chem. Toxicol.* 36, 1053–1060. doi: 10.1016/S0278-6915(98)00086-6
- d'Ávila Farias, M., Oliveira, P. S., Dutra, F. S. P., Fernandes, T. J., De Pereira, C. M. P., De Oliveira, S. Q., et al. (2014). Eugenol derivatives as potential anti-oxidants: is phenolic hydroxyl necessary to obtain an effect? *J. Pharm. Pharmacol.* 66, 733–746. doi: 10.1111/jphp.12197
- Dalcin, A. J. F., Santos, C. G., Gündel, S. S., Roggia, I., Raffin, R. P., Ourique, A. F., et al. (2017). Anti biofilm effect of dihydromyricetin-loaded nanocapsules on urinary catheter infected by *Pseudomonas aeruginosa*. *Colloids Surf. B Biointerfaces* 156, 282–291. doi: 10.1016/j.colsurfb.2017.05.029
- De Paiva, R. K. C., Da Silva, J. F., Moreira, H. A., Pinto, O. G., Camargo, L. T. F. M., Naves, P. L. F., et al. (2019). Synthesis, antimicrobial activity and structure-activity relationship of some 5-Arylidene-thiazolidine-2, 4-dione derivatives. *J. Braz. Chem. Soc.* 30, 164–172. doi: 10.21577/0103-5053.20180167
- De, R., Sarkar, A., Ghosh, P., Ganguly, M., Karmakar, B. C., Saha, D. R., et al. (2018). Antimicrobial activity of ellagic acid against helicobacter pylori isolates from India and during infections in mice. *J. Antimicrob. Chemother.* 73, 1595–1603. doi: 10.1093/jac/dky079
- Defoirdt, T. (2016). Specific Antivirulence activity, a new concept for reliable screening of virulence inhibitors. *Trends Biotechnol.* 34, 527–529. doi: 10.1016/j.tibtech.2016.01.009
- Defoirdt, T., Brackman, G., and Coenye, T. (2013). Quorum sensing inhibitors: how strong is the evidence? *Trends Microbiol.* 21, 619–624. doi: 10.1016/j.tim.2013.09.006
- Dettweiler, M., Lyles, J. T., Nelson, K., Dale, B., Reddinger, R. M., Zurawski, D. V., et al. (2019). American civil war plant medicines inhibit growth, biofilm formation, and quorum sensing by multidrug-resistant bacteria. *Sci. Rep.* 9:7692. doi: 10.1038/s41598-019-44242-y
- Dieltjens, L., Appermans, K., Lissens, M., Lories, B., Kim, W., Van der Eycken, E. V., et al. (2020). Inhibiting bacterial cooperation is an evolutionarily robust anti-biofilm strategy. *Nat. Commun.* 11:107. doi: 10.1038/s41467-019-13660-x
- Dong, G., Liu, H., Yu, X., Zhang, X., Lu, H., Zhou, T., et al. (2018). Antimicrobial and anti-biofilm activity of tannic acid against *Staphylococcus aureus*. *Nat. Prod. Res.* 32, 2225–2228. doi: 10.1080/14786419.2017.1366485
- Dos Santos Goncalves, M., Delattre, C., Balestrino, D., Charbonnel, N., Elbouchfaoui, R., Wadouachi, A., et al. (2014). Anti-biofilm activity: a function of *Klebsiella pneumoniae* capsular polysaccharide. *PLoS One* 9:e99995. doi: 10.1371/journal.pone.0099995
- Ekambaram, S. P., Perumal, S. S., and Balakrishnan, A. (2016). Scope of hydrolysable tannins as possible antimicrobial agent. *Phyther. Res.* 30, 1035–1045. doi: 10.1002/ptr.5616
- Falcão, L., and Araújo, M. E. M. (2011). Tannins characterisation in new and historic vegetable tanned leathers fibres by spot tests. *J. Cult. Herit.* 12, 149–156. doi: 10.1016/j.culher.2010.10.005
- Fang, Y., Lu, Y., Zang, X., Wu, T., Qi, X., Pan, S., et al. (2016). 3D-QSAR and docking studies of flavonoids as potent *Escherichia coli* inhibitors. *Sci. Rep.* 6:23634. doi: 10.1038/srep23634
- Farha, A. K., Yang, Q.-Q., Kim, G., Li, H.-B., Zhu, F., Liu, H.-Y., et al. (2020). Tannins as an alternative to antibiotics. *Food Biosci.* 38:100751. doi: 10.1016/j.fbio.2020.100751
- Gao, L., Li, M., Ehrmann, S., Tu, Z., and Haag, R. (2019). Positively charged nanoaggregates based on Zwitterionic pillar[5]arene that combat planktonic bacteria and disrupt biofilms. *Angew. Chemie – Int. Ed.* 58, 3645–3649. doi: 10.1002/anie.201810314
- García-Contreras, R., Maeda, T., and Wood, T. K. (2016). Can resistance against quorum-sensing interference be selected? *ISME J.* 10, 4–10. doi: 10.1038/ismej.2015.84
- Ge, J., Shi, X., Cai, M., Wu, R., and Wang, M. (2003). A novel biodegradable antimicrobial PU foam from wattle tannin. *J. Appl. Polym. Sci.* 90, 2756–2763. doi: 10.1002/app.12928
- Hemmati, F., Salehi, R., Ghotaslou, R., Samadi Kafil, H., Hasani, A., Gholizadeh, P., et al. (2020). Quorum quenching: a potential target for antipseudomonal therapy. *Infect. Drug Resist.* 13, 2989–3005. doi: 10.2147/IDR.S263196

- Hricoviniová, Z., Mascaretti, Š., Hricoviniová, J., Čížek, A., and Jampilek, J. (2021). New unnatural Gallotannins: a way toward green antioxidants, antimicrobials and antibiofilm agents. *Antioxidants* 10:1288. doi: 10.3390/antiox10081288
- Imperi, F., Fiscarelli, E. V., Visaggio, D., Leoni, L., and Visca, P. (2019). Activity and impact on resistance development of two antivirulence fluoropyrimidine drugs in *Pseudomonas aeruginosa*. *Front. Cell. Infect. Microbiol.* 9, 1–11. doi: 10.3389/fcimb.2019.00049
- Ito, M., Murakami, K., and Yoshino, M. (2005). Antioxidant action of eugenol compounds: role of metal ion in the inhibition of lipid peroxidation. *Food Chem. Toxicol.* 43, 461–466. doi: 10.1016/j.fct.2004.11.019
- Jagani, S., Chelikani, R., and Kim, D. S. (2009). Effects of phenol and natural phenolic compounds on biofilm formation by *Pseudomonas aeruginosa*. *Biofouling* 25, 321–324. doi: 10.1080/08927010802660854
- Janecki, A., and Kolodziej, H. (2010). Anti-adhesive activities of Flavan-3-ols and Proanthocyanidins in the interaction of group A-streptococci and human epithelial cells. *Molecules* 15, 1739–1752. doi: 10.3390/molecules15107139
- Janssens, J. C. A., Steenackers, H., Robijns, S., Gellens, E., Levin, J., Zhao, H., et al. (2008). Brominated furanones inhibit biofilm formation by salmonella enterica serovar typhimurium. *Appl. Environ. Microbiol.* 74, 6639–6648. doi: 10.1128/AEM.01262-08
- Jia, B., Ma, Y. M., Liu, B., Chen, P., Hu, Y., and Zhang, R. (2019). Synthesis, antimicrobial activity, structure-activity relationship, and molecular docking studies of indole Diketopiperazine alkaloids. *Front. Chem.* 7, 1–13. doi: 10.3389/fchem.2019.00837
- Kalia, V. C., Wood, T. K., and Kumar, P. (2014). Evolution of resistance to quorum-sensing inhibitors. *Microb. Ecol.* 68, 13–23. doi: 10.1007/s00248-013-0316-y
- Kemegne, G. A., Mkounga, P., Essia Ngang, J. J., Sado Kamdem, S. L., and Nkengfack, A. E. (2017). Antimicrobial structure activity relationship of five anthraquinones of emodine type isolated from *Vismia laurentii*. *BMC Microbiol.* 17, 41–48. doi: 10.1186/s12866-017-0954-1
- Kim, G., Gan, R.-Y., Zhang, D., Farha, A. K., Habimana, O., Mavumengwana, V., et al. (2020). Large-scale screening of 239 traditional Chinese medicinal plant extracts for their antibacterial activities against multidrug-resistant *Staphylococcus aureus* and cytotoxic activities. *Pathogens* 9:185. doi: 10.3390/pathogens9030185
- Kim, J.-K., Kim, N., and Lim, Y.-H. (2010). Evaluation of the antibacterial activity of Rhapontigenin produced from Rhapontin by biotransformation against *Propionibacterium acnes*. *J. Microbiol. Biotechnol.* 20, 82–87. doi: 10.4014/jmb.0907.07022
- Kim, Y.-G., Lee, J.-H., Lee, S., Lee, Y.-K., Hwang, B. S., and Lee, J. (2021). Antibiofilm activity of Phorbaketal from the marine sponge *Phorbis* sp. against *Staphylococcus aureus*. *Mar. Drugs* 19:301. doi: 10.3390/md19060301
- Klug, T. V., Novello, J., Laranja, D. C., Aguirre, T. A. S., de Oliveira Rios, A., Tondo, E. C., et al. (2017). Effect of tannin extracts on biofilms and attachment of *Escherichia coli* on lettuce leaves. *Food Bioprocess Technol.* 10, 275–283. doi: 10.1007/s11947-016-1812-0
- Koleckar, V., Kubikova, K., Rehakova, Z., Kuca, K., Jun, D., Jahodar, L., et al. (2008). Condensed and hydrolysable tannins as antioxidants influencing the health. *Mini-Reviews Med. Chem.* 8, 436–447. doi: 10.2174/138955708784223486
- Lahiri, D., Dash, S., Dutta, R., and Nag, M. (2019). Elucidating the effect of antibiofilm activity of bioactive compounds extracted from plants. *J. Biosci.* 44:52. doi: 10.1007/s12038-019-9868-4
- Lee, J.-H., Park, J.-H., Cho, H. S., Joo, S. W., Cho, M. H., and Lee, J. (2013). Antibiofilm activities of quercetin and tannic acid against *Staphylococcus aureus*. *Biofouling* 29, 491–499. doi: 10.1080/08927014.2013.788692
- Lin, M. H., Chang, F. R., Hua, M. Y., Wu, Y. C., and Liu, S. T. (2011). Inhibitory effects of 1,2,3,4,6-penta-O-galloyl- β -D-glucopyranose on biofilm formation by *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 55, 1021–1027. doi: 10.1128/AAC.00843-10
- Maura, D., Ballok, A. E., and Rahme, L. G. (2016). Considerations and caveats in anti-virulence drug development. *Curr. Opin. Microbiol.* 33, 41–46. doi: 10.1016/j.mib.2016.06.001
- Mori, A., Nishino, C., Enoki, N., and Tawata, S. (1987). Antibacterial activity and mode of action of plant flavonoids against *Proteus vulgaris* and *Staphylococcus aureus*. *Phytochemistry* 26, 2231–2234. doi: 10.1016/S0031-9422(00)84689-0
- Payne, D. E., Martin, N. R., Parzych, K. R., Rickard, A. H., Underwood, A., and Boles, B. R. (2013). Tannic acid inhibits *Staphylococcus aureus* surface colonization in an IsaA-dependent manner. *Infect. Immun.* 81, 496–504. doi: 10.1128/IAI.00877-12
- Peeters, E., Hooyberghs, G., Robijns, S., Waldrant, K., De Weerd, A., Delattin, N., et al. (2016). Modulation of the substitution pattern of 5-Aryl-2-Aminoimidazoles allows fine-tuning of their Antibiofilm activity Spectrum and toxicity. *Antimicrob. Agents Chemother.* 60, 6483–6497. doi: 10.1128/AAC.00035-16
- Pizzi, A. (2019). Tannins: prospective and actual industrial applications. *Biomol. Ther.* 9:344. doi: 10.3390/biom9080344
- Puljula, E., Walton, G., Woodward, M. J., and Karonen, M. (2020). Antimicrobial activities of Ellagitannins against *Clostridiales perfringens*, *Escherichia coli*, *Lactobacillus plantarum* and *Staphylococcus aureus*. *Molecules* 25:3714. doi: 10.3390/molecules25163714
- Rabin, N., Zheng, Y., Opoku-Temeng, C., Du, Y., Bonsu, E., and Sintim, H. O. (2015). Biofilm formation mechanisms and targets for developing antibiofilm agents. *Future Med. Chem.* 7, 493–512. doi: 10.4155/fmc.15.6
- Roy, R., Tiwari, M., Donelli, G., and Tiwari, V. (2018). Strategies for combating bacterial biofilms: a focus on anti-biofilm agents and their mechanisms of action. *Virulence* 9, 522–554. doi: 10.1080/21505594.2017.1313372
- Salar, R. K., Purewal, S. S., and Sandhu, K. S. (2017). Relationships between DNA damage protection activity, total phenolic content, condensed tannin content and antioxidant potential among Indian barley cultivars. *Biocatal. Agric. Biotechnol.* 11, 201–206. doi: 10.1016/j.bcab.2017.07.006
- Sarabhai, S., Sharma, P., and Capalash, N. (2013). Ellagic acid derivatives from *Terminalia chebula* Retz. Downregulate the expression of quorum sensing genes to attenuate *Pseudomonas aeruginosa* PAO1 virulence. *PLoS One* 8:e53441. doi: 10.1371/journal.pone.0053441
- Scalbert, A. (1991). Antimicrobial properties of tannins. *Phytochemistry* 30, 3875–3883. doi: 10.1016/0031-9422(91)83426-L
- Shukla, V., and Bhatena, Z. (2015). Sustained release of a purified tannin component of *Terminalia chebula* from a titanium implant surface prevents biofilm formation by *Staphylococcus aureus*. *Appl. Biochem. Biotechnol.* 175, 3542–3556. doi: 10.1007/s12010-015-1525-2
- Slobodníková, L., Fialová, S., Rendeková, K., Kováč, J., and Mučaji, P. (2016). Antibiofilm activity of plant polyphenols. *Molecules* 21:1717. doi: 10.3390/molecules21121717
- Song, Z.-M., Zhang, J.-L., Zhou, K., Yue, L.-M., Zhang, Y., Wang, C.-Y., et al. (2021). Anthraquinones as potential Antibiofilm agents against methicillin-resistant *Staphylococcus aureus*. *Front. Microbiol.* 12, 1–16. doi: 10.3389/fmicb.2021.709826
- Steenackers, H. P. L., Ermolatev, D. S., Savaliya, B., Weerd, A. D., Coster, D. D., Shah, A., et al. (2011). Structure-activity relationship of 2-hydroxy-2-aryl-2,3-dihydro-imidazo[1,2-a]pyrimidinium salts and 2N-substituted 4(5)-aryl-2-amino-1H-imidazoles as inhibitors of biofilm formation by *Salmonella typhimurium* and *Pseudomonas aeruginosa*. *Bioorg. Med. Chem.* 19, 3462–3473. doi: 10.1016/j.bmc.2011.04.026
- Taganna, J. C., Quanicco, J. P., Perono, R. M. G., Amor, E. C., and Rivera, W. L. (2011). Tannin-rich fraction from *Terminalia catappa* inhibits quorum sensing (QS) in *Chromobacterium violaceum* and the QS-controlled biofilm maturation and LasA staphylolytic activity in *Pseudomonas aeruginosa*. *J. Ethnopharmacol.* 134, 865–871. doi: 10.1016/j.jep.2011.01.028
- Taguri, T., Tanaka, T., and Kouno, I. (2004). Antimicrobial activity of 10 different plant polyphenols against bacteria causing food-borne disease. *Biol. Pharm. Bull.* 27, 1965–1969. doi: 10.1248/bpb.27.1965
- Tintino, S. R., Oliveira-Tintino, C. D. M., Campina, F. F., Silva, R. L. P., Costa, M. D. S., Menezes, I. R. A., et al. (2016). Evaluation of the tannic acid inhibitory effect against the NorA efflux pump of *Staphylococcus aureus*. *Microb. Pathog.* 97, 9–13. doi: 10.1016/j.micpath.2016.04.003
- Totsika, M. (2016). Benefits and challenges of Antivirulence antimicrobials at the Dawn of the post-antibiotic era. *Drug Deliv. Lett.* 6, 30–37. doi: 10.2127/4/2210303106666160506120057
- Trentin, D. S., Silva, D. B., Amaral, M. W., Zimmer, K. R., Silva, M. V., Lopes, N. P., et al. (2013). Tannins possessing bacteriostatic effect impair *Pseudomonas aeruginosa* adhesion and biofilm formation. *PLoS One* 8:e66257. doi: 10.1371/journal.pone.0066257
- Ulrey, R. K., Barksdale, S. M., Zhou, W., and van Hoek, M. L. (2014). Cranberry proanthocyanidins have anti-biofilm properties against *Pseudomonas aeruginosa*. *BMC Complement. Altern. Med.* 14:499. doi: 10.1186/1472-6882-14-499
- Ultee, A., Bennik, M. H. J., and Moezelaar, R. (2002). The phenolic hydroxyl Group of Carvacrol is Essential for action against the food-borne pathogen *Bacillus cereus*. *Appl. Environ. Microbiol.* 68, 1561–1568. doi: 10.1128/AEM.68.4.1561-1568.2002
- Vattem, D. A., Lin, Y.-T., Labbe, R. G., and Shetty, K. (2004). Antimicrobial activity against select food-borne pathogens by phenolic antioxidants enriched in cranberry pomace by solid-state bioprocessing using the food grade fungus *Rhizopus oligosporus*. *Process Biochem.* 39, 1939–1946. doi: 10.1016/j.procbio.2003.09.032
- Veldhuizen, E. J. A., Tjeerdma-van Bokhoven, J. L. M., Zweijter, C., Burt, S. A., and Haagsman, H. P. (2006). Structural requirements for the antimicrobial activity of Carvacrol. *J. Agric. Food Chem.* 54, 1874–1879. doi: 10.1021/jf052564y

- Versari, A., du Toit, W., and Parpinello, G. P. (2013). Oenological tannins: a review. *Aust. J. Grape Wine Res.* 19, 1–10. doi: 10.1111/ajgw.12002
- Vijayakumar, K., and Thirunanasambandham, R. (2021). 5-Hydroxymethylfurfural inhibits *Acinetobacter baumannii* biofilms: an in vitro study. *Arch. Microbiol.* 203, 673–682. doi: 10.1007/s00203-020-02061-0
- Vu, T. T., Kim, H., Tran, V. K., Vu, H. D., Hoang, T. X., Han, J. W., et al. (2017). Antibacterial activity of tannins isolated from *Sapium baccatum* extract and use for control of tomato bacterial wilt. *PLoS One* 12:e0181499. doi: 10.1371/journal.pone.0181499
- Warraich, A. A., Mohammed, A. R., Perrie, Y., Hussain, M., Gibson, H., and Rahman, A. (2020). Evaluation of anti-biofilm activity of acidic amino acids and synergy with ciprofloxacin on *Staphylococcus aureus* biofilms. *Sci. Rep.* 10:9021. doi: 10.1038/s41598-020-66082-x
- Widsten, P., Cruz, C. D., Fletcher, G. C., Pajak, M. A., and McGhie, T. K. (2014). Tannins and extracts of fruit byproducts: antibacterial activity against foodborne bacteria and antioxidant capacity. *J. Agric. Food Chem.* 62, 11146–11156. doi: 10.1021/jf503819t
- Wu, D., Wu, X.-D., You, X.-F., Ma, X.-F., and Tian, W.-X. (2010). Inhibitory effects on bacterial growth and b-ketoacyl-ACP reductase by different species of maple leaf extracts and tannic acid. *Phyther. Res.* 24, S35–S41. doi: 10.1002/ptr.2873
- Zhen, L., Lange, H., and Crestini, C. (2021a). An analytical toolbox for fast and straightforward structural characterisation of commercially available tannins. *Molecules* 26:2532. doi: 10.3390/molecules26092532
- Zhen, L., Lange, H., Zongo, L., and Crestini, C. (2021b). Chemical derivatization of commercially available condensed and Hydrolyzable tannins. *ACS Sustain. Chem. Eng.* 9, 10154–10166. doi: 10.1021/acssuschemeng.1c02114