

## **OPEN ACCESS**

EDITED AND REVIEWED BY Rustam Aminov, University of Aberdeen, United Kingdom

\*CORRESPONDENCE

Djamel Drider

☑ djamel.drider@univ-lille1.fr

RECEIVED 13 June 2025 ACCEPTED 28 July 2025 PUBLISHED 11 August 2025

### CITATION

Al Atya AK, Belguesmia Y, Chataigne G, Ravallec R, Vachée A, Szunerits S, Boukherroub R and Drider D (2025) Correction: Anti-MRSA activities of enterocins DD28 and DD93 and evidences on their role in the inhibition of biofilm formation. Front. Microbiol. 16:1646639. doi: 10.3389/fmicb.2025.1646639

## COPYRIGHT

© 2025 Al Atya, Belguesmia, Chataigne, Ravallec, Vachée, Szunerits, Boukherroub and Drider. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Correction: Anti-MRSA activities of enterocins DD28 and DD93 and evidences on their role in the inhibition of biofilm formation

Ahmed K. Al Atya<sup>1</sup>, Yanath Belguesmia<sup>1</sup>, Gabrielle Chataigne<sup>1</sup>, Rozenn Ravallec<sup>1</sup>, Anne Vachée<sup>2</sup>, Sabine Szunerits<sup>3</sup>, Rabah Boukherroub<sup>3</sup> and Djamel Drider<sup>1</sup>\*

<sup>1</sup>Université de Lille 1 Sciences et Technologies - Institut Charles Viollette, Lille, France, <sup>2</sup>Hôpital Victor Provo de Roubaix, Roubaix, France, <sup>3</sup>Institut d'Electronique, de Microélectronique et de Nanotechnologie, UMR CNRS 8520, Université Lille 1, Lille, France

## KEYWORDS

enterocins, antibiotics, MRSA, synergism, bacteriocins

# A Correction on

Anti-MRSA activities of enterocins DD28 and DD93 and evidences on their role in the inhibition of biofilm formation

by Al Atya, A. K., Belguesmia, Y., Chataigne, G., Ravallec, R., Vachée, A., Szunerits, S., Boukherroub, R., and Drider, D. (2016). Front. Microbiol. 7:817. doi: 10.3389/fmicb.2016.00817

There was a mistake in Figure 4 as published. A wrong figure appeared.

The corrected Figure 4 and it's caption appear below.

The original version of this article has been updated.

# Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Al Atya et al. 10.3389/fmicb.2025.1646639




FIGURE 4
Biofilms formation by MRSA-S1 on AISI 304L stainless steel slides conditioned with antimicrobial agents for 2 h and then washed and inoculated with 2 ml of  $10^7 \text{ CFU/ml}$  MRSA-S1 culture before removing, washing and adding TSB-YE medium for 0, 3, 6, and 24 h of incubation at  $37^{\circ}$ C. Concentrations of antimicrobial agents used were 8 mg/L for erythromycin, 0.03 mg/L for rifampin, 200 mg/L for enterocin DD28, 50 mg/L + 1 mg/L for enterocin DD28 + erythromycin, respectively. Biofilms were stained with the BacLight Live/Dead Viability Kit and imaged by epifluorescence microscopy after staining pattern for live cells (green) and dead cells (red). The experiments were performed at least twice and representative images are shown.