



Formation of Complexes Between O Proteins and Replication Origin Regions of Shiga Toxin-Converting Bacteriophages

Katarzyna Kozłowska¹, Monika Glinkowska², Lidia Boss², Lidia Gaffke¹, Jakub Deptuła¹ and Grzegorz Węgrzyn^{1*}

OPEN ACCESS

Edited by:

Chew Chieng Yeo,
Sultan Zainal Abidin University,
Malaysia

Reviewed by:

Radosław Pluta,
Institute for Research in Biomedicine,
Spain

Lanying Zeng,

Texas A&M University, United States

Stephen M. Kwong,

The University of Sydney, Australia

*Correspondence:

Grzegorz Węgrzyn
grzegorz.wegrzyn@biol.ug.edu.pl;
wegrzyn@biotech.ug.gda.pl

Specialty section:

This article was submitted to
Molecular Recognition,
a section of the journal
Frontiers in Molecular Biosciences

Received: 09 April 2020

Accepted: 28 July 2020

Published: 19 August 2020

Citation:

Kozłowska K, Glinkowska M,
Boss L, Gaffke L, Deptuła J and
Węgrzyn G (2020) Formation
of Complexes Between O Proteins
and Replication Origin Regions
of Shiga Toxin-Converting
Bacteriophages.
Front. Mol. Biosci. 7:207.
doi: 10.3389/fmolb.2020.00207

¹ Department of Molecular Biology, Faculty of Biology, University of Gdańsk, Gdańsk, Poland, ² Department of Bacterial Molecular Genetics, Faculty of Biology, University of Gdańsk, Gdańsk, Poland

Shiga toxin-converting bacteriophages (or Stx phages) are responsible for virulence of enterohemorrhagic *Escherichia coli* strains. Although they belong to the group of lambdoid phages, which have served as models in studies on DNA replication mechanisms, details of regulation of replication of Stx phage genomes are poorly understood. Despite high similarity of their replication regions to that of phage lambda, considerable differences occur between them. Here, we present a comparison of origins of replication and O proteins of lambda and selected Stx phages (phages P27 and 933W). Stx initiator proteins, similarly to the lambda O protein, exist in the form of dimers. Only 4 iteron sequences are strongly bound *in vitro* by the O proteins, despite the presence of 6 such fragments in the Stx *ori*, while the function of the other two iterons is still crucial for transformation of *E. coli* wild-type strain by the P27-derived lambdoid plasmid. As these sequences are found in the gene coding for Stx O proteins, the sequences of these proteins themselves are also extended compared to lambda phage. Therefore, proteins O of Stx phages P27 and 933W have 13 additional amino acids. They can act as a space barrier, thus affecting the lesser packing of the O-some Stx complex compared to the structure found in lambda. Such structure of the DNA replication initiation complex may determine its lesser dependence on the processes occurring in the host cell, including transcriptional activation of the origin. Differences between molecular processes occurring during formation of replication complexes in lambda and Stx phages may indicate the specialization of the latter phages and their adaptation to specific environmental conditions where quick genetic switches are crucial.

Keywords: Shiga toxin-converting bacteriophages, DNA replication, replication initiator protein, replication complex, protein-DNA interactions

INTRODUCTION

Escherichia coli constitutes crucial component of the natural microbiota of the human and warm-blooded animals intestine (Actis, 2014). Although the vast majority of *E. coli* strains are commensals, some of them can also be pathogenic to humans (Leimbach et al., 2013). Shiga toxin-producing *E. coli* (STEC) strains, synthesizing one of the strongest toxins, are among the most dangerous pathogens (Paletta et al., 2020). Infections with these strains are particularly dangerous, because they can cause severe complications, while available antibacterial therapies are often not effective against those pathogens (Karmali, 2018). The latter problem is caused by two factors: firstly, STEC strains are often resistant to many antibiotics, secondly, the use of antimicrobials may be counterproductive, as some of them may enhance expression of Shiga toxins genes. Thus, even if a STEC strain is susceptible to a particular therapeutic its application may cause production of sufficiently high level of toxins to evoke hemolytic uremic syndrome in patients, resulting in significant morbidity and mortality (Kakoullis et al., 2019; Mir and Kudva, 2019). There are examples of many outbreaks caused by STEC which resulted in severe complications in thousands of patients and relatively many deaths (Devleesschauwer et al., 2019; Kintz et al., 2019).

Shiga toxins, the main virulence factors of STEC, are encoded by the genes (called *stx* genes) present in genomes of Shiga toxin-converting bacteriophages (or Stx bacteriophages) rather than in bacterial chromosome (Łoś et al., 2011). Stx phages occur as prophages in STEC genomes, and *stx* genes, like vast majority of prophage genes, are silent in the lysogenic state (Krüger and Lucchesi, 2015). Under various stress conditions, transition of the phage from lysogenic into lytic development is possible which involves the excision of its genome and replication of phage DNA, followed by progeny virion production and cell lysis (Łoś et al., 2011; Licznarska et al., 2016). Simultaneously, extensive production of Shiga toxin occurs, while inhibition of phage DNA replication impairs expression of *stx* genes (coding for the toxins) (Nejman-Faleńczyk et al., 2012; Nowicki et al., 2013; Balasubramanian et al., 2019). Therefore, detailed knowledge on the regulation of DNA replication of Stx phages may contribute to better understanding of the mechanisms of pathogenicity of STEC strains.

Stx phages belong to the group of lambdoid bacteriophages which include the model virus of molecular biology – bacteriophage λ (Łoś et al., 2011). DNA replication of this phage has been well understood, but its regulation is still not clear (Węgrzyn et al., 2012). It was described as a complex process involving both phage elements and protein machinery of its bacterial host (Weigel and Seitz, 2006). The key role in the regulation of initiation of this process is played by the bacteriophage-encoded O protein and transcriptional activation of the *origin* (Taylor and Węgrzyn, 1995). λ DNA replication requires binding of the phage-encoded O protein to four DNA sequences in the *origin* of replication region, named iterons, and formation of the nucleoprotein structure called O-some. This is the crucial step in the initiation event, and provides the basis for

formation of the replication complex (Taylor and Węgrzyn, 1995; Weigel and Seitz, 2006).

It was demonstrated that the replication regions in genomes of Stx bacteriophages are similar to that of bacteriophage λ (Nejman et al., 2009). The key elements in this region are the *p_R* promoter, and O and P genes (coding for replication proteins) with the *origin* sequence (consisting of O-binding sequences, called iterons, and the AT-rich tract) located in the middle of the former gene (Figure 1). Only a few alterations were found between replication regions of λ and Stx phages, but they are responsible for significant differences in the DNA replication regulation. Particularly, various requirements for host-encoded factors, like DnaA and DksA proteins, and different dependence on the transcriptional activation of the *origin* (the process demonstrated previously as crucial for the control of the replication initiation; see Szalewska-Pałasz et al., 1994; Szambowska et al., 2011) were reported (Nejman et al., 2011). These differences result in various responses of intracellularly developing λ and Stx phages to environmental conditions, including starvation (Nejman et al., 2011).

Replication regions of Stx phages contain 6 iterons, contrary to 4 iterons present in λ , and a few amino acid alterations can be found in O and P proteins between these phages (Figure 1), which nevertheless cause significant differences in the phage DNA replication regulation (Nejman et al., 2011; Nowicki et al., 2015; Kozłowska et al., 2017). Recently, an improved method for overproduction and purification of O proteins of Stx phages has been reported which allowed to identify their specific binding to the *origin* regions (Kozłowska et al., 2017). Nevertheless, details of molecular interactions between these O proteins and iterons remained unknown, especially in the light of the above described differences between λ and Stx phages. Therefore, in this work, we aimed to determine details of formation of complexes between O proteins and *origin* regions of Stx phages and compare them to those of λ .

MATERIALS AND METHODS

Proteins

Overproduction and purification of O proteins derived from phages λ , P27 and 933W used in this study was described previously (Kozłowska et al., 2017).

Bacterial Strains and Source Plasmids

E. coli DH5 α (F⁻ Φ 80lacZ Δ M15 Δ (lacZYA-argF) U169 *recA1 endA1 hsdR17*(r⁻, m⁺) *phoA supE44 thi-1 gyrA96 relA1* λ ⁻) strain (Taylor et al., 1993; Chen et al., 2018), and plasmids pCB104cmr (λ phage-derived), p27cmr (P27 phage-derived) (Nejman et al., 2009), pET24a_wtO_histag, pET24a_P27O_histag, pET24a_933WO_histag, pUC18_ori λ , and pUC18_oriStx (Kozłowska et al., 2017), and pUC18 (Yanisch-Perron et al., 1985) were used for production of proteins and DNAs for *in vitro* experiments. *E. coli* MG1655 and its *dnaA46* and Δ *seqA* derivatives (Nejman et al., 2009)




FIGURE 1 | Differences in the nucleotide sequences of viral genome fragments that are involved in the initiation of DNA replication of bacteriophage λ and Stx phages (P27 and 933). Compositions of iterons and the AT-rich fragment are shown in **(A)** (since sequences of iterons 1 and 1'' are identical, two alternative ways of comparison between λ and Stx *origin* regions are possible, with dashed line indicating the missing fragment; the upper scheme presents the comparison as originally proposed by Nejman et al. (2009), and the lower scheme presents the alternative option). The scheme is not drawn to scale. Nucleotide sequences of iterons are shown in **(B)**, with (–) indicating bases identical to those occurring in iteron 1 and 1'', and letters indicating substitutions. Differences in amino acid sequences of O proteins of bacteriophage λ and Stx phages (P27 and 933W) are depicted in **(C)**.

were used for transformation efficiency analysis, performed as described previously (Nejman et al., 2009).

Construction of DNA Templates and Plasmids

O protein binding to single iterons was tested by EMSA assay, employing DNA fragments obtained by single stranded oligonucleotides hybridization. Respective primer sequences are presented in **Table 1** (oligonucleotides 8–19). Single-stranded

DNA fragments (synthetic oligonucleotides) for hybridization were ordered from Sigma Aldrich. Oligonucleotides were used at a final concentration 1 µg/ml. Hybridization buffer was prepared in a volume of 1 ml. Synthetic oligonucleotide hybridization buffer consisted of 0.1 ml of 1 M Na₃PO₄, 0.03 ml of 5 M NaCl and 0.002 ml of 0.5 M EDTA at pH 8. The hybridization was carried out in a thermocycler starting from temperature 98°C for 2 min, then 2 min at 80°C, another 2 min at 65°C and slow cooling down to 20°C. Following hybridization, double stranded DNA fragments were ligated with plasmid

TABLE 1 | Oligonucleotides.

No.	Oligonucleotide name	Sequence 5'→3'	Assay	PCR template
1.	Cy5_ori_EMSA_ATrich_for	[Cyanine 5]TCAAGCAGCAAGGCGGCATGTTTG	EMSA, forward	pUC18_ori λ and pUC18_oriStx
2.	ori_EMSA_ATrich_for	CACTTGCTGCGCCTCTGAATTGC	EMSA, reverse	
3.	ATrich_57GC_rev	GGATTGCGCCAGAATTCTCGGACGAAGCAT CGCCTCAGCTCGCTTCGGTACTAGTGTCC GATGTGTCCC	EMSA, reverse with AT-rich region replaced with region of 57% GC content	
4.	Cy5_pUC18_iterony_for	[Cyanine 5]GGGCTGGCTTAACTATGCGGCATC	PCR primer, forward	
5.	pUC18_iterony_rev	GTTAGCTCACTCATTAGGCACCC	PCR primer, reverse	
6.	pUC18_liniowy_footprint_for	CTTCCGGCTGGCTGGTTTATTG	Linearization of plasmid for footprinting	
7.	pUC18_liniowy_footprint_rev	CTTTATCCGCCTCCATCCAGTC	Linearization of plasmid for footprinting	
8.	Iteron1_pUC18_for	GATCCCGTAGCAGTCGCTGACATCCCT CAAACGAGGGACGTAGCAGT CGCTGACA	Hybridization, forward	Not applicable
9.	it1_rev	AGCTTGTCAGCGACTGCTACGTCCCTC GTTTTGAGGGATGTCAGCGAC TGCTACGG	Hybridization, reverse	
10.	Iteron1 \times _pUC18_for	GATCCCGTAGCAGTCGCTGAGTTCCC CTAAATGAGGGACGTAGCAGT CGCTGACA	Hybridization, forward	
11.	it1'_rev	AGCTTGTCAGCGACTGCTACGTCCCTC ATTTTAGGGGAACCTCAGCGAC TGCTACGG	Hybridization, reverse	
12.	Iteron2_pUC18_for	GATCCCGTAGCAGTCGCTGAAATCCCCT AAAACGAGGGACGTAGCAGT CGCTGACA	Hybridization, forward	
13.	it2_rev	AGCTTGTCAGCGACTGCTACGTCCCTCG TTTTAGGGGATTTAGCGAC TGCTACGG	Hybridization, reverse	
14.	Iteron3_pUC18_for	GATCCCGTAGCAGTCGCTGACATCCCTC AAATTGGGGACGTAGCAGT CGCTGACA	Hybridization, forward	
15.	it3_rev	AGCTTGTCAGCGACTGCTACGTCCCTC AATTTGAGGGATGTCAGCGAC TGCTACGG	Hybridization, reverse	
16.	Iteron3*_pUC18_for	GATCCCGTAGCAGTCGCTGACATCCCT CAAATTAGGGGACGTAGCAGT CGCTGACA	Hybridization, forward	
17.	it3*_rev	AGCTTGTCAGCGACTGCTACGTCCCTC TAATTTGAGGGATGTCAGCGAC TGCTACGG	Hybridization, reverse	
18.	Iteron4_pUC18_for	GATCCCGTAGCAGTCGCTGAATCCCTC AAAACAGGGGGACGTAGCAGT CGCTGACA	Hybridization, forward	
19.	it4_rev	AGCTTGTCAGCGACTGCTACGTCCCTC CTGTTTTGAGGGATTAGCGAC TGCTACGG	Hybridization, reverse	
20.	EMSA_for	[Cyanine 5]TGCTGCAAGGCGA	EMSA with single iteron sequence, forward	pUC18 with single iterons sequences (depending on iteron)
21.	EMSA_rev	CACTTTATGCTTCCGGCTCGTAT	EMSA with single iteron sequence, reverse	
22.	Footprinting for	AAGCAGCAAGGCGGCATGTTTG	Footprinting	pUC18_ori λ and pUC18_oriStx
23.	Footprinting rev	GCTGGTCAGAGGATTCGC	Footprinting	
24.	FRET	[Cyanine 5]CTGTTCTATTGTGATCTTATTAG [Cyanine 3]	FRET	Not applicable
25.	FRET_Donor_Cy3	CTGTTCTATTGTGATCTTATTAG[Cyanine 3]	FRET	

(Continued)

TABLE 1 | Continued

No.	Oligonucleotide name	Sequence 5'→3'	Assay	PCR template
26.	FRET_methylation	CT(mp)G(mp) T(mp)TC(mp) T(mp)A(mp)T T(mp)G(mp)T(mp) GA(mp)T CTC TTA TTA G	EMSA with methylated ssDNA	
27.	Mut1Fd	GGACCAAATAAAAACATCTCAGAATGGTGC ATTCTCAAACGAAGGAGGTTCCCTAAAATG	Site-directed mutagenesis of iteron sequence 1 (ATCCCTCAAAAACGA GGGAAATTCCTCAAACGAAGGA)	p27R6K
28.	Mut1Rev	GATGTCCCTCATTITAGGGGAACCTCCTTC GTTTGGAGGAATGCACCATTCTGAGATG	Site-directed mutagenesis of iteron sequence 1 (ATCCCTCAAAA ACGAGGGAAATTCCTCAAACGAAGGA)	p27R6K
29.	Mut1 + 1'Fd	CATCTCAGAATGGTGCATTCCTCAAAACG AAGGAGGTTCTCTAAAATGAGTGACATC CCTCAAACGAGGG	Site-directed mutagenesis of iteron sequences 1 and 1' (ATCCCT CAAAAACGAGGGAAATTCCTCAAAA CGAAGGA and TTCCCTAAA ATGAGGGATTCTCTAAAATGAGTGA)	p27mut1
30.	Mut1 + 1'Rev	GATGTTTTATCCCTCGTTTTAGGGGATTTT CCCTCGTTTTGAGGGATGCTACTCATTTT AGGAGAACCTCCT	Site-directed mutagenesis of iteron sequences 1 and 1' (ATCCCTCA AAAACGAGGGAAATTCCTCAAACGA AAGGA and TTCCCTAAAATGAG GATTCTCTAAAATGAGTGA respectively)	p27mut1
31.	Mut1'Fd	CATCTCAGAATGGTGCATCCCTCAAAACG AGGGAGGTTCTCTAAAATGAGTGACATC CCTCAAACGAGGG	Site-directed mutagenesis of iteron sequence 1' (TTCCCTAAAATGAG GGATTCTCTAAAATGAGTGA)	p27R6K
32.	Mut1'Rev	GATGTTTTATCCCTCGTTTTAGGGGATTTT CCTCGTTTTGAGGGATGCTACTCATTTAGGA GAACCTCCC	Site-directed mutagenesis of iteron sequence 1' (TTCCCTAAAATGAG GGATTCTCTAAAATGAGTGA)	p27R6K
33.	p27_IT1_Fd	[Cyanine5]GTGCATCCCTCAAACGAGGGAGGTT	EMSA with wt and mutant variants in iteron 1	
34.	p27_IT1_Rev	AACCTCCCTCGTTTTGAGGGATGCAC	EMSA with wt variant in iteron 1 of P27	
35.	p27_IT1_prim_Fd	[Cyanine5]GAGGTTCCCTCAAATGAGGGACATC	EMSA with wt variant in iteron 1' of P27	
36.	p27_IT1_prim_Rev	[Cyanine5]GTGCATTCCTCAAACGAAGGAGGTT	EMSA with wt variant in iteron 1' of P27	
37.	p27_IT1mut_Rev	AACCTCCTCGTTTTGAGGAATGCAC	EMSA with the mutant variant in iteron 1 of P27	
38.	p27_IT1mut_Rev	AACCTCCTCGTTTTGAGGAATGCAC	EMSA with the mutant variant in iteron 1 of P27	
39.	p27_IT1_prim_mut_Fd	[Cyanine5]GAGGTTCTCTAAAATGAGTGACATC	EMSA with the mutant variant in iteron 1' of P27	
40.	p27IT1_prim_mut_Rev	GATGCTACTCATTTAGGAGAACCTC	EMSA with the mutant variant in iteron 1' of P27	

pUC18, previously linearized with *Bam*HI and *Hind*III (Thermo ScientificTM), and used for transformation of *E. coli* DH5 α competent cells. Obtained plasmids were isolated from bacterial cells using plasmid isolation kit (Sigma Aldrich) and sequenced to confirm their proper construction. DNA fragments for EMSA or footprinting tests were obtained by PCR technique with use of specific primers (see **Table 1** for details).

Derivatives of the p27cmr plasmid, bearing additional *origin* of replication of the R6K plasmid, were constructed as follows. The PCR fragment bearing *origin* R6K was obtained after the reaction with pCAH56 plasmid and primers *OriR6KFd* (5' – GCC GAG CTC CAT CCC TGG CTT GTT GTC C) and *OriR6KRev* (5' – GAT GAG CTC GAT CCG GCC ACG ATG CGT C). The reaction product was cut with *Sac*I, and ligated with the *Sac*I fragment of the p27cmr plasmid. Changes in iteron sequences 1 and/or 1' of p27R6K plasmids, which did

not alter the O amino acid sequence, were introduced by site-directed mutagenesis, producing p27mut1, p27mut1', and p27mut1 + 1' plasmids, respectively, using suitable primers (**Table 1**, primers 27–32). *E. coli* BW25142 (*lacI*^q *rrnB3 lacZ4787 hsdR D(araBAD) D(rhaBAD) phoBR580 rph-1 galU95 endA9 uidA:pir-116 recA1*) strain was used for plasmid replication after site-directed mutagenesis.

Electrophoretic Mobility Shift Assay (EMSA)

All electrophoretic mobility shift assays (EMSA) were performed in the same manner. Indicated amounts of purified O proteins were incubated with 100 ng of different DNA fragments (1 μ g in case of ssDNA fragment – oligonucleotides 26 in **Table 1**) in a freshly prepared O incubation buffer (0.2 M sodium phosphate

pH 7.4, 0.1 M NaCl, 0.1 mM EDTA, 10 mM MgCl₂, 1 mM DTT, 5% glycerol) containing 2 µg of poly dI:dC (final volume of the reaction mixture was 20 µl). Incubation was performed for 15 min at 30°C, and it was stopped by placing samples on ice, and addition 1 µl of 40% saccharose. 1% agarose gel electrophoresis was run in 0.5 × TBE at 4°C, 80 V, for 3.5 h. Gels were visualized with the use of Typhoon FLA 7000 scanner (GE Healthcare).

Dimethyl Sulfide (DMS) Footprinting Labeling of Oligonucleotides With ³²P Isotope

Labeling of oligonucleotides for the primer extension reaction with a ³²P phosphorus isotope was performed using polynucleotide kinase (PNK) (Thermo Scientific). Reagents (oligonucleotides, Buffer A, γ³²P-ATP, T4 PNK – in amounts added according to the PNK manufacturer's instruction) were mixed and incubated for 25 min at 37°C. Then, 10 µl of 50 mM EDTA were added to the mixture and incubation was continued for 10 min at 75°C. Following incubation, the solution was applied to Sephadex G-50 (Sigma Aldrich) resin. The labeled oligonucleotides were purified on the resin by spinning for 5 min at 800 rpm. The efficiency of the labeling reaction was assessed by measuring the radiation of 1 µl of purified oligonucleotide sample with a Geiger counter (minimum 1000 /µl).

Obtaining DNA Template for Primer Extension Reaction

In order to prepare a modified plasmid DNA template for primer extension reactions, 1 µg of plasmid DNA (pCB104cmr or P27cmr) were incubated with proteins λ O, P27 O and 933W O for 15 min at 30°C (to achieve different molar ratios: 1:2, 1:4, 1:8, 1:16, and 1:32) in the incubation buffer [see section “Electrophoretic Mobility Shift Assay (EMSA)”. Following addition of 1 mM dimethyl sulfide (DMS) to the mixture, the incubation was continued for 5 min at 30°C. Reaction was halted by addition of 100 µl ice-cold stopping buffer (3 M ammonium acetate, 1 M β-mercaptoethanol, 20 mM EDTA) and 300 µl of 96% ethanol. DNA samples were then incubated at –80°C for 30 min. Samples were centrifuged for 30 min at 15,000 rpm at 4°C. The pellets were washed gently with 300 µl of 70% ethanol and centrifuged again under above described conditions. Precipitates were dried using a Vacufuge Concentrator (Eppendorf). Pellets were suspended in 100 µl of 1 M piperidine solution and incubated for 30 min at 90°C to break methylated DNA bonds. 100 µl of the solution were applied to Sephacryl S-500 resin (Sigma Aldrich), and centrifuged for 5 min at 500 rpm.

DNA Primer Extension Reaction

Primer extension reaction with Taq polymerase (Thermo Fisher Scientific) was carried out using an isotope-labeled oligonucleotide and modified plasmid DNA template under the conditions specified by Taq polymerase manufacturer. Following incubation scheme was applied: 3 min at 92°C, then 40 cycles of denaturation for 30 s at 92°C, annealing for 50 s at 50°C, and extension for 1 min at 72°C, and then final extension for 2 min at 72°C.

Sanger Sequencing Reaction

The sequencing reaction was carried out using the DNA Cycle Sequencing Kit (Jena Bioscience), according to the manufacturer's instructions.

DNA Electrophoresis

The 8% polyacrylamide gel was prepared using 50 ml of TBE buffer, 100 ml 40% polyacrylamide 19:1 mixture, 240 g urea and 150 ml ultrapure water. The mixture was slightly heated to dissolve urea and degassed using a vacuum pump. To 80 ml of the mixture 640 µl of 10% APS and 30 µl of TEMED were added, and the mixture was poured between glass plates of the electrophoresis apparatus. The gel polymerization lasted 1 h at room temperature. DNA sample after the primer extension reaction and sequencing was dried in a heating block at 99°C for 20–30 min. Obtained pellet was suspended in 12 µl of Loading Buffer (95% formamide, 20 mM EDTA, 0.05% bromophenol blue, 0.05% xylene cyanol FF) and 5 µl of the solution were applied onto the polyacrylamide gel. Electrophoresis was carried out in a previously heated (45–50°C) gel, at 80 W, 1900 V, and 60 mA for 3.0–3.5 h. Following electrophoresis, gel was rinsed for 5 min in water, then 5 min in 10% acetic acid and again for 5 min in water. The gel was then dried, and sealed with an ionizing radiation capture screen (GE Healthcare). The exhibition was carried out overnight. Screen signal was visualized in the Typhoon FLA 7000 scanner (GE Healthcare).

FRET Assay

15 pM of ssDNA fragment with donor (Cy3) and acceptor (Cy5) was incubated with 500 ng of purified O proteins in Incubation buffer (0.2 M sodium phosphate pH 7.4, 0.1 M NaCl, 10 mM MgCl₂, 1 mM DTT, 0.1 mM EDTA, 5% glycerol) in a final volume of 20 µl for 5 min at room temperature. Excitation was conducted at 530 nm (Cy3) and fluorescence (550–700 nm wavelength spectrum) was detected with the use of EnSpire™ Multimode Plate Reader (Perkin Elmer).

RESULTS

The *origin* of replication of lambdoid bacteriophages, located in the middle of the *O* gene, consists of sequences which bind the *O* replication initiator protein, called iterons, and the region containing large proportion of AT base pairs (the AT-rich region). As indicated in **Figure 1A**, the *origin* of bacteriophage λ DNA replication, *ori*λ, contains 4 iterons, while *origins* of the Stx phages (exemplified in this work by phages P27 and 933W) contain 6 iterons. They are numbered as 1, 1', 1'', 2, 3, and 4, where iterons 1, 2, 3, and 4 are present in λ, and iterons 1' and 1'' occur only in Stx phages. There are a few nucleotide alterations between cognate iterons (**Figure 1B**). The sequence alignment is presented in **Supplementary Figure S1**. All iterons are almost the same in all tested *origin* regions (when compared between phages), except one nucleotide difference in iteron 3 of Stx phages relative to that of λ. Thus, this Stx-specific iteron has been named 3*. However, since iterons 1 and 1'' have identical sequences, there are two possible ways of comparing

the replication regions of λ and Stx phages (Figure 1A). The presence of 2 additional iterons in DNA of Stx phages results in appearance of 13 additional amino acid residues in the O protein, relative to phage λ O gene product (Figure 1C). Moreover, the sequence of the O protein of phage P27 contains two alterations relative to λ O protein (Gly17Glu and Leu37Ile), and that of phage 933W contains three such alterations (Gly17Glu, Leu37Ile, and Ser282Gly) (Figure 1C). The O protein sequence alignment with secondary structure assignment is shown in Supplementary Figure S2. Moreover, potential secondary DNA structures formed at the iteron region are presented in Supplementary Figure S3. Considering significant differences in the regulation of DNA replication between λ and Stx phages (Nejman et al., 2009, 2011; Nowicki et al., 2015), we aimed to investigate the effects of the above mentioned alterations on formation of the nucleoprotein complexes by the O proteins of lambdoid phages and their *origin* regions.

The O proteins of phages λ , P27, and 933W were purified as described previously (Kozłowska et al., 2017). Using EMSA assay with DNA fragments containing the replication *origin* regions of phages λ and P27 (with 4 and 6 iterons, respectively), we found that all tested O proteins formed multiple complexes with both templates (Figure 2). However, binding efficiency to *origin* regions varied among the O proteins variants. It was also different for the two Stx O proteins dependent on whether the origin fragment containing 4 or 6 iterons was present in the binding reactions, whereas λ O seemed to bind similarly to both origins. For the four iteron λ *ori* both Stx O proteins formed nucleoprotein complexes of lower mass slightly more efficiently, at lower protein concentrations, than the λ O protein. However, as the protein concentration was increased, λ O formed DNA-protein complexes of higher mass more readily than both the P27 and 933W O proteins. On the other hand, the two latter O proteins formed consecutive nucleoprotein complexes with their cognate origin with lower efficiency than the λ O protein. The difference in binding efficiency in comparison to λ O was higher for the 933 O protein than for its counterpart from the phage P27 (Figure 2). These results strongly suggest that amino acid alterations between various O proteins result in differences in efficiency of formation of their complexes with the *origin* region. However, it was difficult to verify in the EMSA test if all iterons in the Stx *ori* are bound by the O proteins at higher protein concentrations, since nucleoprotein complexes of high mass often formed diffused bands, migrating at variable positions, reflecting their lower stability, various conformations or protein aggregation. Discrete bands representing 4, 5, or 6 dimers of the O proteins bound could not be distinguished.

The differences in interactions of O proteins derived from λ and Stx phages to *origin* regions might result from either various binding cooperativity properties of the initiator proteins or lower affinity of O proteins of Stx phages to iterons. Therefore, we tested efficiency of binding of O proteins to individual iterons, using templates containing single iterons: 1/1'', 1', 2, 3, 3*, 4. We found that all tested O proteins formed complexes with each tested iteron, however, efficiency of their formation were slightly different. Phage λ O protein forms two different complexes with each iteron while O proteins of Stx phages form predominantly

one complex (Figure 3). Again, binding of the λ O protein was the strongest among tested proteins. Moreover, iterons 2 and 3* were bound with the highest efficiency by all tested O proteins, while formation of the nucleoprotein complexes was the weakest for iterons 1' and 4 (Figure 3). Nevertheless, the differences between efficiency of binding of O proteins by individual iterons were not dramatic.

More detailed information about binding of the O proteins to iterons were obtained in footprinting experiments. Results of experiments with both types of *origin* regions (with 4 and 6 iterons) confirmed the efficiency of binding of different variants of the O protein, with λ O being the most effective and the 933W O protein the least effective in interactions with iterons (Figures 4, 5 for results of experiments with the templates containing 4 and 6 iterons, respectively). In experiments with the template containing four iterons, all O proteins bound iterons 2 and 3 with the highest efficiency, then iteron 1, and iteron 4 with the lowest efficiency (Figure 4). Protection of the DNA sequence, indicating the contact sites with the O protein, is evident on both sides of the template, and includes two distal G residues in the G-tracks present in each iteron. Enhancement of the signal, indicating DNA bending, can be seen also at both template sides, between O-bound residues and in the middle of each iteron (Figure 4). These results are in agreement with previously published observations (Zahn and Blattner, 1985). Importantly, when the template containing six iterons was used, only four of them were bound efficiently by O proteins in each experimental system (i.e., with the O protein of phages λ , P27 and 933W). Sequences of two proximal iterons were not clearly protected by any of the tested O proteins, though weak interactions of iteron 1' with O at the highest protein concentrations could be observed (Figure 5). Therefore, we conclude that the arrangement of iterons in the Stx phage *origin* relative to λ should be represented as indicated in the lower part of Figure 1A. At higher protein concentrations, λ O protein bound iteron 1', although with lower efficiency than the four iterons proximal to the AT-rich region, and very weakly interacted with the most distant iteron 1''. On the other hand the P27 and W933 O proteins showed even weaker footprint on iteron 1', with very low level of protection and moderate enhancement at the correct nucleotides. The signal of binding was very weak at protein concentrations which resulted in full occupancy of the consecutive iterons 1/1'', 2, 3, and 4 by both proteins (Supplementary Figure S4). Similar, but even weaker pattern was observed with respect to all O proteins at the 1/1'' iteron situated at the edge of the origin.

Iterons are intrinsically bent sequences (Zahn and Blattner, 1985) and their anisotropy could affect binding parameters of the O protein. To assess whether curvature could underlie slight differences in binding of the O proteins to individual iterons, we performed an analysis of iterons geometry using AA Wedge model (Goodsell and Dickerson, 1994). The data obtained with DNA Curvature Analysis online application¹ showed that iterons 1/1'', 1', 2, and 4 have identical local bending pattern. On the other hand, iterons 3 and 3*, which contain a track of three A, instead of four present in the remaining iterons, have

¹<https://www.lfd.uci.edu/~gohlke/dnacurve/>




FIGURE 2 | Interactions of O proteins of phages λ , P27 and 933W with DNA templates containing four panel (A) or six panel (B) iterons, as assessed by EMSA. Increasing molar ratios of DNA:O proteins, indicated in particular wells, were as follows: 1:2, 1:4, 1:8, 1:16, and 1:32. Lane no. 1 contains no O protein.

slightly different curvature from the rest but indistinguishable from each other (**Supplementary Figure S5**). This suggests that differences in intrinsic DNA bending between individual iterons do not account for the observed preferences of the O proteins to particular sequences present in the origin context.

Using the tool mentioned above, we also analyzed DNA curvature of the entire *origin* regions, both containing 4

and 6 iterons. The results of the analysis (**Supplementary Figures S6, S7**) indicate that both types of origins contain a significantly curved region encompassing iterons 1–3. Both iteron 4 and especially iterons 1'–1'' are separated by regions of low curvature from iterons 1–3. The DNA structure of phage origins could be related to the observed hierarchy of iterons occupation by the O




FIGURE 3 | Interactions of O proteins of phages λ , P27 and 933W with DNA templates containing single itérons: iteron 1/1'' (A), iteron 1' (B), iteron 2 (C), iteron 3 (D), iteron 3* (E), and iteron 4 (F). Increasing molar ratios of DNA:O proteins, indicated in particular wells, were as follows: 1:1, 1:2, 1:4, and 1:8. Lane no. 1 contains no O protein. Note that lanes 10/11 in panel A, and lanes 4/5 in panel D, are swapped.

replication initiators, with itérons 4 and 1'-1'' being the least preferred.

To test if the additional itérons occurring in the *origin* regions of Stx phages are functionally significant, we have constructed plasmids bearing disrupted iteron 1 and/or 1' sequence(s). DNA templates bearing such mutations bound λ O only very weakly at the highest used amounts of the protein, and did not bind the P27 O protein at all (Figure 6). Some differences in results of experiments with wild-type itérons may be due to differences in the DNA fragment lengths (55 vs. 28 bps). The presence of two bands of free DNA for wild-type itérons but only one band for mutated ones is apparently because of formation of bending in wild-type sequence which disappears when the mutation is present. To test functionality of the iteron 1', double-*origin* plasmids have been constructed which

contained lambdoid (P27-derived) replication region and *origin* of plasmid R6K, active only in the presence of the active *pir* gene. Sequences of itérons 1 and 1' were changed as described above, and the alterations of base pairs were chosen to obtain disrupted O-binding properties and unchanged amino acid sequences of the O proteins, encoded by the O genes (as mentioned, the *origin* is located in the middle of the O gene). Replication of plasmids containing either wild-type or mutated itérons was possible in the *E. coli* strain expressing the *pir* gene due to activity of the *ori* R6K, irrespective of the activity of lambdoid *origin*. However, transformation of *E. coli* cells devoid of the *pir* gene was possible only in the presence of the active lambdoid *origin*. We have tested efficiency of transformation by lambdoid plasmids bearing either wild-type itérons or those with mutations in iteron 1 and/or 1' of *E. coli* wild-type host, as well as *dnaA46*, Δ *seqA* or




FIGURE 4 | Interactions of O proteins of phages λ, P27 and 933W with DNA templates containing four iterons, as assessed by DMS footprinting. Increasing molar ratios of DNA:O proteins, indicated in particular wells, were as follows: 1:0 (no O protein), 1:2, 1:4, 1:8, 1:16, and 1:32. Arrows indicate major changes (enhancement) in bands' intensities.





FIGURE 5 | Interactions of O proteins of phages λ, P27 and 933W with DNA templates containing six iterons, as assessed by DMS footprinting. Increasing molar ratios of DNA:O proteins, indicated in particular wells, were as follows: 1:0 (no O protein), 1:2, 1:4, 1:8, 1:16, and 1:32. Arrows indicate major changes (enhancement) in bands' intensities.



double *dnaA46* Δ *seqA* mutants. These mutants were used since it was previously demonstrated that: (i) incompatibility between λ plasmids and *dnaA* mutations arises from both inefficient stimulation of transcription from the *p_R* promoter, responsible for transcriptional activation of the *origin*, by the mutant DnaA protein and the competition between the lambda P protein and the host DnaA and DnaC proteins for DnaB helicase, (ii) both mechanisms must be operative for the incompatibility, and (iii) the incompatibility is abolished in the absence of the *seqA* gene function (Glinkowska et al., 2001) as the *seqA* gene product modulates *p_R* activity (Słomińska et al., 2003a) and interplays with DnaA at this promoter (Słomińska et al., 2003b). We found that disruption of either interon 1 or interon 1' or both abolished ability of lambdoid plasmid to transform *E. coli* wild-type strain.

The presence of mutation(s) in *dnaA*, *seqA* or both these genes did not change the plasmid inability to transform host cells (Table 2). We conclude, that both interon 1 and interon 1' are necessary for efficient replication of lambdoid plasmids derived from Stx phages, despite only weak interactions of the latter interon with the O protein.

In the lambdoid phage *origin* of replication, iterons are flanked by the AT-rich region (see Figure 1A). To test if the AT-rich sequence influence binding of the O protein to iterons, we have replaced the original AT-rich region with that containing 57% GC base pairs. We found that such a change did not influence efficiency of binding of O protein to iterons (Figure 7). Patterns of formed nucleoprotein complexes were also not influenced by replacement of the AT-rich region with a neutral sequence. This

TABLE 2 | Transformation efficiency of *E. coli* wild-type, *dnaA*, *seqA*, and *dnaA seqA* hosts by derivatives of p27cmr plasmids bearing mutated iteron sequences 1 and 1' at 30°C. Results are presented as mean values of three independent experiments ± SD.

Plasmid	Efficiency of transformation (transformants/μg of plasmid DNA)			
	<i>dnaA</i> ⁺ <i>seqA</i> ⁺	<i>dnaA46 seqA</i> ⁺	<i>dnaA</i> ⁺ <i>ΔseqA</i>	<i>dnaA46 ΔseqA</i>
p27R6K	5.38 ± 1.37 × 10 ⁴	3.79 ± 0.39 × 10 ⁴	<10 ¹	7.00 ± 3.49 × 10 ²
p27mut1	<10 ¹	<10 ¹	<10 ¹	<10 ¹
p27mut1'	<10 ¹	<10 ¹	<10 ¹	<10 ¹
p27mut1 + 1'	<10 ¹	<10 ¹	<10 ¹	<10 ¹

was true for both templates, containing either four or six iterons (Figures 7A,B, respectively). The presence of two bands for free DNA (Figure 7B, lines 1–14) and the presence of six protein-DNA bands (line 26) can be explained by formation of secondary structured by ssDNA fragments.

Binding of phage λ O protein to single stranded DNA (ssDNA) has been demonstrated previously (Learn et al., 1997). Therefore, we have asked if all tested O proteins can interact with ssDNA of the AT-rich region. The analysis was performed using FRET technique. ssDNA was labeled with Cy3 and Cy5 fluorescent dyes forming the donor-acceptor pair. For the free ssDNA, donor excitation resulted in effective energy transfer and an increase in acceptor fluorescence emission concomitant with a decrease of the donor fluorescence. Binding of the O proteins caused reduction of FRET efficiency observed as an increase of the donor signal and diminution of the acceptor fluorescence. Results of these experiments indicated that O proteins of phages λ, P27 and 933W can efficiently interact with ssDNA, indeed (Figure 8). The fluorescence signals were similar in all experiments in which O proteins (from λ, P27 and 933W) were present (Figure 8), suggesting that the investigated mechanism does not differ between tested phages.

DISCUSSION


Since DNA replication of Stx phages appears to be necessary for effective production of Shiga toxins by STEC strains (Nejman-Faleńczyk et al., 2012; Nowicki et al., 2013; Balasubramanian et al., 2019), understanding of the regulation of this process is crucial not only for basic science but also for applications related to development of novel anti-STECS treatments. Although replication regions of genomes of Stx phages are highly similar to that of bacteriophage λ, there are important differences (summarized in Figure 1 and Supplementary Figures S1–S3) which result in significantly altered regulation of Stx phage DNA replication relative to λ (Nejman et al., 2009, 2011; Nowicki et al., 2015). Therefore, we decided to investigate details of molecular interactions between the initiator O proteins of Stx phages with the *origin* regions, using purified proteins and nucleic acids.

Previously published results (Zahn and Blattner, 1985, 1987; Schnos et al., 1988) indicated that the 19 bp (imperfect) inverted

repeat of the iteron is the minimal sequence required for efficient binding by the λ O protein. At low protein concentrations, only the inner two iterons were bound, while binding to all four iterons occurred at higher protein concentrations. The O protein binds to the replication origin in both linear and circular (relaxed) forms, but *origin* unwinding requires the negatively supercoiled DNA form introduced by the host gyrase. Results of our experiments indicated that O proteins of Stx bacteriophages bind specifically to iterons in the *origin* regions (Figure 2; see also Kozłowska et al., 2017), however, these interactions are weaker relative to binding of bacteriophage λ O protein to its specific DNA target sequences (Figure 2). Apparently, amino acid alterations in O proteins of Stx phages relative to the λ O protein, listed in Figure 1C and indicated in Supplementary Figure S2), are responsible for these differences in their affinity to iterons. Although each individual iteron can be bound by any of tested O proteins (encoded by λ, P27 or 933W phages) (Figure 3), when the whole *origin* region was considered, only four iterons were involved in effective interactions with O protein molecules, irrespective of the presence of either four or six such DNA sequences in the template (Figures 4, 5). Nevertheless, two additional iterons (1' and 1'') in *origin* of Stx phage can be bound by the O protein and are necessary for effective DNA replication (Figure 6, Table 2, and Supplementary Figure S4).

The AT-rich region appears to be dispensable for formation of the O-some structure (a specific nucleoprotein structure containing the O protein and iterons) (Figure 7). In fact, it was demonstrated (Dodson et al., 1986) that two iterons and the adjacent AT-rich region form the minimal DNA fragment competent for unwinding. Moreover, for *in vitro* replication of *oriλ*-based plasmids, only the two iterons proximal to the AT-rich region were required (Wickner and McKenney, 1987). Since this AT-rich tract is crucial for DNA unwinding during replication initiation, and all tested O protein variants revealed similar ability to interact with single stranded DNA rich in A and T residues (Figure 8), one may suggest that the mechanism of DNA stretching is the same for all tested phages.


The presence of additional two iterons in *origin* sequences of Stx phages relative to λ results also in appearance of additional 13 amino acid residues in corresponding O proteins, as iterons are located in the middle of the O gene (Figure 1). One might ask if such additional protein fragment can influence interaction between O and its DNA target. The secondary structure prediction of O proteins (Supplementary Figure S2) suggests that the additional protein fragments are not involved in interactions with DNA. Regarding the role of these two additional iterons in Stx phages, on one hand, it was reported that three iterons are sufficient for efficient replication of phage λ, as suggested by the normal growth and burst size of the mutant phage (Moore et al., 1981). On the other hand, additional iterons were found here to be necessary for transformation of P27-derived plasmids (Table 2). Therefore, one might propose that iterons 1' and 1'' ensure either elevation of the local concentration of the O protein or stabilization of the O-some which can be crucial due to lower affinity of Stx O proteins to iterons, relative to λ O. Such a model is compatible with the already described mechanism of DNA-protein interactions which usually requires



a proximity of A residue stretch in DNA (Rohs et al., 2009), and such a motif is actually present in iterons located in the *origin* of lambdoid phage replication (see **Figure 1B**).

There is also a question about potential roles of hairpin and cruciform structures, potentially formed by the replication region of lambdoid phages. Roles of such structures in DNA replication have been discussed previously (Bikard et al., 2010). In fact, each iteron in lambdoid *origin* is an inverted repeat, and in the sequences of the entire gene O, the biggest and most stable hairpin structures are encoded by the iterons 2 and 4, 1' and 4, and 2 and 3 (**Supplementary Figure S3**). It appears that duplication of the iterons in Stx phages may influence folding of the potential hairpin/cruciform structures. This may influence efficiency of binding of the O protein and formation of the O-some, and in fact, Stx replication regions are bound less efficiently than that of λ . Therefore changes in secondary DNA structures might be an additional role of additional iterons in Stx phages.

Recent studies indicated that DNA replication may significantly influence the major developmental decision of bacteriophage λ , whether to produce progeny phages and lyse the host cell or to lysogenize it (Shao et al., 2018). Since such decision is crucial for all lambdoid phages, one can speculate that λ and Stx phages evolved in somewhat different ways to produce optimal regulatory systems functioning in environments they occur in. Less stringent binding of the O protein to iterons, and the presence of additional iterons which can modulate formation of the O-some, possibly in response to various environmental conditions, might be profitable for Stx phages which have to quickly change the developmental pathway. Gastrointestinal tract of ruminants is their primary habitat in which STEC strains developed a specific Stx phage-dependent strategy to fight unicellular eukaryotic predators, like ciliates (Łoś et al., 2013). Therefore, ability to adjust the genetic switch between lysogenic and lytic developmental options rapidly may require more flexible regulation of DNA replication initiation, ensured



by modulatory effects of additional iterons and weaker binding of the O protein to them. On the other hand, phage λ that was discovered as a parasite of the *E. coli* K12 strain (Lederberg and Lederberg, 1953), a bacterium originally isolated from human stool and known to occur in human and animal intestine as well as in natural waters (Kuhnert et al., 1995; Na et al., 2006). Hence, this phage might prefer more strict developmental decisions which can be effectively made when employing stronger interactions between the O initiator protein and the replication *origin*. This strategy may ensure more unequivocal lysis-vs.-lysogenization decision which is compatible with the proposal of Shao et al. (2018) about the coupling of phage DNA replication to the λ developmental choice.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation, to any qualified researcher.

AUTHOR CONTRIBUTIONS

KK participated in planning experiments, performed most of them, participated in data interpretation, and drafting the manuscript. MG participated in planning experiments, data analysis, and revising the manuscript. LB participated in constructing of plasmids, performing EMSA, conducting experiments with mutant plasmids, performing DMS footprinting experiments, and revising the manuscript. LG participated in data analysis, preparation of figures, and revising the manuscript. JD assisted in EMSA experiments. GW was principal investigator of the project, supervised the work, and drafted the manuscript. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by the National Science Center (Poland) (project grant no. 2011/02/A/NZ1/00009).

ACKNOWLEDGMENTS

We are grateful to Anna Zawilak-Pawlik and Paweł Donczew for consultations and advices regarding DMS footprinting experiments.

REFERENCES

- Actis, G. C. (2014). The gut microbiome. *Inflamm. Allergy Drug Targets*. 13, 217–223.
- Balasubramanian, S., Osburne, M. S., Brin-Jones, H., Tai, A. K., and Leong, J. M. (2019). Prophage induction, but not production of phage particles, is required for lethal disease in a microbiome-replete murine model of enterohemorrhagic *E. coli* infection. *PLoS Pathog.* 15:e1007494. doi: 10.1371/journal.ppat.1007494
- Bikard, D., Loot, C., Baharoglu, Z., and Mazel, D. (2010). Folded DNA in action: hairpin formation and biological functions in prokaryotes. *Microbiol. Mol. Biol. Rev.* 74, 570–588. doi: 10.1128/MMBR.00026-10
- Chen, J., Li, Y., Zhang, K., and Wang, H. (2018). Whole-genome sequence of phage-resistant strain *Escherichia coli* DH5 α . *Genome Announc.* 6, : e00097–e-18.
- Devleeschauwer, B., Pires, S. M., Young, I., Gill, A., Majowicz, S. E., and the study team. (2019). Associating sporadic, foodborne illness caused by Shiga toxin-producing *Escherichia coli* with specific foods: a systematic review and meta-analysis of case-control studies**. *Epidemiol. Infect.* 147, :e235. doi: 10.1017/S0950268819001183
- Dodson, M., Echols, H., Wickner, S., Alfano, C., Mensa-Wilmot, K., Gomes, B., et al. (1986). Specialized nucleoprotein structures at the origin of replication of bacteriophage lambda: localized unwinding of duplex DNA by a six protein reaction. *Proc. Natl. Acad. Sci. U.S.A.* 83, 7638–7642. doi: 10.1073/pnas.83.20.7638
- Glinkowska, M., Konopa, G., Węgrzyn, A., Herman-Antosiewicz, A., Weigel, C., Seitz, H., et al. (2001). The double mechanism of incompatibility between lambda plasmids and *Escherichia coli* dnaA(ts) host cells. *Microbiology* 147, 1923–1928. doi: 10.1099/0021287-147-7-1923
- Goodsell, D. S., and Dickerson, R. E. (1994). Bending and curvature calculations in B-DNA. *Nucleic Acids Res.* 22, 5497–5503. doi: 10.1093/nar/22.24.5497
- Kakoullis, L., Papachristodoulou, E., Chra, P., and Panos, G. (2019). Shiga toxin-induced haemolytic uraemic syndrome and the role of antibiotics: a global overview. *J. Infect.* 79, 75–94. doi: 10.1016/j.jinf.2019.05.018
- Karmali, M. A. (2018). Factors in the emergence of serious human infections associated with highly pathogenic strains of shiga toxin-producing *Escherichia coli*. *Int. J. Med. Microbiol.* 308, 1067–1072. doi: 10.1016/j.ijmm.2018.08.005
- Kintz, E., Byrne, L., Jenkins, C., McCarthy, N., Vivancos, R., and Hunter, P. (2019). Outbreaks of Shiga toxin-producing *Escherichia coli* linked to sprouted seeds, salad, and leafy greens: a systematic review. *J. Food Prot.* 82, 1950–1958. doi: 10.4315/0362-028X.JFP-19-014
- Kozłowska, K. I., Tymecka-Mulik, J., and Węgrzyn, G. (2017). Purified Stx and λ phage initiator O proteins bind specifically to two different origins of replication in vitro. *Protein Expr. Purif.* 131, 16–26. doi: 10.1016/j.pep.2016.11.002
- Krüger, A., and Lucchesi, P. M. (2015). Shiga toxins and stx phages: highly diverse entities. *Microbiology* 161, 451–462. doi: 10.1099/mic.0.000003
- Kuhnert, P., Nicolet, J., and Frey, J. (1995). Rapid and accurate identification of *Escherichia coli* K-12 strains. *Appl. Environ. Microbiol.* 61, 4135–4139. doi: 10.1128/aem.61.11.4135-4139.1995
- Learn, B. A., Um, S. J., Huang, L., and McMacken, R. (1997). Cryptic single-stranded-DNA binding activities of the phage lambda P and *Escherichia coli* DnaC replication initiation proteins facilitate the transfer of *E. coli* DnaB helicase onto DNA. *Proc. Natl. Acad. Sci. U.S.A.* 94, 1154–1159. doi: 10.1073/pnas.94.4.1154
- Lederberg, E. M., and Lederberg, J. (1953). Genetic studies of lysogenicity in *Escherichia coli*. *Genetics* 38, 51–64.
- Leimbach, A., Hacker, J., and Dobrindt, U. (2013). *E. coli* as an all-rounder: the thin line between commensalism and pathogenicity. *Curr. Top. Microbiol. Immunol.* 358, 3–32. doi: 10.1007/82_2012_303
- Licznarska, K., Nejman-Faleńczyk, B., Bloch, S., Dydecka, A., Topka, G., Gąsior, T., et al. (2016). Oxidative stress in Shiga toxin production by enterohemorrhagic *Escherichia coli*. *Oxid. Med. Cell. Longev.* 2016, :3578368. doi: 10.1155/2016/3578368
- Łoś, J. M., Łoś, M., Węgrzyn, A., and Węgrzyn, G. (2013). Altruism of Shiga toxin-producing *Escherichia coli*: recent hypothesis versus experimental results. *Front. Cell. Infect. Microbiol.* 2:166. doi: 10.3389/fcimb.2012.00166
- Łoś, J. M., Łoś, M., and Węgrzyn, G. (2011). Bacteriophages carrying Shiga toxin genes: genomic variations, detection and potential treatment of pathogenic bacteria. *Future Microbiol.* 6, 909–924. doi: 10.2217/FMB.11.70
- Mir, R. A., and Kudva, I. T. (2019). Antibiotic-resistant Shiga toxin-producing *Escherichia coli*: an overview of prevalence and intervention strategies. *Zoonoses Public Health*. 66, 1–13. doi: 10.1111/zph.12533
- Moore, D. D., Denniston, K. J., and Blattner, F. R. (1981). Sequence organization of the origins of DNA replication in lambdoid coliphages. *Gene* 14, 91–101. doi: 10.1016/0378-1119(81)90151-7
- Na, S. H., Miyayaga, K., Unno, H., and Tanji, Y. (2006). The survival response of *Escherichia coli* K12 in a natural environment. *Appl. Microbiol. Biotechnol.* 72, 386–392. doi: 10.1007/s00253-005-0268-3
- Nejman, B., Łoś, J. M., Łoś, M., Węgrzyn, G., and Węgrzyn, A. (2009). Plasmids derived from lambdoid bacteriophages as models for studying replication of mobile genetic elements responsible for the production of Shiga toxins by pathogenic *Escherichia coli* strains. *J. Mol. Microbiol. Biotechnol.* 17, 211–220. doi: 10.1159/000242447
- Nejman, B., Nadratowska-Wesołowska, B., Szalewska-Pałasz, A., Węgrzyn, A., and Węgrzyn, G. (2011). Replication of plasmids derived from Shiga toxin-converting bacteriophages in starved *Escherichia coli*. *Microbiology* 157, 220–233. doi: 10.1099/mic.0.042820-0
- Nejman-Faleńczyk, B., Golec, P., Maciąg, M., Węgrzyn, A., and Węgrzyn, G. (2012). Inhibition of development of Shiga toxin-converting bacteriophages by either treatment with citrate or amino acid starvation. *Foodborne Pathog. Dis.* 9, 13–19. doi: 10.1089/fpd.2011.0980
- Nowicki, D., Bloch, S., Nejman-Faleńczyk, B., Szalewska-Pałasz, A., Węgrzyn, A., and Węgrzyn, G. (2015). Defects in RNA polyadenylation impair both lysogenization by and lytic development of Shiga toxin-converting bacteriophages. *J. Gen. Virol.* 96, 1957–1968. doi: 10.1099/vir.0.000102
- Nowicki, D., Kobiela, W., Węgrzyn, A., Węgrzyn, G., and Szalewska-Pałasz, A. (2013). ppGpp-dependent negative control of DNA replication of Shiga toxin-converting bacteriophages in *Escherichia coli*. *J. Bacteriol.* 195, 5007–5015. doi: 10.1128/JB.00592-13
- Paletta, A. C. C., Castro, V. S., and Conte-Junior, C. A. (2020). Shiga toxin-producing and enteroaggregative *Escherichia coli* in animal, foods, and humans: pathogenicity mechanisms, detection methods, and epidemiology. *Curr. Microbiol.* 77, 612–620. doi: 10.1007/s00284-019-01842-1
- Rohs, R., West, S. M., Sosinsky, A., Liu, P., Mann, R. S., and Honig, B. (2009). The role of DNA shape in protein-DNA recognition. *Nature* 461, 1248–1253. doi: 10.1038/nature08473
- Schnos, M., Zahn, K., Inman, R. B., and Blattner, F. R. (1988). Initiation protein induced helix destabilization at the lambda origin: a prepriming step in DNA replication. *Cell* 52, 385–395. doi: 10.1016/s0092-8674(88)80031-x
- Shao, Q., Cortes, M. G., Trinh, J. T., Guan, J., Balázs, G., and Zeng, L. (2018). Coupling of DNA replication and negative feedback controls gene expression for cell-fate decisions. *iScience* 6, 1–12. doi: 10.1016/j.isci.2018.07.006
- Słomińska, M., Konopa, G., Barańska, S., Węgrzyn, G., and Węgrzyn, A. (2003b). Interplay between DnaA and SeqA proteins during regulation of bacteriophage lambda pR promoter activity. *J. Mol. Biol.* 329, 59–68. doi: 10.1016/s0022-2836(03)00378-4
- Słomińska, M., Konopa, G., Ostrowska, J., Kedzierska, B., Węgrzyn, G., and Węgrzyn, A. (2003a). SeqA-mediated stimulation of a promoter activity by facilitating functions of a transcription activator. *Mol. Microbiol.* 47, 1669–1679. doi: 10.1046/j.1365-2958.2003.03392.x

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmolb.2020.00207/full#supplementary-material>

- Szalewska-Pałasz, A., Węgrzyn, A., Herman, A., and Węgrzyn, G. (1994). The mechanism of the stringent control of lambda plasmid DNA replication. *EMBO J.* 13, 5779–5785. doi: 10.1002/j.1460-2075.1994.tb06916.x
- Szambowska, A., Pierechod, M., Węgrzyn, G., and Glinkowska, M. (2011). Coupling of transcription and replication machineries in λ DNA replication initiation: evidence for direct interaction of *Escherichia coli* RNA polymerase and the λ O protein. *Nucleic Acids Res.* 39, 168–177. doi: 10.1093/nar/gkq752
- Taylor, K., and Węgrzyn, G. (1995). Replication of coliphage lambda DNA. *FEMS Microbiol. Rev.* 17, 109–119. doi: 10.1111/j.1574-6976.1995.tb00192.x
- Taylor, R. G., Walker, D. C., and McInnes, R. R. (1993). *E. coli* host strains significantly affect the quality of small scale plasmid DNA preparations used for sequencing. *Nucleic Acids Res.* 21, 1677–1678. doi: 10.1093/nar/21.7.1677
- Węgrzyn, G., Licznarska, K., and Węgrzyn, A. (2012). Phage λ - new insights into regulatory circuits. *Adv. Virus Res.* 82, 155–178. doi: 10.1016/B978-0-12-394621-8.00016-9
- Weigel, C., and Seitz, H. (2006). Bacteriophage replication modules. *FEMS Microbiol. Rev.* 30, 321–381. doi: 10.1111/j.1574-6976.2006.00015.x
- Wickner, S., and McKenney, K. (1987). Deletion analysis of the DNA sequence required for the in vitro initiation of replication of bacteriophage lambda. *J. Biol. Chem.* 262, 13163–13167.
- Yanisch-Perron, C., Vieira, J., and Messing, J. (1985). Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* 33, 103–119. doi: 10.1016/0378-1119(85)90120-9
- Zahn, K., and Blattner, F. R. (1985). Binding and bending of the lambda replication origin by the phage O protein. *EMBO J.* 4, 3605–3616. doi: 10.1002/j.1460-2075.1985.tb04124.x
- Zahn, K., and Blattner, F. R. (1987). Direct evidence for DNA bending at the lambda replication origin. *Science* 236, 416–422. doi: 10.1126/science.2951850

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Kozłowska, Glinkowska, Boss, Gaffke, Deptuła and Węgrzyn. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.