



Absence of Cardiolipin From the Outer Leaflet of a Mitochondrial Inner Membrane Mimic Restricts Opa1-Mediated Fusion

Yifan Ge^{1,2†*}, Sivakumar Boopathy^{1,2}, Tran H. Nguyen^{1,2}, Camila Makhoulouta Lugo^{1,2} and Luke H. Chao^{1,2*}

¹Department of Molecular Biology, Massachusetts General Hospital, Boston, MA, United States, ²Department of Genetics, Harvard Medical School, Boston, MA, United States

OPEN ACCESS

Edited by:

James Munro,
University of Massachusetts Medical
School, United States

Reviewed by:

Junjie Hu,
Institute of Biophysics (CAS), China
IVAN LOPEZ-MONTERO,
Complutense University of Madrid,
Spain

*Correspondence:

Yifan Ge
yifange@ntu.edu.cn
Luke H. Chao
chao@molbio.mgh.harvard.edu

†Present address:

Yifan Ge,
Key Laboratory of Neuroregeneration,
Jiangsu Province and Ministry of
education, Nantong University,
Nantong, Jiangsu, China

Specialty section:

This article was submitted to
Biophysics,
a section of the journal
Frontiers in Molecular Biosciences

Received: 01 September 2021

Accepted: 07 December 2021

Published: 22 December 2021

Citation:

Ge Y, Boopathy S, Nguyen TH,
Lugo CM and Chao LH (2021)
Absence of Cardiolipin From the Outer
Leaflet of a Mitochondrial Inner
Membrane Mimic Restricts Opa1-
Mediated Fusion.
Front. Mol. Biosci. 8:769135.
doi: 10.3389/fmolb.2021.769135

Cardiolipin is a tetra-acylated di-phosphatidylglycerol lipid enriched in the matrix-facing (inner) leaflet of the mitochondrial inner membrane. Cardiolipin plays an important role in regulating mitochondria function and dynamics. Yet, the mechanisms connecting cardiolipin distribution and mitochondrial protein function remain indirect. In our previous work, we established an *in vitro* system reconstituting mitochondrial inner membrane fusion mediated by Opa1. We found that the long form of Opa1 (l-Opa1) works together with the proteolytically processed short form (s-Opa1) to mediate fast and efficient membrane fusion. Here, we extend our reconstitution system to generate supported lipid bilayers with asymmetric cardiolipin distribution. Using this system, we find the presence of cardiolipin on the inter-membrane space-facing (outer) leaflet is important for membrane tethering and fusion. We discuss how the presence of cardiolipin in this leaflet may influence protein and membrane properties, and future applications for this approach.

Keywords: cardiolipin, mitochondrial fusion, membrane asymmetry, OPA1, membrane heterogeneity

INTRODUCTION

Mitochondria are important eukaryotic organelles that regulate cell metabolism, signaling and death (McBride et al., 2006; Giacomello et al., 2020). Mitochondria feature a unique double membrane structure. The outer mitochondrial membrane (OMM) hosts machinery for lipid and protein transport, while the inner mitochondrial membrane (IMM), with its cristae invaginations, is the site for ATP synthesis and respiratory function (Kroemer et al., 2007) (Cogliati et al., 2016; Kondadi et al., 2020). Mitochondrial membranes are dynamic, undergoing fusion and fission to distribute metabolites and buffer mutation in mitochondrial DNA, while changing morphology in response to physiological conditions (Chan, 2020; Gao and Hu, 2021).

While phospholipid composition is recognized as a key regulator of membrane properties, the functional roles of phospholipid distribution are challenging to investigate (Harayama and Riezman, 2018). The mitochondrial membrane is distinct in composition from other endomembrane systems, and comprises cardiolipin (CL), phosphocholine (PC), and phosphatidylethanolamine (PE) (Horvath and Daum, 2013). Other lipid species that may play important roles include phosphatidylinositol (PI), phosphatidylserine (PS), phosphatidylglycerol (PG), phosphatidic acid

(PA), lysophospholipids, sterols, sphingomyelin and low levels of cholesterol (CH) (Horvath and Daum, 2013).

In eukaryotes, CL is uniquely found in the mitochondria. Consistent with an endosymbiotic origin, CL is found in bacteria (Friedman and Nunnari, 2014). CL comprises ~20% of total mitochondrial phospholipid content. In the OMM, PC and PE are major membrane components. In contrast, PE and CL makes up nearly 50% of the total phospholipid in the IMM (Horvath and Daum, 2013; Oemer et al., 2018; Oemer et al., 2020). Electron paramagnetic resonance spectroscopy studies of yeast mitochondria show CL enrichment in the inner leaflet of the IMM (Petit et al., 1994). An inner-leaflet IMM distribution of CL was also shown using fluorescent probes that specifically target CL (Krebs et al., 1979; Harb et al., 1981; Petit et al., 1994). These observations are consistent with data indicating CL is synthesized from PA in the matrix-proximal face of the IMM (Schlame, 2008). While the distribution between leaflets may be dynamic, data suggest CL also comprises ~5–15% of the total lipids in the inter-membrane space (IMS)-facing (outer) leaflet (Harb et al., 1981; Gallet et al., 1997).

CL has four acyl chains, forming a unique conical structure expected to have high flexibility. This structure classifies CL as a ‘non-structural’ phospholipid that tends to introduce a hexagonal phase in mixtures (Kagan et al., 2014). The high concentration of PE and CL in the IMM is expected to alter membrane tension (Traïkia et al., 2002), which influences membrane bending. Differential stress between leaflets can change membrane elasticity, stiffness and lateral pressure (Hossein and Deserno, 2020). These changes can favor incorporation of transmembrane proteins that induce curvature, association of peripheral membrane proteins, or self-assembly of proteins on either leaflet (Wilson et al., 2019; Lorent et al., 2020).

In addition to inter-leaflet asymmetry, lipids can exhibit planar segregation (Doktorova et al., 2020; Lorent et al., 2020). Lateral heterogeneity in the plasma membrane has been more extensively studied (Lorent et al., 2020). For example, model plasma membrane experiments show planar segregation coupled to inter-leaflet asymmetry can affect lipid packing and hydrophobic thickness, to sequester transmembrane proteins (Hussain et al., 2013). Similar physical consequences of planar segregation may be at play in the IMM. Although lateral segregation and principles of IMM organization are less well studied in well-controlled biophysical systems, cellular experiments suggest membrane heterogeneity may play a role in organizing respiratory chain super-complexes (Dudek, 2017).

Theory and computation have been applied to investigate the physical consequences of heterogeneity in CL distribution (Lemmin et al., 2013; Boyd et al., 2017; Elias-Wolff et al., 2019; Wilson et al., 2019). Not surprisingly, CL distribution is predicted to strongly shape the electrostatic potential of membrane surface, with cation interactions observed to introduce lateral CL heterogeneity in simulations. (Lemmin et al., 2013). Boyd and coworkers demonstrated concentration-dependent bending and buckling of CL bilayers, with CL mediating regions of high curvature, and CL concentrating in the negatively curved regions (Boyd et al., 2017). This finding was supported by work from Elias-Wolff and coworkers, which also

found CL inter-leaflet asymmetry contributes to membrane buckling (Elias-Wolff et al., 2019).

CL distribution can change in response to environmental stimuli, and has important consequences in disease (Gallet et al., 1997). Recent studies show CL can be externalized by pro-mitophagy stimuli in primary cortical neurons and in cellular models of Parkinson’s disease (Chu et al., 2013). Externalization of CL to the outer leaflet of the OMM activates binding to microtubule-associated-protein-1 light chain 3, which localizes damaged mitochondria to autophagosomes (Chu et al., 2013; Ryan et al., 2018). CL externalization is recognized by macrophages to cause inflammation (Pizzuto and Pelegrin, 2020). These processes are associated with altered fission and fusion dynamics and altered mitochondrial morphology (Kordower et al., 2008; Van Laar and Berman, 2009). Many mitochondrial proteins depend on CL for activity (Ban et al., 2010; Zhang et al., 2020), yet the role of asymmetric lipid distribution on protein activity has been technically challenging to investigate.

In our previous work, we fabricated a model membrane system that mimicked the IMM in composition. We reconstituted Opa1 (the IMM fusogen) into a supported bilayer and noticed CL played a key role in several stages of membrane fusion (in particular, GTP-dependent membrane tethering). In our previous study, we focused on understanding how the full-length ‘long form’ of Opa1 (l-Opa1) worked together with a proteolytically processed ‘short form’ (s-Opa1) to regulate full fusion. We showed that while l-Opa1 can induce low levels of membrane fusion, both s-Opa1 and l-Opa1 are required for fast and efficient fusion. These experiments were performed under conditions containing symmetric lipid composition in the bilayer, with both leaflets containing 20% CL. In these previous experiments, we found that when the model membranes lacked CL entirely, fusion was rare.

Here, we present new data investigating Opa1 activity under asymmetric bilayer conditions. We take advantage of our synthetic model-membrane system to generate supported lipid bilayers with asymmetric CL distribution. Our new results show CL in the outer leaflet (facing the IMS) plays an important role in Opa1-mediated bilayer tethering and pore opening. These data implicate CL-protein interactions during specific steps of membrane fusion. We discuss how changes in bilayer composition may influence membrane properties and protein-protein interactions.

MATERIALS AND METHODS

Preparation of Polymer-Tethered Asymmetric Bilayers and Reconstitution of l-Opa1 Into Asymmetric Bilayers

Polymer-tethered lipid bilayers were fabricated using mixtures of synthesized lipid reagents, including 1,2-dioleoyl-sn-glycero-2-phosphocholine (DOPC); 1,2-dioleoyl-sn-glycero-2-phosphoethanolamine-N-[methoxy (polyethylene glycerol)-2000] (DOPE-PEG2000), L- α -phosphatidylinositol (liver PI)

and 1',3'-bis [1,2-dioleoyl-sn-glycero-3-phospho]-glycerol (18:1 cardiolipin) from Avanti Polar Lipids (AL, United States). Inter-leaflet asymmetric bilayers were assembled in a layer-by-layer manner, using a combination of Langmuir-Blodgett dipping and Langmuir-Schaefer transfer. Bilayers were fabricated on a glass substrate (25 mm diameter cover glass, Fisher Scientific) as previously described (Ge et al., 2020b). For all fusion experiments, the bottom leaflet included 7% (mol) liver PI, 20% (mol) CL, 20% (mol) DOPE, 0.2% (mol) Cy5-PE and 5% (mol) DOPE-PEG2000. For fluorescence anisotropy analysis, the bottom leaflet contained 0.2% (mol) Top-fluor CL (Avanti Polar Lipids), which served as a reporter for CL distribution and to evaluate lipid flip-flop. The lipid mixture was spread carefully on top of the air-water interface on a Langmuir-Blodgett trough (KSV-NIMA, NY, United States). A film was applied at a pressure of 37 mN/m and kept for 30 min, then transferred to the glass substrate, forming the bottom leaflet of the bilayer. The top leaflet was fabricated using a Langmuir-Schaefer transfer method following fabrication of the LB layer, as previously described (Ge et al., 2020b). For l-Opa1 reconstitution experiments, the top leaflet (mimicking the outer, IMS-facing side) of the asymmetric bilayer was completely CL free, whereas the bottom leaflet (facing the inner, matrix side) contains 20% CL. TopFluor CL was used to evaluate bilayer anisotropy. Texas Red was used to monitor hemifusion kinetics. For all bilayers, the lipid mixture of the bottom leaflet was spread on top of the air-water interface, while pressure was kept at 37 mN/m for 30 min before dipping. The bottom leaflet was prepared at 30 mN/m pressured for 30 min before Schaefer transfer and assembly into an asymmetric bilayer. This helped maintain a similar area/molecule ratio in both top and bottom leaflets (Ge et al., 2020b).

Purification and Reconstitution of Opa1 in Asymmetric Bilayers

l- and s-Opa1 were purified following a protocol reported previously (Ge et al., 2020a; Ge et al., 2020b). Human l-Opa1 (isoform 1) and s-Opa1 with Twin-strep-tag at N-terminus and deca-histidine tag at C-terminus (GenScript, NJ, United States) was expressed in *Pichia pastoris* strain SMD1163. The cells were harvested and milled using SPEX 6875D Freezer/Mill. The milled powder was resuspended in buffer containing 50 mM sodium phosphate, 300 mM NaCl, 1 mM 2-mercaptoethanol, pH 7.5, with benzonase nuclease and protease inhibitors. The membrane fraction was collected by ultra-centrifugation at 235,000 xg for 45 min at 4°C, then incubated for 1 h in the same resuspension buffer supplemented with 2% DDM (Anatrace, OH, United States) and 0.1 mg/ml cardiolipin. Membrane fraction solubilized in DDM containing buffer was ultracentrifuged at 100,000 xg for 1 h at 4°C. The supernatant extract containing l-Opa1 was loaded onto a Ni-NTA column (Biorad, CA, USA), washed with 50 mM sodium phosphate, 350 mM NaCl, 1 mM 2-mercaptoethanol, 1 mM DDM, 0.025 mg/ml 18:1 CL, pH 7.5, with 25 mM imidazole followed with 100 mM imidazole. The bound Opa1 was eluted with the above buffer containing 500 mM imidazole. The elution was buffer exchanged into 100 mM Tris-HCl, 150 mM NaCl, 1 mM EDTA,

1 mM 2-mercaptoethanol, 0.15 mM DDM, 0.025 mg/ml 18:1 CL, pH 8.0. The C-terminal His-tag was then cleaved by treatment with TEV protease and applied to Ni-NTA and Strep-Tactin XT Superflow column (IBA life Sciences, Göttingen, Germany) attached in tandem and washed. The Strep-Tactin XT column was detached and the protein eluted with buffer containing 50 mM biotin. The elution fraction was concentrated and subjected to size exclusion chromatography in buffer with 25 mM Bis-Tris Propane, 100 mM NaCl, 1 mM TCEP, 0.025 mg/ml 18:1 CL, pH 7.5.

l-Opa1 was reconstituted into bilayers by mixing purified l-Opa1 with a surfactant cocktail containing DDM and OG, applied as reconstitution reagents, at a concentration of 1.2 and 1.1 nM, respectively. l-Opa1 was reconstituted to total amount of 1.3×10^{-12} mol (protein:lipid 1:100000). The protein was incubated for 2 h in surfactant at room temperature. Extra surfactant was then removed using Bio-Beads SM2 (Bio-Rad, CA, United States), at a concentration of 10 µg/ml.

Preparation of Liposomes and Proteoliposomes


Calcein encapsulated liposomes containing 7% (mol), PI 20% (mol), CL, 20 PE % (mol), 0.2% (mol), TexasRed-PE, and DOPC 52.8% (mol), were prepared as previously described (Ge et al., 2020a; Ge et al., 2020b). Lipids dissolved chloroform were mixed in the appropriate ratio and dried under nitrogen flow for 20–30 min. The resulting lipid film was hydrated with calcein containing Bis-Tris buffer (50 mM calcein, 25 mM Bis-Tris, 150 mM NaCl, pH 7.4) at 70°C for 30 min following vigorous agitation. The resulting vesicles were homogenized using a mini-extruder using 200 nm polycarbonate membrane for 15–20 times extrusion to prepare large unilamellar vesicles (LUVs).

l-Opa1 was reconstituted in the LUVs at a molar ratio of 1:5000. Purified l-Opa1 was added to the LUV suspensions with DDM (final concentration 0.1 µM) and incubated for 2 h at 4°C. After reconstitution, the sample was dialyzed against calcein buffer using a 3.5 kDa dialysis cassette at 4°C overnight. Calcein that was not encapsulated in the liposomes was removed using a PD10 desalting column.

TIRF Microscopy and Fluorescence Anisotropy Analysis

Fluorescence images were taken using a Vector TIRF system (Intelligent Imaging Innovation, Inc, Denver, CO, United States) equipped with a W-view Gemini system (Hamamatsu photonics, Bridgewater, NJ). TIRF images were acquired using a 100X oil-immersion TIRF objective (Zeiss, N.A 1.4). Fluorescent images were taken using a Prime 95B scientific CMOS camera kept at -10°C (Photometrics), recorded at a frame rate of 100 ms.

To validate bilayer asymmetry, fluorescence anisotropy was measured using a modified light path embedded in the microscopy. Under this experimental setting, the 488 nm excitation laser was polarized through a fiber switch system, equipped with internalized polarizer (Intelligent Imaging Innovation, Inc, Denver, CO, United States). The p- and s-emission were separated using a polarized beam splitter



(532 nm, Thorlabs) inserted in the Gemini system. As a result, p-emission and s-emission was separated onto two halves of the camera chip. To ensure a reliable emission result, the p-polarized emission and s-polarized emission was adjusted to be similar intensity and was determined using a photometer. The fluorescence anisotropy of dye-labeled bilayers in both symmetric and asymmetric bilayers was determined by the following formula:

$$r = \frac{I_{\perp} - GI_{\parallel}}{I_{\perp} + 2GI_{\parallel}}$$

Where G represents the correction factor (added to compensate for uneven light distribution caused by the polarized emission beam splitter optics, and potential depolarization effects caused by the high NA objective). The G factor was determined using a sample solution containing 1 mM Calcein dissolved in glycerol, where the fluorescence anisotropy was well defined. The G factor was evaluated for each pixel of the image before every measurement, and the final r of each pixel was obtained and

averaged for the lipid bilayer samples to eliminate potential effects of inner leaflet dye aggregation.

For fusion and membrane tethering experiments, a 543 nm laser was used to analyze TexasRed-PE signal in liposomes and proteoliposomes and to observe lipid demixing. The bilayer was labeled with Cy5 (excited by a 633 nm laser), that reported on homogeneity of the bilayer. A 488 nm laser was applied for recording content release of the encapsulated calcein. Time-lapse images were taken at a frequency of 100 ms to record fusion dynamics.

RESULTS

Bilayers With Asymmetric Cardiolipin Distribution can be Generated Through Layer-by-Layer Assembly

We generated a PEG polymer-tethered lipid bilayer with asymmetric CL distribution that mimics the leaflet distribution

found in the mitochondrial inner membrane. Our approach was inspired by methods successfully used to study the leaflet asymmetry of cholesterol in the plasma membrane (Hussain et al., 2013). These methods have been used to study phase separation in planar lipid bilayers and liposomes (Hussain et al., 2013; Doktorova et al., 2018; London, 2019). Our control symmetric bilayers comprised 15% (mol) CL on both top and bottom leaflets (with 0.1% CL in both leaflets replaced by Top-Fluor CL) (**Figure 1A**). Control asymmetric bilayers comprised 10% (mol) CL on the top leaflet and 20% (mol) CL in the bottom leaflet (Top-Fluor CL only labeling the bottom leaflet) (**Figure 1D**). We noticed symmetric bilayers fabricated in our previous work had to be used immediately, as they acquired defects within 24 h (Ge et al., 2020b). We did not observe this issue with our asymmetric bilayers, which were stable for at least 48 h.

We analyzed TIRF-fluorescence anisotropy images to evaluate lipid asymmetry, as performed previously in cellular membrane and model liposome membranes (Li et al., 2019; St Clair et al., 2020). Both symmetric and asymmetric bilayer were labeled with the same total concentration of fluorophore TopFluor-cardiolipin. EPI-fluorescence images show the same fluorescence intensity (**Figures 1B,E**). Fluorescence anisotropy of the asymmetric bilayer (average $r_{asy} = 0.125$) was approximately four times higher than that of the symmetric bilayer (average $r_{sym} = 0.024$) (**Figures 1C,F,G**). Models of plasma membrane asymmetry show high levels of cholesterol and phospholipid scrambling, which causes membranes to homogenize over time (Doktorova et al., 2020). To evaluate the potential for CL flip-flop and loss of bilayer asymmetry, anisotropy measurements were also performed after 24 and 48 h. There was no statistically significant anisotropy change after 48 h (**Figure 1G**). Finally, to validate bilayer asymmetry, we also measured anisotropy after reconstitution of l-Opa1, and found similar levels of anisotropy to those in the absence of protein (Supplemental Figure 1). These findings are consistent with simulation studies showing that an asymmetric distribution of CL is thermodynamically favorable (Kagan et al., 2015; Elias-Wolff et al., 2019).

l-Opa1 Mediated Homotypic Membrane Tethering is Dependent on CL Distribution and Influenced by Membrane Asymmetry

In our previous studies, we showed CL is important for both l-Opa1 (homotypic and heterotypic) and s-Opa1-mediated membrane tethering (Ge et al., 2020b). l-Opa1 tethering can potentially occur through homotypic protein-protein interactions, or heterotypic protein-lipid interactions. Here, we first investigated homotypic l-Opa1 membrane tethering in a bilayer with asymmetric CL distribution. We compared four conditions varying CL presence in the bilayer and proteoliposome. Initial validation of our system show that we can generate asymmetric bilayers that are stable even with 10% inter-leaflet difference in CL concentration. To evaluate Opa1 function under extreme membrane asymmetry, we completely depleted the CL on the top leaflet in our reconstitution

experiments, while leaving 20% CL in the bottom leaflet. Under homotypic l-Opa1 conditions (reconstituted in both bilayer and liposome), the presence of CL does not significantly stimulate GTP-independent membrane tethering, independent of the location of CL or leaflet distribution (**Figure 2 A-ii apo and A-iii apo compared to A-iv**).

Our previous data, and observations from others (Ban et al., 2010), show that membrane embedded CL can mediate nucleotide-dependent membrane tethering. Interestingly, we observe reduced particle tethering when CL was absent from the top leaflet of the planar bilayer, compared to the symmetrically distributed case (**Figure 2A iii and iv**). In our previous studies, we found that GTP-independent tethering did not depend on the presence of cardiolipin (Ge et al., 2020b). Our quantification here shows that when CL is present in both leaflets of the bilayer (regardless of the CL content in proteoliposomes), addition of GTP induces ~ three-fold increase in proteoliposome tethering to the bilayer (**Figures 2A,B ii and iv**). In contrast, when CL is depleted from only the outer leaflet of the bilayer, GTP triggers ~1.5 fold increase in proteoliposome tethering to the bilayer (**Figure 2 A iii, 2B**).



s-Opa1 Promotes Membrane Tethering in Asymmetric CL Bilayers


Proteolytic processing of the transmembrane anchored l-Opa1 to s-Opa1 is a key cellular regulator of mitochondrial inner membrane fusion and cristae state (Del Dotto et al., 2017; Wang et al., 2021). Our previous *in vitro* reconstitution studies using symmetric membranes, showed that stoichiometric levels of s-Opa1 were optimal for stimulating fast and efficient membrane fusion by l-Opa1 (Ge et al., 2020b). This result reconciled different cellular observations of mitochondrial reticulum phenotypes when modulating the fusion-fission equilibrium (Song et al., 2007; Ehses et al., 2009; Anand et al., 2014; Wang et al., 2021). Here, we investigate the effect of s-Opa1 addition in an asymmetric CL bilayer system. Unexpectedly, depleting CL from the top leaflet of the supported planar bilayer changed the effect of s-Opa1 on proteoliposome tethering.

Upon addition of s-Opa1 at a sub-stoichiometric ratio of 1:10 s-Opa1:l-Opa1, the number of tethered proteoliposomes increased >5 fold, in the presence of GTP (**Figure 3B**). Addition of stoichiometric levels of s-Opa1 to l-Opa1, did not increase proteoliposome tethering to the bilayer. Instead, lipid tubes were observed to form. These tubes were tethered to the bilayer. When s-Opa1 levels were in excess of l-Opa1, fewer proteoliposomes tethered to the bilayer, and the lipid tubes dissociated from the bilayer surface (**Figures 3D,E**).

Cardiolipin in the Outer Leaflet Plays Roles in Efficient l-Opa1-Mediated Homotypic Fusion and s-Opa1 Stimulated Fusion

Model membrane and cellular studies show l-Opa1 is required for membrane fusion, and s-Opa1 plays an important stimulatory role (Song et al., 2007; Ge et al., 2020b). Previously, we developed a three color imaging assay to evaluate the kinetics of membrane





fusion (Ge et al., 2020b). The fluorophores used for monitoring fusion differ from those we used for characterizing anisotropy. Briefly, membrane docking was defined by the FRET signal between Cy-5 (bilayer label) and TexasRed (proteoliposome label). Lipid demixing or hemifusion events were reported by a proteoliposome marker (Texas Red) diffusing into the lipid bilayer. Release of calcein from the proteoliposomes (loaded at quenched concentrations) reported on pore opening events and their kinetics. Using this assay, we observed low levels of hemifusion and full fusion (pore opening) in a homotypic I-Opa1 format (I-Opa1 reconstituted into the supported bilayer and proteoliposomes, both with symmetric CL distribution). Under these conditions, addition of s-Opa1 dramatically stimulated full fusion (Ge et al., 2020b).

Here, in I-Opa1 reconstituted bilayers with asymmetric CL distribution, we observed fewer hemifusion and full fusion events (**Figures 4D,G**, respectively), compared to our previous finding in symmetric bilayers (Ge et al., 2020b), where 20% of the total particles hemifuse with the bilayer. Under asymmetric conditions, addition of s-Opa1 did not increase the efficiency of membrane hemifusion or fusion (**Figures 4D,G**).

We also observed a significant delay in lipid demixing with I-Opa1 alone. Compared to symmetric bilayers, the average time for lipid demixing in asymmetric bilayers changed from 15 s to 20 s (in the ~50 events calculated in three different experiments) (**Figures 4B,C**), whereas the content release time changed from 20 s to 30 s or more (**Figures 4E,F**). The calcein release time was also significantly longer (from milliseconds to several seconds) with asymmetric CL bilayers, possibly suggesting an effect on calcein diffusion due to slow fusion pore expansion. Together

these observations all point to the importance of CL in the outer, IMS-facing leaflet in efficient s-Opa1 stimulated fusion.

DISCUSSION

The mitochondrial inner membrane (IMM) is expected to have an asymmetric distribution of CL under physiological conditions (Kagan et al., 2014; Pizzuto and Pelegrin, 2020). In this study, we used an *in vitro* model IMM to explore Opa1 activity upon selectively depleting CL from the outer leaflet. We found three interesting effects under these asymmetric CL conditions. First, removing CL from the outer leaflet reduces GTP-dependent homotypic I-Opa1 tethering. Second, we observe lipid tube formation when sub-stoichiometric levels of s-Opa1 are added to I-Opa1 tethered proteoliposomes containing CL. Third, asymmetric CL distribution slows pore opening kinetics. These observations all point to the importance of outer leaflet CL in mediating efficient fusion.

Recent structures of s-Opa1 and its yeast orthologue s-Mgm provide candidate interfaces for understanding self-assembly. These structures all show Opa1 has a canonical dynamin-family GTPase domain arrangement, with GTPase domain, bundle signaling element (BSE), stalk domain and paddle region (Faelber et al., 2019; Zhang et al., 2020). Basic and hydrophobic residues in the paddle region directly interact with membranes containing CL (Faelber et al., 2019; Yan et al., 2020; Zhang et al., 2020). Mutation of these residues results in decreased affinity of s-Opa1 to the lipid bilayer and affects GTPase activity (Zhang et al., 2020). In Cryo-ET studies,

interfaces formed by the helical stalk region mediate self-assembly (Faelber et al., 2019).

Our observation of reduced tethering under asymmetric conditions suggests that the presence of CL in the outer leaflet facilitates inter-membrane interactions. We still lack high-resolution structural information for l-Opa1. Existing observations suggest potential modes of CL-influenced tethering may be at play. CL in the bilayer might stabilize a conformation of Opa1 such that it can interact in a homotypic fashion with another copy of Opa1 in the proteoliposome bilayer. It should be noted that although the outer leaflet of the bilayer was fabricated without CL for our assay, recombinant l-Opa1 was purified in the presence of CL. We estimate that during reconstitution, around 10^{-12} mol of CL may be reconstituted into the outer leaflet, resulting in an outer leaflet with 0.01% (mol) CL, which is higher than the total amount of l-Opa1 reconstituted in the planar bilayer. Based on previous GTP-assays, however, this level of CL is unlikely to significantly affect Opa1 activity (Ge et al., 2020b).

Membrane tubes are an intriguing *in vitro* observation of Opa1 behavior, with several candidate *in vivo* roles proposed. In our previous work with symmetric bilayers, we did not observe lipid tube formation with the protein concentrations used under three different conditions, namely, l-Opa1 or s-Opa1 alone, and l-Opa1 mixed with s-Opa1 (Ge et al., 2020b). Interestingly, we do observe tube formation with asymmetric bilayers lacking CL in the outer leaflet, upon addition of s-Opa1. With a bilayer containing symmetric inter-leaflet CL distribution, no lipid tubules were observed previously. In this work (and our previous studies), we used low concentrations of reconstituted protein combined in the bilayers and liposomes, making the total l-Opa1 concentration of 6.5×10^{-12} mol/ml (protein:lipid, 1:5000), and a final s-Opa1 concentration of 6.5×10^{-11} mol/ml (final protein:lipid 1:500) (Ge et al., 2020b). The Daumke and Sun groups determined helical assemblies of s-Mgm1 and s-Opa1, respectively, decorating the outside of lipid tubes and inducing positive curvature (Faelber et al., 2019; Zhang et al., 2020). Faelber and others also observe s-Mgm1 tubes decorating the interior of tubes, with negative curvature (analogous to the topology expected within cristae). Li and co-workers suggest a s-Mgm1 trimer may help 'bud' regions of bilayer, through puckering membranes with positive curvature (Yan et al., 2020). In these reports, s-Opa1 was added at 1:1 (protein: lipid molar) ratio, for a total concentration of s-Opa1 5×10^{-9} mol/ml (Ban et al., 2010; Zhang et al., 2020). In these studies, interaction of s-Opa1 with CL is important for tubulation. In our asymmetric CL experiments, we observe tubulation at even lower concentrations. Due to the lack of CL on the outer (IMS facing) leaflet, we suspect that the major source of membrane for generating the tubes is the liposomes. Whether aspects of tubulation represent an intermediate in fusion will require further investigation. Future studies with asymmetric liposomes will also help discriminate s-Opa1 binding modes. In the asymmetric CL setting, s-Opa1 may interact with l-Opa1, and this heterotypic interaction might explain the observed increase in tethering.

In our previous studies under symmetric CL conditions, we observed fast and efficient membrane fusion upon addition of

stoichiometric levels of s-Opa1 (Ge et al., 2020b). In this study, asymmetric CL distribution (specifically the lack of CL in the outer leaflet) appears to stall progression to fusion, resulting in the growth of tubes. Furthermore, addition of s-Opa1 dissociated tubes from the bilayer, suggesting a hetero-oligomeric l-Opa1:s-Opa1 interaction plays a role in association of these tubes with the supported bilayer. We did not observe any hemifusion events between tubes and the supported bilayer, emphasizing the importance of CL in this leaflet for proceeding through fusion. The lack of stimulatory effect upon addition of s-Opa1 to a homotypic l-Opa1 arrangement also suggests s-Opa1 assembly (or interaction) with the supported bilayer plays a role in fusion.

The *in vitro* system we describe here opens exploration of how membrane lipid composition and distribution influence the interplay of different physical properties to regulate fusion and ultrastructure. For example, fusion efficiency may be influenced by the combination of membrane elastic properties, including monolayer spontaneous curvature and bending modulus. (Fan et al., 2016). Calculations based on physical models suggest spontaneous curvature is a factor regulating the elastic energy of the pre-fusion planar bilayer. In experiments performed here, we investigated supported lipid bilayers with cardiolipin depleted from the outer leaflet. If the unsaturated, four acyl-chain CL is depleted from the outer leaflet, intrinsic negative curvature may be disfavored. Negative spontaneous curvature would be expected to favor the formation of hemifusion stalk, and be important at regions such as the cristae tip. The converse situation, with cardiolipin absent from the inner-leaflet will not only eliminate Opa1-lipid interactions, but also potentially disfavor positive curvature. Lowering the barrier to positive curvature may be important at regions like the cristae junction. Depletion of CL in either leaflet could result in bilayers with a higher bending modulus than that of a symmetric bilayers. Changes in the bending modulus could increase the energy barrier from planar membrane to a highly curved states. Other physical properties likely influenced by cardiolipin distribution include line tension, hydrophobic mismatch and membrane packing. Investigating how these factors regulate membrane rearrangement are of great interest for future studies.

CONCLUSION

In this study we tested the effects of selectively removing CL from one leaflet of a model of the mitochondrial innermembrane. We observed that simply removing CL from the IMS facing leaflet resulted in interesting changes in membrane tethering, and stalled fusion. Furthermore, we observe formation of membrane tubes under these conditions. Low levels of CL are expected to be present in the IMS-facing leaflet under physiological conditions. These observations point to important conformations and assembly states of Opa1 regulated by the presence of CL. These studies also suggest that interesting membrane properties resulting from lipid asymmetry remain to be explored. The specific protein arrangements and membrane states of fusion are key questions for future integrated structural and biophysical studies.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

AUTHOR CONTRIBUTIONS

LC, YG, SB, TN, CL: Conceptualization, Resources, Investigation, Data curation, Validation YG, SB: Formal analysis, Methodology

REFERENCES

- Anand, R., Wai, T., Baker, M. J., Kladt, N., Schauss, A. C., Rugarli, E., et al. (2014). The I-AAA Protease YME1L and OMA1 Cleave OPA1 to Balance Mitochondrial Fusion and Fission. *J. Cel Biol* 204, 919–929. doi:10.1083/jcb.201308006
- Ban, T., Heymann, J. A. W., Song, Z., Hinshaw, J. E., and Chan, D. C. (2010). OPA1 Disease Alleles Causing Dominant Optic Atrophy Have Defects in Cardiolipin-Stimulated GTP Hydrolysis and Membrane Tubulation. *Hum. Mol. Genet.* 19, 2113–2122. doi:10.1093/hmg/ddq088
- Boyd, K. J., Alder, N. N., and May, E. R. (2017). Buckling under Pressure: Curvature-Based Lipid Segregation and Stability Modulation in Cardiolipin-Containing Bilayers. *Langmuir* 33, 6937–6946. doi:10.1021/acs.langmuir.7b01185
- Chan, D. C. (2020). Mitochondrial Dynamics and its Involvement in Disease. *Annu. Rev. Pathol. Mech. Dis.* 15, 235–259. doi:10.1146/annurev-pathmechdis-012419-032711
- Chu, C. T., Ji, J., Dagda, R. K., Jiang, J. F., Tyurina, Y. Y., Kapralov, A. A., et al. (2013). Cardiolipin Externalization to the Outer Mitochondrial Membrane Acts as an Elimination Signal for Mitophagy in Neuronal Cells. *Nat. Cel Biol* 15, 1197–1205. doi:10.1038/ncb2837
- Cogliati, S., Enriquez, J. A., and Scorrano, L. (2016). Mitochondrial Cristae: Where Beauty Meets Functionality. *Trends Biochem. Sci.* 41, 261–273. doi:10.1016/j.tibs.2016.01.001
- Del Dotto, V., Mishra, P., Vidoni, S., Fogazza, M., Maresca, A., Caporali, L., et al. (2017). OPA1 Isoforms in the Hierarchical Organization of Mitochondrial Functions. *Cel Rep.* 19, 2557–2571. doi:10.1016/j.celrep.2017.05.073
- Doktorova, M., Heberle, F. A., Eicher, B., Standaert, R. F., Katsaras, J., London, E., et al. (2018). Preparation of Asymmetric Phospholipid Vesicles for Use as Cell Membrane Models. *Nat. Protoc.* 13, 2086–2101. doi:10.1038/s41596-018-0033-6
- Doktorova, M., Symons, J. L., and Levental, I. (2020). Structural and Functional Consequences of Reversible Lipid Asymmetry in Living Membranes. *Nat. Chem. Biol.* 16, 1321–1330. doi:10.1038/s41589-020-00688-0
- Dudek, J. (2017). Role of Cardiolipin in Mitochondrial Signaling Pathways. *Front. Cel Dev. Biol.* 5, 90. doi:10.3389/fcell.2017.00090
- Ehse, S., Raschke, I., Mancuso, G., Bernacchia, A., Geimer, S., Tondera, D., et al. (2009). Regulation of OPA1 Processing and Mitochondrial Fusion by M-AAA Protease Isoenzymes and OMA1. *J. Cel Biol* 187, 1023–1036. doi:10.1083/jcb.200906084
- Elias-Wolff, F., Lindén, M., Lyubartsev, A. P., and Brandt, E. G. (2019). Curvature Sensing by Cardiolipin in Simulated Buckled Membranes. *Soft Matter* 15, 792–802. doi:10.1039/c8sm02133c
- Faelber, K., Dietrich, L., Noel, J. K., Wollweber, F., Pfitzner, A.-K., Mühleip, A., et al. (2019). Structure and Assembly of the Mitochondrial Membrane Remodelling GTPase Mgm1. *Nature* 571, 429–433. doi:10.1038/s41586-019-1372-3
- Fan, Z.-A., Tsang, K.-Y., Chen, S.-H., and Chen, Y.-F. (2016). Revisit the Correlation between the Elastic Mechanics and Fusion of Lipid Membranes. *Sci. Rep.* 6, 31470. doi:10.1038/srep31470
- Friedman, J. R., and Nunnari, J. (2014). Mitochondrial Form and Function. *Nature* 505, 335–343. doi:10.1038/nature12985

LC, YG: Supervision, Project administration LC: Funding acquisition.

ACKNOWLEDGMENTS

We thank all members of the Chao Lab for helpful discussions and suggestions. This work was supported by funding from the Charles H Hood Foundation Child Health Research Award and the National Institutes of Health (R35GM142553) to LC.

- Gallet, P. F., Petit, J. M., Maftah, A., Zachowski, A., and Julien, R. (1997). Asymmetrical Distribution of Cardiolipin in Yeast Inner Mitochondrial Membrane Triggered by Carbon Catabolite Repression. *Biochem. J.* 324 (Pt 2), 627–634. doi:10.1042/bj3240627
- Gao, S., and Hu, J. (2021). Mitochondrial Fusion: The Machineries in and Out. *Trends Cel Biol.* 31, 62–74. doi:10.1016/j.tcb.2020.09.008
- Ge, Y., Shi, X., Boopathy, S., Mcdonald, J., Smith, A. W., and Chao, L. H. (2020b). Two Forms of Opa1 Cooperate to Complete Fusion of the Mitochondrial Inner-Membrane. *Elife* 9. doi:10.7554/eLife.50973
- Ge, Y., Boopathy, S., Smith, A., and Chao, L. H. (2020a). A Model Membrane Platform for Reconstituting Mitochondrial Membrane Dynamics. *J. Vis. Exp.* doi:10.3791/61620
- Giacomello, M., Pyakurel, A., Glytsou, C., and Scorrano, L. (2020). The Cell Biology of Mitochondrial Membrane Dynamics. *Nat. Rev. Mol. Cel Biol* 21, 204–224. doi:10.1038/s41580-020-0210-7
- Harayama, T., and Riezman, H. (2018). Understanding the Diversity of Membrane Lipid Composition. *Nat. Rev. Mol. Cel Biol* 19, 281–296. doi:10.1038/nrm.2017.138
- Harb, J. S., Comte, J., and Gautheron, D. C. (1981). Asymmetrical Orientation of Phospholipids and Their Interactions with Marker Enzymes in Pig Heart Mitochondrial Inner Membrane. *Arch. Biochem. Biophys.* 208, 305–318. doi:10.1016/0003-9861(81)90153-3
- Horvath, S. E., and Daum, G. (2013). Lipids of Mitochondria. *Prog. Lipid Res.* 52, 590–614. doi:10.1016/j.plipres.2013.07.002
- Hossein, A., and Deserno, M. (2020). Spontaneous Curvature, Differential Stress, and Bending Modulus of Asymmetric Lipid Membranes. *Biophysical J.* 118, 624–642. doi:10.1016/j.bpj.2019.11.3398
- Hussain, N. F., Siegel, A. P., Ge, Y., Jordan, R., and Naumann, C. A. (2013). Bilayer Asymmetry Influences Integrin Sequestering in Raft-Mimicking Lipid Mixtures. *Biophysical J.* 104, 2212–2221. doi:10.1016/j.bpj.2013.04.020
- Kagan, V. E., Chu, C. T., Tyurina, Y. Y., Cheikhi, A., and Bayir, H. (2014). Cardiolipin Asymmetry, Oxidation and Signaling. *Chem. Phys. Lipids* 179, 64–69. doi:10.1016/j.chemphyslip.2013.11.010
- Kagan, V. E., Tyurina, Y. Y., Tyurin, V. A., Mohammadyani, D., Angeli, J. P. F., Baranov, S. V., et al. (2015). Cardiolipin Signaling Mechanisms: Collapse of Asymmetry and Oxidation. *Antioxid. Redox Signaling* 22, 1667–1680. doi:10.1089/ars.2014.6219
- Kondadi, A. K., Anand, R., and Reichert, A. S. (2020). Cristae Membrane Dynamics - A Paradigm Change. *Trends Cel Biol.* 30, 923–936. doi:10.1016/j.tcb.2020.08.008
- Kordower, J. H., Chu, Y., Hauser, R. A., Freeman, T. B., and Olanow, C. W. (2008). Lewy Body-like Pathology in Long-Term Embryonic Nigral Transplants in Parkinson's Disease. *Nat. Med.* 14, 504–506. doi:10.1038/nm1747
- Krebs, J. J., Hauser, H., and Carafoli, E. (1979). Asymmetric Distribution of Phospholipids in the Inner Membrane of Beef Heart Mitochondria. *J. Biol. Chem.* 254, 5308–5316. doi:10.1016/s0021-9258(18)50596-2
- Kroemer, G., Galluzzi, L., and Brenner, C. (2007). Mitochondrial Membrane Permeabilization in Cell Death. *Physiol. Rev.* 87, 99–163. doi:10.1152/physrev.00013.2006
- Lemmin, T., Bovigny, C., Lançon, D., and Dal Peraro, M. (2013). Cardiolipin Models for Molecular Simulations of Bacterial and Mitochondrial Membranes. *J. Chem. Theor. Comput.* 9, 670–678. doi:10.1021/ct300590v
- Li, G., Kakuda, S., Suresh, P., Canals, D., Salamone, S., and London, E. (2019). Replacing Plasma Membrane Outer Leaflet Lipids with Exogenous Lipid

- without Damaging Membrane Integrity. *PLoS One* 14, e0223572. doi:10.1371/journal.pone.0223572
- London, E. (2019). Membrane Structure-Function Insights from Asymmetric Lipid Vesicles. *Acc. Chem. Res.* 52, 2382–2391. doi:10.1021/acs.accounts.9b00300
- Lorent, J. H., Levental, K. R., Ganesan, L., Rivera-Longworth, G., Sezgin, E., Doktorova, M., et al. (2020). Plasma Membranes Are Asymmetric in Lipid Unsaturation, Packing and Protein Shape. *Nat. Chem. Biol.* 16, 644–652. doi:10.1038/s41589-020-0529-6
- Mcbride, H. M., Neuspiel, M., and Wasiak, S. (2006). Mitochondria: More Than Just a Powerhouse. *Curr. Biol.* 16, R551–R560. doi:10.1016/j.cub.2006.06.054
- Oemer, G., Koch, J., Wohlfarter, Y., Alam, M. T., Lackner, K., Sailer, S., et al. (2020). Phospholipid Acyl Chain Diversity Controls the Tissue-specific Assembly of Mitochondrial Cardiolipins. *Cel Rep.* 30, 4281–4291. doi:10.1016/j.celrep.2020.02.115
- Oemer, G., Lackner, K., Muigg, K., Krumshnabel, G., Watschinger, K., Sailer, S., et al. (2018). Molecular Structural Diversity of Mitochondrial Cardiolipins. *Proc. Natl. Acad. Sci. USA* 115, 4158–4163. doi:10.1073/pnas.1719407115
- Petit, J.-M., Huet, O., Gallet, P. F., Maftah, A., Ratinaud, M.-H., and Julien, R. (1994). Direct Analysis and Significance of Cardiolipin Transverse Distribution in Mitochondrial Inner Membranes. *Eur. J. Biochem.* 220, 871–879. doi:10.1111/j.1432-1033.1994.tb18690.x
- Pizzuto, M., and Pelegrin, P. (2020). Cardiolipin in Immune Signaling and Cell Death. *Trends Cel Biol.* 30, 892–903. doi:10.1016/j.tcb.2020.09.004
- Ryan, T., Bamm, V. V., Stykel, M. G., Coackley, C. L., Humphries, K. M., Jamieson-Williams, R., et al. (2018). Cardiolipin Exposure on the Outer Mitochondrial Membrane Modulates α -synuclein. *Nat. Commun.* 9, 817. doi:10.1038/s41467-018-03241-9
- Schlame, M. (2008). Thematic Review Series: Glycerolipids. Cardiolipin Synthesis for the Assembly of Bacterial and Mitochondrial Membranes. *J. Lipid Res.* 49, 1607–1620. doi:10.1194/jlr.r700018-jlr200
- Song, Z., Chen, H., Fiket, M., Alexander, C., and Chan, D. C. (2007). OPA1 Processing Controls Mitochondrial Fusion and Is Regulated by mRNA Splicing, Membrane Potential, and Yme1L. *J. Cel Biol* 178, 749–755. doi:10.1083/jcb.200704110
- St. Clair, J. W., Kakuda, S., and London, E. (2020). Induction of Ordered Lipid Raft Domain Formation by Loss of Lipid Asymmetry. *Biophysical J.* 119, 483–492. doi:10.1016/j.bpj.2020.06.030
- Traïkia, M., Warschawski, D. E., Lambert, O., Rigaud, J. L., and Devaux, P. F. (2002). Asymmetrical Membranes and Surface Tension. *Biophys. J.* 83, 1443–1454. doi:10.1016/S0006-3495(02)73915-5
- Van Laar, V. S., and Berman, S. B. (2009). Mitochondrial Dynamics in Parkinson's Disease. *Exp. Neurol.* 218, 247–256. doi:10.1016/j.expneurol.2009.03.019
- Wang, R., Mishra, P., Garbis, S. D., Moradian, A., Sweredoski, M. J., and Chan, D. C. (2021). Identification of New OPA1 Cleavage Site Reveals that Short Isoforms Regulate Mitochondrial Fusion. *MBoC* 32, 157–168. doi:10.1091/mbc.e20-09-0605
- Wilson, B. A., Ramanathan, A., and Lopez, C. F. (2019). Cardiolipin-Dependent Properties of Model Mitochondrial Membranes from Molecular Simulations. *Biophysical J.* 117, 429–444. doi:10.1016/j.bpj.2019.06.023
- Yan, L., Qi, Y., Ricketson, D., Li, L., Subramanian, K., Zhao, J., et al. (2020). Structural Analysis of a Trimeric Assembly of the Mitochondrial Dynamin-like GTPase Mgm1. *Proc. Natl. Acad. Sci. USA* 117, 4061–4070. doi:10.1073/pnas.1919116117
- Zhang, D., Zhang, Y., Ma, J., Zhu, C., Niu, T., Chen, W., et al. (2020). Cryo-EM Structures of S-OPA1 Reveal its Interactions with Membrane and Changes upon Nucleotide Binding. *Elife* 9, e50294. doi:10.7554/eLife.50294

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Ge, Boopathy, Nguyen, Lugo and Chao. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.