



# Purines released from astrocytes inhibit excitatory synaptic transmission in the ventral horn of the spinal cord

Eva Meier Carlsen and Jean-François Perrier\*

Laboratory of Neuronal Signaling, Department of Neuroscience and Pharmacology, University of Copenhagen, Copenhagen, Denmark

## Edited by:

Brian R. Noga, University of Miami, USA

## Reviewed by:

David Parker, Cambridge University, UK

Klaus Ballanyi, University of Alberta, Canada

## \*Correspondence:

Jean-François Perrier, Laboratory of Neuronal Signaling, Department of Neuroscience and Pharmacology, University of Copenhagen, Blegdamsvej 3, 2200 Copenhagen N, Denmark  
e-mail: perrier@sund.ku.dk

Spinal neuronal networks are essential for motor function. They are involved in the integration of sensory inputs and the generation of rhythmic motor outputs. They continuously adapt their activity to the internal state of the organism and to the environment. This plasticity can be provided by different neuromodulators. These substances are usually thought of being released by dedicated neurons. However, in other networks from the central nervous system synaptic transmission is also modulated by transmitters released from astrocytes. The star-shaped glial cell responds to neurotransmitters by releasing gliotransmitters, which in turn modulate synaptic transmission. Here we investigated if astrocytes present in the ventral horn of the spinal cord modulate synaptic transmission. We evoked synaptic inputs in ventral horn neurons recorded in a slice preparation from the spinal cord of neonatal mice. Neurons responded to electrical stimulation by monosynaptic EPSCs (excitatory monosynaptic postsynaptic currents). We used mice expressing the enhanced green fluorescent protein under the promoter of the glial fibrillary acidic protein to identify astrocytes. Chelating calcium with BAPTA in a single neighboring astrocyte increased the amplitude of synaptic currents. In contrast, when we selectively stimulated astrocytes by activating PAR-1 receptors with the peptide TFLLR, the amplitude of EPSCs evoked by a paired stimulation protocol was reduced. The paired-pulse ratio was increased, suggesting an inhibition occurring at the presynaptic side of synapses. In the presence of blockers for extracellular ectonucleotidases, TFLLR did not induce presynaptic inhibition. Puffing adenosine reproduced the effect of TFLLR and blocking adenosine A<sub>1</sub> receptors with 8-Cyclopentyl-1,3-dipropylxanthine prevented it. Altogether our results show that ventral horn astrocytes are responsible for a tonic and a phasic inhibition of excitatory synaptic transmission by releasing ATP, which gets converted into adenosine that binds to inhibitory presynaptic A<sub>1</sub> receptors.

**Keywords:** astrocyte, ATP, adenosine, purine, spinal cord, synapse, motor control

## INTRODUCTION

Neuronal networks located in the ventral horns of the spinal cord are essential for motor control (Kjaerulff and Kiehn, 1996; Prut and Perlmutter, 2003; Lanuza et al., 2004; Dai et al., 2005; Brocard et al., 2010; Mui et al., 2012). Their architecture provides the necessary stability for generating stereotyped movements. In order to adapt to internal and external changes, the organism adjusts neuronal activity by different cellular mechanisms providing high degrees of flexibility on time scales ranging from milliseconds to hours. The activity of spinal networks is regulated by descending pathways releasing neuromodulators such as serotonin, noradrenaline, glutamate or GABA (Arvidsson et al., 1990; Jacobs and Azmitia, 1992; Ono and Fukuda, 1995; Schmidt and Jordan, 2000; Conn et al., 2005; Farrant and Nusser, 2005; Perrier and Delgado-Lezama, 2005; Cotel et al., 2013; Perrier et al., 2013). Flexibility is also ensured by intrinsic modulatory systems releasing transmitters such as endocannabinoids (Ameri, 1999; El Manira and Kyriakatos, 2010) or purines (Dale and Gilday, 1996). The modulation provided by purines is of particular interest because adenosine triphosphate (ATP) and its metabolic product adenosine exert opposite effects on neurons. In the spinal cord of

embryonic *Xenopus*, ATP promotes swimming by inhibiting voltage gated potassium conductances (Dale and Gilday, 1996; Brown and Dale, 2002). Similarly, in the brainstem, ATP activates neurons from the retrotrapezoid nucleus, involved in breathing (Gourine et al., 2010). Once in the extracellular space, ATP is converted by ectonucleotidases into adenosine diphosphate (ADP), adenosine monophosphate (AMP), and adenosine (Dunwiddie et al., 1997). The binding of adenosine to metabotropic A<sub>1</sub> receptors lowers the excitability of spinal neurons belonging to motor circuits by inhibiting Ca<sup>2+</sup> currents (Dale and Gilday, 1996; Brown and Dale, 2000). In agreement, (Taccola et al., 2012) and (Brockhaus and Ballanyi, 2000) showed that A<sub>1</sub> receptor activation depresses bicuculline-evoked seizure-like bursting in newborn rat spinal cords. In other systems such as the hippocampus or the cerebellum, the activation of presynaptic A<sub>1</sub> receptors decreases neurotransmitter release by inhibiting Ca<sup>2+</sup> channels (Hollins and Stone, 1980; Wu and Saggau, 1994; Zhang et al., 2003).

Sources of ATP in the nervous system are multiple. ATP is co-released from neurons with neurotransmitters such as acetylcholine, noradrenaline, or GABA (Silinsky and Redman,

1996; Jo and Schlichter, 1999; Burnstock, 2007). In addition, ATP is secreted from astrocytes (Guthrie et al., 1999; Fields and Burnstock, 2006; Gourine et al., 2010; Christensen et al., 2013; Lalo et al., 2014). In the spinal cord, some of the modulatory actions induced by ATP are blocked by the glial metabolic poison fluoroacetate (Witts et al., 2012). It was therefore suggested that spinal glial cells also release ATP. However, a direct demonstration that ATP released from spinal astrocytes modulates synaptic transmission is still lacking.

In this study, we investigated if purines released from spinal astrocytes have any effect on synaptic transmission between neurons from the ventral horn. We found that astrocytes release ATP, which after being converted to adenosine produces both tonic and phasic inhibition of excitatory synaptic transmission by decreasing the probability of neurotransmitter release.

## MATERIALS AND METHODS


Experiments were performed on neonatal (P4–P22) wild type (C7BL/6; Taconic) and transgenic mice expressing enhanced green fluorescent protein (E-GFP) under the promoter of glial fibrillary acidic protein [GFAP; line Tgn(hgFAPEGEP)] GFEC 335 (Figures 1A,B). Transgenic mice were kindly provided by Professor Frank Kirchhoff. The astrocytes of these mice were identified by their fluorescence (Nolte et al., 2001). The surgical procedures complied with Danish legislation. Mice were killed by decapitation.

### SLICE PREPARATION

The lumbar enlargement of the spinal cord was removed and placed in cold artificial cerebrospinal fluid containing NaCl 125 mM, KCl 2.5 mM, NaHCO<sub>3</sub> 26 mM, CaCl<sub>2</sub> 2 mM, MgCl<sub>2</sub> 1 mM, NaH<sub>2</sub>PO<sub>4</sub> 1.25 mM, Glucose 25 mM. Ringer's solution was continuously carbogenated by gassing with 95% O<sub>2</sub> plus 5% CO<sub>2</sub>. 300 μm thick slices were obtained with a vibratome (MicroM slicer HM 650V; Microm International GmbH, Germany) equipped with cooling unit CU65 set at 2°C. Slices were then positioned in a recording chamber and continuously perfused with Ringer's solution at room temperature.

### PATCH CLAMP RECORDING

Visual guided patch clamp recording was done with a Multi-clamp 700B amplifier (Molecular Devices, USA). Neurons were visualized by means of a BW51WI microscope (Olympus, Japan) equipped with an oblique illumination condenser. Astrocytes were identified by means of fluorescence illumination obtained with a monochromator (Polychrome V; Till Photonics, Germany) tuned at 488 nm. The pipette solution (in mM): 122 K-gluconate, 2.5 MgCl<sub>2</sub>, 0.0003 CaCl<sub>2</sub>, 5.6 Mg-gluconate, 5 K-HEPES, 5 H-HEPES, 5 Na<sub>2</sub>ATP, 1 EGTA, 2.5 biocytine, 0.01 Alexa 488 hydrazide, sodium salt (Life Technologies, USA), and KOH to adjust the pH to 7.4. Calcium-clamp pipette solution contained (in mM): 40 K-gluconate, 30 K<sub>4</sub>-BAPTA, 50 sucrose, 2.5 MgCl<sub>2</sub>, 0.0003 CaCl<sub>2</sub>, 5.6 Mg-gluconate, 5 K-HEPES, 5 H-HEPES, 5 Na<sub>2</sub>ATP, 1 EGTA, 5 biocytine, 0.068 Alexa 568 hydrazide sodium salt (Life Technologies, USA) and the necessary amount of KOH to adjust the pH to 7.4. Electrodes had a resistance ranging from 4 to 8 MΩ. Recordings were sampled at 10 kHz with a 16-bit analog-to-digital



**FIGURE 1 | Confocal microscopy of GFAP-EGFP astrocytes in spinal cord slices.** (A) Image of a lumbar spinal cord slice from a neonatal mouse (P7) scanned with confocal microscopy. The red dashed line indicates the limits between gray and white matter. Note the presence of multiple protoplasmic fluorescent astrocytes in the gray matter. Scale bar: 100 μm. (B) High magnification picture (x63) of ventral horn E-GFP positive astrocytes. Twenty-four consecutive parallel confocal plans were superimposed. Scale bar: 10 μm. (C) Relative positions of a fraction of the neurons recorded for the study. Upward triangles indicate responses paired pulse stimulation protocol with a PPR > 1. Downward triangles correspond to a PPR < 1. Squares: PPR ≈ 1.

converter (DIGIDATA 1440; Molecular Devices, USA) and displayed by means of Clampex 10.2 software (Molecular Devices, USA).

### ELECTRICAL STIMULATION

Local electric stimulation was performed with a bipolar concentric electrode (TM33CCNON; World Precision Instruments, Sarasota, FL, USA) connected to an isolation unit (Isolator 11, Axon Instruments, USA) triggered by an external signal. The electrode was positioned in areas devoid of cell bodies, in the vicinity of neurons recorded from. Different positions were tried until a reliable response was induced by stimulation. Responses to paired pulse stimulations (duration from 10 to 40 μs; amplitude from 0.1 to 1 mA; interval from 10 to 30 ms) were recorded in neurons in voltage-clamp mode. They were considered as putative excitatory monosynaptic postsynaptic currents (EPSCs) when all the following criteria were satisfied: (1) fixed latency and variable amplitude; (2) decrease of amplitude upon depolarization, as one would expect for EPSCs in contrast with IPSCs; (3) absence of response when reverting the polarity of stimulation to exclude possible artifacts caused by direct electrical

stimulation. For a fraction of neurons, we also tested responses to stimulation at high frequencies (10, 20, and 50 Hz). The neurons that did not follow these frequencies were discarded from the sample.

### INPUT RESISTANCE

The input resistance of the neurons recorded from was calculated as the ratio of voltage to current measured during small depolarizing pulses generated in voltage-clamp mode.

### FOCAL APPLICATION OF DRUGS

Electrodes made from borosilicate capillaries (tip diameter ranging from 1.5–2  $\mu\text{m}$ ; BF150-86-7.5, Sutter Instrument, USA) were either filled with TFLLR-NH<sub>2</sub> (10  $\mu\text{M}$ ; Tocris Bioscience, UK), ATP (1 mM; Sigma–Aldrich, St. Louis, MO, USA) or adenosine (1 mM; Sigma–Aldrich, St. Louis, MO, USA). Drugs were puff applied at 14–35 Pa by a homemade time-controlled pressure device.

### DRUGS

The following drugs were used: BAPTA (30 mM; Invitrogen, Carlsbad, CA, USA), 6-*N,N*-Diethyl-D- $\beta,\gamma$ -dibromomethylene ATP trisodium salt (ARL 67156 trisodium salt, 50  $\mu\text{M}$ ; Tocris Bioscience, UK), 8-Cyclopentyl-1,3-dipropylxanthine (DPCPX, 5  $\mu\text{M}$ ; Tocris Bioscience, UK).

### CONFOCAL MICROSCOPY

Confocal microscopy images were obtained at the core facility for integrated microscopy (CFIM) of the Faculty of Health and Medical Sciences of the University of Copenhagen. Pictures were taken with a LSM 700 confocal microscope (Zeiss, Germany) equipped with Plan-Neofluar, X5 (N.A. 0.15) and Plan-Apochromat X63 (N.A. 1.4) objectives. E-GFP positive cells were excited with a 488 nm diode laser (10 mW).

### DATA ANALYSIS

Statistical analyses were performed offline by means of Origin software (version 8.6; MicroCal Inc., USA). The normality distribution of each sample was tested with Shapiro–Wilk test. Non-parametric tests were used for small samples ( $n < 10$ ) and when Gaussian distribution could not be approximated. Data are represented as mean  $\pm$  standard deviation of the mean.

## RESULTS


### ASTROCYTES PRODUCE TONIC PRESYNAPTIC INHIBITION IN THE VENTRAL HORN

We recorded ventral horn neurons (Figure 1C) from the spinal cord of neonatal mice with the whole-cell patch-clamp technique in voltage-clamp mode. Most of the neurons recorded from had a high input resistance ( $> 500 \text{ M}\Omega$ ) and were located in the medial part of the ventral horn. They were likely interneurons. In few instances putative motoneurons located in the lateral part of the ventral horn were also recorded. We did not investigate their identity further. Since all these neurons responded similarly to the different tests of our study, they were pooled in a single sample. We evoked pairs of monosynaptic EPSCs by stimulating a nearby region with a bipolar electrode (Figures 2A,B; see Materials and Methods). The second EPSC was bigger, similar or smaller

than the first. Consequently, the paired pulse ratio (PPR), calculated as the relative value of the second and the first EPSCs was either below or above 1 (Figure 1C). To investigate if and how astrocytes modulate synaptic transmission, we used a second pipette filled with patch solution enriched with a high concentration of the Ca<sup>2+</sup> chelator BAPTA (30 mM; see Materials and Methods). We approached it close to a neighboring fluorescent astrocyte (Figures 1A,B). Recording the astrocyte in cell-attached mode did not affect the evoked EPSCs (Figure 2D). However, few minutes (3–21) after breaking-in to whole-cell configuration, the amplitude of the first EPSC was significantly increased (Figures 2B–D;  $n = 6$ ; significant increase for five of six cells considered individually;  $6.8 \cdot 10^{-13} < p < 0.033$ ; two sample *t*-test and two sample Kolmogorov–Smirnov; significant increase for the mean values of all cells pooled together;  $105 \pm 142 \text{ pA}$  to  $135 \pm 179 \text{ pA}$ ,  $p = 0.03$ ; Wilcoxon signed-rank test). This result suggests that astrocytes exert a tonic inhibition on ventral horn neurons by releasing a gliotransmitter through a Ca<sup>2+</sup> dependent mechanism. To determine if the inhibition occurred pre- or postsynaptically, we considered the response induced by the second stimulation. On average, chelating Ca<sup>2+</sup> in the astrocyte had no significant effect on the second EPSC (Figure 2E;  $95 \pm 63 \text{ pA}$  to  $97 \pm 68 \text{ pA}$ ;  $n = 6$ ;  $p = 0.69$ ; Wilcoxon signed-rank test). Consequently the PPR was decreased in all the cells tested (Figures 2F–H; significant decrease for all the cells pooled together;  $1.4 \pm 0.5$  to  $1.0 \pm 0.3$ ,  $p = 0.03$  Wilcoxon signed-rank test). The input resistance of the recorded neurons was not changed after chelating Ca<sup>2+</sup> in astrocytes ( $514 \pm 298 \text{ M}\Omega$  in control;  $507 \pm 284 \text{ M}\Omega$  after Ca<sup>2+</sup> chelating,  $n = 5$ ;  $p = 0.3$ ; Wilcoxon signed-rank test). Altogether these results suggest that astrocytes produce a tonic inhibition in the ventral horn by releasing a gliotransmitter that inhibits excitatory transmission at the presynaptic level.

### SPINAL ASTROCYTES GENERATE PHASIC PRESYNAPTIC INHIBITION


Next, we investigated if a phasic activation of astrocytes also produced an inhibition of synaptic transmission. For this purpose, we used a pipette filled with the peptide TFLLR, an agonist for the G-protein coupled PAR-1 receptor that is preferentially expressed in astrocytes (Lee et al., 2007; Shigetomi et al., 2008). TFLLR induces a Ca<sup>2+</sup> increase in astrocytes (Lee et al., 2007; Shigetomi et al., 2008; Wang et al., 2013). When we puffed TFLLR between the stimulation and recording electrodes (Figure 3A), the amplitude of the first EPSC evoked by a paired stimulation protocol was significantly decreased (Figures 3B–D;  $n = 25$ ; significant decrease for 20/25 cells tested individually;  $3.3 \cdot 10^{-6} < p < 0.05$ ; paired sample *t*-test, Kolmogorov–Smirnov test and two sample *t*-test; significant decrease for all the cells pooled together;  $p = 1.9 \cdot 10^{-7}$ , paired sample *t*-test). The second EPSC was not significantly affected by TFLLR (Figure 3E;  $n = 25$ ;  $p = 0.06$ ; paired *t*-test). Consequently the PPR was increased (Figures 3F–H;  $n = 25$ ; significant increase for all cells taken together;  $p = 1.7 \cdot 10^{-5}$ , paired *t*-test). The effect of TFLLR was correlated with the PPR observed under control conditions: the inhibition was stronger for synapses with a high release probability (PPR  $< 1$ ; significant correlation;  $p = 0.04$ ;  $R = -0.4$ ; Pearson correlation test;  $n = 25$ ). Since neither the input resistance of the postsynaptic



neuron, nor the holding current were affected by TFLLR (control input resistance:  $730 \pm 531 \text{ M}\Omega$ ; after TFLLR:  $741 \pm 572 \text{ M}\Omega$ ,  $n = 23$ ;  $p = 0.36$ , Wilcoxon signed-rank test; control holding current:  $-63 \pm 69 \text{ pA}$ ; after TFLLR:  $-67 \pm 75 \text{ pA}$ ,  $n = 25$ ;  $p = 0.11$ , Wilcoxon signed-rank test), our results suggest that activation of astrocytes triggers the release of a substance that produces presynaptic inhibition of excitatory synaptic transmission in the ventral horn of the spinal cord. To identify the gliotransmitter involved, we performed a range of pharmacological tests.

**DUAL MODULATION OF SYNAPTIC TRANSMISSION BY ATP**

Since purines are major intrinsic modulators in the spinal cord (Dale and Gilday, 1996) that can be released by astrocytes (Pascual et al., 2005; Fields and Burnstock, 2006; Panatier et al., 2011), we tested the effect of ATP on synaptic transmission (Figure 4A). A 10 s Puff of ATP (1 mM) between the stimulation and recording electrodes reduced the amplitude of evoked pairs EPSCs (Figure 4B;  $n = 6$ ). A careful inspection of the recordings revealed that the effects induced by ATP evolved with time. During the puff itself, the amplitude of the first EPSC was reduced of 18% (Figure 4C;




from  $131 \pm 72$  pA to  $108 \pm 37$  pA; non-significant reduction,  $p = 0.22$ , Wilcoxon signed-rank test) while the second EPSC was decreased of 30% (Figure 4D; from  $132 \pm 71$  pA to  $93 \pm 45$  pA; significant reduction for all the cells tested together;  $n = 6$ ;  $p = 0.03$ ; Wilcoxon signed-rank test). Consequently, the PPR was significantly reduced (Figures 4E,F; from  $1.08 \pm 0.5$  in control conditions to  $0.85 \pm 0.3$  during ATP;  $p = 0.03$ , Wilcoxon signed-rank test). The effects initially induced by ATP were concomitant with a decrease in input resistance of postsynaptic neurons (from  $480 \pm 318$  M $\Omega$  to  $442 \pm 295$  M $\Omega$ ;  $p = 0.03$ , Wilcoxon signed-rank test), suggesting a mixture of pre- and postsynaptic effects. However, 1 min after the puff, the effects induced by ATP were different. The response to the first shock was more decreased (Figures 4B,C; from  $131 \pm 72$  pA to  $81 \pm 35$  pA;  $n = 6$ ; significant decrease for 4/6 cells,  $0.0015 < p < 0.019$ , two sample  $t$ -test; significant decrease for all cells pooled together,  $p = 0.03$ , Wilcoxon signed-rank test), while the response to the second shock was less affected

(Figure 4D; from  $132 \pm 71$  pA to  $109 \pm 59$  pA, significant decrease for all cells pooled together,  $p = 0.03$ , Wilcoxon signed-rank test). Thus, after an initial decrease, ATP induced an increase of the PPR (Figures 4E,F; from  $1.08 \pm 0.5$  to  $1.35 \pm 0.5$ ,  $p = 0.03$ , Wilcoxon signed-rank test). Since 1 min after the puff, the input resistance of the recorded neuron was not significantly different from control conditions ( $480 \pm 317$  M $\Omega$  in control;  $500 \pm 363$  M $\Omega$  1 min after ATP;  $p = 0.6$ , Wilcoxon signed-rank test), our results suggest that ATP induced presynaptic inhibition of transmitter release.

#### ADENOSINE INDUCES PRESYNAPTIC INHIBITION

The latency of the effects induced by ATP suggests that presynaptic inhibition is not induced by ATP itself, but by one of its metabolic products. Since ATP gets converted into adenosine by extracellular ectonucleotidases (Fields and Burnstock, 2006), we tested the effect of a puff of adenosine (1 mM), using the



same protocol (Figure 5A). We found that adenosine induced a strong inhibition of the first EPSC (Figure 5C;  $n = 6$ ; from  $133 \pm 134$  pA to  $64 \pm 52$  pA; significant decrease for 5/6 cells;  $5.9 \cdot 10^{-17} < p < 0.002$ ; two sample *t*-test and two sample Kolmogorov–Smirnov; significant decrease for all cells pooled together,  $p = 0.03$ , Wilcoxon signed rank test) and a weaker inhibition of the second EPSC (Figure 5D;  $n = 6$ ; from  $141 \pm 170$  pA to  $85 \pm 76$  pA; significant decrease for 4/6 cells;  $1.2 \cdot 10^{-12} < p < 0.008$ , two sample *t*-test and two sample Kolmogorov–Smirnov; non-significant decrease for all cells pooled together,  $p = 0.16$ , Wilcoxon signed rank test). Here also, the PPR was significantly increased (Figure 5E; from  $1.0 \pm 0.3$  to  $1.4 \pm 0.4$ ;  $p = 0.03$ , Wilcoxon signed-rank test). In contrast with ATP, the effects of adenosine were homogeneous and occurred without significant change in input resistance ( $1007 \pm 794$  M $\Omega$  in control;  $949 \pm 680$  M $\Omega$  after adenosine;  $p = 0.8$ ; Wilcoxon signed-rank test). Altogether, these results suggest that adenosine inhibits EPSCs via a presynaptic mechanism.


#### THE SELECTIVE BLOCKADE OF ECTONUCLEOTIDASES PREVENTS THE EFFECTS OF TFLLR

We tested further the possible involvement of ATP in presynaptic inhibition by means of the very selective ectonucleotidase

inhibitor ARL 67156 (Levesque et al., 2007). We selected neurons for which TFLLR induced presynaptic inhibition (Figures 6A,B). After addition of ARL 67156 (50  $\mu$ M) to the extracellular medium, the first EPSC was not affected anymore by a puff of TFLLR (Figure 6E;  $n = 6$ ; EPSC in normal Ringer:  $-76.2 \pm 30$  pA; after TFLLR:  $-52.0 \pm 31.6$  pA; significant decrease for 4/6 cells,  $3.3 \cdot 10^{-6} < p < 0.047$ , two sample *t*-test and two sample Kolmogorov–Smirnov; significant decrease for all cells pooled together,  $p = 0.03$ , Wilcoxon signed-rank test; EPSC in ARL 67156:  $70.6 \pm 44.3$  pA; after TFLLR:  $73.2 \pm 48.9$  pA; no significant decrease for cells considered individually  $0.11 < p < 0.81$ , two sample *t*-test and Kolmogorov–Smirnov; no significant decrease for all cells pooled together,  $p = 0.7$ , Wilcoxon signed-rank test). Moreover, the PPR was not significantly affected (Figures 6D,F;  $n = 6$ ; PPR in normal Ringer:  $1.0 \pm 0.2$ ; TFLLR:  $1.7 \pm 0.4$ ;  $p = 0.03$  Wilcoxon signed-rank test; PPR in ARL 67156:  $1.2 \pm 0.4$ ; TFLLR:  $1.0 \pm 0.2$ ;  $p = 0.2$  Wilcoxon signed-rank test). We therefore conclude that presynaptic inhibition is not mediated by ATP, but rather by one of its metabolic products.

#### TFLLR INDUCES PRESYNAPTIC INHIBITION BY ACTIVATING ADENOSINE A<sub>1</sub> RECEPTORS

Since adenosine induced presynaptic inhibition (Figure 5), we tested if blocking adenosine A<sub>1</sub> receptors prevented the induction



of presynaptic inhibition by TFLLR (Figure 7A). We found that in the presence of the selective adenosine A<sub>1</sub> receptor antagonist DPCPX (5 μM), puffing TFLLR did not affect the amplitude of evoked EPSCs anymore (Figures 7B,E; n = 6; effect of TFLLR in control conditions: from 153 ± 63 pA to 111 ± 47; significant decrease for 6/6 cells, 0.0017 < p < 0.043, two sample t-test and two sample Kolmogorov–Smirnov; significant decrease for all cells pooled together, p = 0.03, Wilcoxon signed-rank test; effect of TFLLR in DPCPX: from 135 ± 63 pA to 131 ± 58 pA; no significant decrease for cells considered individually: 0.18 < p < 0.71, two sample t-test and Kolmogorov–Smirnov; no significant decrease for all cells pooled together; p = 0.3, Wilcoxon signed-rank test). The PPR, which was significantly increased in control conditions (Figures 7C,D), was not altered anymore (Figures 7D,F; n = 6; increase of PPR in control conditions: from 1.1 ± 0.3 to 1.4 ± 0.5; p = 0.03, Wilcoxon signed-rank test; no change of PPR in DPCPX: from 1.2 ± 0.4 to 1.1 ± 0.4; p = 0.3, Wilcoxon signed-rank test). Our results demonstrate that the metabolic product of ATP that induces presynaptic inhibition of excitatory synaptic transmission is likely to be adenosine acting on A<sub>1</sub> receptors.

**DISCUSSION**


Our study suggests that astrocytes located in the ventral horn of the spinal cord induce both tonic and phasic presynaptic inhibition of

excitatory synaptic transmission. Our results indicate that extracellular ectonucleotidases convert ATP released by astrocytes into adenosine, which binds to A<sub>1</sub> inhibitory receptors located on the presynaptic side of excitatory synapses (Figure 8).

**THE MODULATION INDUCED BY ASTROCYTES IS BOTH TONIC AND PHASIC**

Chelation of Ca<sup>2+</sup> in one astrocyte induced a decrease of the PPR suggesting a removal of presynaptic inhibition (Figure 2). The most likely mechanism is an increase of the residual Ca<sup>2+</sup> present in presynaptic terminals since both the release probability (p) and the pool of available quanta (n) are Ca<sup>2+</sup> dependent (Zucker and Regehr, 2002). Even though the conductance measured in postsynaptic neurons was unaffected by the BAPTA treatment, we cannot rule out postsynaptic mechanisms (for example if there are two or more conductances affected in opposite ways).

This result suggests that astrocytes continuously release purines that produce presynaptic inhibition that might contribute to lower the activity of the spinal motor network. Tonic release of vesicular ATP from astrocytes was previously reported in the hippocampus (Pascual et al., 2005). Released ATP gets converted into adenosine by extracellular ectonucleotidases. This mechanism explains the persistent synaptic suppression mediated by adenosine (Pascual



et al., 2005). The central nervous system uses different strategies to decrease its excitability, the most common being a general increase of membrane conductance provided by an ambient concentration of GABA (Farrant and Nusser, 2005; Bautista et al., 2010).


Puffing TFLLR produced a transient inhibition of excitatory synaptic transmission (Figure 3). This suggests that the level of presynaptic inhibition provided by astrocytes is not only tonic but can also be regulated by the activity of the network. TFLLR is an exogenous molecule and does not activate astrocytes under physiological conditions. We did not investigate the nature of endogenous neurotransmitters responsible for the activation of astrocytes. The most likely scenario is that glutamate released from excitatory synapses binds to receptors located in the membrane of neighboring astrocytes that would in turn release ATP. Such a mechanism has been described in slices from the hippocampus where glutamate binding to metabotropic glutamate subtype 5 receptors triggers the release of ATP from astrocytes (Zhang et al., 2003; Panatier et al., 2011). Another possibility could be a change in pH. Brain hypoxia and ischemia lead to acidosis (Tombaugh and Sapolsky, 1993), which induces the release of ATP from astrocytes in the brainstem (Gourine et al., 2010). This process is essential for adjusting respiratory responses. These possibilities will be tested during future experiments.

after puffing TFLLR as a function of PPR before puffing TFLLR. In control conditions, the PPR was significantly increased (green open circles). In the presence of ARL 67156, the PPR was not significantly increased anymore (brown open circles). (E) Same neuron as in (B), recorded after addition of ARL 67156 to the bath. Puffing TFLLR did not affect the amplitude of the EPSC anymore (brown trace). (F) Mean values of the PPR before and after puffing TFLLR in the presence of ARL 67156.

The interpretation of our results relies on the hypothesis that the number of axons recruited by the electrical stimulation remained constant during each experiment. Puffing drug with pressure could in principle induce a mechanical artifact resulting in a change of the excitability of stimulated fibers. However, if it was the case, the number of recruited axons would either increase or decrease in a random manner. In contrast, puffing TFLLR induced either no effect or a reliable decrease of synaptic transmission. In addition, when blocking ectonucleotidases or A<sub>1</sub> receptors abolished the effect of TFLLR puff (see Figures 6 and 7), which rules out the possibility of a substantial change in the number of recruited axons.

#### IDENTITY OF THE PURINES AND OF THE RECEPTORS RESPONSIBLE FOR PRESYNAPTIC INHIBITION

Our results demonstrate that presynaptic inhibition is triggered by ATP, which, after enzymatic hydrolysis to adenosine binds to presynaptic receptors. Several arguments support this interpretation. First, presynaptic inhibition induced by TFLLR was both reproduced by ATP and adenosine (Figures 4 and 5). However, in contrast to adenosine, the inhibition produced by ATP was complex since, during the puff application, it was characterized by a decrease in PPR, which is not compatible with presynaptic inhibition. After 1 min, the inhibition produced by ATP was characterized by an increase in PPR, like the effect




induced by adenosine. This suggests that adenosine rather than ATP is responsible for presynaptic inhibition. This interpretation is supported by the fact that the conversion of ATP to adenosine occurs within a range of 200 ms (Dunwiddie et al., 1997). Second, blocking the hydrolysis of ATP to adenosine by inhibiting extracellular ectonucleotidases with ARL 67156 suppressed the presynaptic inhibition triggered by TFLLR (Figure 6). This confirms that ATP itself does not induce presynaptic inhibition and suggests instead that a metabolic product of ATP such as ADP, AMP, or adenosine is involved (Fields and Burnstock, 2006). Third, blocking adenosine  $A_1$  receptors with DPCPX suppressed the effects of TFLLR (Figure 7). This demonstrates that the purine responsible for presynaptic inhibition is actually adenosine.

Adenosine inhibits synaptic release of transmitter in other systems such as the dentate gyrus (Dolphin and Archer, 1983) and the CA1 region of the hippocampus (Corradetti et al., 1984; Wu and Saggau, 1994), the basal forebrain (Hawryluk et al., 2012), and the cerebral cortex (Hollins and Stone, 1980). In most cases, presynaptic adenosine  $A_1$  receptors decrease transmitter release by inhibiting a  $Ca^{2+}$  current (Hollins and Stone, 1980; Wu and Saggau, 1994; Dale and Gilday, 1996; Brown and Dale, 2000; Zhang et al., 2003; Fields and Burnstock, 2006). In contrast, adenosine  $A_{2A}$  receptors facilitate synaptic transmission (Panatier

et al., 2011). In agreement, we found that  $A_1$  receptors mediate presynaptic inhibition in the ventral horn of the spinal cord (Figure 7).

#### FUNCTIONAL RELEVANCE

Regulating the excitability motor networks is essential for adjusting the motor command. Hyperexcitability of motor networks leads to involuntary movements and spasms. Tonic presynaptic inhibition of excitatory transmission may provide a general decrease of the activity of highly active networks and act as a homeostatic mechanism. In other regions of the central nervous system such as the hippocampus, adenosine released from astrocytes has neuroprotective effects and the activation of presynaptic  $A_1$  receptors acts as an endogenous anticonvulsant (Boison, 2012). In the spinal cord, artificial spinal rhythmic activities induced by cocktails of pharmacological agents are modulated by adenosine. The activation of  $A_1$  receptors depresses bicuculline-evoked seizures in newborn rats (Brockhaus and Ballanyi, 2000; Taccola et al., 2012). This suggests that adenosine also has anticonvulsant effects in the spinal cord. Indeed, in the spinal cord of young rats, acute hypoxia triggers the release of adenosine, which in turn depresses monosynaptic reflexes (Otsuguro et al., 2011). Such a neuroprotective role of adenosine is well documented in patients suffering from temporal lobe epilepsy. Seizures trigger a massive release of adenosine,



which in turn activate A<sub>1</sub> receptors and thereby terminate the activity (Boison, 2012).

Adenosine might also be involved in the tuning of locomotor activity. Antagonists for adenosine receptors from the Methylxanthine family (such as caffeine or theophylline) increase locomotor activity (Snyder et al., 1981). During fictive locomotion induced in neonatal mice by a mixture of serotonin, NMDA and dopamine, bath application of adenosine slows down rhythmic activity by around 30% (Witts et al., 2012). To figure out the actual physiological role of purines during locomotion in adult mammals, one will have to measure their concentrations. The recent development of probes allowing the detection of ATP and adenosine in real time (Frenguelli et al., 2007) may allow figuring out if the concentration of purines changes tonically or rhythmically during movement. Interestingly, ATP increases the excitability of spinal locomotor networks (Dale and Gilday, 1996). Since the hydrolysis of ATP to adenosine occurs one range of magnitude faster than the locomotor cycle (Dunwiddie et al., 1997), a possible scenario could be that astrocytes release ATP cyclically, in phase with the rhythm. The fast degradation of ATP into adenosine would then induce a rhythmic inhibition of excitatory transmission. However, high-speed locomotion could be too fast for rhythmic

mediated inhibition. We will test these hypotheses during future experiments.

## ACKNOWLEDGMENTS

The authors wish to thank Professor Frank Kirchhoff and Dr. Anja Scheller (University of Saarland, Germany) for providing the transgenic mouse and Dr Laure Plantard from the CFIM for her help with confocal microscopy. The project was funded by Owensenske Fond, Simon Fougner Hartmanns Familiefond, Agnes and Poul Friis Fond, Novo Scholarship Programme, Augustinus Foundation, Oticon Foundation, Molecular Biomedicine Travel Stipend, Lundbeck Foundation and Carlsberg-fondet.

## REFERENCES

- Ameri, A. (1999). The effects of cannabinoids on the brain. *Prog. Neurobiol.* 58, 315–348. doi: 10.1016/S0301-0082(98)00087-2
- Arvidsson, U., Cullheim, S., Ulfhake, B., Bennett, G. W., Fone, K. C., Cuello, A. C., et al. (1990). 5-Hydroxytryptamine, substance P, and thyrotropin-releasing hormone in the adult rat spinal cord segment L7: immunohistochemical and chemical studies. *Synapse* 6, 237–270. doi: 10.1002/syn.890060305
- Bautista, W., Aguilar, J., Loeza-Alcocer, J. E., and Delgado-Lezama, R. (2010). Pre- and postsynaptic modulation of monosynaptic reflex by GABA<sub>A</sub> receptors on turtle spinal cord. *J. Physiol.* 588, 2621–2631. doi: 10.1113/jphysiol.2010.188979
- Boison, D. (2012). Adenosine dysfunction in epilepsy. *Glia* 60, 1234–1243. doi: 10.1002/glia.22285
- Brocard, F., Tazerart, S., and Vinay, L. (2010). Do pacemakers drive the central pattern generator for locomotion in mammals? *Neuroscientist* 16, 139–155. doi: 10.1177/1073858409346339
- Brockhaus, J., and Ballanyi, K. (2000). Anticonvulsant A(1) receptor-mediated adenosine action on neuronal networks in the brainstem-spinal cord of newborn rats. *Neuroscience* 96, 359–371. doi: 10.1016/S0306-4522(99)00544-8
- Brown, P., and Dale, N. (2000). Adenosine A1 receptors modulate high voltage-activated Ca<sup>2+</sup> currents and motor pattern generation in the xenopus embryo. *J. Physiol.* 525(Pt 3), 655–667. doi: 10.1111/j.1469-7793.2000.00655.x
- Brown, P., and Dale, N. (2002). Modulation of K(+) currents in Xenopus spinal neurons by p2y receptors: a role for ATP and ADP in motor pattern generation. *J. Physiol.* 540, 843–850. doi: 10.1113/jphysiol.2001.013192
- Burnstock, G. (2007). Physiology and pathophysiology of purinergic neurotransmission. *Physiol. Rev.* 87, 659–797. doi: 10.1152/physrev.00043.2006
- Christensen, R. K., Petersen, A. V., and Perrier, J. F. (2013). How do glial cells contribute to motor control? *Curr. Pharm. Des.* 19, 4385–4399. doi: 10.2174/13816128113199990384
- Conn, P. J., Battaglia, G., Marino, M. J., and Nicoletti, F. (2005). Metabotropic glutamate receptors in the basal ganglia motor circuit. *Nat. Rev. Neurosci.* 6, 787–798. doi: 10.1038/nrn1763
- Corradetti, R., Lo Conte, G., Moroni, F., Passani, M. B., and Pepeu, G. (1984). Adenosine decreases aspartate and glutamate release from rat hippocampal slices. *Eur. J. Pharmacol.* 104, 19–26. doi: 10.1016/0014-2999(84)90364-9
- Cotel, F., Exley, R., Cragg, S. J., and Perrier, J. F. (2013). Serotonin spillover onto the axon initial segment of motoneurons induces central fatigue by inhibiting action potential initiation. *Proc. Natl. Acad. Sci. U.S.A.* 110, 4774–4779. doi: 10.1073/pnas.1216150110
- Dai, X., Noga, B. R., Douglas, J. R., and Jordan, L. M. (2005). Localization of spinal neurons activated during locomotion using the c-fos immunohistochemical method. *J. Neurophysiol.* 93, 3442–3452. doi: 10.1152/jn.00578.2004
- Dale, N., and Gilday, D. (1996). Regulation of rhythmic movements by purinergic neurotransmitters in frog embryos. *Nature* 383, 259–263. doi: 10.1038/383259a0
- Dolphin, A. C., and Archer, E. R. (1983). An adenosine agonist inhibits and a cyclic AMP analogue enhances the release of glutamate but not GABA from slices of rat dentate gyrus. *Neurosci. Lett.* 43, 49–54. doi: 10.1016/0304-3940(83)90127-1
- Dunwiddie, T. V., Diao, L., and Proctor, W. R. (1997). Adenine nucleotides undergo rapid, quantitative conversion to adenosine in the extracellular space in rat hippocampus. *J. Neurosci.* 17, 7673–7682.

- El Manira, A., and Kyriakatos, A. (2010). The role of endocannabinoid signaling in motor control. *Physiology (Bethesda)* 25, 230–238. doi: 10.1152/physiol.00007.2010
- Farrant, M., and Nusser, Z. (2005). Variations on an inhibitory theme: phasic and tonic activation of GABA(A) receptors. *Nat. Rev. Neurosci.* 6, 215–229. doi: 10.1038/nrn1625
- Fields, R. D., and Burnstock, G. (2006). Purinergic signalling in neuron-glia interactions. *Nat. Rev. Neurosci.* 7, 423–436. doi: 10.1038/nrn1928
- Freguelli, B. G., Wigmore, G., Llaudet, E., and Dale, N. (2007). Temporal and mechanistic dissociation of ATP and adenosine release during ischaemia in the mammalian hippocampus. *J. Neurochem.* 101, 1400–1413. doi: 10.1111/j.1471-4159.2007.04425.x
- Gourine, A. V., Kasymov, V., Marina, N., Tang, F., Figueiredo, M. F., Lane, S., et al. (2010). Astrocytes control breathing through pH-dependent release of ATP. *Science* 329, 571–575. doi: 10.1126/science.1190721
- Guthrie, P. B., Knappenberger, J., Segal, M., Bennett, M. V. L., Charles, A. C., and Kater, S. B. (1999). ATP released from astrocytes mediates glial calcium waves. *J. Neurosci.* 19, 520–528.
- Hawryluk, J. M., Ferrari, L. L., Keating, S. A., and Arrigoni, E. (2012). Adenosine inhibits glutamatergic input to basal forebrain cholinergic neurons. *J. Neurophysiol.* 107, 2769–2781. doi: 10.1152/jn.00528.2011
- Hollins, C., and Stone, T. W. (1980). Adenosine inhibition of gamma-aminobutyric acid release from slices of rat cerebral cortex. *Br. J. Pharmacol.* 69, 107–112. doi: 10.1111/j.1476-5381.1980.tb10888.x
- Jacobs, B. L., and Azmitia, E. C. (1992). Structure and function of the brain serotonin system. *Physiol. Rev.* 72, 165–229.
- Jo, Y. H., and Schlichter, R. (1999). Synaptic corelease of ATP and GABA in cultured spinal neurons. *Nat. Neurosci.* 2, 241–245. doi: 10.1038/6344
- Kjaerulff, O., and Kiehn, O. (1996). Distribution of networks generating and coordinating locomotor activity in the neonatal rat spinal cord in vitro: a lesion study. *J. Neurosci.* 16, 5777–5794.
- Lalo, U., Palygin, O., Rasooli-Nejad, S., Andrew, J., Haydon, P. G., and Pankratov, Y. (2014). Exocytosis of ATP from astrocytes modulates phasic and tonic inhibition in the neocortex. *PLoS Biol.* 12:e1001747. doi: 10.1371/journal.pbio.1001747
- Lanuza, G. M., Gosgnach, S., Pierani, A., Jessell, T. M., and Goulding, M. (2004). Genetic identification of spinal interneurons that coordinate left-right locomotor activity necessary for walking movements. *Neuron* 42, 375–386. doi: 10.1016/S0896-6273(04)00249-1
- Lee, C. J., Mannaioni, G., Yuan, H., Woo, D. H., Gingrich, M. B., and Traynelis, S. F. (2007). Astrocytic control of synaptic NMDA receptors. *J. Physiol.* 581, 1057–1081. doi: 10.1113/jphysiol.2007.130377
- Levesque, S. A., Lavoie, E. G., Lecka, J., Bigonnesse, F., and Sevigny, J. (2007). Specificity of the ecto-ATPase inhibitor ARL 67156 on human and mouse ectonucleotidases. *Br. J. Pharmacol.* 152, 141–150. doi: 10.1038/sj.bjp.0707361
- Mui, J. W., Willis, K. L., Hao, Z. Z., and Berkowitz, A. (2012). Distributions of active spinal cord neurons during swimming and scratching motor patterns. *J. Comp. Physiol. A Neuroethol. Sens. Neural. Behav. Physiol.* 198, 877–889. doi: 10.1007/s00359-012-0758-6
- Nolte, C., Matyash, M., Pivneva, T., Schipke, C. G., Ohlemeyer, C., Hanisch, U. K., et al. (2001). GFAP promoter-controlled EGFP-expressing transgenic mice: a tool to visualize astrocytes and astrogliosis in living brain tissue. *Glia* 33, 72–86. doi: 10.1002/1098-1136(20010101)33:1<72::AID-GLIA1007>3.0.CO;2-A
- Ono, H., and Fukuda, H. (1995). Pharmacology of descending noradrenergic systems in relation to motor function. *Pharmacol. Ther.* 68, 105–112. doi: 10.1016/0163-7258(95)02001-2
- Otsuguro, K., Wada, M., and Ito, S. (2011). Differential contributions of adenosine to hypoxia-evoked depressions of three neuronal pathways in isolated spinal cord of neonatal rats. *Br. J. Pharmacol.* 164, 132–144. doi: 10.1111/j.1476-5381.2011.01333.x
- Panatier, A., Vallee, J., Haber, M., Murai, K. K., Lacaille, J. C., and Robitaille, R. (2011). Astrocytes are endogenous regulators of basal transmission at central synapses. *Cell* 146, 785–798. doi: 10.1016/j.cell.2011.07.022
- Pascual, O., Casper, K. B., Kubera, C., Zhang, J., Revilla-Sanchez, R., Sul, J. Y., et al. (2005). Astrocytic purinergic signaling coordinates synaptic networks. *Science* 310, 113–116. doi: 10.1126/science.1116916
- Perrier, J. F., and Delgado-Lezama, R. (2005). Synaptic release of serotonin induced by stimulation of the raphe nucleus promotes plateau potentials in spinal motoneurons of the adult turtle. *J. Neurosci.* 25, 7993–7999. doi: 10.1523/JNEUROSCI.1957-05.2005
- Perrier, J. F., Rasmussen, H. B., Christensen, R. K., and Petersen, A. V. (2013). Modulation of the intrinsic properties of motoneurons by serotonin. *Curr. Pharm. Des.* 19, 4371–4384. doi: 10.2174/13816128113199990341
- Prut, Y., and Perlmutter, S. I. (2003). Firing properties of spinal interneurons during voluntary movement. I. State-dependent regularity of firing. *J. Neurosci.* 23, 9600–9610.
- Schmidt, B. J., and Jordan, L. M. (2000). The role of serotonin in reflex modulation and locomotor rhythm production in the mammalian spinal cord. *Brain Res. Bull.* 53, 689–710. doi: 10.1016/S0304-9230(00)00402-0
- Shigetomi, E., Bowser, D. N., Sofroniew, M. V., and Khakh, B. S. (2008). Two forms of astrocyte calcium excitability have distinct effects on NMDA receptor-mediated slow inward currents in pyramidal neurons. *J. Neurosci.* 28, 6659–6663. doi: 10.1523/JNEUROSCI.1717-08.2008
- Silinsky, E. M., and Redman, R. S. (1996). Synchronous release of ATP and neurotransmitter within milliseconds of a motor nerve impulse in the frog. *J. Physiol.* 492(Pt 3), 815–822.
- Snyder, S. H., Katims, J. J., Annau, Z., Bruns, R. F., and Daly, J. W. (1981). Adenosine receptors and behavioral actions of methylxanthines. *Proc. Natl. Acad. Sci. U.S.A.* 78, 3260–3264. doi: 10.1073/pnas.78.5.3260
- Taccola, G., Olivieri, D., D'Angelo, G., Blackburn, P., Secchia, L., and Balanyi, K. (2012). A(1) adenosine receptor modulation of chemically and electrically evoked lumbar locomotor network activity in isolated newborn rat spinal cords. *Neuroscience* 222, 191–204. doi: 10.1016/j.neuroscience.2012.07.030
- Tombaugh, G. C., and Sapolsky, R. M. (1993). Evolving concepts about the role of acidosis in ischemic neuropathology. *J. Neurochem.* 61, 793–803. doi: 10.1111/j.1471-4159.1993.tb03589.x
- Wang, F., Smith, N. A., Xu, Q., Goldman, S., Peng, W., Huang, J. H., et al. (2013). Photolysis of caged Ca<sup>2+</sup> but not receptor-mediated Ca<sup>2+</sup> signaling triggers astrocytic glutamate release. *J. Neurosci.* 33, 17404–17412. doi: 10.1523/JNEUROSCI.2178-13.2013
- Witts, E. C., Panetta, K. M., and Miles, G. B. (2012). Glial-derived adenosine modulates spinal motor networks in mice. *J. Neurophysiol.* 107, 1925–1934. doi: 10.1152/jn.00513.2011
- Wu, L. G., and Saggau, P. (1994). Adenosine inhibits evoked synaptic transmission primarily by reducing presynaptic calcium influx in area CA1 of hippocampus. *Neuron* 12, 1139–1148. doi: 10.1016/0896-6273(94)90321-2
- Zhang, J. M., Wang, H. K., Ye, C. Q., Ge, W., Chen, Y., Jiang, Z. L., et al. (2003). ATP released by astrocytes mediates glutamatergic activity-dependent heterosynaptic suppression. *Neuron* 40, 971–982. doi: 10.1016/S0896-6273(03)00717-7
- Zucker, R. S., and Regehr, W. G. (2002). Short-term synaptic plasticity. *Annu. Rev. Physiol.* 64, 355–405. doi: 10.1146/annurev.physiol.64.092501.114547

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 17 March 2014; accepted: 20 May 2014; published online: 04 June 2014.

Citation: Carlsen EM and Perrier J-F (2014) Purines released from astrocytes inhibit excitatory synaptic transmission in the ventral horn of the spinal cord. *Front. Neural Circuits* 8:60. doi: 10.3389/fncir.2014.00060

This article was submitted to the journal *Frontiers in Neural Circuits*.

Copyright © 2014 Carlsen and Perrier. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.