



Corrigendum: Reconsidering Tonotopic Maps in the Auditory Cortex and Lemniscal Auditory Thalamus in Mice

Hiroaki Tsukano^{1*}, Masao Horie^{2†}, Shinpei Ohga¹, Kuniyuki Takahashi³, Yamato Kubota³, Ryuichi Hishida¹, Hirohide Takebayashi² and Katsuei Shibuki¹

¹ Department of Neurophysiology, Brain Research Institute, Niigata University, Niigata, Japan, ² Division of Neurobiology and Anatomy, Graduate School of Medicine and Dental Sciences, Niigata University, Niigata, Japan, ³ Division of Otolaryngology, Graduate School of Medicine and Dental Sciences, Niigata University, Niigata, Japan

Keywords: brain map, auditory cortex, medial geniculate body, tonotopy, topology, thalamocortical pathway, multiple compartments, mice

OPEN ACCESS

Edited and reviewed by:

Takao K. Hensch,
Harvard University, United States

*Correspondence:

Hiroaki Tsukano
tsukano-nii@umin.ac.jp

† Present Address:

Masao Horie,
Department of Morphological
Sciences, Graduate School of Medical
and Dental Sciences, Kagoshima
University, Kagoshima, Japan

Received: 14 May 2017

Accepted: 22 May 2017

Published: 31 May 2017

Citation:

Tsukano H, Horie M, Ohga S,
Takahashi K, Kubota Y, Hishida R,
Takebayashi H and Shibuki K (2017)
Corrigendum: Reconsidering
Tonotopic Maps in the Auditory Cortex
and Lemniscal Auditory Thalamus in
Mice. *Front. Neural Circuits* 11:39.
doi: 10.3389/fncir.2017.00039

A corrigendum on

Reconsidering Tonotopic Maps in the Auditory Cortex and Lemniscal Auditory Thalamus in Mice

by Tsukano, H., Horie, M., Ohga, S., Takahashi, K., Kubota, Y., Hishida, R., et al. (2017). *Front. Neural Circuits* 11:14. doi: 10.3389/fncir.2017.00014

We noticed that there was an error in the illustration in **Figure 2B**. In particular, we inadvertently drew the red oval in the opposite direction to the original data in the references (Tsukano et al., 2015, 2017). There are no relevant errors in the text part.

REFERENCES

- Tsukano, H., Horie, M., Bo, T., Uchimura, A., Hishida, R., Kudoh, M., et al. (2015). Delineation of a frequency-organized region isolated from the mouse primary auditory cortex. *J. Neurophysiol.* 113, 2900–2920. doi: 10.1152/jn.00932.2014
- Tsukano, H., Horie, M., Hishida, R., Takahashi, K., Takebayashi, H., and Shibuki, K. (2017). Independent tonotopy and thalamocortical projection patterns in two adjacent parts of the classical primary auditory cortex in mice. *Neurosci. Lett.* 637, 26–30. doi: 10.1016/j.neulet.2016.11.062

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Tsukano, Horie, Ohga, Takahashi, Kubota, Hishida, Takebayashi and Shibuki. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

