



Mechanisms Underlying Serotonergic Excitation of Callosal Projection Neurons in the Mouse Medial Prefrontal Cortex

Emily K. Stephens[†], Arielle L. Baker[†] and Allan T. Gulledge*

Department of Molecular and Systems Biology, Geisel School of Medicine at Dartmouth College, Hanover, NH, United States

Serotonin (5-HT) selectively excites subpopulations of pyramidal neurons in the neocortex via activation of 5-HT_{2A} (2A) receptors coupled to G_q subtype G-protein alpha subunits. G_q-mediated excitatory responses have been attributed primarily to suppression of potassium conductances, including those mediated by K_v7 potassium channels (i.e., the M-current), or activation of non-specific cation conductances that underlie calcium-dependent afterdepolarizations (ADPs). However, 2A-dependent excitation of cortical neurons has not been extensively studied, and no consensus exists regarding the underlying ionic effector(s) involved. In layer 5 of the mouse medial prefrontal cortex, we tested potential mechanisms of serotonergic excitation in commissural/callosal (COM) projection neurons, a subpopulation of pyramidal neurons that exhibits 2A-dependent excitation in response to 5-HT. In baseline conditions, 5-HT enhanced the rate of action potential generation in COM neurons experiencing suprathreshold somatic current injection. This serotonergic excitation was occluded by activation of muscarinic acetylcholine (ACh) receptors, confirming that 5-HT acts via the same G_q-signaling cascades engaged by ACh. Like ACh, 5-HT promoted the generation of calcium-dependent ADPs following spike trains. However, calcium was not necessary for serotonergic excitation, as responses to 5-HT were enhanced (by >100%), rather than reduced, by chelation of intracellular calcium with 10 mM BAPTA. This suggests intracellular calcium negatively regulates additional ionic conductances gated by 2A receptors. Removal of extracellular calcium had no effect when intracellular calcium signaling was intact, but suppressed 5-HT response amplitudes, by about 50%, when BAPTA was included in patch pipettes. This suggests that 2A excitation involves activation of a non-specific cation conductance that is both calcium-sensitive and calcium-permeable. M-current suppression was found to be a third ionic effector, as blockade of K_v7 channels with XE991 (10 μM) reduced serotonergic excitation by ~50% in control conditions, and by ~30% with intracellular BAPTA present. Together, these findings demonstrate a role for at least three distinct ionic effectors, including K_v7 channels, a calcium-sensitive and calcium-permeable non-specific cation conductance, and the calcium-dependent ADP conductance, in mediating serotonergic excitation of COM neurons.

Keywords: serotonin, cortex, pyramidal neuron, 5-HT_{2A} receptor, calcium, M-current, K_v7 channels, afterdepolarization

OPEN ACCESS

Edited by:

Michael E. Hasselmo,
Boston University, United States

Reviewed by:

John J. Woodward,
Medical University of South Carolina,
United States

Laura Schrader,
Tulane University, United States

*Correspondence:

Allan T. Gulledge
allan.gulledge@dartmouth.edu

[†]These authors have contributed
equally to this work.

Received: 03 October 2017

Accepted: 04 January 2018

Published: 18 January 2018

Citation:

Stephens EK, Baker AL and
Gulledge AT (2018) Mechanisms
Underlying Serotonergic Excitation
of Callosal Projection Neurons
in the Mouse Medial Prefrontal
Cortex. *Front. Neural Circuits* 12:2.
doi: 10.3389/fncir.2018.00002

INTRODUCTION

In the prefrontal cortex (PFC), serotonin (5-HT) acts as a neurotransmitter to regulate diverse cognitive processes, including working memory (Williams et al., 2002; Liy-Salmeron and Meneses, 2008), cognitive flexibility (Clarke et al., 2004, 2005), impulsivity (Harrison et al., 1997; Winstanley et al., 2003), attention (Winstanley et al., 2003), and fear (Almada et al., 2015; Marinho et al., 2015; Leon et al., 2017). 5-HT regulates prefrontal circuits by acting on a variety of receptor subtypes differentially expressed in subpopulations of cortical neurons. In pyramidal cells, which comprise heterogeneous subpopulations of glutamatergic neurons, 5-HT acts primarily via two G-protein-coupled receptors that have opposing influences on neuronal excitability. 5-HT_{1A} (1A) receptors are expressed in perisomatic or axonal compartments (Azmitia et al., 1996; DeFelipe et al., 2001; Czyrak et al., 2003; Cruz et al., 2004) and inhibit neurons by enhancing potassium conductances downstream of G_{i/o}-associated G-protein β subunits (Andrade and Nicoll, 1987; Colino and Halliwell, 1987; Davies et al., 1987; Tanaka and North, 1993; Spain, 1994; Luscher et al., 1997; Goodfellow et al., 2009). On the other hand, 5-HT_{2A} (2A) receptors coupled to G_q subtype G-protein α subunits are expressed in dendritic compartments (Willins et al., 1997; Jakab and Goldman-Rakic, 1998; Xu and Pandey, 2000; Miner et al., 2003), and act generally to enhance pyramidal neuron excitability. However, the ionic mechanisms responsible for 2A-dependent excitation of pyramidal neurons have been less well studied (Araneda and Andrade, 1991; Spain, 1994; Villalobos et al., 2005, 2011), largely due to the fact that the 2A receptor is not universally expressed in adult neurons. As a result, much of our understanding of 2A signaling is inferred from studies of cholinergic excitation, which is mediated by the more ubiquitous G_q-coupled M1-subtype muscarinic acetylcholine (ACh) receptor. Classically, G_q-coupled excitation in pyramidal neurons is associated with suppression of potassium conductances, including those mediating the “M” (muscarine-suppressed) current and those responsible for afterhyperpolarizations (AHPs) in pyramidal neurons following bouts of action potential generation (Krnjevic et al., 1971; Schwindt et al., 1988; McCormick and Williamson, 1989; Araneda and Andrade, 1991; Wang and McCormick, 1993; Villalobos et al., 2005, 2011). However, reduction of potassium conductances does not fully account for G_q-mediated excitation in neocortical pyramidal neurons (Andrade, 1991;

Haj-Dahmane and Andrade, 1996), as G_q-coupled muscarinic ACh and 5-HT receptors also engage inward currents generated by calcium-dependent (Haj-Dahmane and Andrade, 1998; Yan et al., 2009; Lei et al., 2014) and independent (Haj-Dahmane and Andrade, 1996; Shalinsky et al., 2002; Egorov et al., 2003; Magistretti et al., 2004; Baker et al., 2018) non-specific cation channels.

Here, to better understand the mechanisms underlying 2A-mediated excitation in the cerebral cortex, we take advantage of the selective and robust 2A excitation that occurs in a subpopulation of pyramidal neurons that project to the contralateral cerebral hemisphere (Avesar and Gullledge, 2012). Using fluorescent retrograde tracers injected unilaterally in the mouse medial PFC (mPFC), we challenged 2A excitation in retrograde-labeled commissural/callosal (COM) projection neurons with a combination of pharmacological and ionic substitution approaches. Our results demonstrate that serotonin 2A receptors engage multiple ionic effectors, including the M-current, a calcium-sensitive but calcium-permeable non-specific cation conductance, and the calcium-dependent non-specific cation conductance underlying the ADP, to promote action potential generation in COM neurons.

MATERIALS AND METHODS

Animals

Experiments involved adult (6-to-10-week-old) C57BL/6J male and female mice housed in 12:12 h light:dark cycle and provided food and water *ad libitum*. All experiments were carried out according to methods approved by the Institutional Animal Care and Use Committee of Dartmouth College. No significant sex-dependent differences in cellular physiology or 5-HT responses were observed (Table 1).

Retrograde Labeling

Red or green fluorescent microbeads (Retrobeads, Lumafuor Inc.) were injected unilaterally into the left prelimbic cortex using age-appropriate coordinates (Paxinos and Franklin, 2004) to label COM neurons in the contralateral mPFC. Animals were anesthetized throughout surgeries with vaporized isoflurane (~2%). Following craniotomy, a microsyringe was lowered into the mPFC and 300 nL of undiluted Retrobead solution was injected over a 10 min period. Animals were allowed to recover from surgery for at least 72 h before use in electrophysiological

TABLE 1 | Physiology and serotonin responses in neurons from female and male mice.

Sex	n	RMP (mV)	R _N (M Ω)	I _h (%sag)	Serotonin responses	
					% Increase in firing rate	Response integral (Hz \times s)
Female	17	-80 \pm 5	171 \pm 45	8 \pm 4	112 \pm 56	114 \pm 90
Male	37	-81 \pm 5	182 \pm 67	9 \pm 4	94 \pm 49	145 \pm 117
P-value		0.64	0.51	0.70	0.28	0.30
All	54	-81 \pm 5	179 \pm 60	8 \pm 4	100 \pm 52	135 \pm 110

Data presented as means \pm standard deviations.

experiments. Locations of microbead injections were confirmed *post hoc* in coronal sections of the mPFC.

Slice Preparation

Following isoflurane anesthesia and decapitation, brains were removed and submerged in artificial cerebral spinal fluid (aCSF) containing, in mM: 125 NaCl, 25 NaHCO₃, 3 KCl, and 1.25 NaH₂PO₄, 0.5 CaCl₂, 5 MgCl₂, 25 glucose, and saturated with 95% O₂ / 5% CO₂. Coronal brain slices (250 μm thick) of the mPFC were cut using a Leica VT 1200 slicer and stored in a chamber filled with aCSF containing 2 mM CaCl₂ and 1 mM MgCl₂ at 35°C for ~45 min, then kept at room temperature (~26°C) until use in experiments.

Electrophysiology

Slices were transferred to a recording chamber continuously perfused with oxygenated aCSF at 35–36°C and visualized with an Olympus BX51WI microscope. Unless otherwise noted, whole-cell current-clamp recordings of Retrobead-labeled COM neurons in layer 5 were made with patch pipettes (~5 MΩ) filled with our standard intracellular solution containing, in mM: 135 K-gluconate, 2 NaCl, 2 MgCl₂, 10 HEPES, 3 Na₂ATP, and 0.3 NaGTP (pH 7.2 with KOH). Epifluorescence illumination using 470 nm or 530 nm LEDs was used to target Retrobead-labeled layer 5 COM neurons in the prelimbic cortex for whole-cell recording. Data were acquired with Axograph software (Axograph Scientific) using BVC-700 amplifiers (Dagan Corporation) and ITC-18 digitizers (HEKA Instruments). Membrane potentials were sampled at 25 kHz, filtered at 10 kHz, and corrected for a junction potential of +12 mV. As 2A-mediated excitation is activity-dependent (Stephens et al., 2014), responses to 5-HT were measured during periods of action potential generation (~5 Hz) produced by somatic DC current injection. Mean currents injected, and the resulting baseline firing frequencies, for different experimental conditions are listed in **Table 2**.

5-HT (100 μM) was dissolved in aCSF, loaded into a patch pipette placed ~50 μm from the targeted soma, and

pneumatically applied for 1 s at ~15 psi. COM neurons were classified as “COM-excited” or “COM-biphasic” based on their initial response to 5-HT (see Avesar and Gullledge, 2012). At 100 μM, all excitatory effects of 5-HT on COM neurons can be attributed to 2A receptors, as excitation is eliminated in the presence of 500 nM MDL 11939 (Avesar and Gullledge, 2012; Stephens et al., 2014). To preserve focus on 2A-driven signaling, only COM-excited neurons were used for experiments. Serotonergic excitation was quantified as the average instantaneous spike frequency (ISF) over the first 500 ms following the peak post-5-HT increase in ISF, as measured relative to the mean baseline ISF (i.e., as the percent increase above baseline firing rates), and as the integral of the increased ISF over time.

Pharmacological Manipulations

Carbamylcholine chloride (carbachol, 50–100 μM; Sigma–Aldrich) was used to activate G_q-coupled M1 muscarinic acetylcholine receptors, which mediate cholinergic excitation of pyramidal neurons in the mPFC and hippocampus (Gullledge et al., 2009; Dasari and Gullledge, 2011). XE991 (10 μM; Tocris Bioscience or Cayman Chemicals) was used to block K_{V7} channels. 1,2-Bis(2-Aminophenoxy)ethane-N,N,N',N'-tetraacetic acid (BAPTA, 10 mM; Sigma–Aldrich) was included in the patch-pipette solution in some experiments to chelate intracellular calcium. For experiments using nominally calcium-free aCSF, CaCl₂ was replaced with either equimolar or 5 mM MgCl₂ (no differences were observed between these two substitution conditions). In some experiments, extracellular KCl was reduced to 0.5 mM, with NaCl being raised to 127.5 mM.

Statistical Analyses

Unless otherwise noted, data are presented as mean ± SEM, and were assessed with either Student's *t*-test (two-tailed, paired or unpaired) or a one-way ANOVA (two-tailed, repeated measures where appropriate) using Wizard for Mac version 1.9 (Evan Miller). Significance was defined as *p* < 0.05.

TABLE 2 | Current injections and firing frequencies across experimental conditions.

Internal solution	Pharmacological manipulation	<i>n</i>	Baseline		+ Pharmacological manipulation		<i>p</i> -values (<i>I/f</i>)
			<i>I</i> (nA)	<i>f</i> (Hz)	<i>I</i> (nA)	<i>f</i> (Hz)	
Standard ↓	Aggregate baseline	54	0.12 ± 0.06	4.68 ± 1.41	—	—	—
	+ carbachol	9	0.14 ± 0.07	5.23 ± 2.59	0.07 ± 0.08	4.98 ± 1.36	0.03/0.76
	None (ADP experiments)	16	0.10 ± 0.07	4.00 ± 0.87	—	—	—
	+ Ca ²⁺ -free aCSF	9	0.11 ± 0.04	5.05 ± 0.97	0.10 ± 0.07	6.56 ± 1.61	0.47/ 0.04
	+ XE991	10	0.12 ± 0.05	4.03 ± 0.77	0.12 ± 0.06	3.65 ± 0.84	0.10/0.31
10 mM BAPTA ↓	+ low K ⁺ aCSF	10	0.13 ± 0.06	5.57 ± 0.66	0.12 ± 0.06	5.61 ± 0.66	0.12/0.81
	Aggregate baseline	43	0.28 ± 0.12	5.54 ± 1.51	vs. aggregate standard internal solution →		<0.001/0.004
	None	13	0.31 ± 0.13	6.34 ± 1.90	vs. 16 ADP cells →		<0.001/<0.001
	+ Ca ²⁺ -free aCSF	12	0.13 ± 0.05	4.52 ± 1.04	0.13 ± 0.05	5.24 ± 1.63	0.86/0.21
	+ XE991	10	0.23 ± 0.08	5.33 ± 0.81	0.28 ± 0.13	5.21 ± 0.87	0.03/0.72
+ low K ⁺ aCSF	8	0.24 ± 0.13	6.04 ± 1.24	0.25 ± 0.16	5.76 ± 1.91	0.69/0.51	

Data presented as means ± standard deviations. *p*-values below 0.05 are indicated in boldface type.

RESULTS


To characterize the ionic mechanisms underlying 2A-receptor-mediated serotonergic excitation in the cerebral cortex, we made electrical recordings from labeled callosal/commissural (COM) projection neurons in layer 5 of the mouse mPFC. Because 2A-mediated excitation of COM neurons requires coincident exogenous excitatory drive (Stephens et al., 2014), 5-HT was focally applied during periods of sustained action potential generation (~ 5 Hz) produced by somatic DC-current injection (**Figure 1A**). Under these conditions, 5-HT (1 s at ~ 15 psi) generated robust increases in instantaneous spike frequency (ISF) that persisted for many seconds. In these same neurons, 5-HT delivered at resting membrane potentials had little, if any, effect (**Figure 1A**). Across all COM neurons recorded with our normal whole-cell pipette solution ($n = 54$), the magnitude of serotonergic excitation, as measured by the peak increase in ISF following 5-HT application, was negatively correlated with baseline firing rate ($p = 0.004$, $R = -0.39$, **Figure 1B**), suggesting a ceiling effect of serotonergic excitation at high baseline firing rates. Thus, for each experimental manipulation, we adjusted current intensities to maintain similar baseline firing frequencies across experimental conditions (**Table 2**).

Cholinergic Activation of L5PNs Occludes Serotonergic Excitation


Since serotonergic responses are qualitatively similar to cholinergic responses mediated by M1 muscarinic ACh receptors (Araneda and Andrade, 1991; Haj-Dahmane and Andrade, 1999; Gullledge et al., 2009), we first tested whether 5-HT and ACh share common intracellular signaling cascades by measuring responses to 5-HT before and after tonically activating muscarinic ACh receptors with carbachol (50–100 μM , bath applied). In baseline conditions, focal application of 5-HT produced mean increases in peak ISF of $117 \pm 23\%$ ($n = 9$; $p < 0.001$, paired Student's *t*-test), with response integrals being $168 \pm 34 \text{ Hz}\cdot\text{s}$ (**Figure 2A**). Bath application of carbachol depolarized COM neurons (by $6 \pm 2 \text{ mV}$) and lowered the current necessary to evoke sustained action potential generation by $54 \pm 16\%$, from $138 \pm 24 \text{ pA}$ to $71 \pm 26 \text{ pA}$ ($n = 6$ of 9 neurons). Carbachol induced spontaneous firing in 3 of 9 neurons. Across all 9 neurons, bath-applied carbachol fully occluded serotonergic excitation, as 5-HT failed to enhance action potential frequency during cholinergic stimulation ($n = 9$; **Figure 2**). Instead, 5-HT application in the presence of carbachol reduced the firing frequency of COM neurons (peak change in ISF = $-40 \pm 10\%$; $n = 9$, $p = 0.003$, paired Student's *t*-test; **Figures 2B,C**), resulting in negative response integrals ($-103 \pm 39 \text{ Hz}\cdot\text{s}$). These results confirm significant overlap of G_q -signaling cascades downstream of M1 and 2A receptors in COM neurons.

5-HT Generates Afterdepolarizations in COM Neurons

A prominent feature of muscarinic excitation of pyramidal neurons is the generation of afterdepolarizations (ADPs) that occur after pairing M1 receptor activation with brief



suprathreshold current injections (Andrade, 1991; Haj-Dahmane and Andrade, 1998; Gullledge et al., 2009; Dasari et al., 2013). Similarly, 5-HT acting via 2A receptors can initiate ADPs in neocortical pyramidal neurons (Araneda and Andrade, 1991; Spain, 1994; Zhang and Arseneault, 2005). We confirmed that 5-HT promotes ADPs in COM neurons by evoking trains of ten action potentials using brief (2 ms) high-amplitude (3 nA) current steps presented at 25 Hz (**Figure 3A**). On some trials, 5-HT was focally applied 1 s before current pulses, and peak depolarization relative to the resting membrane potential (RMP) was measured in the first 1 s following the spike train. Pairing 5-HT with current pulses generated small ADPs with mean



amplitudes of 1.6 ± 0.2 mV ($n = 16$; $p < 0.001$, paired Student's *t*-test; **Figure 3B**). Serotonergic ADPs were absent when the calcium chelating agent BAPTA (10 mM) was included in the patch-pipette solution ($n = 6$; **Figure 3B**), confirming that they are gated by intracellular calcium signaling (Spain, 1994; Haj-Dahmane and Andrade, 1998; Zhang and Arsenault, 2005; Dasari et al., 2013). These results suggest that the calcium-dependent and G_q -triggered ADP may represent a common ionic mechanism contributing to serotonergic and cholinergic excitation in COM neurons.

Role of Calcium in Serotonergic Excitation of COM Neurons

If the ADP current is the major contributor to serotonergic excitation in COM neurons, chelation of intracellular calcium with BAPTA should reduce or eliminate excitatory responses to 5-HT. Instead, addition of BAPTA (10 mM) to the pipette solution ($n = 13$) enhanced serotonergic excitation, with mean 5-HT response amplitudes being twice as large as in COM neurons patched with our standard intracellular solution (**Figure 4**). In control neurons ($n = 16$), peak increases in ISF were $113 \pm 13\%$ above baseline values, while in BAPTA-filled neurons ($n = 13$) peak ISF increases were $226 \pm 23\%$ above baseline values ($p < 0.001$ vs. control; Student's *t*-test). While response integrals in these BAPTA-filled COM neurons (249 ± 56 Hz \cdot s) were not significantly larger than those in

control neurons (176 ± 26 Hz \cdot s; $p = 0.22$; **Figure 4C**), in three additional independent experimental groups (see below) we observed significant enhancement of both amplitudes and integrals of 5-HT responses in BAPTA-filled neurons relative to controls. These findings demonstrate that serotonergic excitation, as a whole, is not calcium-dependent, and suggest that the calcium-dependent ADP may not represent the major ionic effector mediating 5-HT responses in COM neurons. Instead, the enhancement of responses in the presence of intracellular BAPTA suggests that intracellular calcium acts as a negative regulator for additional ionic effectors contributing to 2A-mediated excitation.

If activity- and/or 5-HT-driven calcium entry negatively regulates serotonergic excitation of COM neurons, removal of extracellular calcium should enhance 5-HT responses in a manner similar to inclusion of intracellular BAPTA. To test this possibility, we measured responses to 5-HT in COM neurons in baseline conditions and after replacing extracellular calcium with magnesium (see Materials and Methods; **Figure 5**). Successful elimination of extracellular calcium was verified in a subset of neurons ($n = 4$) by measuring electrically evoked EPSPs before, during, and after calcium replacement (**Figure 5A**). While removal of extracellular calcium eliminated synaptic responses, it did not affect the magnitude or integrals of serotonergic responses ($n = 9$; **Figures 5B–D**). Peak increases in ISFs were $78 \pm 11\%$ and $75 \pm 12\%$ above initial firing frequencies in baseline and calcium-free conditions, respectively




FIGURE 3 | 5-HT induces a calcium-dependent afterdepolarization (ADP) in COM neurons. **(A)** Voltage traces from a COM neuron experiencing a train of ten high-amplitude (3 nA, 2 ms, 25 Hz) current injections in baseline conditions (blue) and following focal application of 5-HT (red). Action potentials are truncated. Black arrows indicate current step applications. **(B)** Average voltage responses following spike trains paired with 5-HT, resampled to 50 Hz and plotted as mean \pm SEM, for 16 neurons in baseline conditions (blue) and after 5-HT (red). Gray symbols show responses following 5-HT application in a different group of COM neurons filled with 10 mM BAPTA ($n = 6$).

($p = 0.83$, paired Student's t -test), while response integrals for the two conditions were 141 ± 29 Hz \bullet s and 164 ± 44 Hz \bullet s, respectively ($p = 0.58$). This consistency in 5-HT responses across experimental conditions occurred in spite of a small (1.5 ± 0.6 Hz; $p = 0.043$, paired Student's t -test) increase in baseline firing frequency that would otherwise predict a $22 \pm 9\%$ decrease in peak excitation (see **Figure 1** and **Table 2**). Thus, with normal intracellular calcium signaling intact, removal of extracellular calcium had little, if any, effect on 5-HT response amplitudes or integrals.


The lack of effect of extracellular calcium removal on 5-HT responses likely reflects competing cellular processes. For instance, even as removal of calcium may enhance some 5-HT-gated conductances, it will also eliminate the calcium-dependent ADP conductance and the calcium component of any calcium-permeable cation conductances contributing to excitation. To further explore the impact of extracellular calcium on serotonergic signaling in COM neurons, we filled additional neurons with BAPTA (10 mM) and measured serotonergic responses before and after replacement of extracellular calcium with magnesium (**Figures 5E,F**). As in our earlier experiments, the presence of intracellular BAPTA led to robust serotonergic excitation in baseline

(i.e., normal extracellular calcium) conditions, with peak increases in ISF being $213 \pm 36\%$ above initial firing rates, and response integrals being 163 ± 34 Hz \bullet s ($n = 12$). Subsequent removal of extracellular calcium greatly diminished both peak excitation (to $105 \pm 11\%$; $p = 0.007$, paired Student's t -test) and response integrals (to 63 ± 16 Hz \bullet s; $p = 0.004$, **Figure 5G**). Because intracellular calcium signaling was blocked with BAPTA throughout these experiments, the reduction in response amplitudes and integrals observed after removal of extracellular calcium cannot be explained by effects on the ADP or other calcium-dependent conductances. Instead, these results point to involvement of a calcium-permeable cation conductance that, under normal conditions, is suppressed by intracellular calcium, but which is enhanced after chelation of intracellular calcium. Because responses in the presence of both intracellular BAPTA (which itself doubles 5-HT response amplitudes) and calcium-free aCSF (which in the absence of BAPTA has no effect on 5-HT responses amplitudes) results in responses comparable in size to those in control conditions (i.e., with normal intracellular and extracellular calcium levels), this calcium-sensitive calcium conductance appears to be balanced with the ADP and other calcium-dependent conductances gated by 2A receptors.

Role of M-current in Serotonergic Excitation

A classic ionic effector associated with G_q -coupled receptors that may exhibit calcium-sensitivity is the M- ("muscarinic") current mediated by voltage-dependent K_V7 (KCNQ) potassium channels (Delmas and Brown, 2005). Does M-current contribute to serotonergic excitation of COM neurons? If so, does intracellular calcium reduce the contribution of M-current to excitatory responses? To test the role of M-current suppression in generating serotonergic excitation, we measured 5-HT responses in COM neurons before and after bath applying the selective M-current blocker XE991 (10 μ M; **Figures 6A,B**). XE991 depolarized RMPs, from -80 ± 2 to -76 ± 1 mV ($n = 10$; $p = 0.014$, paired Student's t -test), and increased input resistances (R_N), from 155 ± 14 to 185 ± 16 M Ω ($p = 0.004$, paired Student's t -test), indicating that the M-current contributes to the resting membrane conductance of layer 5 COM neurons in the mouse mPFC. XE991 also reduced both the magnitude (by $49 \pm 5\%$) and integral (by $56 \pm 15\%$) of serotonergic excitation. Peak increases in ISF in response to 5-HT dropped from $94 \pm 15\%$ in baseline conditions to $50 \pm 11\%$ after application of XE991 ($p = 0.001$, paired Student's t -test), while response integrals dropped from 132 ± 23 Hz \bullet s to 48 ± 16 Hz \bullet s ($p = 0.009$, **Figures 6E,F**). The substantial reduction in response amplitudes and integrals in the presence of XE991 suggests that suppression of the M-current is a key mediator of serotonergic excitation in COM neurons.

If calcium sensitivity of M-current (see, for instance, Selyanko and Brown, 1996) accounts for the enhanced serotonergic excitation observed after intracellular calcium chelation, 5-HT responses after blockade of K_V7 channels should be similar in both control and BAPTA-filled neurons. We tested this prediction




by recording from a second group of neurons ($n = 10$) filled with 10 mM BAPTA (Figures 6C,D). Consistent with our earlier experiments using intracellular BAPTA, baseline responses to 5-HT were much larger in amplitude (mean increase in ISF was $230 \pm 22\%$; $p < 0.001$ vs. neurons lacking BAPTA; Student's t -test, $n = 10$ per group), and integral (319 ± 107 Hz \bullet s; $p = 0.10$) in BAPTA-filled neurons relative to responses in neurons with intracellular calcium signaling intact. However, intracellular BAPTA did not enhance the impact of K_V7 blockade with 10 μ M XE991. With BAPTA present, serotonergic excitation in XE991 ($+164 \pm 30\%$ peak increase in ISF) remained three times larger than in control neurons treated with XE991 ($p = 0.002$, Student's t -test; Figures 6E,F). However, mean integrals (122 ± 43 Hz \bullet s), while generally larger than in control neurons treated with XE991, were not statistically different in magnitude ($p = 0.13$). Overall, peak response amplitudes were reduced by $29 \pm 9\%$ ($p = 0.022$ vs. baseline responses; paired Student's t -test) and integrals by $57 \pm 11\%$ ($p = 0.024$ vs. baseline responses) in BAPTA-filled neurons. Thus, the XE991-sensitive portions of serotonin responses in control and BAPTA recording conditions were similar, suggesting that M-current alone does not account for the calcium-sensitivity of 5-HT excitation in COM neurons.

The results above suggest that 5-HT acts via at least three distinct mechanisms (K_V7 suppression, the ADP conductance, and a calcium-sensitive calcium conductance) to enhance the excitability of COM neurons. To test whether M-current is the dominant potassium conductance contributing to serotonergic excitation, we enhanced the driving force for potassium by lowering the external potassium concentration ($[K^+]_o$)

six-fold to 0.5 mM (replaced with equimolar sodium; Figure 7). By increasing the driving force for potassium, this manipulation will enhance the impact of M-current suppression by 5-HT, but will also act to reduce the net current through potassium-permeable non-specific cation conductances. In neurons recorded with control intracellular solution, lowering $[K^+]_o$ revealed a brief inhibition occurring immediately after 5-HT application that was absent in control conditions (Figures 7A,C); these inhibitory responses are likely G_q -driven hyperpolarizations (mediated by SK-type potassium channels) that occur regularly in pyramidal neurons following M1 muscarinic receptor activation (Gulledge et al., 2009), but which are only rarely observed in response to 5-HT in control conditions. Lowering $[K^+]_o$ enhanced this early potassium conductance, and reduced the magnitude of serotonergic excitation by $31 \pm 9\%$ ($n = 10$, paired). In control conditions (e.g., 3 mM $[K^+]_o$), 5-HT generated peak responses of $82 \pm 14\%$ with integrals of 157 ± 44 Hz \bullet s. After reducing extracellular potassium to 0.5 mM, peak excitation was $61 \pm 15\%$ ($p = 0.003$ relative to control conditions) with integrals of 117 ± 47 Hz \bullet s ($p = 0.057$, Figure 7D). Because the larger driving force for potassium is expected to increase 5-HT excitation by enhancing the contribution of M-current suppression, the observed reductions in response magnitudes and integrals suggest the participation of potassium-permeable non-specific cation conductances, such as the ADP conductance (Haj-Dahmane and Andrade, 1998).


To further test the impact of potassium manipulations on 5-HT responses in the absence of the non-specific cation conductance underlying the ADP, and calcium-activated



potassium conductances, we repeated experiments in COM neurons filled with 10 mM BAPTA (Figures 7B,C). With intracellular calcium chelated, baseline responses were again large, with $254 \pm 34\%$ increases in ISF and 295 ± 61 Hz \cdot s response integrals ($n = 8$). When $[K^+]_o$ was lowered to 0.5 mM, response amplitudes were reduced by $42 \pm 7\%$ (to $156 \pm 36\%$ above baseline firing rates; $p < 0.001$; paired Student's *t*-test), while response integrals were decreased by $63 \pm 12\%$ (to 114 ± 55 Hz \cdot s; $p = 0.029$; Figure 7D). The percent reductions in both amplitudes ($p = 0.342$) and integrals ($p = 0.559$) after lowering $[K^+]_o$ in BAPTA filled neurons were similar to those observed in control neurons, suggesting that the calcium-sensitive conductance enhanced by intracellular BAPTA must also be a potassium-permeable non-specific cation conductance.

DISCUSSION

Activation of G_q -coupled receptors, including serotonergic 2A and muscarinic M1 receptors, enhances the intrinsic excitability of many neurons throughout the nervous system. As common and robust as these responses are, it is surprising that there is not yet consensus on their underlying mechanisms. Data regarding cortical 2A signaling are particularly limited, in part due to the selective expression of these receptors in subpopulations of cortical neurons in adult animals (Weber and Andrade, 2010; Avesar and Gullledge, 2012). Therefore, much of our understanding of 2A receptor signaling has been inferred from studies of the more ubiquitous G_q -coupled muscarinic ACh receptors. Here, we selectively targeted 2A-responsive COM neurons to probe the mechanisms underlying serotonergic




excitation. Our results confirm that 2A and M1 receptors share downstream signaling pathways, and point to at least three distinct ionic effectors contributing to serotonergic excitation: suppression of K_v7 conductances, activation of a non-specific cation channel that is both calcium-sensitive and calcium-permeable, and activation of the calcium-dependent non-specific cation conductance responsible for the ADP (**Figure 8**).

Ionic Conductances Contributing to Serotonergic Excitation of COM Neurons

Both M-current suppression (Caulfield et al., 1994) and activation of non-specific cation conductances (Haj-Dahmane and Andrade, 1996) have been proposed as mechanisms for G_q -mediated excitation. We found that under control conditions, with intracellular calcium signaling intact, suppression of

M-current accounted for about half of the total serotonergic response. While K_v7 channels are reported to be calcium-sensitive (Selyanko and Brown, 1996), the impact of M-current suppression was similar in control neurons and those filled with BAPTA, with responses to 5-HT in the presence of both BAPTA and XE991 remaining much larger than responses measured in control conditions with intracellular calcium signaling and M-current intact. This suggests that the calcium-sensitivity of K_v7 channels cannot account for the larger 5-HT responses observed after chelation of intracellular calcium.


We also confirmed that 5-HT generates calcium-dependent ADPs in COM neurons. However, quantifying the relative contribution of the ADP conductance to enhanced action potential generation is problematic, as the ion channel mediating the ADP remains a matter of debate



(Kuzmiski and MacVicar, 2001; Yan et al., 2009; Dasari et al., 2013; Lei et al., 2014), and attempts to block the conductance with intracellular BAPTA, or by removal of extracellular calcium, paradoxically enhanced the magnitude of serotonergic excitation. Indeed, in four independent experimental groups, baseline responses to 5-HT in the presence of intracellular BAPTA were twice as large as responses in control neurons with normal intracellular calcium signaling. These findings demonstrate that the ADP conductance is not required for serotonergic excitation, and that the net effect of normal levels of intracellular calcium is moderation, rather than amplification, of serotonergic excitation.

Two observations suggest that the calcium-sensitivity of 5-HT responses reflects a non-specific cation conductance that itself is calcium permeable. First, the involvement of non-specific cation conductances is strongly suggested by the effect

of reducing extracellular potassium. While this manipulation enhances selective potassium currents (e.g., the M-current), it also lowers the equilibrium potential for potassium-permeable non-specific cation conductances. Thus, the reduction in serotonergic excitation observed after lowering extracellular potassium suggests the involvement of non-specific cation conductances, such as the ADP conductance (Haj-Dahmane and Andrade, 1998). Since we continued to observe a reduction in response amplitudes after ADP currents were blocked by BAPTA, the remaining calcium-sensitive calcium conductance, which was amplified by chelation of intracellular calcium, is likely also a potassium-permeable non-specific cation conductance. Second, we found that the effect of removing extracellular calcium was dependent on the baseline state of intracellular calcium signaling within neurons. In control conditions, with calcium-signaling intact, removal of extracellular calcium had no impact



on 5-HT response amplitudes or integrals. Yet, when intracellular calcium signaling was blocked with BAPTA, and baseline responses to 5-HT were twice as large as in control neurons, removal of extracellular calcium greatly reduced the amplitudes and integrals of 5-HT responses, such that they became comparable in size to control 5-HT responses observed in the absence of BAPTA. These findings suggest the involvement of a calcium-permeable non-specific cation conductance that is itself suppressed by intracellular calcium (Figure 8). Under normal conditions, with intracellular calcium signaling intact, removal of extracellular calcium is expected to have two opposing effects: it will eliminate a charge carrier for the calcium-permeable non-specific cation conductance, thereby hyperpolarizing the effective equilibrium potential for the conductance, but will also enhance the maximum conductance by relieving the channel from negative regulation by intracellular calcium. Our results suggest these two competing influences are fairly balanced with the calcium-dependent ADP conductance, such that calcium-removal in control conditions has little impact on serotonergic responses. However, when the conductance is maximized by chelating intracellular calcium with BAPTA, subsequent removal of extracellular calcium acts only to reduce net current, such that 5-HT responses become smaller and equivalent to responses observed in control conditions.

These findings regarding calcium regulation of serotonergic excitation are most comparable to those of Magistretti et al. (2004), who found G_q -mediated cholinergic responses in layer 2 pyramidal neurons in the entorhinal cortex to be bidirectionally influenced by calcium influx. They found that modest increases in intracellular calcium enhanced net inward current (attributed to a calcium-dependent ADP-like conductance), while larger increases in intracellular calcium, including normal levels of calcium accumulation during periods of action potential generation, acted to suppress cholinergic excitation (Magistretti et al., 2004). While Magistretti et al. (2004) concluded that their results reflected bidirectional calcium regulation of the

ADP conductance, results from our experiments using BAPTA and calcium-free aCSF dissociate the calcium-activated ADP conductance from the additional calcium-sensitive non-specific cation conductance. Co-activation of two non-specific cation conductances with opposing calcium regulation may contribute to the robustness of G_q -mediated excitation by maintaining a consistent net current amplitude independent of intracellular calcium levels, while also promoting calcium-influx when intracellular calcium levels are low. This negative feedback of intracellular calcium on the calcium-permeable conductance should act to stabilize intracellular calcium levels, and could provide a mechanism for calcium store refilling after G_q receptor signaling (Dasari et al., 2017).

G_q -Signaling in Cortical Projection Neurons

The ionic mechanisms contributing to serotonergic excitation of COM neurons appear to be conserved across cortical neuron subtypes and transmitter signaling systems. We have recently found that phasic activation of the G_q -coupled M1 muscarinic ACh receptor enhances the excitability of pyramidal neurons in layer 5 of the mouse mPFC, an enhancement more robust and longer lasting in corticopontine (CPn) neurons relative to COM neurons (Baker et al., 2018; see also Dembrow et al., 2010). Further investigation of the mechanisms underlying this robust excitation in CPn neurons revealed that the M1 mAChR pathway involves the same combination of ionic effectors (M-current suppression, the ADP conductance, and a calcium-sensitive and calcium-permeable non-specific cation conductance) as described here in relation to serotonergic 2A signaling in COM neurons.

Some notable differences do exist, however, in G_q -mediated excitation of COM and CPn neurons by 5-HT and ACh, respectively. For instance, although stimulation of muscarinic ACh receptors robustly generated transient SK-channel mediated inhibition in both COM and CPn neurons (Baker et al., 2018),

in normal conditions (3 mM $[K^+]_o$) we rarely observed 5-HT-triggered SK-dependent inhibition in mouse COM neurons, even as ACh and 5-HT generate similar increases in ISF (~100% above baseline levels) in these same neurons. Even after enhancing the driving force for potassium channels with lowered extracellular potassium, SK-like responses to 5-HT were observed in only 7 of 10 neurons, and had shorter durations (0.7 ± 0.2 s) than SK responses generated by ACh in either COM (~1.5 s) or CPn (~1 s) neurons (Baker et al., 2018). This may reflect differential subcellular localization of M1 muscarinic receptors relative to serotonergic 2A receptors in COM neurons, or subtle differences in G_q -coupling to second messenger systems. Another difference is the relative contribution of ionic effectors in generating excitatory responses. In CPn neurons, suppression of M-current accounted for only ~20% of total cholinergic excitation, while chelation of intracellular calcium enhanced responses by only ~25% (Baker et al., 2018). This suggests that the ADP conductance may play a larger role in mediating cholinergic excitation in CPn neurons relative to COM neurons. Thus, M-current suppression and the calcium-sensitive non-specific cation conductance appear more important for serotonergic excitation of COM neurons, relative to cholinergic excitation of CPn neurons.

In addition to 2A and M1 receptors, neocortical pyramidal neurons express a wide variety of aminergic, glutamatergic, purinergic, and peptidergic receptors that couple to G_q alpha subunits. Our results reveal multiple ionic effectors contributing to serotonergic 2A signal transduction in COM neurons, and point to significant overlap of 2A signaling with M1 muscarinic receptors in these same neurons (see also Baker et al., 2018), suggesting that a common set of ionic effectors is targeted by a wide variety of transmitter systems. While parallel G_q signaling cascades enhance the fidelity and robustness of excitatory responses, targeted expression of G_q -coupled receptors and/or downstream ionic effectors in specific cellular compartments may allow fine-tuning of local responses to neuromodulatory input. Expression of additional, non- G_q -coupled receptors (e.g.,

5-HT_{1A} receptors), and the potential formation of heteromeric receptor complexes (McCormick and Botta, 2017), further adds to the diversity of G_q signaling. A detailed understanding of G_q signaling in the cerebral cortex will clarify our understanding of how neuromodulators facilitate cognition, and, especially in regard to 2A receptors, may facilitate development of pharmacological interventions for psychiatric disorders that are more efficacious and have fewer side effects than approaches that target G-protein-coupled receptors directly.

ETHICS STATEMENT

This study was carried out in accordance with the recommendations of the Association for Assessment and Accreditation of Laboratory Animal Care International (AAALAC). The protocol was approved by the Institutional Animal Care and Use Committee of Dartmouth College.

AUTHOR CONTRIBUTIONS

AG and ES designed the research. AG, AB, and ES conducted the research and analyzed the data. AG, ES, and AB wrote the manuscript.

FUNDING

This work was supported by a grant from the National Institute for Mental Health (R01 MH099054; AG) and a Frank and Myra Weiser Scholar Award, AG.

ACKNOWLEDGMENTS

The authors thank Saiko Ikeda for technical assistance.

REFERENCES

- Almada, R. C., Coimbra, N. C., and Brandao, M. L. (2015). Medial prefrontal cortex serotonergic and GABAergic mechanisms modulate the expression of contextual fear: intratelencephalic pathways and differential involvement of cortical subregions. *Neuroscience* 284, 988–997. doi: 10.1016/j.neuroscience.2014.11.001
- Andrade, R. (1991). Cell excitation enhances muscarinic cholinergic responses in rat association cortex. *Brain Res.* 548, 81–93. doi: 10.1016/0006-8993(91)91109-E
- Andrade, R., and Nicoll, R. A. (1987). Pharmacologically distinct actions of serotonin on single pyramidal neurones of the rat hippocampus recorded in vitro. *J. Physiol.* 394, 99–124. doi: 10.1113/jphysiol.1987.sp016862
- Araneda, R., and Andrade, R. (1991). 5-Hydroxytryptamine₂ and 5-hydroxytryptamine_{1A} receptors mediate opposing responses on membrane excitability in rat association cortex. *Neuroscience* 40, 399–412. doi: 10.1016/0306-4522(91)90128-B
- Avesar, D., and Gullledge, A. T. (2012). Selective serotonergic excitation of callosal projection neurons. *Front. Neural Circuits* 6:12. doi: 10.3389/fncir.2012.00012
- Azmitia, E. C., Gannon, P. J., Kheck, N. M., and Whitaker-Azmitia, P. M. (1996). Cellular localization of the 5-HT_{1A} receptor in primate brain neurons and glial cells. *Neuropsychopharmacology* 14, 35–46. doi: 10.1016/S0893-133X(96)80057-1
- Baker, A. L., O'Toole, R. J., and Gullledge, A. T. (2018). Preferential cholinergic excitation of corticopontine neurons. *J. Physiol.* (in Press). doi: 10.1113/JP275194
- Caulfield, M. P., Jones, S., Vallis, Y., Buckley, N. J., Kim, G. D., Milligan, G., et al. (1994). Muscarinic M-current inhibition via G alpha q/11 and alpha-adrenoceptor inhibition of Ca²⁺ current via G alpha o in rat sympathetic neurones. *J. Physiol.* 477, 415–422. doi: 10.1113/jphysiol.1994.sp020203
- Clarke, H. F., Dalley, J. W., Crofts, H. S., Robbins, T. W., and Roberts, A. C. (2004). Cognitive inflexibility after prefrontal serotonin depletion. *Science* 304, 878–880. doi: 10.1126/science.1094987
- Clarke, H. F., Walker, S. C., Crofts, H. S., Dalley, J. W., Robbins, T. W., and Roberts, A. C. (2005). Prefrontal serotonin depletion affects reversal learning but not attentional set shifting. *J. Neurosci.* 25, 532–538. doi: 10.1523/JNEUROSCI.3690-04.2005
- Colino, A., and Halliwell, J. V. (1987). Differential modulation of three separate K-conductances in hippocampal CA1 neurons by serotonin. *Nature* 328, 73–77. doi: 10.1038/328073a0
- Cruz, D. A., Eggen, S. M., Azmitia, E. C., and Lewis, D. A. (2004). Serotonin_{1A} receptors at the axon initial segment of prefrontal pyramidal neurons in schizophrenia. *Am. J. Psychiatry* 161, 739–742. doi: 10.1176/appi.ajp.161.4.739

- Czyrak, A., Czepiel, K., Mackowiak, M., Chocyk, A., and Wedzony, K. (2003). Serotonin 5-HT_{1A} receptors might control the output of cortical glutamatergic neurons in rat cingulate cortex. *Brain Res.* 989, 42–51. doi: 10.1016/S0006-8993(03)03352-3
- Dasari, S., Abramowitz, J., Birnbaumer, L., and Gullledge, A. T. (2013). Do canonical transient receptor potential channels mediate cholinergic excitation of cortical pyramidal neurons? *Neuroreport* 24, 550–554. doi: 10.1097/WNR.0b013e3283621344
- Dasari, S., and Gullledge, A. T. (2011). M1 and M4 receptors modulate hippocampal pyramidal neurons. *J. Neurophysiol.* 105, 779–792. doi: 10.1152/jn.00686.2010
- Dasari, S., Hill, C., and Gullledge, A. T. (2017). A unifying hypothesis for M1 muscarinic receptor signalling in pyramidal neurons. *J. Physiol.* 595, 1711–1723. doi: 10.1113/JP273627
- Davies, M. F., Deisz, R. A., Prince, D. A., and Peroutka, S. J. (1987). Two distinct effects of 5-hydroxytryptamine on single cortical neurons. *Brain Res.* 423, 347–352. doi: 10.1016/0006-8993(87)90861-4
- DeFelipe, J., Arellano, J. I., Gomez, A., Azmitia, E. C., and Munoz, A. (2001). Pyramidal cell axons show a local specialization for GABA and 5-HT inputs in monkey and human cerebral cortex. *J. Comp. Neurol.* 433, 148–155. doi: 10.1002/cne.1132
- Delmas, P., and Brown, D. A. (2005). Pathways modulating neural KCNQ/M (Kv7) potassium channels. *Nat. Rev. Neurosci.* 6, 850–862. doi: 10.1038/nrn1785
- Dembrow, N. C., Chitwood, R. A., and Johnston, D. (2010). Projection-specific neuromodulation of medial prefrontal cortex neurons. *J. Neurosci.* 30, 16922–16937. doi: 10.1523/JNEUROSCI.3644-10.2010
- Egorov, A. V., Angelova, P. R., Heinemann, U., and Müller, W. (2003). Ca²⁺-independent muscarinic excitation of rat medial entorhinal cortex layer V neurons. *Eur. J. Neurosci.* 18, 3343–3351. doi: 10.1111/j.1460-9568.2003.03050.x
- Goodfellow, N. M., Benekareddy, M., Vaidya, V. A., and Lambe, E. K. (2009). Layer II/III of the prefrontal cortex: Inhibition by the serotonin 5-HT_{1A} receptor in development and stress. *J. Neurosci.* 29, 10094–10103. doi: 10.1523/JNEUROSCI.1960-09.2009
- Gullledge, A. T., Bucci, D. J., Zhang, S. S., Matsui, M., and Yeh, H. H. (2009). M1 receptors mediate cholinergic modulation of excitability in neocortical pyramidal neurons. *J. Neurosci.* 29, 9888–9902. doi: 10.1523/JNEUROSCI.1366-09.2009
- Haj-Dahmane, S., and Andrade, R. (1996). Muscarinic activation of a voltage-dependent cation nonselective current in rat association cortex. *J. Neurosci.* 16, 3848–3861.
- Haj-Dahmane, S., and Andrade, R. (1998). Ionic mechanism of the slow afterdepolarization induced by muscarinic receptor activation in rat prefrontal cortex. *J. Neurophysiol.* 80, 1197–1210. doi: 10.1152/jn.1998.80.3.1197
- Haj-Dahmane, S., and Andrade, R. (1999). Muscarinic receptors regulate two different calcium-dependent non-selective cation currents in rat prefrontal cortex. *Eur. J. Neurosci.* 11, 1973–1980. doi: 10.1046/j.1460-9568.1999.00612.x
- Harrison, A. A., Everitt, B. J., and Robbins, T. W. (1997). Central 5-HT depletion enhances impulsive responding without affecting the accuracy of attentional performance: interactions with dopaminergic mechanisms. *Psychopharmacology (Berl.)* 133, 329–342. doi: 10.1007/s002130050410
- Jakab, R. L., and Goldman-Rakic, P. S. (1998). 5-Hydroxytryptamine_{2A} serotonin receptors in the primate cerebral cortex: possible site of action of hallucinogenic and antipsychotic drugs in pyramidal cell apical dendrites. *Proc. Natl. Acad. Sci. U.S.A.* 95, 735–740. doi: 10.1073/pnas.95.2.735
- Krnjevic, K., Pumain, R., and Renaud, L. (1971). The mechanism of excitation by acetylcholine in the cerebral cortex. *J. Physiol.* 215, 247–268. doi: 10.1113/jphysiol.1971.sp009467
- Kuzmiski, J. B., and MacVicar, B. A. (2001). Cyclic nucleotide-gated channels contribute to the cholinergic plateau potential in hippocampal CA1 pyramidal neurons. *J. Neurosci.* 21, 8707–8714.
- Lei, Y. T., Thuault, S. J., Launay, P., Margolskee, R. F., Kandel, E. R., and Siegelbaum, S. A. (2014). Differential contribution of TRPM4 and TRPM5 nonselective cation channels to the slow afterdepolarization in mouse prefrontal cortex neurons. *Front. Cell. Neurosci.* 8:267. doi: 10.3389/fncel.2014.00267
- Leon, L. A., Castro-Gomes, V., Zarate-Guerrero, S., Corredor, K., Mello Cruz, A. P., Brandao, M. L., et al. (2017). Behavioral effects of systemic, infralimbic and prelimbic injections of a serotonin 5-HT_{2A} antagonist in caracara high- and low-conditioned freezing rats. *Front. Behav. Neurosci.* 11:117. doi: 10.3389/fnbeh.2017.00117
- Liy-Salmeron, G., and Meneses, A. (2008). Effects of 5-HT drugs in prefrontal cortex during memory formation and the ketamine amnesia-model. *Hippocampus* 18, 965–974. doi: 10.1002/hipo.20459
- Luscher, C., Jan, L. Y., Stoffel, M., Malenka, R. C., and Nicoll, R. A. (1997). G protein-coupled inwardly rectifying K⁺ channels (GIRKs) mediate postsynaptic but not presynaptic transmitter actions in hippocampal neurons. *Neuron* 19, 687–695. doi: 10.1016/S0896-6273(00)80381-5
- Magistretti, J., Ma, L., Shalinsky, M. H., Lin, W., Klink, R., and Alonso, A. (2004). Spike patterning by Ca²⁺-dependent regulation of a muscarinic cation current in entorhinal cortex layer II neurons. *J. Neurophysiol.* 92, 1644–1657. doi: 10.1152/jn.00036.2004
- Marinho, A. L., Vila-Verde, C., Fogaca, M. V., and Guimaraes, F. S. (2015). Effects of intra-infralimbic prefrontal cortex injections of cannabidiol in the modulation of emotional behaviors in rats: contribution of 5HT(1)A receptors and stressful experiences. *Behav. Brain Res.* 286, 49–56. doi: 10.1016/j.bbr.2015.02.023
- McCormick, D. A., and Williamson, A. (1989). Convergence and divergence of neurotransmitter action in human cerebral cortex. *Proc. Natl. Acad. Sci. U.S.A.* 86, 8098–8102. doi: 10.1073/pnas.86.20.8098
- McCormick, P. J., and Botta, J. (2017). “Heteromers form novel signaling complexes,” in *G-Protein-Coupled Receptor Dimers*, eds K. Herrick-Davis, G. Milligan, and G. Di Giovanni (Cham: Springer), 467–475. doi: 10.1007/978-3-319-60174-8_19
- Miner, L. A., Backstrom, J. R., Sanders-Bush, E., and Sesack, S. R. (2003). Ultrastructural localization of serotonin_{2A} receptors in the middle layers of the rat prelimbic prefrontal cortex. *Neuroscience* 116, 107–117. doi: 10.1016/S0306-4522(02)00580-8
- Paxinos, G., and Franklin, K. B. J. (2004). *The Mouse Brain in Stereotaxic Coordinates*. San Diego, CA: Elsevier.
- Schwindt, P. C., Spain, W. J., Foehring, R. C., Chubb, M. C., and Crill, W. E. (1988). Slow conductances in neurons from cat sensorimotor cortex in vitro and their role in slow excitability changes. *J. Neurophysiol.* 59, 450–467. doi: 10.1152/jn.1988.59.2.450
- Selyanko, A. A., and Brown, D. A. (1996). Intracellular calcium directly inhibits potassium M channels in excised membrane patches from rat sympathetic neurons. *Neuron* 16, 151–162. doi: 10.1016/S0896-6273(00)80032-X
- Shalinsky, M. H., Magistretti, J., Ma, L., and Alonso, A. A. (2002). Muscarinic activation of a cation current and associated current noise in entorhinal-cortex layer-II neurons. *J. Neurophysiol.* 88, 1197–1211. doi: 10.1152/jn.2002.88.3.1197
- Spain, W. J. (1994). Serotonin has different effects on two classes of Betz cells from the cat. *J. Neurophysiol.* 72, 1925–1937. doi: 10.1152/jn.1994.72.4.1925
- Stephens, E. K., Avesar, D., and Gullledge, A. T. (2014). Activity-dependent serotonergic excitation of callosal projection neurons in the mouse prefrontal cortex. *Front. Neural Circuits* 8:97. doi: 10.3389/fncir.2014.00097
- Tanaka, E., and North, R. A. (1993). Actions of 5-hydroxytryptamine on neurons of the rat cingulate cortex. *J. Neurophysiol.* 69, 1749–1757. doi: 10.1152/jn.1993.69.5.1749
- Villalobos, C., Beique, J. C., Gingrich, J. A., and Andrade, R. (2005). Serotonergic regulation of calcium-activated potassium currents in rodent prefrontal cortex. *Eur. J. Neurosci.* 22, 1120–1126. doi: 10.1111/j.1460-9568.2005.04307.x
- Villalobos, C., Foehring, R. C., Lee, J. C., and Andrade, R. (2011). Essential role for phosphatidylinositol 4,5-bisphosphate in the expression, regulation, and gating of the slow afterhyperpolarization current in the cerebral cortex. *J. Neurosci.* 31, 18303–18312. doi: 10.1523/JNEUROSCI.3203-11.2011
- Wang, Z., and McCormick, D. A. (1993). Control of firing mode of corticotectal and corticopontine layer V burst-generating neurons by norepinephrine, acetylcholine, and 1S,3R-ACPD. *J. Neurosci.* 13, 2199–2216.

- Weber, E. T., and Andrade, R. (2010). *Htr2a* gene and 5-HT_{2A} receptor expression in the cerebral cortex studied using genetically modified mice. *Front. Neurosci.* 4:36. doi: 10.3389/fnins.2010.00036
- Williams, G. V., Rao, S. G., and Goldman-Rakic, P. S. (2002). The physiological role of 5-HT_{2A} receptors in working memory. *J. Neurosci.* 22, 2843–2854.
- Willins, D. L., Deutch, A. Y., and Roth, B. L. (1997). Serotonin 5-HT_{2A} receptors are expressed on pyramidal cells and interneurons in the rat cortex. *Synapse* 27, 79–82. doi: 10.1002/(SICI)1098-2396(199709)27:1<79::AID-SYN8<3.0.CO;2-A
- Winstanley, C. A., Chudasama, Y., Dalley, J. W., Theobald, D. E., Glennon, J. C., and Robbins, T. W. (2003). Intra-prefrontal 8-OH-DPAT and M100907 improve visuospatial attention and decrease impulsivity on the five-choice serial reaction time task in rats. *Psychopharmacology* 167, 304–314. doi: 10.1007/s00213-003-1398-x
- Xu, T., and Pandey, S. C. (2000). Cellular localization of serotonin_{2A} (5HT_{2A}) receptors in the rat brain. *Brain Res. Bull.* 51, 499–505. doi: 10.1016/S0361-9230(99)00278-6
- Yan, H. D., Villalobos, C., and Andrade, R. (2009). TRPC channels mediate a muscarinic receptor-induced afterdepolarization in cerebral cortex. *J. Neurosci.* 29, 10038–10046. doi: 10.1523/JNEUROSCI.1042-09.2009
- Zhang, Z. W., and Arsenault, D. (2005). Gain modulation by serotonin in pyramidal neurones of the rat prefrontal cortex. *J. Physiol.* 566, 379–394. doi: 10.1113/jphysiol.2005.086066

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Stephens, Baker and Gullledge. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.