



Effects of Isosorbide Mononitrate and/or Cilostazol on Hematological Markers, Platelet Function, and Hemodynamics in Patients With Lacunar Ischaemic Stroke: Safety Data From the Lacunar Intervention-1 (LACI-1) Trial

OPEN ACCESS

Edited by:

Andreas Charidimou,
Massachusetts General Hospital,
Harvard Medical School,
United States

Reviewed by:

Vincent Chung Tong Mok,
The Chinese University of
Hong Kong, China
Joan Marti Fabregas,
Hospital de la Santa Creu i Sant
Pau, Spain

*Correspondence:

Joanna M. Wardlaw
joanna.wardlaw@ed.ac.uk

†These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Stroke,
a section of the journal
Frontiers in Neurology

Received: 08 April 2019

Accepted: 18 June 2019

Published: 03 July 2019

Citation:

Appleton JP, Blair GW, Flaherty K,
Law ZK, May J, Woodhouse LJ,
Doubal F, Sprigg N, Bath PM and
Wardlaw JM (2019) Effects of
Isosorbide Mononitrate and/or
Cilostazol on Hematological Markers,
Platelet Function, and Hemodynamics
in Patients With Lacunar Ischaemic
Stroke: Safety Data From the Lacunar
Intervention-1 (LACI-1) Trial.
Front. Neurol. 10:723.
doi: 10.3389/fneur.2019.00723

Jason P. Appleton^{1,2†}, Gordon W. Blair^{3,4,5†}, Katie Flaherty¹, Zhe Kang Law^{1,2,6}, Jane May¹,
Lisa J. Woodhouse¹, Fergus Doubal^{3,4,5}, Nikola Sprigg^{1,2}, Philip M. Bath^{1,2†} and
Joanna M. Wardlaw^{3,4,5†}

¹ Stroke Trials Unit, Division of Clinical Neuroscience, University of Nottingham, Nottingham, United Kingdom, ² Stroke, Nottingham University Hospitals NHS Trust, Nottingham, United Kingdom, ³ Brain Research Imaging Centre, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, United Kingdom, ⁴ Edinburgh Dementia Research Centre in the UK Dementia Research Initiative, Edinburgh, United Kingdom, ⁵ Edinburgh Imaging, University of Edinburgh, Edinburgh, United Kingdom, ⁶ Department of Medicine, National University of Malaysia, Kuala Lumpur, Malaysia

Background: Cilostazol and isosorbide mononitrate (ISMN) are candidate treatments for cerebral small vessel disease and lacunar ischaemic stroke. As both drugs may influence hemoglobin and platelet count, and hemodynamics, we sought to assess their effects in the lacunar intervention-1 (LACI-1) trial.

Methods: Fifty-seven lacunar ischaemic stroke patients were randomized to immediate ISMN, cilostazol, or their combination for 9 weeks in addition to guideline stroke prevention. A fourth group received both drugs with a delayed start. Full blood count, platelet function, peripheral blood pressure (BP), heart rate and central hemodynamics (Augmentation index, Buckberg index) were measured at baseline, and weeks 3 and 8. Differences were assessed by multiple linear regression adjusted for baseline and key prognostic variables. Registration ISRCTN 12580546.

Results: At week 8, platelet count was higher with cilostazol vs. no cilostazol (mean difference, MD 35.73, 95% confidence intervals, 95% CI 2.81–68.66, $p = 0.033$), but no significant differences were noted for hemoglobin levels or platelet function. At week 8, BP did not differ between the treatment groups, whilst heart rate was higher in those taking cilostazol vs. no cilostazol (MD 6.42, 95% CI 1.17–11.68, $p = 0.017$). Buckberg index (subendocardial perfusion) was lower in those randomized to cilostazol vs. no cilostazol and in those randomized to both drugs vs. either drug. Whilst ISMN significantly increased unadjusted augmentation index (arterial stiffness, MD 21.19, 95% CI 9.08–33.31, $p = 0.001$), in isolation both drugs non-significantly reduced augmentation index adjusted for heart rate.

Conclusions: Cilostazol increased heart rate and platelet count, and reduced Buckberg index, whilst both drugs may individually reduce arterial stiffness adjusted for heart rate. Neither drug had clinically significant effects on hemoglobin or platelet function over 8 weeks. Further assessment of the safety and efficacy of these medications following lacunar ischaemic stroke is warranted.

Keywords: cilostazol, isosorbide mononitrate, blood pressure, safety, platelets, lacunar stroke, randomized clinical trial

INTRODUCTION

Cerebral small vessel disease (SVD) is a common cause of stroke—both “lacunar” ischaemic and haemorrhagic stroke—cognitive impairment and dementia (1). SVD is a disorder of the small perforating arterioles of the brain involving endothelial dysfunction, inflammation (2), and blood-brain barrier breakdown (3). However, there are no specific treatments for either primary or secondary prevention of SVD and its associated clinical manifestations (4). Two medications with potential beneficial mechanisms of action have been highlighted for potential repurposing: cilostazol (a phosphodiesterase 3' inhibitor used in stroke prevention in the Asia-Pacific Region) and isosorbide mononitrate (ISMN, a nitric oxide donor widely used in ischaemic heart disease). Little is known about the safety and efficacy of these drugs in SVD, particularly in combination, and yet their effects may be synergistic (4). As a mild antiplatelet, cilostazol can cause bleeding sufficient to result in anemia. Further, cilostazol can rarely cause thrombocytopenia (5). Both drugs can lower peripheral blood pressure (BP) and increase heart rate (5, 6), but data regarding their effects on central hemodynamics following ischaemic lacunar stroke are lacking.

The Lacunar Intervention Trial-1 (LACI-1) tested cilostazol and ISMN, alone and in combination, for tolerability in patients with prior lacunar ischaemic stroke (7). Overall, the medications were well-tolerated by participants (8). As both drugs may influence platelet count and hemoglobin level in addition to hemodynamics, we sought to assess these important safety outcomes.

METHODS

Population

LACI-1 was a phase IIa, partial factorial, dose-escalation, prospective, randomized, open-label, blinded endpoint (PROBE) trial. Details regarding the trial protocol and statistical analysis plan are available (7), and the main results have been presented and published (8, 9). In summary, LACI-1 recruited 57 patients from stroke centers in Edinburgh and Nottingham, UK, with clinically confirmed lacunar ischaemic stroke, without dependency and able to consent themselves, and randomized them to one of four groups for 9 weeks: ISMN 25 mg twice daily; cilostazol 100 mg twice daily; both ISMN and cilostazol

Abbreviations: Bpm, beats per minute; BP, blood pressure; CI, confidence interval; FBC, full blood count; ISMN, isosorbide mononitrate; MD, mean difference; SVD, small vessel disease.




FIGURE 1 | P-selectin surface expression in response to arachidonic acid (Aspirin test). AA, Arachidonic acid; ISMN, isosorbide mononitrate; MF, median fluorescence.

started immediately; or both medications with a 3-week delayed start, thus providing a drug-free control period (see **Figure 1** in main paper) (8). Patients were contacted on alternate weeks to enquire about symptoms (headache, nausea, diarrhea, vomiting, and bleeding) using a structured questionnaire. All patients continued to take guideline secondary stroke prevention medications including statins and antihypertensive treatment where appropriate. Hematology and haemodynamic measures were assessed blind to treatment allocation.

All patients provided written informed consent. LACI-1 was approved by the Scotland A Research Ethics Committee (15/SS/0154) and registered (ISRCTN 12580546).

Hematology and Platelet Function Testing

Full blood counts were taken at baseline, and at weeks 3 and 8: hemoglobin (g/L); and platelet count ($\times 10^9/L$). Platelet function was assessed with measurement of surface expression of P-selectin (CD63P) using kits sensitive to aspirin (arachidonic acid) or clopidogrel (adenosine diphosphate) at the same timepoints (10). An unstimulated sample provided baseline expression data at each timepoint. Median fluorescence (MF) was recorded for platelet surface expression of P-selectin for each sample using flow cytometry. Platelet P-selectin expression was chosen since blood samples can be obtained and fixed at multiple clinical sites then transported for measurement at a central core laboratory.

Haemodynamic Measures

Hemodynamics were measured at baseline and at weeks 3 and 8. Peripheral blood pressure (BP, mmHg) and heart rate (bpm)

TABLE 1 | Baseline characteristics of LACI-1 participants.

	All	Both delayed	ISMN	Cilostazol	Both immediate
Patients	57	15	15	13	14
Age (years)*	66.1 (11.1)	63.4 (11.5)	62.2 (11.0)	75.8 (8.7)	64.4 (8.1)
Sex, female	18 (31.6%)	6 (40.0%)	4 (26.7%)	5 (38.5%)	3 (21.4%)
Onset to randomization (days)	202.6 (256.2)	153.3 (180.4)	137.5 (214.2)	279.2 (324.7)	254.0 (290.7)
Systolic BP (mmHg)	147.3 (20.5)	141.8 (19.6)	150.9 (19.0)	146.1 (21.3)	150.4 (22.9)
Diastolic BP (mmHg)	83.3 (12.6)	82.9 (14.3)	86.6 (11.5)	79.3 (12.0)	83.7 (12.6)
Past medical history (%)					
Treated hypertension	42 (73.7%)	11 (73.3%)	11 (73.3%)	12 (92.3%)	8 (57.1%)
Treated hyperlipidaemia	48 (84.2%)	11 (73.3%)	12 (80.0%)	13 (100.0%)	12 (85.7%)
Diabetes	11 (19.3%)	3 (20.0%)	3 (20.0%)	2 (15.4%)	3 (21.4%)
Atrial fibrillation	0	0	0	0	0
Myocardial infarction	1 (1.8%)	1 (6.7%)	0	0	0
Previous stroke	6 (10.5%)	1 (6.7%)	2 (13.3%)	1 (7.7%)	2 (14.3%)
Smoking	26 (45.6%)	10 (66.7%)	6 (40.0%)	6 (46.2%)	4 (28.6%)
Patient status					
mRS, baseline [6]	1.0 [0.0, 1.0]	1.0 [0.0, 2.0]	1.0 [1.0, 1.0]	1.0 [1.0, 1.0]	1.0 [0.0, 1.0]
Current NIHSS [42]	0.0 [0.0, 1.0]	0.0 [0.0, 1.0]	0.0 [0.0, 1.0]	0.0 [0.0, 1.0]	0.5 [0.0, 1.0]

BP, blood pressure; ISMN, isosorbide mononitrate; mRS, modified Rankin Scale; NIHSS, National Institute of Health Stroke Scale.

*Significant difference between the treatment groups ($p < 0.05$), comparisons done and Kruskal-Wallis tests.

were measured three times sitting and three times standing at each visit using a validated monitor. Central hemodynamics were measured in duplicate using the SphygmoCor device at the same timepoints: mean arterial pressure (MAP); Augmentation index (%)—a measure of arterial stiffness; Buckberg index (%)—a measure of subendocardial perfusion; and pulse wave velocity (m/s)—a measure of arterial stiffness, assessed using non-invasive tonometry measurements taken at the radial and carotid arteries.

Sample Size and Statistics

The sample size calculation for LACI-1 was based on the ability to detect a difference of 90 vs. 55% (i.e., an absolute difference of 35%) between those reaching target dose on one drug vs. both drugs. For 80% power, significance 0.05, a sample size of 55 was needed.

Data are number (%), mean (standard deviation, SD), and median [interquartile quartiles, IQR]. Baseline differences between treatment groups were assessed using chi-square and Kruskal-Wallis tests. Differences in hemodynamics and hematological measures were assessed by multiple linear regression, adjusted for baseline values, age, SVD imaging score and time to randomization. Haemodynamic measures were also adjusted for baseline systolic BP. As heart rate can lead to spurious results, Augmentation index (%) was measured unadjusted and normalized to heart rate 75 bpm as is standard practice. The resultant mean difference (MD) with 95% confidence intervals (CI) are presented with significance set at $p < 0.05$. Analyses were performed using SAS.

RESULTS

In total, 57 patients were recruited at mean 203 [median 256] days after their index stroke with a mean age of 66 (11) years

and 18 (32%) were female (Table 1). Most participants (97%) were on clopidogrel for stroke secondary prevention (guideline therapy in the UK), the remainder on aspirin. Apart from the cilostazol group being slightly older, baseline characteristics were well-balanced across the treatment groups. One participant in the delayed start group withdrew during follow-up.

Hematology and Platelet Function Testing

There were no differences in hemoglobin at week 8 between treatment groups (Table 2). Platelet count was slightly higher in those on cilostazol $286.8 \times 10^9/L$ vs. no cilostazol $249.1 \times 10^9/L$ (MD 35.73, 95% CI 2.81 to 68.66, $p = 0.033$). No differences were noted in platelet function between treatment groups in aspirin or clopidogrel tests (Figures 1, 2). There was no difference in bruising or bleeding with either medication in isolation or combination during the study.

Hemodynamics

Full haemodynamic data were available for 56 participants. Baseline peripheral BP and heart rate did not differ between treatment groups (Table 1). At week 8, BP did not significantly differ between the treatment groups, whilst heart rate was significantly higher in those taking cilostazol 82.8 bpm vs. no cilostazol 74.5 bpm (MD 6.42, 95% CI 1.17 to 11.68, $p = 0.017$, Table 3). Central MAP did not differ between treatment groups at week 8. Buckberg index (subendocardial perfusion) was reduced in those randomized to cilostazol 142.2% vs. no cilostazol 160.7% (MD -10.81 , 95% CI -21.15 to -0.47 , $p = 0.040$) and in those randomized to both drugs 148.9% vs. either drug 159.9% (MD -11.37 , 95% CI -21.01 to -1.72 , $p = 0.021$). Unadjusted Augmentation index was significantly higher (i.e., increased arterial stiffness) in those randomized to ISMN 123.8% vs. no ISMN 119.8% (MD 21.19, 95% CI 9.08 to 33.31, $p = 0.001$).

TABLE 2 | Hematology and platelet function testing at week 8 by treatment group.

Week 8	Both delayed N = 14		ISMN N = 15		Cilostazol N = 13		Both immediate N = 14		Cilostazol vs. none (2 + 3 vs. 1 + 4*)		ISMN vs. none (1 + 3 vs. 2 + 4*)		Cil + ISMN vs. one or other (3 + 4 vs. 1 + 2)	
	Week 3 (4*)	Week 8 (4)	Week 8 (1)	Week 8 (2)	Week 8 (3)	MD (95% CI)	p-value	MD (95% CI)	p-value	MD (95% CI)	p-value			
Full blood count, adjusted for baseline														
Hemoglobin (g/L), mean (SD)	133.0 (12.2)	136.2 (13.5)	146.9 (12.0)	130.8 (9.5)	137.1 (14.6)	-3.7 (-8.4, 1.0)	0.12	-1.0 (-6.9, 4.9)	0.73	-2.4 (-6.8, 2.1)	0.30			
Platelet count (x10 ⁹ /L), mean (SD)	260.3 (47.1)	258.4 (56.1)	238.6 (45.5)	289.5 (61.4)	284.3 (57.8)	35.7 (2.8, 68.7)	0.033	3.9 (-32.4, 40.2)	0.83	-7.4 (-36.8, 21.9)	0.62			
Platelet function, adjusted for baseline														
Unstimulated median fluorescence, mean (SD)	87.0 (29.0)	80.4 (29.1)	86.1 (29.0)	75.9 (27.3)	83.7 (33.8)	-2.9 (-15.1, 9.3)	0.64	-8.2 (-20.7, 4.2)	0.20	7.5 (-4.0, 19.0)	0.20			
AA median fluorescence, mean (SD)	741.0 (395.0)	775.4 (474.4)	675.2 (364.1)	636.4 (272.4)	679.6 (376.6)	-83.6 (-256.5, 89.4)	0.34	-14.1 (-192.5, 164.2)	0.88	109.7 (-54.0, 273.5)	0.19			
ADP median fluorescence, mean (SD)	519.3 (234.5)	502.1 (242.8)	454.9 (238.8)	518.9 (179.5)	451.7 (176.5)	-1.3 (-77.6, 75.0)	0.97	-0.5 (-80.1, 79.1)	0.99	111.9 (-60.2, 84.1)	0.75			

AA, arachidonic acid (aspirin test); ADP, adenosine diphosphate (clopidogrel test); CI, confidence intervals; ISMN, isosorbide mononitrate; MD, mean difference; SD, standard deviation.




FIGURE 2 | P-selectin surface expression in response to adenosine diphosphate (Clopidogrel test). ADP, adenosine diphosphate; ISMN, isosorbide mononitrate; MF, median fluorescence.

In contrast, when adjusted for heart rate there was a tendency toward lower Augmentation index (i.e., less arterial stiffness) in those randomized to cilostazol 18.7% vs. no cilostazol 20.6% (MD -4.79, 95% CI -9.72 to 0.15, *p* = 0.057) and ISMN 16.1% vs. no ISMN 23.5% (MD -4.74, 95% CI -9.78 to 0.30, *p* = 0.066). No differences in pulse wave velocity were seen.

DISCUSSION

In this secondary analysis of the LACI-1 trial—the first trial to assess cilostazol and/or ISMN in lacunar ischaemic stroke patients, and the first trial to assess both together in any ischaemic stroke—we have demonstrated that cilostazol and ISMN in isolation and in combination do not influence hemoglobin levels or platelet function in the short-term when given in addition to usual secondary stroke prevention therapy. Platelet count was significantly higher in those randomized to cilostazol. Cilostazol alone and both drugs in combination, reduced Buckberg index (subendocardial perfusion), whilst cilostazol increased heart rate. Both drugs in isolation may reduce arterial stiffness when adjusted for heart rate.

Due to its mild antiplatelet effects, cilostazol can uncommonly ($\geq 1/100$ to $< 1/10$) cause bleeding sufficient to lead to anemia according to the summary of product characteristics (SmPC) (5). Although we have a small population, it is reassuring that hemoglobin was stable between treatment groups during the trial given that all participants were already on guideline antiplatelet therapy. The increase in platelet count seen with cilostazol at 8 weeks needs further investigation, especially since thrombocythaemia, is known to occur rarely ($\geq 1/10,000$ to $< 1/1,000$) (5). The lack of any difference in platelet function between the treatment groups is of further reassurance.

ISMN is known to lower BP and increase heart rate and so can commonly lead to postural dizziness ($\geq 1/100$ to $< 1/10$) and rarely syncope (6). Cilostazol can increase heart rate and, if co-administered with BP-lowering medication, an additive hypotensive effect with reflex tachycardia is reported (5). Neither symptom were apparent across the trial population (8). Importantly, despite cilostazol increasing heart rate in

TABLE 3 | Peripheral and central hemodynamics by treatment group.

	Both delayed N = 14		ISMN N = 15		Cilostazol N = 13		Both immediate N = 14		Cilostazol vs. none (2 + 3 vs. 1 + 4*)		ISMN vs. none (1 + 3 vs. 2 + 4*)		Cil + ISMN vs. one or other (3 + 4 vs. 1 + 2)	
	Week 3 (4*)	Week 8 (4)	Week 8 (1)	Week 8 (2)	Week 8 (3)	MD (95% CI)	p-value	MD (95% CI)	p-value	MD (95% CI)	p-value			
Peripheral hemodynamics, adjusted for baseline														
Systolic (mmHg), mean (SD)	134.8 (12.4)	131.4 (19.0)	144.1 (16.5)	146.7 (19.4)	139.9 (20.7)	-1.11 (-8.38, 6.16)	0.765	-1.06 (-8.52, 6.40)	0.780	-4.91 (-11.71, 1.88)	0.156			
Diastolic (mmHg), mean (SD)	80.4 (9.7)	78.0 (10.8)	84.9 (7.6)	81.3 (11.2)	79.8 (10.5)	-1.63 (-5.84, 2.58)	0.448	-0.65 (-4.96, 3.66)	0.768	-2.56 (-6.50, 1.38)	0.202			
Heart rate, mean (SD)	74.3 (9.3)	79.5 (11.2)	74.7 (12.5)	80.6 (15.0)	84.8 (15.1)	6.42 (1.17, 11.68)	0.017	0.06 (-5.50, 5.62)	0.984	3.13 (-1.93, 8.20)	0.225			
Central hemodynamics, adjusted for baseline														
Pulse wave velocity (m/s), mean (SD)	8.4 (1.3)	8.7 (1.4)	8.7 (1.6)	8.6 (1.3)	8.2 (1.3)	-0.34 (-0.99, 0.30)	0.296	-0.14 (-0.84, 0.55)	0.686	0.00 (-0.64, 0.64)	0.996			
Augmentation index 75 (%), mean (SD)	24.0 (10.6)	16.2 (12.4)	17.5 (12.9)	23.0 (10.6)	14.7 (10.5)	-4.79 (-9.72, 0.15)	0.057	-4.74 (-9.78, 0.30)	0.066	0.09 (-4.76, 4.94)	0.970			
Augmentation index (%), mean (SD)	118.6 (28.3)	123.3 (23.2)	131.1 (16.8)	121.1 (18.9)	133.1 (25.0)	3.46 (-8.65, 15.57)	0.575	21.19 (9.08, 33.31)	0.001	0.97 (-10.77, 12.71)	0.871			
Central blood pressure (mmHg), mean (SD)	99.0 (14.6)	95.4 (12.7)	101.2 (12.4)	103.1 (11.8)	98.8 (11.7)	-2.03 (-8.23, 4.17)	0.521	-4.74 (-10.67, 1.19)	0.117	-3.36 (-9.12, 2.40)	0.253			
Buckberg index (%), mean (SD)	154.4 (21.5)	164.8 (22.2)	166.6 (25.2)	152.2 (31.4)	132.9 (22.7)	-10.81 (-21.15, -0.47)	0.040	6.75 (-3.59, 17.09)	0.201	-11.37 (-21.01, -1.72)	0.021			

CI, confidence intervals; ISMN, isosorbide mononitrate; MD, mean difference; SD, standard deviation. Augmentation index 75: normalized to heart rate 75 bpm.

our population, no significant hypotensive effect was seen in combination with participants' pre-prescribed antihypertensives (74% were on antihypertensives at randomization) or ISMN. Central haemodynamic data regarding cilostazol and/or ISMN in the context of lacunar ischaemic stroke are lacking. Buckberg index is a marker of subendocardial perfusion, which is reduced with increased heart rate (11). Cilostazol and both drugs in combination reduced Buckberg index, which may be due to their underlying effects on heart rate, or may reflect that the patients allocated to cilostazol were older than those in other allocated groups. ISMN was associated with increased Augmentation index (a marker of arterial stiffness) unadjusted for heart rate, but when Augmentation index was adjusted for heart rate, cilostazol and ISMN in isolation both non-significantly reduced arterial stiffness. Given that small vessel stiffness is implicated in the pathophysiology of cerebral SVD (12), medications that reduce arterial stiffness may be of benefit.

The strengths of this LACI-1 substudy include using data from the first randomized controlled trial assessing both cilostazol and ISMN in lacunar ischaemic stroke patients. There are, however, limitations. First, the sample size was small and the study was not powered to look for differences in hematological or haemodynamic secondary outcomes, therefore some findings may represent chance. Second, the treatment duration was relatively short and so we cannot comment on the longer-term effects of these medications on hematological markers or hemodynamics. Third, those who were randomized to cilostazol were older which may explain some of the reported differences in these outcomes. Last, there was no formal control group throughout the trial, rather the delayed start group represented a drug-free control period. This gave the opportunity to establish whether starting one drug first over the other impacted upon side-effects, whilst still providing a control group for the first 3 weeks of the study.

In summary, cilostazol and ISMN were safe in isolation and combination in the short- to medium-term in this small population following lacunar ischaemic stroke. Further data from larger trials are required to establish the longer term effects of these medications on safety and efficacy following lacunar ischaemic stroke. As such, the LACI-2 study is testing their effects on stroke recurrence, cognitive function, imaging features of SVD and safety when given for 12 months in patients with lacunar ischaemic stroke (ISRCTN14911850).

DATA AVAILABILITY

The datasets generated for this study are available on request to the corresponding author.

ETHICS STATEMENT

LACI-1 was approved by the Scotland A Research Ethics Committee (15/SS/0154) and registered (ISRCTN 12580546).

AUTHOR CONTRIBUTIONS

JA: recruitment, data collection, analysis, and manuscript preparation. GB and ZL: recruitment, data collection, analysis, and manuscript editing. KF and LW: statistical analysis, manuscript preparation, and editing. JM: platelet function testing. FD and NS: study design, supervision, study set up, data collection, SAE adjudication, and manuscript editing. PB: trial conception, design, management, supervision, data collection and analysis, and manuscript editing. JW: trial conception, design, funding, supervision, data collection, analysis, manuscript editing, and overall guarantor.

FUNDING

LACI-1 is funded by the Alzheimer's Society Ref: 252 (AS-PG-14-033). Additional support is provided by the European Union Horizon 2020 project No 666881, SVDs@Target (GB), the Stroke Association Princess Margaret Research Development

REFERENCES

1. Wardlaw JM, Smith C, Dichgans M. Small vessel disease: mechanisms and clinical implications. *Lancet Neurol.* (2019) 18: 684–96. doi: 10.1016/S1474-4422(19)30079-1
2. Wiseman S, Marlborough F, Doubal F, Webb DJ, Wardlaw J. Blood markers of coagulation, fibrinolysis, endothelial dysfunction and inflammation in lacunar stroke versus non-lacunar stroke and non-stroke: systematic review and meta-analysis. *Cerebrovasc Dis.* (2014) 37:64–75. doi: 10.1159/000356789
3. Wardlaw JM, Makin SJ, Valdés Hernández MC, Armitage PA, Heye AK, Chappell FM, et al. Blood-brain barrier failure as a core mechanism in cerebral small vessel disease and dementia: evidence from a cohort study. *Alzheimer Dement.* (2017) 13:634–43. doi: 10.1016/j.jalz.2016.09.006
4. Bath P, Wardlaw J. Pharmacological treatment and prevention of cerebral small vessel disease: a review of potential interventions. *Stroke.* (2015) 10:469–78. doi: 10.1111/ijvs.12466
5. European Medicines Agency. *Cilostazol—Summary of Product Characteristics.* (2016). Available online at: <https://www.medicines.org.uk/emc/product/3379/smpc> (accessed September 1, 2018).
6. SmPC. *Isosorbide Mononitrate—Summary of Product Characteristics.* (2014). Available online at: <https://www.medicines.org.uk/emc/product/6115/smpc> (accessed September 1, 2018).
7. Blair GW, Appleton JP, Law ZK, Doubal F, Flaherty K, Dooley R, et al. Preventing cognitive decline and dementia from cerebral small vessel disease: the LACI-1 trial. protocol and statistical analysis plan of a phase IIa dose escalation trial testing tolerability, safety and effect on intermediary endpoints of isosorbide mononitrate and cilostazol, separately and in combination. *Int J Stroke.* (2018) 13:530–8. doi: 10.1177/1747493017731947
8. Blair GW, Appleton JP, Flaherty K, Doubal F, Sprigg N, Dooley R, et al. Tolerability, safety and intermediary pharmacological effects of cilostazol and isosorbide mononitrate, alone and combined, in patients with lacunar ischaemic stroke: the LACunar Intervention-1 (LACI-1) trial, a randomised clinical trial. *EclinicalMedicine.* (in press). doi: 10.1016/j.eclinm.2019.04.001

Fellowship scheme (GB), the Stroke Association Garfield Weston Foundation Stroke Association Senior Clinical Lectureship (FD), NHS Research Scotland (FD), NHS Lothian Research and Development Office and the Scottish Funding Council through the Scottish Imaging Network, A Platform for Scientific Excellence (SINAPSE) Collaboration. Funding is gratefully acknowledged from the Foundation Leducq (ref no. 16 CVD 05) and Edinburgh and Lothians Health Foundation. JA and LW are supported, in part, by the National Institutes of Health Research (NIHR) HTA TARDIS and BHF RIGHT-2 trials. ZL and KF are supported, in part, by the NIHR HTA TICH-2 trial. PB is Stroke Association Professor of Stroke Medicine, and is a NIHR Senior Investigator. LACI-1 was adopted by the NIHR CRN.

ACKNOWLEDGMENTS

Data contributing to this manuscript were presented, in part, as posters at the European Stroke Organization Conference (ESOC) in Gothenburg, Sweden May 2018: Appleton et al. (13, 14).

9. Blair GW, Appleton JP, Flaherty K, Sprigg N, Doubal F, Boyd J, et al. Preventing cognitive decline and dementia from cerebral small vessel disease: the LACI-1 trial. *Int J Stroke.* (2017) 12 (5 Suppl. 2):15.
10. Bath PM, May J, Flaherty K, Woodhouse LJ, Dovlatova N, Fox SC, et al. Remote assessment of platelet function in patients with acute stroke or transient ischaemic attack. *Stroke Res Treat.* (2017) 2017:7365684. doi: 10.1155/2017/7365684
11. Buckberg GD, Fixler DE, Archie JP, Hoffman JL. Experimental subendocardial ischemia in dogs with normal coronary arteries. *Circ Res.* (1972) 30:67–81. doi: 10.1161/01.RES.30.1.67
12. Del Bene A, Makin SD, Doubal FN, Wardlaw JM. Do risk factors for lacunar ischaemic stroke vary with the location or appearance of the lacunar infarct? *Cerebrovasc Dis.* (2012) 33:21.
13. Appleton JP, Flaherty K, Blair GW, Law ZK, Doubal F, Sprigg N, et al. Platelet and haemoglobin levels in patients on isosorbide mononitrate and/or cilostazol with lacunar ischaemic stroke: data from the LACI-1 trial. *Eur Stroke J.* (2018) 3:488–9.
14. Appleton JP, Blair GW, Flaherty K, Law ZK, Doubal F, Sprigg N, et al. Peripheral and central haemodynamics in patients on isosorbide mononitrate and/or cilostazol with lacunar ischaemic stroke: data from the LACI-1 trial. *Eur Stroke J.* (2018) 3:489.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Appleton, Blair, Flaherty, Law, May, Woodhouse, Doubal, Sprigg, Bath and Wardlaw. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.